

This is a repository copy of *Density-functional theory of polar insulators*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/3995/>

---

**Article:**

Godby, R.W. [orcid.org/0000-0002-1012-4176](http://orcid.org/0000-0002-1012-4176), Gonze, X. and Ghosez, P. (1997)  
Density-functional theory of polar insulators. *Physical Review Letters*. pp. 294-297. ISSN  
1079-7114

<https://doi.org/10.1103/PhysRevLett.78.294>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

*promoting access to White Rose research papers*



**Universities of Leeds, Sheffield and York**  
**<http://eprints.whiterose.ac.uk/>**

---

White Rose Research Online URL for this paper:  
**<http://eprints.whiterose.ac.uk/3995>**

---

**Published paper**

**Gonze, X, Ghosez, P, Godby, RW (1995)** *Density-functional theory of polar insulators*

**Physical Review Letters 78 (294-297)**

---

# Density-Functional Theory of Polar Insulators

X. Gonze\*, Ph. Ghosez\*, and R.W. Godby†

\* *Unité PCPM, Université Catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium*

† *Department of Physics, University of York, Heslington, York YO1 5DD, U.K.*

(February 1, 2008)

## Abstract

We examine the density-functional theory of macroscopic insulators, obtained in the large-cluster limit or under periodic boundary conditions. For *polar* crystals, we find that the two procedures are not equivalent. In a large-cluster case, the exact exchange-correlation potential acquires a homogeneous “electric field” which is absent from the usual local approximations, and the Kohn-Sham electronic system becomes metallic. With periodic boundary conditions, such a field is forbidden, and the polarization deduced from Kohn-Sham wavefunctions is incorrect even if the exact functional is used.

PACS numbers: 71.15.Mb; 71.10.-w; 77.22.Ej

Although the density-functional theory (DFT) introduced by Hohenberg, Kohn and Sham [1,2] has become the standard method for first-principles calculations of the ground-state properties of solids, to our knowledge [3], the implications of applying DFT to infinite, insulating, crystals have not been fully appreciated. In part, this reflects the fact that the key theorems of DFT [1,2] were proved for arbitrarily large, but not infinite, systems. In the present paper, we show that the exact DFT treatment of *polar* crystals (a) with the usual Born-von Karman (BvK) boundary conditions, or (b) from the macroscopic limit of large clusters, will generally give different macroscopic polarizations. Only (b) is correct.

Investigating the *response* of periodic insulators to a homogeneous electric field, we recently revealed [4] the polarization-dependence of the exchange-correlation energy, and its consequences on the dielectric response. Aulbur, Jönsson and Wilkins [5] quantified this effect for real materials, while Resta [6] discussed the origin of such a behaviour in connection with long-range correlation effects. The present study emphasizes a more basic role of the polarization in DFT: careful handling of the polarization is mandatory for polar solids, even under zero electric field. In polar materials, the spontaneous polarization computed from the Kohn-Sham (KS) wavefunctions will be correct only if an exchange-correlation homogeneous electric field is allowed throughout the material, in which case the KS electronic system becomes *metallic*. This field will appear in the exact DFT treatment of a finite cluster but is forbidden when using BvK conditions. Approximate density-functionals such as the Local Density Approximation (LDA) and the Generalized Gradient Approximation (GGA) always fail to yield an exchange-correlation electric field: within these approximations, using BvK boundary conditions or finite clusters incorrectly provide the *same* value of the polarization [7]. Any improvement to these functionals which retains a dependence only on the periodic density will be similarly flawed. We will exhibit our results for a one-dimensional model semiconductor.

The correct definition of a macroscopic crystal is clearly as the limit of a finite crystal of increasing size. Fig. 1(a) shows schematically the total electrostatic potential  $V_{\text{elec}} = V_e + V_N + V_{\text{appl}}$  in such a finite crystal, where  $V_e$  is the electrostatic potential due to the

ground-state electron density,  $V_N$  is the potential due to the nuclei, and  $V_{\text{appl}}$  is an applied potential, created by an external short-circuited capacitor, that maintains equality of the electrostatic potential on the two sides [8]. The sum  $V_{\text{ext}} = V_N + V_{\text{appl}}$  is referred to as the external potential. The total electrostatic potential in the bulk region is periodic and, crucially for a non-zero polarization, non-centrosymmetric. The potential just outside the surface is fixed by the electrostatic potential of the capacitor plates. The corresponding ground-state electron density is also shown. In the bulk region, it is periodic, with the same periodicity as the local potential [9]. Close to the surface, the density deviates from perfect periodicity, although this effect decreases exponentially with the distance from the surface [10].

The macroscopic polarization of such a finite solid is directly linked to the total surface charge [11,12]. Its value is equal to zero (modulo a half-quantum) if the crystal is centrosymmetric [12,13], but otherwise can have any value and must be calculated. For a long time the macroscopic polarization was only accessible from the surface charge and was a well-defined concept only for finite solids. Recent theoretical advances have shown that it can also conveniently be determined, up to a quantum, from a Berry phase of the correlated many-body wavefunction of the *bulk* [12–16]. Within this approach the macroscopic polarization appears as a bulk property and is unambiguously defined even for the infinite periodic solid, which is of practical interest in solid state *ab initio* calculations.

The breakthrough of King-Smith and Vanderbilt [14], leading to the modern theory of the polarization, was actually carried out in the context of DFT. Later, they argued [12] that the Berry phase of the occupied KS wavefunctions possesses an exact physical meaning since the surface charge must be exactly reproduced within DFT [12]. We now show that the justification of Vanderbilt and King-Smith apply to exact DFT *only* when considering *finite* solids, and not when applying BvK periodic boundary conditions.

In the context of DFT, it is shown that the density  $n(\mathbf{r})$  of the ground state of a system uniquely determines the external potential up to a constant. Following Kohn and Sham [2], one can introduce a fictitious system of non-interacting electrons in an effective potential

$V_{\text{eff}} = V_{\text{ext}} + V_{\text{H}} + V_{\text{xc}}$  (where  $V_{\text{xc}}$  is the exchange correlation potential) that reproduces the ground-state electron density of the real system. In the particular case where *periodic boundary conditions* are imposed, although the KS effective potential  $V_{\text{eff}}$  is constrained to reproduce the correct *periodic* bulk density of the polar solid, there is no guarantee that it will reproduce the correct *polarization*, since this information is not contained in the criterion for the effective potential to be correct [18]. Such a periodic DFT is based only on the periodic part of the density, while the polarization is a completely independent quantity [13,4,6] that depends on the phase of the correlated wavefunctions. The polarization will be correct only in those solids where a fundamental symmetry (such as centrosymmetry) constrains the polarization, or where external parameters, such as the pressure, are fortuitously chosen.

We illustrate this for a one-dimensional model semiconductor [19]. In this model the electrostatic potential is periodic and asymmetric:  $V_{\text{elec}}(x) = V_c \cos \frac{2\pi x}{a} + V_s \sin \frac{4\pi x}{a}$ . A non-local self-energy operator, intended to mimic the relevant many-body effects, has the same non-local form as in Ref. [4]:  $\Sigma(x, x', \omega) = \frac{f(x)+f(x')}{2}g(|x - x'|)$  where  $f(x) = F_o[1 - \cos \frac{2\pi x}{a}]$  is a negative function with the periodicity of one unit cell and  $g(y)$  is a normalized gaussian of width  $w$ .

First, the Schrödinger equation containing the self-energy operator is solved by direct diagonalization using a plane-wave basis set. The density is deduced from the sum of the squares of the eigenfunctions. From this result, using standard iterative optimization techniques, we construct an *exact* density-functional theory by determining the local potential  $V_{\text{eff}}(x)$  which, when filled with *non-interacting* electrons (no self-energy operator), reproduces the same electron density as in the self-energy calculation. Fig. 2 presents the function  $V_{\text{elec}}(x)$ , as well as the density  $n(x)$ , and the effective potential  $V_{\text{eff}}(x)$ , for the following set of parameters :  $a_0 = 4$  a.u.,  $V_c = V_s = 2.72$  eV,  $F_o = -4.08$  eV,  $w = 2$  a.u.

Using the Berry-phase approach [13,14], we then compute the polarization [20]. In the self-energy calculation, the polarization is  $22.68 \cdot 10^{-3}$  electrons with respect to the centrosymmetric system with  $V_s = 0$ , while that calculated from the Berry phase of the Kohn-Sham wave-functions is  $21.99 \cdot 10^{-3}$  electrons. The two polarizations differ by 3%, well outside

the calculational error bar. This value may be taken as an order of magnitude estimation of the effect in real materials, and is compatible with the observed (and often satisfactory) accuracy of LDA polarization calculations for real ferroelectric materials [17].

The deficiency in the periodic-boundary approach, reflected in the Berry phase of the KS wavefunctions and hence in the polarization, is that the exchange-correlation potential is prevented from having a component which is linear in space [21]. The KS theorem demonstrates that there is only one periodic effective potential  $V_{\text{eff}}$  that reproduces a particular periodic density. However, once an additional linear component is allowed, there exists an infinite family of KS potentials that gives the same periodic density but different polarizations [4]. Imposing BvK conditions on the potential thus arbitrarily constrains the polarization to a specific, usually incorrect, value. This restriction does not apply for the finite cluster, where application of the KS theorem shows that there exists a unique effective potential that, when used in the effective Hamiltonian, will generate the exact ground-state density *everywhere*: not only in the bulk region (as in the BvK case), but *also* in the surface region, resulting in the correct polarization. Fig. 1(c) sketches the behaviour of such an effective potential. The linear part is necessary to yield the correct polarization in polar crystals. This “exchange-correlation electric field” originates in the ultra non-local dependence of the exchange-correlation energy in the surface charge pointed out in Ref. [4].

In the small cluster shown in Fig. 1(c), the magnitude of the exchange-correlation field is approximately independent of the cluster size, since the polarization correction relative to periodic DFT is constant. As the cluster is made larger, a point will be reached where the variation in potential from one side of the cluster to the other, due to the homogeneous exchange-correlation electric field, reaches the DFT band gap of the material. Beyond this point, the KS electronic system is *metallic* and the band edges will “pin” the effective potential (Fig. 3). As the cluster is made still larger, charge will flow freely from one face to the other in order to maintain the correct macroscopic polarization. The magnitude of the homogeneous electric field will now change with the size of the cluster in order to maintain the potential drop: in the limit of large cluster size, the effective homogeneous electric field

will be infinitesimally small, although non-zero. As mentioned in Ref. [22], an infinite system cannot sustain a finite homogeneous electric field in its ground-state. Here, an infinitesimal field appears naturally in the DFT treatment of polar solids.

There is a strong similarity between this behaviour and that of a system of two distant, different, open-shell atoms [23], in which the exact exchange-correlation potential exhibits a long-range spatial variation to align the Kohn-Sham eigenvalues. There is also a connection with the DFT metal/insulator paradox [24] in which an insulating system may be described as metallic in DFT.

In summary, for a polar insulator, when Born-von Karman periodic boundary conditions are used, the polarizations calculated from the Berry phase of the Kohn-Sham wavefunctions and from the Berry phase of the correlated wavefunction will differ, because the DFT effective potential is prevented from acquiring a linear part. When a large cluster is used for the DFT calculation, a homogeneous effective exchange-correlation “electric field” develops in order to correctly reproduce the polarization. The Kohn-Sham system becomes metallic.

The authors thank R. Resta, R. Martin, H. Krakauer, R.E. Cohen and I. Mazin for interesting discussions, and acknowledge financial support from FNRS-Belgium (X.G.), the Concerted Actions program (X.G.), The Royal Society and the Engineering and Physical Sciences Research Council (R.W.G.), the European Union (Human Capital and Mobility Program Contracts CHRX-CT940462 and CHRX-CT930337), and the Academic Research Collaboration Programme between the FNRS, the CGRI, and the British Council.



## REFERENCES

- [1] P. Hohenberg and W. Kohn, Phys. Rev. 136, B864 (1964).
- [2] W. Kohn and L.J. Sham, Phys. Rev. 140, A1133 (1965).
- [3] D.D. Koelling, Rep. Prog. Phys. 44, 139 (1981); *Theory of the Inhomogeneous Electron Gas*, edited by S. Lundqvist and N.H. March (Plenum Press, New York, 1983); *Highlights of Condensed-Matter Theory*, edited by F. Bassani, F. Fumi and M.P. Tosi (North-Holland, New York, 1985); *Density Functional Methods in Physics*, edited by R.M. Dreizler and J. da Providencia (Plenum Press, New York, 1985); G.P. Srivastava and D. Weaire, Adv. Phys. 36, 463 (1987); R.O. Jones and O. Gunnarsson, Rev. Mod. Phys. 61, 689 (1989); W.E. Pickett, Comp. Phys. Rep. 9, 115 (1989); E.S. Kryachko and E.V. Ludena, *Energy Density Functional Theory of Many-Electron Systems* (Kluwer Academic Publishers, Dordrecht, 1990).
- [4] X. Gonze, Ph. Ghosez and R. W. Godby, Phys. Rev. Lett. 74, 4035 (1995).
- [5] W.G. Aulbur, L. Jönsson, and J.W. Wilkins, Phys. Rev. B (to be published).
- [6] R. Resta, Phys. Rev. Lett. 77, 2265 (1996).
- [7] This can be viewed as a practical advantage of LDA or GGA: the computation of the polarization under BvK conditions (from a Berry phase of the KS wavefunctions) is intrinsically easier than from the limit of large clusters and constitutes a consistent procedure within this approximation.
- [8] For the simplicity of the argument, we imagine that the capacitor is a classical device, in which case the analysis of electrostatics performed in Ref. [6] applies.
- [9] In this work, we do not consider symmetry-breaking effects like charge- or spin-density waves.
- [10] W. Kohn, Phys. Rev. 133, A171 (1964).




- [11] L.D. Landau and E.M. Lifshits, *Electrodynamics of Continuous Media* (Pergamon Press, New York, 1960).
- [12] D. Vanderbilt and R.D. King-Smith, Phys. Rev. B 48, 4442 (1993). In the present context, see especially Section VI of this reference.
- [13] R. Resta, Rev. Mod. Phys. 66, 899 (1994).
- [14] R.D. King-Smith and D. Vanderbilt, Phys. Rev. B 47, 1651 (1993).
- [15] G. Ortiz and R.M. Martin, Phys. Rev. B 49, 14202 (1994).
- [16] R. Resta and S. Sorella, Phys. Rev. Lett. 74, 4738 (1995).
- [17] R. Resta, M. Posternak, and A. Baldereschi, Phys. Rev. Lett. 70, 1010 (1993).
- [18] In some cases, even with BvK conditions, the Kohn-Sham system will be a metal [24] in which case the computation of the Berry phase is inappropriate. Henceforth we will suppose that the KS system obtained under BvK boundary conditions is insulating.
- [19] This model is a generalization of that used in R.W. Godby and L.J. Sham, Phys. Rev. B 49, 1849 (1994) and Ref. [4].
- [20] The standard Berry phase analysis [10-14] is applicable to our system, showing the exact polarization to be proportional to the Berry phase of the eigenfunctions of the non-local Hamiltonian. This is distinct from the Berry phase of the Kohn-Sham wavefunctions.
- [21] The electrostatic potential ( $V_{\text{ext}} + V_e$ ) is the same periodic potential as in the many-body case, thus constraining  $V_{\text{xc}}$  to be periodic also.
- [22] R.W. Nunes and D. Vanderbilt, Phys. Rev. Lett. 73, 712 (1994).
- [23] C.O. Almbladh and U. von Barth, in “Density functional methods in physics”, ed. R.M. Dreizler and J. da Providencia, (Plenum Press, New York, 1985), p. 209.
- [24] R.W. Godby and R.J. Needs, Phys. Rev. Lett. 62, 1169 (1989).


## FIGURE CAPTIONS

Fig.1 (a) The local electrostatic potential (external plus Hartree) of an insulator, and the corresponding ground-state density. In the bulk region the potential is periodic. Short-circuited capacitor plates are also present. (b) The effective potential that, when used in Kohn-Sham equations, is able to reproduce the periodic part of the density shown in (a), under Born-von Karman periodic boundary conditions. The macroscopic polarization is not correct. (c) The effective potential that, when used in Kohn-Sham equations, is able to reproduce the density shown in (a), in all the regions of space. The macroscopic polarization is correct (in contrast to (b)).


Fig.2 The electrostatic potential  $V_{\text{elec}}(x)$ , the electron density  $n(x)$  and the Kohn-Sham effective potential  $V_{\text{eff}}(x)$  of the model one-dimensional semiconductor are shown when periodic boundary conditions are imposed (corresponding to Fig. 1(b)). The Kohn-Sham electrons correctly reproduce the electron density, but not the macroscopic polarization.

Fig.3 For sufficiently large clusters, the exchange-correlation field will cause band overlap and hence metallization. Further increase in cluster size leaves the band edges pinned as shown, and charge transfer occurs between the two surfaces.






Conduction States



Fermi Energy



Valence States