

This is a repository copy of *Diagrammatic self-energy approximations and the total particle number*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/4013/>

Article:

Schindlmayr, A, Garcia-Gonzalez, P and Godby, R W orcid.org/0000-0002-1012-4176
(2001) Diagrammatic self-energy approximations and the total particle number. *Physical Review B*. 235106. -. ISSN 2469-9969

<https://doi.org/10.1103/PhysRevB.64.235106>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

promoting access to White Rose research papers



Universities of Leeds, Sheffield and York
<http://eprints.whiterose.ac.uk/>

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/4013>

Published paper

Schindlmayr, A, Garc'ia-González, P, Godby, RW (2001) *Diagrammatic self-energy approximations and the total particle number*

Physical Review B 64 (235106 (6 pages))

Diagrammatic self-energy approximations and the total particle number

Arno Schindlmayr,^{1,*} P. García-González,² and R. W. Godby³

¹*Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, 14195 Berlin-Dahlem, Germany*

²*Departamento de Física Fundamental, Universidad Nacional de Educación a Distancia, Apartado 60141, 28080 Madrid, Spain*

³*Department of Physics, University of York, Heslington, York YO10 5DD, United Kingdom*
(February 1, 2008)

There is increasing interest in many-body perturbation theory as a practical tool for the calculation of ground-state properties. As a consequence, unambiguous sum rules such as the conservation of particle number under the influence of the Coulomb interaction have acquired an importance that did not exist for calculations of excited-state properties. In this paper we obtain a rigorous, simple relation whose fulfilment guarantees particle-number conservation in a given diagrammatic self-energy approximation. Hedin's G_0W_0 approximation does not satisfy this relation and hence violates the particle-number sum rule. Very precise calculations for the homogeneous electron gas and a model inhomogeneous electron system allow the extent of the nonconservation to be estimated.

71.45.Gm,71.15.Qe

I. INTRODUCTION

Many-body perturbation theory is a powerful method for studying interacting electron systems, because the partial summation of self-energy diagrams allows an efficient and systematically converging description of the dominant scattering mechanisms.¹ In solid-state physics, Hedin's GW approximation² includes dynamic screening in the random-phase approximation and has been applied with great success to a large range of materials.³ While calculations have long focused on electronic excitations, such as band structures,^{4,5} that are not normally accessible by variational mean-field schemes, there is now increasing interest in using many-body perturbation theory also to obtain ground-state properties like the charge density⁶ or the total energy⁷⁻¹⁰ in order to circumvent well-known limitations of standard approximations in density-functional theory.¹¹ Unlike the calculation of excited states, which are given immediately by the pole structure of the spectral function, this generally requires a multidimensional integration over the hole part of the Green function, as in Galitskii and Migdal's expression for the total energy.¹² As a consequence, sum rules that could hitherto be ignored have gained new prominence. The most important of these is the conservation of particle number, i.e., the requirement that the integral

$$N = \frac{1}{2\pi i} \sum_{\sigma} \int d^3r \int d\omega G_{\sigma\sigma}(\mathbf{r}, \mathbf{r}; \omega) e^{i\omega\eta} \quad (1)$$

over the diagonal elements of the Green function G equals the true number of electrons. Here σ denotes the spin variable and η is a positive infinitesimal that forces the frequency contour to be closed across the upper complex half plane.

In a seminal paper Baym and Kadanoff¹³ investigated the evolution of nonequilibrium Green functions and derived a set of symmetry relations for diagrammatic many-body approximations that guarantee the overall

conservation of particle number, total energy, and momentum under time-dependent external perturbations. Baym¹⁴ later showed that a self-energy satisfying all of these relations can be represented as the derivative $\Sigma = \delta\Phi/\delta G$ of a generating functional Φ , and that in this case the Green function obtained self-consistently from Dyson's equation¹⁵ moreover yields the exact particle number. In particular, this applies to the fully self-consistent GW approximation, in which both the Green function G and the screened Coulomb interaction W are dressed by self-energy insertions in accordance with a self-consistent solution of Dyson's equation.¹³ However, it has since become clear that Φ derivability is a sufficient but not a necessary requirement for the fulfilment of the particle-number sum rule. For instance, the partially self-consistent GW_0 approximation, in which only the Green function is updated self-consistently but the screened Coulomb interaction remains undressed, is *not* Φ derivable but nevertheless produces the exact particle number.¹⁶ On the other hand, without self-consistency even in the Green function, the particle number is not, in general, given correctly,¹⁷ but this computationally efficient G_0W_0 approach still remains the preferred method for most practical applications.

More complicated self-energy expressions like the cumulant expansion¹⁸ or the T matrix¹⁹ have already been successfully applied to solids, leading to an improved description of satellite resonances. Like the GW approximation, these schemes are typically implemented without full self-consistency and are hence not Φ derivable. In order to avoid expensive numerical tests in such situations, it would be desirable to have clear diagrammatic criteria for the fulfilment of the particle-number sum rule that could be checked *a priori* without actual calculations.

Unfortunately, Baym's proof, which is based on Luttinger's examination of the exact theory²⁰ and determines the volume of the Fermi sea directly, cannot easily be extended, because it relies explicitly on the existence of the generating functional Φ . We therefore take

a different approach by describing the switching on of the Coulomb potential as a time-dependent process that connects the noninteracting and the corresponding interacting electron system on a finite time scale. In this way we can examine the differential conservation laws and deduce a diagrammatic symmetry relation for particle-number conservation *before* taking the adiabatic limit. The theoretical framework is developed in Sec. II. The non-self-consistent G_0W_0 approximation, which violates this symmetry relation and does not conserve the particle number when the interaction is switched on, deserves special attention owing to its pre-eminent role in practical implementations. In Sec. III we therefore present very precise numerical calculations of the particle number for the homogeneous electron gas and a model inhomogeneous system in order to assess the quantitative deviation. Finally, in Sec. IV we summarize our conclusions. Atomic units are used throughout.

II. PARTICLE-NUMBER CONSERVATION

In order to connect the interacting electron system with the corresponding noninteracting system, whose properties are supposedly known exactly, we consider the Hamiltonian

$$\hat{H}(t) = \hat{H}_0 + e^{-\epsilon|t|}\hat{H}_1. \quad (2)$$

The one-body part \hat{H}_0 contains the kinetic energy as well as the external potential V_{ext} , and the Coulomb interaction \hat{H}_1 is switched on exponentially with $\epsilon > 0$. At large times, both in the past and in the future, the Hamiltonian reduces to \hat{H}_0 , which constitutes a solvable problem. The noninteracting Green function G_0 is readily constructed from the solutions of the single-particle Schrödinger equation and yields the correct number of particles. On the other hand, at $t = 0$ the full Coulomb interaction is effective, and the Green function is defined as¹

$$G(x, x') = -i \frac{\langle \Psi | T[\hat{\psi}(x)\hat{\psi}^\dagger(x')] | \Psi \rangle}{\langle \Psi | \Psi \rangle}, \quad (3)$$

where the shorthand notation $x \equiv (\mathbf{r}, \sigma, t)$ indicates a set of spatial, spin, and temporal coordinates, $|\Psi\rangle$ denotes the ground-state wave function of the interacting electron system in the Heisenberg picture, and T is Wick's time-ordering operator that rearranges the subsequent symbols in ascending order from right to left with a sign change for every pair commutation. Furthermore, $\hat{\psi}^\dagger(x')$ and $\hat{\psi}(x)$ represent the electron creation and annihilation operator in the Heisenberg picture, respectively.

The unknown many-body wave function $|\Psi\rangle$ evolves from the noninteracting ground state $|\Psi_0\rangle$ and can formally be expressed as $|\Psi\rangle = \hat{U}_\epsilon(0, -\infty)|\Psi_0\rangle$, where¹

$$\begin{aligned} \hat{U}_\epsilon(t, t') &= \sum_{\nu=0}^{\infty} \frac{(-i)^\nu}{\nu!} \int_{t'}^t dt_1 \cdots \int_{t'}^t dt_\nu e^{-\epsilon(|t_1| + \cdots + |t_\nu|)} \\ &\quad \times T[\hat{H}_1(t_1) \cdots \hat{H}_1(t_\nu)] \end{aligned} \quad (4)$$

represents the time-development operator. With this definition the Green function may be rewritten as

$$G(x, x') = -i \frac{\langle \Psi_0 | T[\hat{S}_\epsilon \hat{\psi}(x)\hat{\psi}^\dagger(x')] | \Psi_0 \rangle}{\langle \Psi_0 | \hat{S}_\epsilon | \Psi_0 \rangle} \quad (5)$$

with $\hat{S}_\epsilon = \hat{U}_\epsilon(\infty, -\infty)$. At this stage the Gell-Mann and Low theorem²¹ asserts that it is, in general, permissible to take the adiabatic limit $\epsilon \rightarrow 0$. However, in the following we continue to perform a time-dependent perturbation analysis for finite ϵ and only take the adiabatic limit after establishing the conservation criteria that apply during the transition. The time-ordered products in Eq. (5) may be evaluated in the usual way by invoking Wick's theorem,²² because the exponentials are scalar functions and commute with the field operators. Hence the perturbative treatment generates the standard series of connected and topologically distinct Feynman diagrams,²³ made up of the noninteracting Green function G_0 and the two-body Coulomb potential, but the latter now acquires an additional prefactor and is given by

$$v(x, x') = \frac{e^{-\epsilon|t|}}{|\mathbf{r} - \mathbf{r}'|} \delta(t - t'). \quad (6)$$

The formal identity of the perturbation expansion in the time-dependent and the adiabatic, time-independent case is a crucial result that forms the basis of our discussion in this section.

For an analysis of the conservation properties we now follow Ref. 13 and write the perturbation series as

$$\begin{aligned} &\int G_0^{-1}(x, x_1) G(x_1, x') dx_1 \\ &= \delta(x - x') - i \int v(x, x_1) G_2(x, x_1; x', x_1^+) dx_1, \end{aligned} \quad (7)$$

invoking the two-particle Green function G_2 . The superscript x^+ indicates that a positive infinitesimal is added to the time variable to ensure the proper ordering. An equivalent form is the adjoint equation of motion

$$\begin{aligned} &\int G(x, x_1) G_0^{-1}(x_1, x') dx_1 \\ &= \delta(x - x') - i \int G_2(x, x_1; x', x_1^+) v(x', x_1) dx_1. \end{aligned} \quad (8)$$

The inverse noninteracting Green function is identical to the operator

$$\begin{aligned} G_0^{-1}(x, x') &= \left(i \frac{\partial}{\partial t} + \frac{1}{2} \nabla^2 - V_{\text{ext}}(\mathbf{r}) \right) \delta(x - x') \\ &= \left(-i \frac{\partial}{\partial t'} + \frac{1}{2} \nabla'^2 - V_{\text{ext}}(\mathbf{r}') \right) \delta(x - x'), \end{aligned} \quad (9)$$

so that after subtracting Eq. (8) from Eq. (7) we obtain

$$\begin{aligned}
& \left[i \left(\frac{\partial}{\partial t} + \frac{\partial}{\partial t'} \right) + \frac{1}{2} (\nabla + \nabla') \cdot (\nabla - \nabla') \right] G(x, x') \\
& = [V_{\text{ext}}(\mathbf{r}) - V_{\text{ext}}(\mathbf{r}')] G(x, x') \\
& - i \int [v(x, x_1) - v(x', x_1)] G_2(x, x_1; x', x_1^+) dx_1.
\end{aligned} \tag{10}$$

When we set $x' = x^+$, the terms on the right-hand side cancel, while the left-hand side reduces to the differential conservation law for the particle number

$$\frac{\partial n(\mathbf{r}, t)}{\partial t} + \nabla \cdot \mathbf{j}(\mathbf{r}, t) = 0 \tag{11}$$

with the electron density $n(\mathbf{r}, t) = -i \sum_{\sigma} G(x, x^+)$ and current $\mathbf{j}(\mathbf{r}, t) = -\frac{1}{2} \sum_{\sigma} [(\nabla - \nabla') G(x, x')]_{x'=x^+}$. Thus whenever Eqs. (7) and (8) are satisfied simultaneously, the total particle number is conserved while the interaction is switched on. This does not depend on the value of ϵ and, in particular, remains true in the adiabatic limit, which can now be taken, turning G into the equilibrium Green function of the interacting electron system.

By multiplying Eqs. (7) and (8) with G from the left and right, respectively, and then subtracting one from the other, their mutual consistency can be stated in the more convenient form,

$$\begin{aligned}
& \int G(x, x_2) v(x_2, x_1) G_2(x_2, x_1; x', x_1^+) dx_1 dx_2 \\
& = \int G_2(x, x_1; x_2, x_1^+) v(x_2, x_1) G(x_2, x') dx_1 dx_2, \tag{12}
\end{aligned}$$

that may easily be verified by visual inspection of a given diagrammatic approximation for the two-particle Green function. Evidently it is the same criterion as derived by Baym and Kadanoff¹³ for particle-number conservation under time-dependent external perturbations. This is no coincidence, of course, because the termwise cancellation of diagrams on the right-hand side of Eq. (10) is of purely topological origin and does not depend on the mathematical properties of the constituent propagators. Hence it is inconsequential whether, as in Refs. 13 and 14, G_0 contains a time-dependent perturbation while the interaction is constant or, as in the physical situation considered here, the noninteracting Green function is invariant under temporal translations while the Coulomb potential instead acquires a time-dependent prefactor.

As an example we now consider the GW approximation. The self-energy, when applied with full self-consistency, is given by

$$\Sigma(x, x') = iG(x, x')W(x^+, x'), \tag{13}$$

where the screened Coulomb interaction W takes the mathematical form of the random-phase approximation but is evaluated using the dressed Green function self-consistently derived from Dyson's equation. The diagrammatic representation of Σ is shown in Fig. 1(a). The corresponding two-particle Green function is obtained by




FIG. 1. Diagrammatic representation of the self-energy $\Sigma(x_1, x_2)$ and the corresponding two-particle Green function $G_2(x_1, x_3; x_2, x_4)$ in (a) the fully self-consistent GW approximation, (b) the partially self-consistent GW_0 approximation, and (c) the non-self-consistent G_0W_0 approximation.

comparing Dyson's equation with the equation of motion (7), which yields the identity

$$\begin{aligned}
& -i \int v(x, x_1) G_2(x, x_1; x', x_1^+) dx_1 \\
& = V_H(x) G(x, x') + \int \Sigma(x, x_1) G(x_1, x') dx_1, \tag{14}
\end{aligned}$$

where $V_H(x) = -i \int v(x, x_1) G(x_1, x_1^+) dx_1$ indicates the Hartree potential. The two-particle Green function corresponding to the GW approximation for the self-energy is also displayed in Fig. 1(a). It is easily seen that it satisfies the symmetry relation (12), which is essentially a horizontal left-right symmetry for the building blocks of G_2 , and hence conserves the total particle number when the Coulomb interaction is switched on. Of course, this result also follows from the existence of the generating functional Φ .¹⁴

The partially self-consistent GW_0 approximation

$$\Sigma(x, x') = iG(x, x')W_0(x^+, x'), \tag{15}$$

in which the screened Coulomb interaction W_0 is evaluated with the noninteracting Green function G_0 , is *not* Φ derivable, which would require an additional vertical

mirror symmetry $G_2(x_1, x_3; x_2, x_4) = G_2(x_3, x_1; x_4, x_2)$ in the diagrammatic structure of the two-particle Green function that has been lost in the transition from full to partial self-consistency. Nevertheless, G_2 , shown in Fig. 1(b), still obeys the consistency relation (12) and hence guarantees the correct total particle number, as previously confirmed by explicit integration of the spectral function.¹⁶ In contrast, the non-self-consistent G_0W_0 approximation

$$\Sigma(x, x') = iG_0(x, x')W_0(x^+, x') \quad (16)$$

leads to a two-particle Green function with lower internal symmetry, displayed in Fig. 1(c), that no longer satisfies Eq. (12), implying an incorrect total particle number. The quantitative deviation is investigated in the following section. In a similar manner, the conservation properties of other diagrammatic self-energy approximations are easily established by an inspection of the underlying two-particle Green function.

III. NUMERICAL RESULTS

In the previous section we proved that the GW and GW_0 approximations conserve the particle number for an arbitrary electron system when the Coulomb interaction is switched on, in contrast to G_0W_0 . This, coupled with their superior performance in ground-state total-energy calculations,^{7,10} might be thought to suggest that the G_0W_0 approach is useless if one is interested in ground-state properties. However, a many-body calculation at only the G_0W_0 level is already sufficient to correct typical limitations of mean-field density-functional theories, such as their inaccuracy in highly inhomogeneous systems or their failure to describe van der Waals forces.²⁴ Moreover, the Green function arising from a G_0W_0 calculation may be used as input in the variational Luttinger and Ward functional,²⁵ and prospective calculations suggest that this is an excellent approach for calculating total energies.²⁶ Since these methods are more amenable to applications in complex systems than the fully or partially self-consistent GW approximations, it is important to determine whether the underlying violation of the particle-number sum rule in the G_0W_0 framework is small enough to be safely ignored.

There are some indications that such an error is indeed fairly small for the homogeneous electron gas at metallic densities,¹⁶ a Hubbard model system,¹⁷ and typical semiconductors.⁶ Here, bearing in mind that many-body total-energy calculations are intended to be used in extreme situations where standard implementations of density-functional theory fail, we present numerical results for thin jellium slabs, whose most relevant feature is the strong inhomogeneity of the electron-density profile, as well as for the homogeneous electron gas over a wide range of densities.

Our concern is the evaluation of the particle-number difference,

$$\delta N = \frac{-i}{\pi} \int_{-\infty}^{+\infty} d\omega \operatorname{tr} [G(\omega) - G_0(\omega)] , \quad (17)$$

where tr denotes the spatial trace (we omit the explicit spatial variables for clarity and also consider only spin-unpolarized systems). Since both G and G_0 behave as $1/\omega$ for large frequencies, we can apply Cauchy's theorem and write Eq. (17) alternatively as

$$\delta N = \frac{1}{\pi} \int_{-\infty}^{+\infty} d\omega \operatorname{tr} [G(\mu + i\omega) - G_0(\mu_0 + i\omega)] , \quad (18)$$

where μ and μ_0 are the chemical potentials of the interacting and the noninteracting system, respectively, which correspond, by definition, to the position of the pole of the Green function at the Fermi surface. As the characteristic sharp structure of $G(\omega)$ (quasiparticle peaks and satellites) does not appear in the analytic continuation $G(\mu + i\omega)$, Eq. (18) is preferred for numerical integration. We hence follow some of the ideas suggested by Rojas *et al.*²⁷ and work exclusively in an imaginary time and frequency representation. An accurate evaluation of Eq. (18) furthermore requires a treatment of the high-frequency tails of G , which can be done easily with the numerical procedures described in Ref. 28.

For the homogeneous electron gas, an analytic expression exists for the noninteracting Green function $G_0(r, i\tau)$ in real space and imaginary time,²⁹ while the screened Coulomb interaction $W_0(k, i\omega)$ in the random-phase approximation is given analytically in reciprocal space by the dynamic Lindhard function.³⁰ The evaluation of the self-energy according to $\Sigma(r, i\tau) = iG_0(r, i\tau)W_0(r, i\tau)$ therefore only requires the numerical Fourier transform $W_0(k, i\omega) \rightarrow W_0(r, i\tau)$. It is this largely analytic approach that makes the present calculation especially precise.

At this stage we remark that the self-energy given by Eq. (16) has the same analytic structure as the underlying Green function G_0 , i.e., the poles of $\Sigma(\omega)$ are located in the upper (lower) complex half-plane for energies smaller (larger) than $\mu_0 = \frac{1}{2}k_F^2$, where k_F denotes the Fermi wave vector. As a consequence, an inconsistency arises because the true self-energy should have a polar structure identical to the *interacting* Green function with the chemical potential μ . The self-energy must therefore be appropriately shifted along the real frequency axis. In the imaginary time/frequency representation, this shift is automatically included in the backward transform

$$\Sigma(\mu + i\omega) = -i \int_{-\infty}^{+\infty} d\tau \Sigma(i\tau) e^{-i\omega\tau} . \quad (19)$$

The calculation of $\Sigma(\mu + i\omega)$, therefore, does not require an advance knowledge of μ , which can now be obtained from the relation $\mu = \mu_0 + \Sigma(k_F, \mu)$.




FIG. 2. Violation of the particle-number sum rule for the homogeneous electron gas in the G_0W_0 approximation. The relative error in the density is always negative and of the order of 0.1% in the range of metallic densities.

Finally, the interacting Green function is calculated in reciprocal space according to

$$G(k, \mu + i\omega) = \frac{1}{i\omega - \frac{1}{2}k^2 - \Sigma(k, \mu + i\omega) + \mu}. \quad (20)$$

In the same representation, the noninteracting Green function is given by

$$G_0(k, \mu_0 + i\omega) = \frac{1}{i\omega - \frac{1}{2}k^2 + \mu_0}, \quad (21)$$

so that the density variation is readily obtained from

$$\delta n = \int \frac{d^3k}{(2\pi)^3} \int_{-\infty}^{+\infty} \frac{d\omega}{\pi} [G(k, \mu + i\omega) - G_0(k, \mu_0 + i\omega)]. \quad (22)$$

In Fig. 2 the relative deviation $\delta n/n_0$ from the exact density is displayed as a function of the Wigner-Seitz radius r_s . In the high-density region $r_s < 1.8$ the particle number is slightly overestimated ($< 0.01\%$), while it is underestimated for lower densities. In the range of metallic densities this underestimation is of the order of 0.1%, but the error becomes increasingly important in the dilute limit (-1.7% for $r_s = 10$ and -6.1% for $r_s = 20$).

As pointed out above, it is also of interest to investigate the error resulting from the G_0W_0 method in the total number of particles for a strongly inhomogeneous system. The model we have chosen is a thin jellium slab with a background density $n_0 = (\frac{4}{3}\pi r_s^3)^{-1}$ and width L . The slab is bounded by two infinite planar walls, so that, if charge neutrality is assumed, the system is fully characterized by the lengths r_s and L . In this case, G_0 corresponds to the Kohn-Sham system obtained self-consistently with the local-density approximation (LDA) for the exchange-correlation potential V_{xc} , as is typically done in practical *ab initio* calculations.

With z chosen as the coordinate perpendicular to the planar walls, the translational symmetry of the system in the xy plane allows an efficient semianalytic evaluation of the relevant propagators. The screened Coulomb interaction is given by $W_0 = \epsilon_0^{-1}v$, where ϵ_0 denotes the dielectric function in the random-phase approximation. The latter is calculated as $\epsilon_0(k, i\omega)_{\alpha\beta}$ in the basis $\zeta_\alpha(z) \exp(i\mathbf{k} \cdot \boldsymbol{\rho})/\sqrt{S}$. Here $\zeta_\alpha(z)$ is a set of cosine functions, $\mathbf{k} = (k_x, k_y)$ and $\boldsymbol{\rho} = (x, y)$ denote the two-dimensional momentum and the position vector in the xy plane, respectively, and S is the slab surface. The matrix elements can be calculated analytically in terms of the scalar products $\langle \zeta_\alpha \phi_n | \phi_m \rangle$,^{24,31} where $\phi_n(z) \exp(i\mathbf{k} \cdot \boldsymbol{\rho})/\sqrt{S}$ are the single-particle eigenstates of the Kohn-Sham Hamiltonian h_{KS} . The matrix elements $v(k)_{\alpha\beta}$ of the Coulomb potential are likewise obtained analytically. The screened Coulomb interaction is then easily calculated by a matrix inversion for each value of k , and the real-space representation is given by expanding

$$W_0(\boldsymbol{\rho}, z, z'; i\tau) = i \sum_{\alpha, \beta} \int \frac{d^2k}{(2\pi)^2} \int_{-\infty}^{+\infty} \frac{d\omega}{2\pi} e^{i(\omega\tau + \mathbf{k} \cdot \boldsymbol{\rho})} \times \zeta_\alpha(z) \zeta_\beta(z') W_0(k, i\omega)_{\alpha\beta}. \quad (23)$$

The Green function $G_0(\boldsymbol{\rho}, z, z'; i\tau)$ is readily calculated from the Kohn-Sham eigenstates, and by employing Eq. (16) we obtain the self-energy in real space and imaginary time as well as, eventually, its representation $\Sigma(k, \mu + i\omega)_{nm}$ in the Kohn-Sham basis set. The presence of infinite confining walls implies a quick convergence with respect to the number of cosine and Kohn-Sham wave functions used in the calculation. The convergence is further accelerated by the analytic treatment of the asymptotic time and frequency tails of all operators.

The Green function is calculated in the basis of Kohn-Sham eigenstates according to

$$G(k, \mu + i\omega) = [i\omega - h_{KS}(k) - \Sigma(k, \mu + i\omega) + V_{xc}(k) + \mu]^{-1} \quad (24)$$

by a matrix inversion in the indices nm . Finally, the variation of the number of particles per surface unit is given by

$$\frac{\delta N}{S} = \sum_m \int \frac{d^2k}{(2\pi)^2} \int_{-\infty}^{+\infty} \frac{d\omega}{\pi} [G(k, \mu + i\omega)_{mm} - G_0(k, \mu_0 + i\omega)_{mm}], \quad (25)$$

where we have used the invariance of the trace with respect to any wave-function representation.

In Fig. 3 we plot the relative deviation of the particle number $\delta N/N$ in the G_0W_0 approximation for several configurations of the model system, keeping the exact number of particles per surface unit $n^{2D} = n_0 L = L/(\frac{4}{3}\pi r_s^3)$ constant. The limit $L \rightarrow 0$ thus corresponds to a two-dimensional (2D) homogeneous electron gas with density n^{2D} . Over the wide variation of the degree of




FIG. 3. Relative violation of particle number in the G_0W_0 approximation for thin jellium slabs of fixed 2D density $n^{2D} = 3/4\pi$ as a function of their thickness L (and the corresponding 3D density parameter r_s). A typical error bar is reported.

homogeneity shown in the figure, it is seen that $\delta N/N$ remains of similar magnitude as in the homogeneous case ($\lesssim 0.2\%$). This observation remains true for other 2D densities inside the range $[0.1, 1]$.

IV. CONCLUSIONS

In this paper we have rigorously obtained a general criterion which allows, by simple inspection, to verify whether a diagrammatic self-energy approximation satisfies the particle-number sum rule for an interacting electron system. As an application, we have demonstrated that the so-called G_0W_0 method does not yield the correct particle number, generalizing the conclusions of a previous analytic study for a Hubbard model Hamiltonian defined only on a discrete lattice.¹⁷ Thus this limitation of the G_0W_0 approximation has been fully confirmed for arbitrary electron systems. By performing a very precise integration of the spectral function, we have furthermore calculated the size of the error in the G_0W_0 particle number in two simple, but very distinct, families of electron systems. The error becomes large only outside the range of densities of physical interest.

ACKNOWLEDGMENTS

The authors thank Professor C.-O. Almbladh and Dr. J. E. Alvarellos for a thorough reading of the manuscript and for valuable discussions. This work was funded in part by the EU through the NANOPHASE Research Training Network (Contract No. HPRN-CT-2000-00167), the Spanish Education Ministry DGESIC Grant

No. PB97-1223-C02-02, and by the Deutscher Akademischer Austauschdienst and the British Council under the British-German Academic Research Collaboration (ARC) program.

* Electronic address: schindlmayr@fhi-berlin.mpg.de

- ¹ A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York, 1971).
- ² L. Hedin, Phys. Rev. **139**, A796 (1965).
- ³ F. Aryasetiawan and O. Gunnarsson, Rep. Prog. Phys. **61**, 237 (1998).
- ⁴ M. S. Hybertsen and S. G. Louie, Phys. Rev. Lett. **55**, 1418 (1985); Phys. Rev. B **34**, 5390 (1986).
- ⁵ R. W. Godby, M. Schlüter, and L. J. Sham, Phys. Rev. Lett. **56**, 2415 (1986); Phys. Rev. B **35**, 4170 (1987).
- ⁶ M. M. Rieger and R. W. Godby, Phys. Rev. B **58**, 1343 (1998).
- ⁷ B. Holm, Phys. Rev. Lett. **83**, 788 (1999).
- ⁸ B. Holm and F. Aryasetiawan, Phys. Rev. B **62**, 4858 (2000).
- ⁹ P. Sánchez-Friera and R. W. Godby, Phys. Rev. Lett. **85**, 5611 (2000).
- ¹⁰ P. García-González and R. W. Godby, Phys. Rev. B **63**, 075112 (2001).
- ¹¹ P. Hohenberg and W. Kohn, Phys. Rev. **136**, B864 (1964); W. Kohn and L. J. Sham, *ibid.* **140**, A1133 (1965).
- ¹² V. M. Galitskii and A. B. Migdal, Zh. Eksp. Teor. Fiz. **34**, 139 (1958) [Sov. Phys. JETP **7**, 96 (1958)].
- ¹³ G. Baym and L. P. Kadanoff, Phys. Rev. **124**, 287 (1961).
- ¹⁴ G. Baym, Phys. Rev. **127**, 1391 (1962).
- ¹⁵ F. J. Dyson, Phys. Rev. **75**, 486 (1949); **75**, 1736 (1949).
- ¹⁶ B. Holm, Ph.D. thesis, Lund University, 1997.
- ¹⁷ A. Schindlmayr, Phys. Rev. B **56**, 3528 (1997).
- ¹⁸ F. Aryasetiawan, L. Hedin, and K. Karlsson, Phys. Rev. Lett. **77**, 2268 (1996).
- ¹⁹ M. Springer, F. Aryasetiawan, and K. Karlsson, Phys. Rev. Lett. **80**, 2389 (1998).
- ²⁰ J. M. Luttinger, Phys. Rev. **119**, 1153 (1960).
- ²¹ M. Gell-Mann and F. Low, Phys. Rev. **84**, 350 (1951).
- ²² G. C. Wick, Phys. Rev. **80**, 268 (1950).
- ²³ R. P. Feynman, Phys. Rev. **76**, 749 (1949); **76**, 769 (1949).
- ²⁴ P. García-González and R. W. Godby (unpublished).
- ²⁵ J. M. Luttinger and J. C. Ward, Phys. Rev. **118**, 1417 (1960).
- ²⁶ C.-O. Almbladh, U. von Barth, and R. van Leeuwen, Int. J. Mod. Phys. B **13**, 535 (1999).
- ²⁷ H. N. Rojas, R. W. Godby, and R. J. Needs, Phys. Rev. Lett. **74**, 1827 (1995).
- ²⁸ L. Steinbeck, A. Rubio, L. Reining, M. Torrent, I. D. White, and R. W. Godby, Comput. Phys. Commun. **125**, 105 (2000).
- ²⁹ A. Schindlmayr, Phys. Rev. B **62**, 12 573 (2000).
- ³⁰ J. Lindhard, K. Dan. Vidensk. Selsk. Mat. Fys. Medd. **28**, No. 8 (1954).
- ³¹ A. G. Eguiluz, Phys. Rev. B **31**, 3303 (1985); J. M. Pitarke and A. G. Eguiluz, *ibid.* **63**, 045116 (2001).