

This is a repository copy of *The necessity of the Hadamard Condition*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/84763/>

Version: Accepted Version

Article:

Verch, Rainer and Fewster, Chris orcid.org/0000-0001-8915-5321 (2013) The necessity of the Hadamard Condition. *Classical and Quantum Gravity*. 235027. ISSN 1361-6382

<https://doi.org/10.1088/0264-9381/30/23/235027>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The Necessity of the Hadamard Condition

Christopher J Fewster^{(1)*} and Rainer Verch^{(2)†}

⁽¹⁾ Department of Mathematics, University of York, Heslington, York YO10 5DD, U.K.

⁽²⁾ Institut für Theoretische Physik, Universität Leipzig, 04009 Leipzig, Germany

October 3, 2013

Abstract

Hadamard states are generally considered as the physical states for linear quantized fields on curved spacetimes, for several good reasons. Here, we provide a new motivation for the Hadamard condition: for “ultrastatic slab spacetimes” with compact Cauchy surface, we show that the Wick squares of all time derivatives of the quantized Klein-Gordon field have finite fluctuations only if the Wick-ordering is defined with respect to a Hadamard state. This provides a converse to an important result of Brunetti and Fredenhagen. The recently proposed “S-J (Sorkin-Johnston) states” are shown, generically, to give infinite fluctuations for the Wick square of the time derivative of the field, further limiting their utility as reasonable states. Motivated by the S-J construction, we also study the general question of extending states that are pure (or given by density matrices relative to a pure state) on a double-cone region of Minkowski space. We prove a result for general quantum field theories showing that such states cannot be extended to any larger double-cone without encountering singular behaviour at the spacelike boundary of the inner region. In the context of the Klein-Gordon field this shows that even if an S-J state is Hadamard within the double cone, this must fail at the boundary.

1 Introduction

One of the fundamental difficulties in the theory of quantum fields on curved spacetimes is that generic spacetimes possess no symmetries that could serve to distinguish a preferred vacuum state. Instead, for linear fields, experience has led to the delineation of the class of Hadamard states [29] whose short-distance structure approximates that of states with finite energy density in Minkowski space, motivated by the equivalence principle. The

*chris.fewster@york.ac.uk

†verch@itp.uni-leipzig.de

two-point function of a Hadamard state ω of a Klein–Gordon field is required to take the form

$$W_\omega(x, y) = \lim_{\epsilon \rightarrow 0^+} \frac{U(x, y)}{\sigma_\epsilon(x, y)} + V(x, y) \ln(\sigma_\epsilon(x, y)) + H_\omega(x, y)$$

upon writing the distribution W_ω formally as a function of spacetime points x and y . Here, $\sigma_\epsilon(x, y)$ denotes the squared geodesic distance between x and y , together with a suitable regularization, U and V are C^∞ functions determined by spacetime metric and Klein-Gordon equation, while H_ω is a C^∞ function containing the state-dependence. (For full details of the definition, see [29]). It is worth noting that for static spacetimes, ground states or thermal equilibrium states of the quantized Klein-Gordon field are quasifree Hadamard states [33]. Hadamard states play an important role in computations, for they permit the evaluation of Wick polynomials, including the stress-energy tensor, and time-ordered products [11]. An elegant reformulation of the Hadamard condition using microlocal analysis, due to Radzikowski [32], has provided sufficiently good control on the class of Hadamard states that many results can be proved for general Hadamard states on general spacetime backgrounds. Examples include Quantum Energy Inequalities [17, 18], no-go results concerning chronology-violating spacetimes [28] and existence results for the semi-classical Friedmann equations [30]. Hadamard states also play an important role in understanding why black holes display the Hawking temperature [29]. It is also worth emphasizing that the class of Hadamard states is defined in a fully local and covariant way [13].

Nonetheless, one might wonder whether a criterion based on ultrashort distance behaviour is physically well-founded, especially in view of the widespread expectation that the continuum model of spacetime should break down at very small scales. An alternative approach might be to start with a model of fundamental spacetime structure and extrapolate macroscopic state selection criteria from that. Just such a proposal was made recently by Afshordi, Aslanbeigi and Sorkin [1] based on ideas developed in the causal set programme [26, 36], and giving a novel proposal for assigning a distinguished ‘‘S-J state’’ of the Klein–Gordon field to suitable spacetimes or spacetime regions. Further results appear in [2, 10, 5]. The idea is the following: consider a globally hyperbolic spacetime M with metric g such that \mathbf{E} , the difference of the advanced and retarded Green functions, extends to a bounded operator on $L^2(M, d\text{vol}_M)$. Then the operator $A = i\mathbf{E}$ is symmetric and [its closure] possesses the polar decomposition $A = U|A|$. One can form the positive part of A , given by $A^+ = (1/2)(|A| + A)$, and define a two-point function

$$W_{SJ}(f, h) = \langle \bar{f}, A^+ h \rangle, \quad f, h \in C_0^\infty(M), \quad (1.1)$$

where the bar denotes complex conjugation and $\langle f, h \rangle = \int_M \bar{f} h d\text{vol}_M$ denotes the L^2 -scalar product; this defines the two-point function of the S-J state [1].

In [19], we analyzed the S-J prescription and showed that W_{SJ} is indeed the 2-point function of a pure quasifree state on the algebra of smeared Klein–Gordon fields on M whenever the prescription is defined, and that this certainly holds whenever M can be isometrically embedded as a globally hyperbolic and relatively compact sub-spacetime inside a larger globally hyperbolic spacetime. However, we also showed that the S-J states

suffer from serious problems. In particular, they generically fail to be Hadamard when computed for ultrastatic ‘slab’ spacetimes with compact spatial section; moreover, they fail to be defined in a locally covariant way. This brings into sharp focus the question of how seriously the Hadamard condition should be taken in the context of models inspired by discontinuous fundamental spacetime structure.

In the present paper, we aim to bring further clarity to a number of issues surrounding the desirability of the Hadamard condition and the problems that must be faced if one aims to define a state on a local region of spacetime. We emphasize that our focus is much broader than S-J states, although they have provided a motivation for some of our results.

First, we consider general pure quasifree states of the Klein–Gordon field on general ultrastatic slab spacetimes with compact spatial section. Any such state ω can be used as the basis of a normal ordering prescription and the construction of Wick powers. Of course, all Wick powers (other than the zero’th order) will necessarily have vanishing expectation value in the basic state ω ; this would apply to the stress-energy tensor as well. However, in order to use ω as the basis for a fully fledged perturbative construction of a self-interacting theory, or for an analysis of the semiclassical Einstein equations, one would need the Wick polynomials to have finite (and preferably small) fluctuations in the state ω . Our main result of Section 2 is that ω has finite fluctuations for all Wick polynomials only if it is Hadamard (the converse statement is established, for all spacetimes, in [11]). In particular, we show that the Wick square of the first time-derivative of the field has infinite fluctuations in the S-J state. However, the main force of our result is to provide a motivation for the Hadamard condition that is less tightly bound to the ultrashort distance structure.

While these results cast serious doubt on the utility of S-J states defined on spatially compact ultrastatic slab spacetimes, strengthening the findings of our previous paper [19], we have not yet addressed the properties of S-J states defined on bounded regions of Minkowski space. Our second result concerns this question, indeed, we will address it in the model-independent context of operator-algebraic quantum field theory. In this setting, we will show that no state that is pure (or even normal to a pure state, i.e. arising as a density matrix state in the GNS Hilbert space representation of a pure state) on the algebra of observables of a double cone region can extend to the observable algebra of a strictly larger double cone region without developing pathologies at the spacelike boundary of the smaller double cone: the extended state necessarily fails to admit a stable short-distance scaling limit at any point of this boundary. To borrow a term from recent controversial discussions of black hole evaporation [3], one might say that the state on the inner region is protected by a firewall (or, ‘energetic curtain’ [9]). In the particular example of the free Klein–Gordon field, the S-J state of any double cone is pure [19] and our result entails that any extension to a larger double cone must fail to be Hadamard at all points of the spacelike boundary, because the Hadamard condition implies the existence of a stable scaling limit. The argument relies on the type of local von Neumann algebras of the double cones. It is well known that under standard assumptions the local von Neumann algebras in quantum field theory are of type III, while the von Neumann algebra induced by a pure state is of type I. In our case, we draw on results taken from [22, 7] (see also [15]) establishing that the type III property follows from the existence of suitable stable scaling limits which

are expected to correspond to the theory admitting an ultra-violet renormalization group fixed point. Our result is consistent with the computation performed in [2], where a two-dimensional diamond region was considered for the massless free field – near the spacelike boundary the two-point function is approximated by that of the ground state of the field in the presence of a mirror at the boundary, so extension through the boundary will not be possible while maintaining the Hadamard form.

The result just described indicates that S-J states defined on extendible spacetime regions have undesirable features and the case of double cones seems to suggest that this has to do with the ‘sharp cut-off’ at the boundary of the spacetime region on which they are defined. It has been suggested to us by various people (we thank Jorma Louko and Rafael Sorkin in particular) that perhaps the Hadamard condition could be restored for a modified version of the S-J state prescription in which the ‘sharp boundary’ is softened. We study two simple such variants gained by averaging procedures, in the ultrastatic slab situation, and show explicitly that neither leads to Hadamard states. Very recently, however, Brum and Fredenhagen [10] have suggested an altered definition of S-J states on certain globally hyperbolic slab spacetimes which corresponds to a different ‘softening-the-boundary’ procedure, and their proposal leads to Hadamard states. This indeed indicates that the failure of the original definition of S-J states to produce Hadamard states stems from the presence of the ‘sharp boundary’ in timelike direction in the case of ultrastatic slab spacetimes considered in [19]. However, the altered definition of [10] involves a dependence on a smoothing function as an extra parameter — which is in the very nature of smoothing procedures — and this is at variance with the original idea that a unique S-J state can be assigned to any spacetime without the need of specifying other parameters or possible choices [1]. Moreover, it is emphasized in [10] that the modified states are unlikely to carry the interpretation of vacuum states. Nevertheless, the proposal of [10] does lead to a new method of constructing Hadamard states for certain static or expanding slab spacetimes, and it would be of interest to see if that could be extended to a larger class of spacetimes.

As mentioned in [10], the definition of S-J states for the quantized scalar Klein-Gordon field seems to have had a forerunner for the case of the quantized Dirac field in the form of the ‘Fermionic projector’ proposed some time ago by Felix Finster [20], see also the recent paper [21]. (We would like to thank Felix Finster for some comments on that point.)

The organization of the present work is as follows. In Section 2, we establish that the requirement of finite variance for the Wick squares of time-derivatives of any order of the quantized Klein-Gordon field on an ultrastatic slab spacetime with compact Cauchy surface implies that the states with respect to which the Wick-ordering is defined must be Hadamard. We also show that S-J states do not comply with this requirement as Wick squares of the time-derivative of the field have infinite fluctuations for S-J states.

In Section 3, we show for general quantum field theories on Minkowski spacetime that a state on the algebra of observables of double cone whose induced GNS Hilbert space representation is of type I cannot be extended to a state on the observable algebra of a larger spacetime region unless the state fails to have a stable short-distance scaling limit at every point of the double cone’s (spacelike) boundary. Such an assertion may be known to some experts, but would surely be hard to trace in this form in the literature.

We then apply this result to show for the case of the quantized Klein-Gordon field on Minkowski spacetime that S-J states defined for double cones cannot be extended to states on observable algebras of larger spacetime regions which are normal to Hadamard states. (The result generalizes, as we shall indicate, also to certain curved spacetime situations.)

In Section 4, we present two simple averaging procedures for S-J states defined for the quantized Klein-Gordon field on ultrastatic slab spacetimes, and we show that these procedures do not produce Hadamard states. We also show that our previous results in [19] on S-J states failing to be Hadamard for ultrastatic slabs can be strengthened by invoking a local-to-global result on the Hadamard property of two-point functions of states due to Radzikowski [31]. In particular, if the Cauchy-surface of the ultrastatic slab spacetime is acted on by a transitive group of isometries, then the corresponding S-J state does not only fail to be Hadamard on the ultrastatic slab spacetime as a whole (i.e., ‘somewhere’), but it fails to be Hadamard everywhere.

The article is concluded by a discussion in Sec. 5.

2 Fluctuations in Wick polynomials and Hadamard property

In this section, we investigate how the requirement that certain Wick polynomials have finite fluctuations enforces the Hadamard condition. In passing, the S-J states on ultrastatic slabs are shown to produce infinite fluctuations for smearings of $:\dot{\phi}^2:$: except for at most a set of measure zero in τ . Throughout the section, we will say that a field has finite fluctuation in a state if all of its smearings with test functions have finite fluctuations in that state.

We consider the massive scalar field $(\square + m^2)\phi = 0$, $m > 0$, on d -dimensional ultrastatic slab spacetimes of the form $M = (-\tau, \tau) \times \Sigma$ with metric $ds^2 = dt^2 - h_{ij}dx^i dx^j$, where Σ is a compact manifold with smooth Riemannian metric h and $\tau \in (0, \infty]$. Then there exists a complete orthonormal basis ψ_j ($j \in J$) for $L^2(\Sigma)$ (with the volume measure induced by h), comprising eigenfunctions of $K = -\Delta + m^2$, where Δ is the Laplacian on (Σ, h) and $K\psi_j = \omega_j^2\psi_j$ with $\omega_j > 0$. The index set J is countable and we assume (without loss of generality) that there exists, for each $j \in J$, a (unique) $\bar{j} \in J$ such that $\psi_{\bar{j}} = \overline{\psi_j}$; we allow for the possibility that \bar{j} might equal j . Clearly, $\omega_{\bar{j}} = \omega_j$ for all $j \in J$.

We will consider the class of quasifree pure states of the quantized Klein-Gordon field on M – equivalently, irreducible Fock space representations of the theory, with the Fock vacuum vector representing the state. Given the simple structure of M , the general theory (see, e.g., [44], or [29]) reduces to familiar constructions using mode functions. As usual, the aim is to identify a set of mode solutions to the field equation of the form $T_j(t)\psi_j(\underline{x})$, which, together with their complex conjugates, form a basis for the space of complex valued solutions in a suitable sense. The most general choice is, of course, $T_j(t) = \alpha_j e^{-i\omega_j t} + \beta_j e^{i\omega_j t}$, for complex-valued α_j and β_j (obeying certain conditions described below). The choice of basis amounts to the choice of a state: for example, the ultrastatic vacuum results from

the choice $\alpha_j = 1$, $\beta_j = 0$ for all $j \in J$. For $\tau < \infty$, the S-J prescription also takes this form [19],¹ with

$$\alpha_j = \sqrt{\frac{\|S_j\|}{\|C_j\|}} \left(1 - \frac{\delta_j}{2}\right), \quad \beta_j = \sqrt{\frac{\|S_j\|}{\|C_j\|}} \frac{\delta_j}{2},$$

where $\|S_j\|$ and $\|C_j\|$ denote the $L^2(-\tau, \tau)$ norms of the functions $S_j(t) = \sin \omega_j t$ and $C_j(t) = \cos \omega_j t$, i.e.,

$$\|C_j\|^2 = \tau(1 + \text{sinc } 2\omega_j \tau), \quad \|S_j\|^2 = \tau(1 - \text{sinc } 2\omega_j \tau),$$

and

$$\delta_j = 1 - \frac{\|C_j\|}{\|S_j\|} = 1 - \sqrt{1 + \frac{2 \text{sinc } 2\omega_j \tau}{1 - \text{sinc } 2\omega_j \tau}} = -\text{sinc } 2\omega_j \tau + O((\omega_j \tau)^{-2}). \quad (2.1)$$

Here, we have used the notation $\text{sinc } x = \sin x/x$.

Returning to the general case, one then constructs a Fock space, in which the field will be given as

$$\phi(t, \underline{x}) = \sum_{j \in J} \frac{1}{\sqrt{2\omega_j}} \left((\alpha_j e^{-i\omega_j t} + \beta_j e^{i\omega_j t}) \psi_j(\underline{x}) a_j + \text{h.c.} \right), \quad (2.2)$$

where $[a_j, a_k^*] = \delta_{jk} \mathbf{1}$ and the Fock vacuum, annihilated by all a_j , will be denoted by Ω . The corresponding two-point function is given formally by

$$W(t, \underline{x}; t', \underline{x}') = \sum_{j \in J} \frac{1}{2\omega_j} (\alpha_j e^{-i\omega_j t} + \beta_j e^{i\omega_j t}) \left(\overline{\alpha_j} e^{i\omega_j t'} + \overline{\beta_j} e^{-i\omega_j t'} \right) \psi_j(\underline{x}) \psi_{\bar{j}}(\underline{x}').$$

As mentioned above, the α_j and β_j are subject to various conditions. We will require that the series for W should converge in the sense of distributions, which means that if the summands are individually smeared against $F(t, \underline{x})G(t', \underline{x}')$ for test functions $F, G \in C_0^\infty((-\tau, \tau) \times \Sigma)$, then the resulting series should converge in \mathbb{C} , with the sum defining the value of $W(F, G)$. Furthermore, the commutation relations of the field require that

$$\begin{aligned} W(t, \underline{x}; t', \underline{x}') - W(t', \underline{x}'; t, \underline{x}) &= iE(t, \underline{x}; t', \underline{x}') \\ &= \sum_{j \in J} \frac{1}{2\omega_j} \left(e^{i\omega_j(t-t')} - e^{-i\omega_j(t-t')} \right) \psi_j(\underline{x}) \psi_{\bar{j}}(\underline{x}'), \end{aligned}$$

which leads to the relations

$$|\alpha_j|^2 - |\beta_{\bar{j}}|^2 = 1, \quad \alpha_j \overline{\beta_{\bar{j}}} = \alpha_{\bar{j}} \overline{\beta_j} \quad (j \in J) \quad (2.3)$$

on considering smearings against test functions of the form $f(t)g(t')\psi_{\bar{j}}(\underline{x})\psi_j(\underline{x}')$. Taking absolute values in the second set of relations in Eq. (2.3), and using the first, we obtain

¹The S-J prescription fails in the case $\tau = \infty$. If one takes a suitable limit of S-J states as $\tau \rightarrow \infty$, however, the ultrastatic ground state is obtained in the limit [19].

$|\alpha_j|^2(1 + |\alpha_{\bar{j}}|^2) = |\alpha_{\bar{j}}|^2(1 + |\alpha_j|^2)$ for all j and hence $|\alpha_j| = |\alpha_{\bar{j}}|$, $|\beta_j| = |\beta_{\bar{j}}|$ for all $j \in J$. As a rephrasing

$$\alpha_j \mapsto \alpha_j e^{i\chi_j}, \quad \beta_j \mapsto \beta_j e^{i\chi_j} \quad (\chi_j \in \mathbb{R})$$

leaves W invariant, and corresponds to a simple unitary transformation on Fock space, we may assume without loss of generality that $\alpha_j \geq 1$ for all $j \in J$, whereupon the class of representations considered is parameterized by the choice of each $\beta_j \in \mathbb{C}$ subject to $\beta_j = \beta_{\bar{j}}$; we then have $\alpha_j = \sqrt{1 + |\beta_j|^2}$. As mentioned, it is additionally required that the series for W should converge in the sense of distributions, so the β_j cannot be too badly behaved. Rather than study the convergence question in detail, we assume for simplicity that there is a polynomial Q such that $|\beta_j| \leq Q(\omega_j)$ for all $j \in J$, which entails a similar polynomial bound on the growth of α_j . In fact, as we will soon see, the β_j must actually decay if the fluctuations of $:\dot{\phi}^2:(F)$ are to be finite. For simplicity of presentation, we proceed with formal computation, though our conclusions would also be reached by a more careful analysis.

Defining normal ordering as usual in the Fock space, we find (formally)

$$\begin{aligned} :\dot{\phi}^2(t, \underline{x}):\Omega &= -\frac{1}{2} \sum_{j, j' \in J} \sqrt{\omega_j \omega_{j'}} \overline{(\alpha_j e^{-i\omega_j t} - \beta_j e^{i\omega_j t}) (\alpha_{j'} e^{-i\omega_{j'} t} - \beta_{j'} e^{i\omega_{j'} t})} \psi_j(\underline{x}) \psi_{j'}(\underline{x}) a_j^* a_{j'}^* \Omega \\ &= -\frac{1}{2} \sum_{j, j' \in J} \sqrt{\omega_j \omega_{j'}} (\alpha_j e^{i\omega_j t} - \bar{\beta}_j e^{-i\omega_j t}) (\alpha_{j'} e^{i\omega_{j'} t} - \bar{\beta}_{j'} e^{-i\omega_{j'} t}) \overline{\psi_j(\underline{x})} \overline{\psi_{j'}(\underline{x})} a_j^* a_{j'}^* \Omega \end{aligned}$$

where, in the second step, we have used the fact that the α_j are real, and relabelled the j' sum by $j' \mapsto \bar{j}'$ (recalling also that $\beta_{\bar{j}'} = \beta_{j'}$ and $\overline{\psi_{\bar{j}'}} = \psi_{j'}$). We smear against test functions so that

$$:\dot{\phi}^2:(f \otimes g) = \int_{-\tau}^{\tau} dt \int_{\Sigma} d\text{vol}_{\Sigma}(\underline{x}) :\dot{\phi}^2(t, \underline{x}): f(t) g(\underline{x}),$$

with g smooth and compactly supported on Σ , and f smooth and compactly supported in $(-\tau, \tau)$.

As $:\dot{\phi}^2:(f \otimes g)$ has vanishing expectation in the state Ω , its fluctuation (or dispersion) is simply the square root of the expectation of its square; in short, the norm of $:\dot{\phi}^2:(f \otimes g)\Omega$. For simplicity, we will take $g(\underline{x}) \equiv 1$ (recall that Σ is compact) and f to be real and even, so that its Fourier transform

$$\hat{f}(\omega) = \int dt f(t) e^{i\omega t}$$

is also real-valued and even, and decays faster than polynomially as $\omega \rightarrow \infty$. Together with the orthonormality of the ψ_j , our assumptions reduce the sum over j and j' to those terms for which $j' = j$, giving

$$:\dot{\phi}^2:(f \otimes 1)\Omega = -\frac{1}{2} \sum_{j \in J} \omega_j \left((\alpha_j^2 + \bar{\beta}_j^2) \hat{f}(2\omega_j) - 2\alpha_j \bar{\beta}_j \hat{f}(0) \right) a_j^* a_j^* \Omega,$$

where we have also used that \hat{f} is real and even. Hence

$$\|:\dot{\phi}^2:(f \otimes 1)\Omega\|^2 = \frac{1}{2} \sum_{j \in J} \omega_j^2 \left| (\alpha_j^2 + \overline{\beta_j}^2) \hat{f}(2\omega_j) - 2\alpha_j \overline{\beta_j} \hat{f}(0) \right|^2. \quad (2.4)$$

While the above computation was rather formal, a more careful analysis leads to the same identity provided that the right-hand side converges; if it diverges, this may be interpreted as showing that $:\dot{\phi}^2:(f \otimes 1)$ has infinite fluctuations in the state Ω .²

In order to estimate this sum, we will use two facts. First, general results on the spectrum of the Laplacian [41, Ch. 8, §3] permit us to choose $N \in \mathbb{N}$ for which $\sum_{j \in J} (\omega_j^2 + m^2)^{-2N}$ converges. Second, as f is smooth and compactly supported, $\sup_{j \in J} |\hat{f}(\omega_j) P(\omega_j)|$ is finite for any polynomial P . Expanding the absolute value in (2.4), we find two terms involving $\hat{f}(2\omega_j)$, both of which lead to convergent sums on combining the two facts just mentioned with the polynomial growth bounds on α_j and β_j . Hence $:\dot{\phi}^2:(f \otimes 1)\Omega$ has finite norm if and only if

$$\sum_{j \in J} \omega_j^2 |\alpha_j \beta_j|^2 < \infty.$$

Introducing parameters $\zeta_j = \tanh^{-1}(|\beta_j|/|\alpha_j|)$, the condition is equivalent to

$$\sum_{j \in J} \omega_j^2 \sinh^2(2\zeta_j) < \infty, \quad (2.5)$$

which implies that $\omega_j \zeta_j \rightarrow 0$ as $j \rightarrow \infty$ in some (and hence any) ordering of J by the natural numbers.

In the case of the S-J state on a slab with $\tau < \infty$, for instance, we have $\alpha_j \sim 1$ and $\beta_j \sim -\text{sinc } 2\omega_j \tau$ as $j \rightarrow \infty$, so $\omega_j \zeta_j \sim (2\tau)^{-1} \sin 2\omega_j \tau$. The following is now essentially immediate, and provides a further pathology of S-J states that casts substantial doubt on their physical relevance.

Theorem 2.1 *For general (Σ, h) , the set of $\tau \in (0, \infty)$ for which $:\dot{\phi}^2:$ has finite fluctuations in the corresponding S-J state is contained in a set of measure zero. If (Σ, h) is either a flat 3-torus or round 3-sphere, then there is no $\tau \in (0, \infty)$ for which $:\dot{\phi}^2:$ has finite fluctuations in the corresponding S-J state.*

Proof: It was shown in [19] that the set of $\tau \in (0, \infty)$ for which $\sin 2\omega_j \tau \rightarrow 0$ has vanishing Lebesgue measure for general (Σ, h) and is empty in the two special cases mentioned. \square

This result can be seen in the same vein as the observation made by Brunetti, Fredenhagen and Hollands that the ‘ α -vacua’ states on de Sitter spacetime yield infinite fluctuations for the Wick-ordered stress-energy tensor, which casts serious doubts on their utility as physical states [12].

²The point is that expressions such as $\langle a_j^* a_k^* \Omega | : \dot{\phi}^2 : (f \otimes 1) \Omega \rangle$ may be defined *in the sense of quadratic forms* by a point-splitting prescription relative to the two-point function W ; the question is whether these matrix elements can be interpreted as literal Fock space inner products, which amounts to the convergence or otherwise of the right-hand side of (2.4).

Returning to the general case, the above argument is easily modified to deal with Wick squares of higher time derivatives of the field, with consequent increase in the power of ω_j in the bound Eq. (2.5). Moreover, one sees easily that sufficient decay of the ζ_j implies that $:(\partial^k \phi / \partial t^k)^2:(f \otimes 1)\Omega$ has finite norm for any $f \in C_0^\infty(-\tau, \tau)$, not just those that are real and even. This proves the following result.

Theorem 2.2 *In any Fock representation of the field taking the form Eq. (2.2), in which the β_j obey a polynomial bound $|\beta_j| \leq Q(\omega_j)$ for some polynomial Q and all $j \in J$, the following are equivalent:*

1. $:(\partial^k \phi / \partial t^k)^2:(f \otimes 1)\Omega$ has finite norm for all $k \in \mathbb{N}$, where $f \in C_0^\infty(-\tau, \tau)$;
2. $\sup_j |P(\omega_j) \sinh 2\zeta_j| < \infty$ for all polynomials P .

In other words, the requirement of finite fluctuations for the above operators forces $\alpha_j \rightarrow 1$ and $\beta_j \rightarrow 0$ with rapid convergence (faster than polynomially in $1/\omega_j$).

We now show that the state constructed under these conditions is, in fact, Hadamard. This requires some use of microlocal techniques to compute the wave-front set $\text{WF}(u)$ of the Hilbert-space-valued distribution $u(F) = \phi(F)\Omega$.³ The procedure is as follows; references include [38, 35]. Fix any $p = (t_p, \underline{x}_p)$ in the slab and consider local coordinates y^i ($1 \leq i \leq d-1$) for Σ near \underline{x}_p and use $y^0 = t$ to give spacetime coordinates $y^\mu(t, \underline{x})$ ($0 \leq \mu \leq n$) near p .⁴ Let $k' \in T_p^*M$ be a nonzero covector at p , with components k'_μ in the chosen coordinates, and define a function $e_{\lambda, k'}(t, \underline{x}) = e^{i\lambda k'_\mu y^\mu(t, \underline{x})}$ on the y^μ coordinate chart. Then the pair $(p, k) \in T_p^*M$ is said to be a *regular direction* for u if there exists $F \in C_0^\infty(M)$ with $F(p) \neq 0$ such that $\|u(Fe_{\lambda, k'})\| = O(\lambda^{-M})$ as $\lambda \rightarrow +\infty$ for every $M \in \mathbb{N}$ uniformly for all k' in an open neighbourhood of k . The *wave-front set* $\text{WF}(u)$ is the set of all $(p, k) \in T^*M$ with $k \neq 0$ that are *not* regular directions. It is known [38] that the state Ω is Hadamard if and only if $\text{WF}(u) \subset \mathcal{N}^-$, the bundle of past-directed null covectors on M .

To start, we will show that every $(p, k) \in T^*M$ with $k_0 > 0$ is a regular direction for u . Observe that

$$\phi(f \otimes g)\Omega = \sum_{j \in J} \frac{1}{\sqrt{2\omega_j}} \left(\alpha_j \hat{f}(\omega_j) + \overline{\beta_j} \hat{f}(-\omega_j) \right) \langle \psi_j | g \rangle a_j^* \Omega$$

for any $f \in C_0^\infty(-\tau, \tau)$ and $g \in C_0^\infty(\Sigma)$, and hence

$$\|\phi(f \otimes g)\Omega\|^2 \leq \|g\|^2 \sum_{j \in J} \frac{1}{2\omega_j} \left| \alpha_j \hat{f}(\omega_j) + \overline{\beta_j} \hat{f}(-\omega_j) \right|^2. \quad (2.6)$$

Now fix f and g so that $f(t_p) \neq 0$ and $g(\underline{x}_p) \neq 0$, with g supported inside the coordinate chart of the y^i . The inequality (2.6) implies

$$\|\phi((f \otimes g)e_{\lambda, k'})\Omega\|^2 \leq \|g\|^2 \Upsilon(\lambda k'_0), \quad (2.7)$$

³The original microlocal formulation of the Hadamard condition used the two-point function [32].

⁴The specific choice of coordinates is irrelevant.

where

$$\Upsilon(\mu) = \sum_{j \in J} \frac{1}{2\omega_j} \left| \alpha_j \hat{f}(\mu + \omega_j) + \overline{\beta_j} \hat{f}(\mu - \omega_j) \right|^2$$

decays faster than any inverse power as $\mu \rightarrow +\infty$, assuming that $\beta_j \rightarrow 0$ faster than polynomially in $1/\omega_j$, as will be shown in Appendix B. Thus for any k with $k_0 > 0$, the right-hand side of (2.7) decays rapidly as $\lambda \rightarrow +\infty$, uniformly for k' in the open neighbourhood $\{\ell \in \mathbb{R}^4 : \ell_0 > k_0/2\}$ of k ; it follows that all points of the form $(p, k) \in T^*M$ with $k_0 > 0$ are regular directions for the vector-valued distribution u , and therefore $\text{WF}(u) \subset \{(p, k) \in T^*M : k_0 < 0\}$. However, as u is a weak solution to the Klein–Gordon equation, its wave-front set must also be contained in the characteristic set of the Klein–Gordon operator [16], i.e., the bundle of null co-vectors. Therefore, $\text{WF}(\phi(\cdot)\Omega) \subset \mathcal{N}^-$, the bundle of past-directed null covectors, and this establishes that the state Ω is in fact Hadamard. On the other hand, all Wick polynomials have finite fluctuations in Hadamard states by the analysis in [11] (in any globally hyperbolic spacetime) so we have proved:

Theorem 2.3 *The state Ω is Hadamard if and only if $(\partial^k \phi / \partial t^k)^2$ has finite fluctuations in Ω for every $k \in \mathbb{N}_0$.*

3 Locally pure states and boundary singularities

The S-J prescription assigns a pure state to bounded regions of spacetime [1, 19]. In this section, we explain why any such prescription can be expected to result in singular behaviour (of a type described below) at the boundary of the region. In fact, the same applies to all states on the bounded region that are defined by a density matrix in the GNS representation of a pure state. We work in an algebraic framework of quantum field theory on $1 + d$ -dimensional Minkowski space [23] and for the most part, our considerations are not restricted to the particular example of the free scalar field. A brief summary of how the Klein–Gordon field fits into the general operator algebraic framework can be found in Appendix A. Our results are consistent with (but go much further than) the computations performed in [2] where the two-point function of the massless Klein–Gordon field is studied near the boundary of a two-dimensional Minkowski diamond.

The starting assumption is that, to each open relatively compact set O of Minkowski space, there is a corresponding unital C^* -algebra $\mathcal{A}(O)$, consisting of the observables of the theory localised in O . These algebras obey the following conditions: (a) if $O \subset \tilde{O}$ then $\mathcal{A}(O)$ is assumed to be a subalgebra of $\mathcal{A}(\tilde{O})$ and they share a common unit; (b) $\mathcal{A}(O)$ and $\mathcal{A}(\tilde{O})$ are assumed to commute elementwise if O and \tilde{O} are spacelike separated. We will not make any assumptions about the action of spacetime symmetries at this level.

Let M be a double-cone in Minkowski space, with corresponding algebra $\mathcal{A}(M)$. Recall that a state on $\mathcal{A}(M)$ is a linear map $\omega : \mathcal{A}(M) \rightarrow \mathbb{C}$ that is normalised ($\omega(\mathbf{1}) = 1$) and positive ($\omega(A^*A) \geq 0$ for all $A \in \mathcal{A}(M)$). Any state on $\mathcal{A}(M)$ induces a Hilbert space GNS representation; this representation is irreducible if and only if the state is pure [8, II.6.4.8]. Further, any state ω on $\mathcal{A}(M)$ induces a whole class of states that are defined by

density matrices in the GNS representation of ω ; these states are said to be *normal* to ω (in particular, this includes all vector states in the representation, and ω itself).

We wish to show that no state normal to a pure state of the algebra $\mathcal{A}(M)$ can be extended to any strictly larger double cone N without encountering singular behaviour at the boundary of M . Our argument actually proceeds in reverse, by showing that singularity-free states of $\mathcal{A}(N)$ do not restrict to $\mathcal{A}(M)$ as pure states, or even states that are normal to pure states.

Let ω be any state of $\mathcal{A}(N)$. It induces a GNS representation $(\mathcal{H}, \pi, \Omega)$ of $\mathcal{A}(N)$ so that $\langle \Omega | \pi(A)\Omega \rangle = \omega(A)$ for all $A \in \mathcal{A}(N)$, with Ω as a cyclic vector for the representation.⁵ Thus, a state ω' of $\mathcal{A}(N)$ is normal to ω if there is a density matrix ϱ on \mathcal{H} such that $\omega'(A) = \text{Tr}(\varrho\pi(A))$ holds for all $A \in \mathcal{A}(N)$. We assume, as a requirement on ω , that the Hilbert space \mathcal{H} is separable. If O is any open bounded region contained in N , we obtain a subalgebra $\pi(\mathcal{A}(O))$ of the bounded operators $\mathcal{B}(\mathcal{H})$ of \mathcal{H} , and a corresponding von Neumann algebra $\mathcal{N}(O) := \pi(\mathcal{A}(O))''$, where the prime denotes the operation of forming the commutant.⁶ By von Neumann's bicommutant theorem, $\mathcal{N}(O)$ coincides with the weak closure of $\pi(\mathcal{A}(O))$ [8, I.9.1.1].

To explain the singular behaviour we have in mind, we assume in addition that the theory has a description involving field operators. Specifically, we assume that there is a dense domain $\mathcal{D} \subset \mathcal{H}$ containing the GNS vector Ω and, to every real-valued test function $f \in C_0^\infty(N)$, there is a corresponding hermitian smeared field operator $\phi(f)$ leaving \mathcal{D} invariant, depending linearly on f and so that the operator closures $\overline{\phi(f)}$ exist and are affiliated to the local algebras.⁷ We may form n -point functions of the field ϕ in state ω by setting

$$W_\omega^{(n)}(f_1, \dots, f_n) = \langle \Omega | \phi(f_1) \cdots \phi(f_n)\Omega \rangle$$

for any test functions $f_k \in C_0^\infty(N)$ and we require these to determine distributions on N .

The next step is to consider scalings of the n -point functions. Let p be a point in N and define the scaling maps

$$\beta_{p,\lambda}f(x) = f((x-p)/\lambda), \quad f \in C_0^\infty(\mathbb{R}^{1+d}), \quad x \in N, \quad 1 > \lambda > 0$$

which contract the supports of test-functions to the point p as $\lambda \rightarrow 0$. The following definition is based on a condition in [7, §16.2.4].

Definition 3.1 *The state ω has a regular scaling limit at p if, in addition to the conditions above, the following two properties hold:*

(i) *The intersection of the local von Neumann algebras in the GNS representation induced by ω is trivial at p , i.e.*

$$\bigcap_{O \ni p} \mathcal{N}(O) = \mathbb{C}\mathbf{1};$$

⁵Cyclicity means that the set $\pi(\mathcal{A}(N))\Omega$ is dense in \mathcal{H} .

⁶The commutant of any subset \mathcal{X} of $\mathcal{B}(\mathcal{H})$ is defined as $\mathcal{X}' = \{B \in \mathcal{B}(\mathcal{H}) : AB = BA \text{ for all } A \in \mathcal{X}\}$, and the bicommutant is defined as $\mathcal{X}'' = (\mathcal{X}')'$.

⁷That is, when one takes a polar decomposition $\overline{\phi(f)} = U_f|\overline{\phi(f)}|$, the partial isometry U_f , and every bounded function of $|\overline{\phi(f)}|$, belong to $\mathcal{N}(O)$ for any O containing the support of f .

(ii) There is some monotone, positive-valued function $\lambda \mapsto \nu(\lambda)$ so that the limits

$$W_0^{(n)}(f_1, \dots, f_n) = \lim_{\lambda \rightarrow 0} \nu(\lambda)^n W_\omega^{(n)}(\beta_{p,\lambda} f_1, \dots, \beta_{p,\lambda} f_n)$$

exist for all $n \in \mathbb{N}$ and all $f_j \in C_0^\infty(\mathbb{R}^{1+d})$ and define (non-trivial) n -point functions satisfying the Wightman axioms for a quantum field theory⁸ in its vacuum representation [37].

The existence of regular scaling limits is expected for quantum field theories possessing an ultraviolet fixed point [14]. We regard the failure of a regular scaling limit as indicating a pathology of the state ω . For our purposes, the importance of scaling limits is that they permit us to determine the *type* of the local von Neumann algebras. We will only need to consider algebras of types I and III. A von Neumann algebra \mathcal{N} is said to be of type I if it is isomorphic [as a von Neumann algebra] to the algebra of bounded operators on some Hilbert space. A von Neumann algebra \mathcal{N} is of type III if it contains no finite projections, which can also be expressed as follows: If $P \in \mathcal{N}$ is any non-zero projection operator, then there is a proper sub-projection $Q \in \mathcal{N}$ (i.e. $QP = PQ = Q$ and $Q \neq P$) together with an operator $V \in \mathcal{N}$ such that $V^*V = P$ and $VV^* = Q$. The type is preserved under von Neumann algebra isomorphisms. Moreover, in cases where the von Neumann algebra is obtained by taking the weak closure of a representation of a C^* -algebra, we describe the representation as having the same type as the von Neumann algebra.

The following is a restatement of [7, Theorem 16.2.18], which is a development of a seminal result of Fredenhagen [22].

Proposition 3.2 *Let M be a double cone with compact closure contained in N and let p be a point in the spacelike boundary of M . Suppose that the state ω of $\mathcal{A}(N)$ has a regular scaling limit at p . Then the local von Neumann algebra $\mathcal{N}(M) = \pi(\mathcal{A}(M))''$ is of type III, i.e., the restriction $\pi|_{\mathcal{A}(M)}$ is a type III representation of $\mathcal{A}(M)$.*

(In fact, even type III₁ is proved in the references given.) Our result is a simple consequence.

Corollary 3.3 (a) *Under the above conditions, the GNS representation induced by $\omega|_{\mathcal{A}(M)}$ is of type III. In particular, $\omega|_{\mathcal{A}(M)}$ is neither a pure state, nor is it normal to the GNS representation of a pure state of $\mathcal{A}(M)$. (b) If, in addition, the von Neumann algebra $\mathcal{N}(M)$ is a factor⁹ and ω' is normal to ω , then $\omega'|_{\mathcal{A}(M)}$ also induces a type III representation.*

Proof. (a) The restriction $\pi|_{\mathcal{A}(M)}$ defines a representation of $\mathcal{A}(M)$ on \mathcal{H} such that $\omega|_{\mathcal{A}(M)}(A) = \langle \Omega | \pi|_{\mathcal{A}(M)}(A)\Omega \rangle$ for all $A \in \mathcal{A}(M)$. Defining \mathcal{H}' to be the closure of $\pi|_{\mathcal{A}(M)}(\mathcal{A}(M))\Omega$, it follows that $(\mathcal{H}', \pi|_{\mathcal{A}(M)}|_{\mathcal{H}'}, \Omega)$ has all the properties of the GNS representation of $\mathcal{A}(M)$ induced by $\omega|_{\mathcal{A}(M)}$ and may therefore be identified with it. Thus, the GNS representation induced by $\omega|_{\mathcal{A}(M)}$ may be identified with a subrepresentation of $\pi|_{\mathcal{A}(M)}$. Applying Prop. 3.2 and [8, III.5.1.7], we have shown that $\omega|_{\mathcal{A}(M)}$ induces a type III representation.

⁸This ‘scaling limit theory’ will typically differ from the original theory [14].

⁹That is, $\mathcal{N}(M)$ intersects its commutant only in multiples of the identity operator.

On the other hand, any pure state ω_0 of $\mathcal{A}(M)$ induces an irreducible representation, which is therefore of type I. Irreducibility implies that ω_0 is *primary*;¹⁰ by [27, 10.3.14], all its normal states induce GNS representations that are quasi-equivalent to that of ω_0 , so they therefore have the same type [8, III.5.1.7]. Hence any (state normal to a) pure state of $\mathcal{A}(M)$ has a type I representation.

(b) As $\mathcal{N}(M)$ is a factor, it follows, first, that $\omega'|_{\mathcal{A}(M)}$ is normal to $\omega|_{\mathcal{A}(M)}$ (see, e.g., Appendix b) of [13]) and, second, that $\omega|_{\mathcal{A}(M)}$ is primary (this is a consequence e.g. of [8, I.9.1.5]). By the argument just used in part (a), $\omega'|_{\mathcal{A}(M)}$ induces a representation of the same type as $\omega|_{\mathcal{A}(M)}$, i.e., type III. \square

Turning this around, if a (state normal to a) pure state of $\mathcal{A}(M)$ is extended to the algebra of any larger region N then the extended state cannot be associated with a regular scaling limit at *any* point of the spacelike boundary of M . Note that although we have described scaling limits in terms of fields, there is an intrinsically operator-algebraic version of these results which would lead to essentially the same conclusion [15].

Let us now apply these results for the case of the linear scalar field with particular regard to S-J states for double cones. Therefore, $\mathcal{A}(O)$ will now denote the C^* -Weyl algebra of the linear scalar field defined for a double cone O in Minkowski spacetime (of arbitrary dimension, not less than 2); the formal definition can be found in Appendix A. By $\omega_{S,J,O}$ we denote the S-J state for the double cone O , i.e. the quasifree state on $\mathcal{A}(O)$ determined by the two-point function $W_{S,J}$ of (1.1), for the case that the spacetime with respect to which $W_{S,J}$ is defined is the double cone O with the induced Minkowski metric.

Proposition 3.4

(a) *Let $M \subset N$ be a strict inclusion of double cones (so that $\overline{M} \subset N$, where both M and N are open double cones), and let $\omega_{S,J,M}$ be the S-J state for the smaller double cone M . Then neither $\omega_{S,J,M}$, nor any state normal to it, can admit an extension to a Hadamard state, or in fact any state normal to a Hadamard state, on $\mathcal{A}(N)$.*

(b) *Supposing again that $M \subset N$ is a strict inclusion of double cones, then the following statements are mutually exclusive.*

(i) $\omega_{S,J,N}$ has a regular scaling limit at some point p in the spacelike boundary of M .

(ii) $\omega_{S,J,M}$ (or a state normal to it) extends to a state ω on $\mathcal{A}(N)$ which is normal to $\omega_{S,J,N}$.

Proof.

(a) Suppose first that ω is a Hadamard state on $\mathcal{A}(N)$. Since Hadamard states possess regular scaling limits at all points in spacetime [34], Corollary 3.3(a) implies that $\omega|_{\mathcal{A}(M)}$ induces a type III representation. Moreover, the von Neumann algebra $\mathcal{N}(M)$ [formed, as above, with respect to the GNS-representation of ω], is a factor [42]. Applying Corollary 3.3(b), if ω' is any state normal to ω , then $\omega'|_{\mathcal{A}(M)}$ induces a type III representation and

¹⁰A state of a C^* -algebra is primary if the von Neumann algebra formed in the corresponding GNS representation is a factor.

cannot coincide with the S-J state $\omega_{S,J,M}$, which is pure, or any state normal to it.

(b) If $\omega_{S,J,N}$ has a regular scaling limit at any point in the spacelike boundary of M , then the von Neumann algebra $\mathcal{N}_{S,J,N}(M) = \pi_{S,J,N}(\mathcal{A}(M))''$ formed in the GNS representation of $\omega_{S,J,N}$ is of type III. On the other hand, since $\omega_{S,J,N}$ is a pure state, and thus primary, any state ω on $\mathcal{A}(N)$ which is normal to $\omega_{S,J,N}$ is quasiequivalent to it by [27, 10.3.14]. This implies that there is an isomorphism $\phi : \pi_{S,J,N}(\mathcal{A}(N))'' \rightarrow \pi_\omega(\mathcal{A}(N))''$ between the von Neumann algebras formed in the GNS representations of the respective states which also satisfies $\phi \circ \pi_{S,J,N} = \pi_\omega$. Consequently, ϕ restricts to a von Neumann algebra isomorphism between $\pi_{S,J,N}(\mathcal{A}(M))''$ and $\pi_\omega(\mathcal{A}(M))''$, and therefore the latter von Neumann algebra is of type III, too. By arguments used in Corollary 3.3(a), it follows that $\omega|_{\mathcal{A}(M)}$ has a GNS representation of type III, and cannot coincide with $\omega_{S,J,M}$ (or any state normal to it) since the latter state is pure. \square

Remarks.

(A) The statements of Prop. 3.4 carry over to curved spacetime, upon making the following modifications: N is a globally hyperbolic spacetime such that the S-J state is defined for the quantized linear scalar field on N (e.g. if N is isometrically embedded into a larger spacetime, see [19]), and M is globally hyperbolic sub-spacetime of N having a sufficiently regular spacelike boundary, and whose closure is properly contained in N . A class of sub-spacetimes M having the required properties is e.g. given by those of the form $M = \text{int}(D(B))$ where $D(B)$ denotes the domain of dependence of B , and B is a coordinate ball of any Cauchy surface Σ of N (with $\Sigma \setminus B$ having a non-void open interior). This is based on the following facts: (i) Proposition 3.2 generalizes to curved spacetimes by [45],[7, Theorem 16.2.18] (also in purely operator-algebraic setting, see [43]), (ii) Hadamard states of the quantized linear scalar field have regular scaling limits also in curved spacetime [34] and (iii) the local von Neumann algebras in their GNS representations are factors for spacetime regions M of the said form [42].

(B) The results clearly indicate that a state which is pure on the algebra of observables of a double cone has a singular behaviour at the double cone's spacelike boundary in the sense that the state cannot be extended to an ambient spacetime so as to have regular scaling limits at the spacelike boundary of the double cone. Statement (b) points at problems if one were to take the point of view that physical states are those which are normal to S-J states (the natural definition of physical states once one is given candidate "vacuum states" [23]). Then the set of physical states depends sensitively on the particular choice of a double cone — similar to the findings we made for S-J states for ultrastatic slab spacetimes in [19] — or there are points in spacetime where S-J states fail to have regular scaling limits. However, as mentioned before, that would commonly be viewed as a pathological short-distance behaviour of a state, particularly for a linear quantized field.

(C) The type III property of the local von Neumann algebras of observables is a typical feature of relativistic quantum field theory which is not encountered in quantum mechanics. A consequence (in combination with the assumption of regular scaling limits) is that the

von Neumann algebras of local observables associated with tangent double cones don't admit product states which are (locally) normal to the vacuum [40]. It also makes the concept of local particle number operators problematic, see e.g. [39] and literature cited there for discussion.

4 Further remarks on S-J States

In this section we make some further observations about the specific case of S-J states. First, we consider whether the failure of S-J states on ultrastatic slab spacetimes, which we have discussed in this paper and in [19], can be removed by ‘softening the boundary’ of the slab by an averaging procedure. Second, we show that the S-J states on such spacetimes (and their averaged variants) fail to be locally Hadamard. We mention, however, that recently Brum and Fredenhagen have considered a somewhat different method of “softening the boundary” of the ultrastatic slab, leading to an altered definition of S-J states which turns out to result in Hadamard states [10]. While this result indicates that it is really the “sharp boundaries” of an ultrastatic slab in time-direction which causes the failing Hadamard property of the associated S-J states, our present results show that Hadamard states cannot be gained by straightforward averaging procedures on the S-J states of ultrastatic slabs.

4.1 Averaging procedures

One could imagine many different prescriptions: here, we study (a) the effect of averaging with respect to a shift in the slab’s absolute time and (b) the effect of averaging with respect to the slab duration parameter τ . As we will see, neither prescription results in a Hadamard state.

We recall from [19] that, on a slab spacetime $M = (-\tau, \tau) \times \Sigma$ as in Section 2, the normal ordered SJ two-point function is

$$:W_{SJ}: (t, \underline{x}; t', \underline{x}') = \sum_{j \in J} \left\{ \frac{\delta_j^2 \cos \omega_j(t - t')}{4\omega_j(1 - \delta_j)} + \frac{\delta_j(2 - \delta_j)}{4\omega_j(1 - \delta_j)} \cos \omega_j(t + t') \right\} \psi_j(\underline{x}) \overline{\psi_j(\underline{x}')}, \quad (4.1)$$

where normal ordering is performed with respect to the standard ultrastatic ground state (which is Hadamard) and the δ_j are given in (2.1). In [19], we showed that $:W_{SJ}:$ cannot be smooth except, perhaps, for a set of τ with measure zero.

For our first averaging prescription, let $\rho \in C_0^\infty(-\epsilon, \epsilon)$ be a nonnegative and even function, with $\epsilon \ll \tau_0$ and $\int d\tau \rho(\tau) = 1$. We consider the quasifree state with two-point function

$$\widehat{W}_{SJ}(t, \underline{x}; t', \underline{x}') = \int d\tau \rho(\tau) W_{SJ, \tau_0}(t + \tau, \underline{x}; t' + \tau, \underline{x}')$$

on the slab $M = (-\tau_0 + \epsilon, \tau_0 - \epsilon) \times \Sigma$. As the ultrastatic ground state is time-translation

invariant, a simple computation gives

$$:\widehat{W}_{SJ}:(t, \underline{x}; t', \underline{x}') = \sum_{j \in J} \left\{ \frac{\delta_j^2 \cos \omega_j(t - t')}{4\omega_j(1 - \delta_j)} + \frac{\delta_j(2 - \delta_j)}{4\omega_j(1 - \delta_j)} \hat{\rho}(2\omega_j) \cos \omega_j(t + t') \right\} \psi_j(\underline{x}) \overline{\psi_j(\underline{x}')}.$$

In order to show that the new state is not globally Hadamard, it is enough to show that this normal ordered two-point function is not smooth. The argument is much as in [19, §4.2]. If $:\widehat{W}_{SJ}:$ were smooth, then the derivative $\partial^{4k}:\widehat{W}_{SJ}:/\partial t^{2k}\partial t'^{2k}$ would have to be the integral kernel of a Hilbert–Schmidt operator on $L^2((-\tau_0 - 2\epsilon, \tau_0 + 2\epsilon) \times \Sigma)$ for all $k \in \mathbb{N}$. Given that $\hat{\rho}(2\omega_j)$ decays rapidly as $j \rightarrow \infty$, it is easily seen that our condition requires that

$$\sum_{j \in J} \frac{\omega_j^{4k} \delta_j^2 \cos \omega_j(t - t')}{4\omega_j(1 - \delta_j)} \psi_j(\underline{x}) \overline{\psi_j(\underline{x}')}$$

is also the integral kernel of such a Hilbert–Schmidt operator. Proceeding as in [19, §4.2] this implies

$$\sum_{j \in J} \frac{\omega_j^{8k-2} \delta_j^4}{(1 - \delta_j)^2} < \infty$$

Given that $\delta_j \rightarrow 1$ we deduce, taking $k = 1$, that $\omega_j \delta_j \rightarrow 0$ as $j \rightarrow \infty$. This can happen for at most a measure zero set of τ_0 (see the proof of [19, Theorem 4.2]).

Turning to the second averaging prescription, let $\rho \in C_0^\infty(\tau_0 - \epsilon, \tau_0 + \epsilon)$ be nonnegative, with $\epsilon \ll \tau_0$ and $\int d\tau \rho(\tau) = 1$. We consider the quasifree state with two-point function

$$\widetilde{W}_{SJ}(t, \underline{x}; t', \underline{x}') = \int d\tau \rho(\tau) W_{SJ,\tau}(t, \underline{x}; t', \underline{x}')$$

which is easily seen to be the two-point function of a state on a slab $M = (-\tau_0 + \epsilon, \tau_0 - \epsilon) \times \Sigma$. Here, $W_{SJ,\tau}$ is the two-point function of the SJ state on the slab $(-\tau, \tau) \times \Sigma$ (cf. [19, §4.5]). The normal-ordered two-point function is

$$:\widetilde{W}_{SJ}:(t, \underline{x}; t', \underline{x}') = \sum_{j \in J} \{A_j \cos \omega_j(t - t') + B_j \cos \omega_j(t + t')\} \psi_j(\underline{x}) \overline{\psi_j(\underline{x}')},$$

where

$$A_j = \int d\tau \rho(\tau) \frac{\delta_j(\tau)^2}{4\omega_j(1 - \delta_j(\tau))}, \quad B_j = \int d\tau \rho(\tau) \frac{\delta_j(\tau)(2 - \delta_j(\tau))}{4\omega_j(1 - \delta_j(\tau))}$$

and we have written the τ -dependence of δ_j explicitly. It is elementary, using the Riemann–Lebesgue lemma, to show that

$$A_j \sim \frac{\text{const}}{\omega_j^3} \sim -B_j$$

Following a similar integral kernel argument to the one just sketched, we see that $:\widetilde{W}_{SJ}:$ cannot be smooth (for any τ_0 or ρ); accordingly, \widetilde{W}_{SJ} is not globally Hadamard.

4.2 Failure to be locally Hadamard

Our results in [19] and in the previous section showing that S-J states and some “smoothed” variants cannot be globally Hadamard on ultrastatic slabs can be strengthened to concluding that they are not even locally Hadamard by applying Radzikowski’s local-to-global theorem for smooth differences of 2-point functions for the KG-field [31], which can be re-stated as follows (where also the propagation-of-singularities theorem [16, Sec. 6.1] is implicitly used):

Proposition 4.1 *Let W be the two-point function of a state defined for a globally hyperbolic spacetime M (assuming W is a bi-distribution). Suppose that W is locally Hadamard in the following sense: there is a Cauchy-surface Σ for M , an open neighbourhood N of Σ and an open covering $\{O_k\}_{k \in \mathcal{K}}$ of N (where \mathcal{K} is a suitable index set) such that $W|_{C_0^\infty(O_k \times O_k)}$ is of Hadamard form for all $k \in \mathcal{K}$. Then W is globally Hadamard.*

This statement has two Corollaries which show that generically the S-J two-point functions $W_{S,J,\tau}$ of the KG-field on ultrastatic slab spacetimes $(-\tau, \tau) \times \Sigma$ cannot be locally Hadamard. The first is an immediate consequence.

Corollary 4.2 *Let $\tau > 0$ be such that $W_{S,J,\tau}$ is not globally Hadamard, let $\tilde{\Sigma}$ be any Cauchy-surface for the ultrastatic slab spacetime $(-\tau, \tau) \times \Sigma$, and let $\{O_{\tilde{k}}\}_{\tilde{k} \in \tilde{\mathcal{K}}}$ be an open cover of (an open neighbourhood of) $\tilde{\Sigma}$. Then there must be some indices \tilde{k} in $\tilde{\mathcal{K}}$ such that $W_{S,J,\tau}|_{C_0^\infty(O_{\tilde{k}} \times O_{\tilde{k}})}$ fails to be Hadamard.*

The statement becomes considerably sharper if the model Cauchy surface Σ of the ultrastatic slab spacetime $(-\tau, \tau) \times \Sigma$ carries a transitive group G of isometries.

Corollary 4.3 *Let $\tau > 0$ be such that $W_{S,J,\tau}$ is not globally Hadamard, and assume that Σ possesses a transitive group G of isometries. Then for any open subset O of $(-\tau, \tau) \times \Sigma$, $W_{S,J,\tau}|_{C_0^\infty(O \times O)}$ fails to be Hadamard.*

Proof: To see this, note that we may without restriction of generality assume that $O = \text{int}(D(\{\tau_0\} \times S))$ for some open subset S of Σ and some $\tau_0 \in (-\tau, \tau)$, as open sets of this type form a base for the topology of an ultrastatic slab. Then we can move S and hence O around with the group G and produce an open cover of $\{\tau_0\} \times \Sigma$ in this way. On the other hand, by construction, $W_{S,J,\tau}$ is invariant under this group action. Thus, assuming that $W_{S,J,\tau}|_{C_0^\infty(O \times O)}$ were Hadamard, we would conclude that $W_{S,J,\tau}$ is locally Hadamard on an open cover of a Cauchy-surface in the sense of Proposition 4.1 and thus globally Hadamard, which again results in a contradiction. \square

The final observation is entirely obvious.

Corollary 4.4 *The statements of Cor. 4.2 and Cor. 4.3 hold likewise for the smoothed-by-averaging versions of the S-J two-point functions on ultrastatic slabs.*

5 Discussion

The Hadamard property for states of quantized linear fields on curved spacetimes is well-motivated by various considerations, e.g. its intimate connection to quantum energy inequalities and stable thermodynamical behaviour. Moreover, it permits a systematic and, more importantly, local and covariant definition of renormalized quantities such as the stress-energy tensor and Wick-ordered and time-ordered product. These properties are instrumental for setting up a local covariant perturbative construction of interacting quantum fields which has hence been carried out, see e.g. [11, 24, 25]. In the present paper, we have shown that, for ultrastatic slabs, the Hadamard property is itself a consequence of imposing finite variance for Wick squares of time-derivatives of field operators — a prerequisite for the quantum mechanical interpretation. Therefore, the failure of S-J states to be Hadamard states cannot, in our view, be taken lightly. In particular, the divergence of the variance of the Wick square of the field’s time-derivative for S-J states on ultrastatic slab spacetimes is a strong argument against considering such states as physical, or even as surrogate vacuum states. We have also seen, as a consequence of more general, model-independent arguments that S-J states defined for double cones must be singular at the spacelike boundary in the sense of not admitting extensions to states on larger spacetime regions having regular scaling limits at the spacelike boundary of the double cones. Furthermore, in the ultrastatic slab situation, the Hadamard property cannot be reached at by simple averaging procedures on S-J states. The altered definition suggested in [10] is different and indicates, in conjunction with our results for double cones, that the problems of S-J states defined for extendible spacetimes stem prominently from a ‘sharp cut-off’ at the ‘edges’ of a spacetime.

The following can, in our view, be concluded from the results. Firstly, the Hadamard property is essentially inevitable for physical states of linear quantized fields on a curved spacetime. Secondly, the definition of S-J states for extendible spacetimes suffers from ultraviolet problems at the ‘spacetime edges’. There is an interesting aspect to it, however. It might be that S-J states (supplemented by a limiting procedure) yield Hadamard states on inextendible, geodesically complete spacetimes, while the Hadamard property fails for S-J states generically on geodesically incomplete spacetimes. If a variant of such a statement could be established, then the deviation of S-J states from the Hadamard property can be taken as a signal for geodesic incompleteness, and at the level of quasi-free representations of the Weyl-algebra, one can associate an algebraic index-like quantity to it. This would be of interest also because failure of geodesic completeness is an indication for singularities in general relativity, and the possibility of associating such a situation with a kind of index of a physically interpretable quantum field theory on the spacetime in question has attractive features from a mathematical point of view, potentially with wider implications.

Finally, the definition of S-J states is a continuum extrapolation of a proposed vacuum-like state for a system of discrete degrees of freedom within the ‘causal sets’ programme [26, 36], a certain approach to quantum gravity. There may be several reasons why this extrapolation does not always lead to Hadamard states. One origin may be that the S-J prescription for continuum quantum fields is simply not an appropriate continuum limit

of the discrete theory. For instance, in the light of our second conclusion, it might be that the limiting procedure ought to exclude geodesically incomplete spacetimes. Another possibility worth contemplating is that the causal set framework might not yet possess sufficient structure to uncover the features relevant for physical states in a continuum limit, like the microlocal spectrum condition (or any other expression of dynamical stability roughly asserting that physical states are mixtures of ‘small perturbations of a ground state or thermal equilibrium state’). Therefore, it might be that our negative results concerning S-J states indicate that there is some structural element missing in the causal sets programme which pertains to information about the dynamics.

The work reported here was partly conducted during the workshop “New Trends in Algebraic Quantum Field Theory”, Frascati, September 2012, and the authors thank the organisers for financial support. We are also grateful to Klaus Fredenhagen for suggesting the idea of computing the fluctuations of Wick polynomials in S-J states. RV thanks the Department of Mathematics of the University of York for hospitality and financial support on occasion of a visit in May 2013 during which major work on the manuscript was done. We also thank Felix Finster, Jorma Louko and Rafael Sorkin for comments.

A The Quantized Linear Scalar Field in the Operator Algebraic Setting

Let us now explain how the linear Klein-Gordon field fits into the framework described in Section 3. Let (M, g) be a globally hyperbolic spacetime, for example (a globally hyperbolic subspacetime of) Minkowski space. Then there are uniquely defined advanced $(-)$ and retarded $(+)$ Green’s operators \mathbf{E}^\pm , defined on $C_0^\infty(M, \mathbb{R})$, for the Klein-Gordon operator $\square + m^2$, where \square is the d’Alembert operator on (M, g) and $m \geq 0$ is a constant, and $\text{supp } \mathbf{E}^\pm f \subset J^\pm(\text{supp } f)$. (See e.g., [6] for details, though our conventions differ.) Their difference $\mathbf{E} = \mathbf{E}^- - \mathbf{E}^+$ is the causal Green’s operator. Factorizing the space $C_0^\infty(M, \mathbb{R})$ by the kernel of \mathbf{E} gives the real vector space $K = C_0^\infty(M, \mathbb{R})/\ker(\mathbf{E})$ of equivalence classes $[f] = f + \ker(\mathbf{E})$ ($f \in C_0^\infty(M, \mathbb{R})$) and it is known that

$$\sigma([f], [h]) = \int_M f(p)(\mathbf{E}h)(p) \, d\text{vol}_M(p)$$

supplies a symplectic form on K (where $d\text{vol}_M$ is the metric-induced volume form of (M, g)). Therefore, (K, σ) is a symplectic space. To any symplectic space there is uniquely associated its *Weyl algebra* $\mathcal{W}(K, \sigma)$, the unique C^* algebra generated by a unit $\mathbf{1}$ and a family of elements $\mathbf{W}([f])$, $[f] \in K$, subject to the relations

$$\mathbf{W}([0]) = \mathbf{1}, \quad \mathbf{W}([f])^* = \mathbf{W}(-[f]), \quad \mathbf{W}([f])\mathbf{W}([h]) = e^{i\sigma([f],[h])/2}\mathbf{W}([f] + [h]),$$

which are the canonical commutation relations in exponentiated form, also called the Weyl relations. One can now set $\mathcal{A}(M) = \mathcal{W}(K, \sigma)$ and define $\mathcal{A}(O)$ as the C^* -subalgebra

generated by all $W([f])$ with $\text{supp}(f) \subset O$. The assignment $(M, g) \rightarrow \mathcal{A}(M)$ is, in fact, functorial [13].

A two-point function is a distribution W in $\mathcal{D}'(M \times M)$ with the property that $W(\bar{f}, f) \geq 0$ ¹¹ for all test-functions $f \in C_0^\infty(M)$ and such that there is some real scalar product μ on K with

$$W(f, h) = \mu([f], [h]) + \frac{i}{2}\sigma([f], [h])$$

for all real-valued test-functions $f, h \in C_0^\infty(M, \mathbb{R})$, and extension by requiring complex linearity in both entries of W . Any two-point function W determines a quasifree state ω on $\mathcal{A}(M) = \mathcal{W}(K, \sigma)$ by setting

$$\omega(W([f])) = e^{-W(f, f)/2} \quad (f \in C_0^\infty(M, \mathbb{R}))$$

and extension by linearity. For such a quasifree state ω , one obtains quantum field operators $\phi(f)$ in the corresponding GNS representation $(\mathcal{H}, \pi, \Omega)$ as the generators of the 1-parameter unitary groups $t \mapsto \pi(W(t[f]))$. That is, to each $f \in C_0^\infty(M, \mathbb{R})$ there is a self-adjoint operator $\phi(f)$ such that

$$\phi(f)\psi = \left. \frac{1}{i} \frac{d}{dt} \pi(W(t[f]))\psi \right|_{t=0}$$

for all ψ in the domain of $\phi(f)$; the subspace \mathcal{D} of \mathcal{H} generated by Ω and all $\phi(f_1) \cdots \phi(f_n)\Omega$ (the ‘‘Wightman domain’’) turns out to be an invariant common domain of essential self-adjointness for all $\phi(f)$ [4]. Furthermore, the $\phi(f)$ are affiliated to the local von Neumann algebras $\mathcal{N}(O)$ in the GNS representation if $\text{supp}(f) \subset O$ since their unitary groups $e^{it\phi(f)} = \pi(W(t[f]))$, $t \in \mathbb{R}$, are clearly contained in $\mathcal{N}(O)$ for real-valued test-functions f supported in O .

It is then easy to check that the quantum field thus obtained in the GNS representation of ω fulfills the following conditions:

- (a) $\phi(f)$ is complex linear in f (by complex linear extension of $\phi(f)$ for real f)
- (b) $\phi(f)^* = \phi(\bar{f})$
- (c) $[\phi(f), \phi(h)] = i\sigma([f], [h])\mathbf{1}$
- (d) $\phi((\square + m^2)f) = 0$

where these relations hold for all $f, h \in C_0^\infty(M)$ as operator identities on \mathcal{D} . Moreover, it holds that $W(f, h) = \langle \Omega | \phi(f)\phi(h)\Omega \rangle$.

A slightly different formulation of the quantized Klein-Gordon field was used in [19] and, implicitly, also in Sections 2 and 4 of the present paper. There, we considered the abstract unital $*$ -algebra $\mathcal{F}(M)$ generated by elements $\phi(f)$ ($f \in C_0^\infty(M)$) subject to relations (a)–(d) above. Evidently, the Hilbert space \mathcal{H} constructed above carries a representation

¹¹We write $W(f, h)$ instead of the technically more correct $W(f \otimes h)$.

of $\mathcal{F}(M)$ in an obvious way, and Ω determines a quasifree state on $\mathcal{F}(M)$ with two-point function W . We also see that, in this sense, the algebra of fields obtained from the Weyl algebra is independent of the quasifree state used in the construction. Moreover, for quasi-free states, one may use the Weyl and field algebras almost interchangeably.

B Decay estimate for the smeared field

We show, subject to the hypotheses of Theorem 2.2, that the function

$$\Upsilon(\mu) = \sum_{j \in J} \frac{1}{2\omega_j} \left| \alpha_j \hat{f}(\mu + \omega_j) + \overline{\beta_j} \hat{f}(\mu - \omega_j) \right|^2$$

decays rapidly as $\mu \rightarrow +\infty$, for all $M \in \mathbb{N}$.

Let $N \in \mathbb{N}$ be chosen so that $C := \sum_{j \in J} (\omega_j^2 + m^2)^{-2N} < \infty$. As $\alpha_j \rightarrow 1$, and f is smooth and compactly supported there is, for each $M \in \mathbb{N}$, a constant C_M so that

$$|\alpha_j \hat{f}(\mu + \omega_j)| \leq \frac{C_M}{((\mu + \omega_j)^2 + m^2)^{2N+M}} \leq \frac{C_M}{(\omega_j^2 + m^2)^{2N} \mu^{2M}}$$

for any $\mu > 0$. Thus we have

$$\sum_{j \in J} \frac{1}{2\omega_j} \left| \alpha_j \hat{f}(\mu + \omega_j) \right|^2 = O(\mu^{-4M}), \quad \sum_{j \in J} \frac{1}{2\omega_j} \left| \overline{\beta_j} \hat{f}(\mu - \omega_j) \right|^2 = O(\mu^{-2M})$$

As $\sup_{j \in J} |\beta_j \hat{f}(\mu - \omega_j)| \leq \sup_{j \in J} |\beta_j| \sup_{\omega} |\hat{f}(\omega)| < \infty$, it follows that

$$\Upsilon(\mu) \leq \sum_{j \in J} \frac{1}{2\omega_j} \left| \overline{\beta_j} \hat{f}(\mu - \omega_j) \right|^2 + O(\mu^{-2M}) \quad (\mu \rightarrow +\infty). \quad (\text{B.1})$$

Now because of the rapid decrease of the β_j and \hat{f} , there are positive constants C'_M and C''_M such that

$$\begin{aligned} (\omega_j^2 + m^2)^N (\omega_j + m)^M |\beta_j| &\leq C'_M & (j \in J) \\ (m^2 + \omega^2)^M |\hat{f}(\omega)| &\leq C''_M & (\omega \in \mathbb{R}). \end{aligned}$$

Hence

$$\begin{aligned} \sum_{j \in J} \frac{1}{2\omega_j} \left| \overline{\beta_j} \hat{f}(\mu - \omega_j) \right|^2 &\leq \frac{(C'_M C''_M)^2}{2m} \sum_{j \in J} \frac{1}{(\omega_j^2 + m^2)^{2N}} \frac{1}{(\omega_j + m)^{2M} ((\omega_j - \mu)^2 + m^2)^{2M}} \\ &\leq \frac{(C'_M C''_M)^2 C}{2m} \sup_{j \in J} \frac{1}{(\omega_j + m)^{2M} ((\omega_j - \mu)^2 + m^2)^{2M}} \\ &\leq \frac{2^{2M-1} (C'_M C''_M)^2 C}{\mu^{2M} m^{4M+1}} \end{aligned} \quad (\text{B.2})$$

for all $\mu > m$. In the last step, we used the following elementary observations. Let $K_{x_0}(x) = (1+x)(1+(x-x_0)^2)$, for $x_0 > 1$. Then, on $[0, \infty)$, K_{x_0} is easily seen to have a local maximum at $x = x_-$ and global minimum at x_+ , where

$$x_{\pm} = \frac{1}{3} \left(2x_0 - 1 \pm \sqrt{(x_0 + 1)^2 - 3} \right).$$

Then $K_{x_0}(x_+) = (1+x_+)(1+(x-x_+)^2) \geq 1+x_+ > 2x_0/3$, so, in particular, $\inf_{\mathbb{R}^+} K_{x_0} > x_0/2$. Moreover, as K_{x_0} is strictly positive on \mathbb{R}^+ , the same classification of critical points applies to positive integer powers of K_{x_0} . In particular, the minimum value of $K_{x_0}(x)^{2M}$ is $K_{x_0}(x_+)^{2M} > (x_0/2)^{2M}$. This estimate is precisely what is needed to obtain (B.2). In conjunction with (B.1), we have shown that $\Upsilon(\mu) = O(\mu^{-2M})$ as $\mu \rightarrow +\infty$, for all $M \in \mathbb{N}$, as desired.

References

- [1] Afshordi, N., Aslanbeigi, S., Sorkin, R. D.: A distinguished vacuum state for a quantum field in a curved spacetime: Formalism, features, and cosmology. *JHEP* **08**, 137 (2012)
- [2] Afshordi, N., Buck, M., Dowker, F., Rideout, D., Sorkin, R. D., Yazaman, K. Y.: A ground state for the causal diamond in 2 dimensions. *JHEP* **10**, 088 (2012)
- [3] Almheiri, A., Marolf, D., Polchinski, J., Sully, J.: Black holes: complementarity or firewalls? *JHEP* **02**, 062 (2013)
- [4] Araki, H., Yamagami, S.: On quasi-equivalence of quasifree states of the canonical commutation relations. *Publ. Res. Inst. Math. Sci.* **18**, 703–758 (283–338) (1982)
- [5] Aslanbeigi, S., Buck, M.: A preferred ground state for the scalar field in de Sitter space, E-print arXiv:1306.3231 (2013)
- [6] Bär, C., Ginoux, N., Pfäffle, F.: Wave equations on Lorentzian manifolds and quantization. European Mathematical Society (EMS), Zürich (2007)
- [7] Baumgärtel, H., Wollenberg, M.: Causal nets of operator algebras. Akademie-Verlag, Berlin (1992)
- [8] Blackadar, B.: Operator algebras, *Encyclopaedia of Mathematical Sciences*, vol. 122. Springer-Verlag, Berlin (2006). Theory of C^* -algebras and von Neumann algebras, Operator Algebras and Non-commutative Geometry, III
- [9] Braunstein, S.L.: Black hole entropy as entropy of entanglement, or it’s curtains for the equivalence principle (2009). ArXiv:0907.1190v1. Published as: Braunstein, S. L., Pirandola, S. and Życzkowski, K., *Physical Review Letters* **110**, 101301 (2013); see arXiv:0907.1190v3
- [10] Brum, M. and Fredenhagen, K.: Modified Sorkin-Johnston states on static and expanding spacetimes. (2013). ArXiv:1307.0482 [gr-qc]
- [11] Brunetti, R., Fredenhagen, K.: Microlocal analysis and interacting quantum field theories: Renormalization on physical backgrounds. *Commun. Math. Phys.* **208**, 623–661 (2000)
- [12] Brunetti, R., Fredenhagen, K., Hollands, S.: A remark on alpha vacua for quantum field theories on de Sitter space. *JHEP* **0505**, 063 (2005)
- [13] Brunetti, R., Fredenhagen, K., Verch, R.: The generally covariant locality principle – a new paradigm for local quantum physics. *Commun. Math. Phys.* **237**, 31–68 (2003)
- [14] Buchholz, D.: Quarks, gluons, colour: facts or fiction? *Nuclear Phys. B* **469**, 333–353 (1996)

- [15] Buchholz, D., Verch, R.: Scaling algebras and renormalization group in algebraic quantum field theory. *Rev. Math. Phys.* **7**, 1195–1239 (1995)
- [16] Duistermaat, J., Hörmander, L.: Fourier integral operators II. *Acta Mathematica* **128**, 183–269 (1972)
- [17] Fewster, C.J.: A general worldline quantum inequality. *Class. Quant. Grav.* **17**, 1897–1911 (2000)
- [18] Fewster, C.J., Verch, R.: A quantum weak energy inequality for Dirac fields in curved spacetime. *Commun. Math. Phys.* **225**, 331–359 (2002)
- [19] Fewster, C.J., Verch, R.: On a recent construction of ‘vacuum-like’ quantum field states in curved spacetime. *Classical Quantum Gravity* **29**, 205017 (2012)
- [20] Finster, F.: Definition of the Dirac sea in the presence of external fields. *Adv. Theor. Math. Phys.* **2**, 963–985 (1998)
- [21] Finster, F., Reintjes, M.: A non-perturbative construction of the Fermionic projector on globally hyperbolic manifolds I - Space-times of finite lifetime (2013). [ArXiv:1301.5420](https://arxiv.org/abs/1301.5420)
- [22] Fredenhagen, K.: On the modular structure of local algebras of observables. *Comm. Math. Phys.* **97**, 79–89 (1985)
- [23] Haag, R.: *Local Quantum Physics: Fields, Particles, Algebras*. Springer-Verlag, Berlin (1992)
- [24] Hollands, S., Wald, R.M.: Local Wick polynomials and time ordered products of quantum fields in curved spacetime. *Commun. Math. Phys.* **223**, 289–326 (2001)
- [25] Hollands, S., Wald, R.M.: Existence of local covariant time ordered products of quantum fields in curved spacetime. *Commun. Math. Phys.* **231**, 309–345 (2002)
- [26] Johnston, S.: Feynman propagator for a free scalar field on a causal set. *Physical Review Letters* **103**, 180401 (2009), [arXiv:0909.0944](https://arxiv.org/abs/0909.0944)
- [27] Kadison, R.V., Ringrose, J.R.: *Fundamentals of the theory of operator algebras. Vol. II*, vol. 100. Academic Press Inc., Orlando, FL (1986)
- [28] Kay, B.S., Radzikowski, M.J., Wald, R.M.: Quantum field theory on spacetimes with a compactly generated Cauchy horizon. *Commun. Math. Phys.* **183**, 533–556 (1997)
- [29] Kay, B. S., Wald, R. M.: Theorems on the uniqueness and thermal properties of stationary, nonsingular, quasifree states on space-times with a bifurcate Killing horizon. *Phys. Rept.* **207**, 49–136 (1991)
- [30] Pinamonti, N.: On the initial conditions and solutions of the semiclassical Einstein equations in a cosmological scenario. *Commun. Math. Phys.* **305**, 563–604 (2011)
- [31] Radzikowski, M.: A local to global singularity theorem for quantum field theory on curved space-time. *Commun.Math.Phys.* **180**, 1–22 (1996)
- [32] Radzikowski, M. J.: Micro-local approach to the Hadamard condition in quantum field theory on curved space-time. *Commun. Math. Phys.* **179**, 529–553 (1996)
- [33] Sahlmann, H., Verch, R.: Passivity and microlocal spectrum condition. *Commun. Math. Phys.* **214**, 705–731 (2000)
- [34] Sahlmann, H., Verch, R.: Microlocal spectrum condition and Hadamard form for vector valued quantum fields in curved space-time. *Rev.Math.Phys.* **13**, 1203–1246 (2001)
- [35] Sanders, K.: Equivalence of the (generalised) Hadamard and microlocal spectrum condition for (generalised) free fields in curved spacetime. *Comm. Math. Phys.* **295**, 485–501 (2010)
- [36] Sorkin, R.D.: Scalar field theory on a causal set in histories form. *Journal of Physics Conference Series* **306**, 012017 (2011)

- [37] Streater, R.F., Wightman, A.S.: PCT, spin and statistics, and all that. Princeton Landmarks in Physics. Princeton University Press, Princeton, NJ (2000)
- [38] Strohmaier, A., Verch, R., Wollenberg, M.: Microlocal analysis of quantum fields on curved spacetimes: Analytic wavefront sets and Reeh-Schlieder theorems. *J. Math. Phys.* **43**, 5514–5530 (2002)
- [39] Summers, S.J.: Subsystems and independence in relativistic microscopic physics. *Stud. Hist. Phil. Mod. Physics* **40**, 133–141 (2009)
- [40] Summers, S.J. , Werner, R.: Maximal violation of Bell’s inequalities for algebras of observables in tangent spacetime regions. *Ann. Inst. H. Poincaré* **49**, 215–243 (1988)
- [41] Taylor, M.E.: Partial differential equations. II, *Applied Mathematical Sciences*, vol. 116. Springer-Verlag, New York (1996). Qualitative studies of linear equations
- [42] Verch, R.: Continuity of symplectically adjoint maps and the algebraic structure of Hadamard vacuum representations for quantum fields on curved space-time. *Rev.Math.Phys.* **9**, 635–674 (1997)
- [43] Verch, R.: Scaling algebras, the renormalization group and the principle of local stability in algebraic quantum field theory. In: Doplicher, S., Guido, D., Longo, R., Zsido, L. (ed.) Proceedings of the Conference on Operator Algebras and Quantum Field Theory, Rome, Italy, 1-6 July 1996, International Press (1996)
- [44] Wald, R.M.: Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics. University of Chicago Press, Chicago (1994)
- [45] Wollenberg, M.: Scaling limits and type of local algebras over curved space-time. In: W.B. Arveson, A.S. Mishchenko, M. Putinar, M.A. Rieffel, Ş. Strătilă (eds.) Operator algebras and topology. Proceedings of the OATE 2 Conference on Operator Algebras and their Connections with Topology and Ergodic Theory held in Craiova, Summer 1989, *Pitman Res. Notes Math. Ser.*, vol. 270, pp. 179–196. Longman Sci. Tech., Harlow (1992)