

This is a repository copy of *American options with gradual exercise under proportional transaction costs*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/89270/>

Version: Accepted Version

Article:

Roux, Alet orcid.org/0000-0001-9176-4468 and Zastawniak, Tomasz (2014) American options with gradual exercise under proportional transaction costs. *International Journal of Theoretical and Applied Finance*. 1450052. ISSN 0219-0249

<https://doi.org/10.1142/S0219024914500526>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

International Journal of Theoretical and Applied Finance
© World Scientific Publishing Company

AMERICAN OPTIONS WITH GRADUAL EXERCISE UNDER PROPORTIONAL TRANSACTION COSTS*

ALET ROUX

*Department of Mathematics, University of York
Heslington, York YO10 5DD, UK
alet.roux@york.ac.uk*

TOMASZ ZASTAWNIAK

*Department of Mathematics, University of York
Heslington, York YO10 5DD, UK
tomasz.zastawniak@york.ac.uk*

American options in a multi-asset market model with proportional transaction costs are studied in the case when the holder of an option is able to exercise it gradually at a so-called mixed (randomised) stopping time. The introduction of gradual exercise leads to tighter bounds on the option price when compared to the case studied in the existing literature, where the standard assumption is that the option can only be exercised instantly at an ordinary stopping time. Algorithmic constructions for the bid and ask prices and the associated superhedging strategies and optimal mixed stopping times for an American option with gradual exercise are developed and implemented, and dual representations are established.

Keywords: American options; transaction costs; mixed stopping times; superhedging; dual representation.

1. Introduction

This work on pricing American options under proportional transaction costs goes back to the seminal discovery by Chalasani & Jha (2001) that to hedge against a buyer who can exercise the option at any (ordinary) stopping time, the seller must in effect be protected against all mixed (randomised) stopping times. This was followed by Bouchard & Temam (2005), who established a dual representation for the set of strategies superhedging the seller's (though not the buyer's) position in an American option under transaction costs in discrete time. Moreover, dual representation of seller's superhedging strategies in continuous time was tackled by Bouchard & Chassagneux (2009) and De Vallière & Denis & Kabanov (2009). Iterative algorithms for computing the upper and lower hedging prices of the option,

*Electronic version of an article published in *International Journal of Theoretical and Applied Finance* **17** (8), 2014, 1450052 DOI 10.1142/S0219024914500526 ©World Scientific Publishing Company <http://www.worldscientific.com/worldscinet/ijtaf>

the hedging strategies, optimal stopping times as well as dual representations (with the first constructive proofs) for both the seller and the buyer of an American option under transaction costs were developed by Roux & Zastawniak (2009) in a discrete model with two assets, and Roux & Zastawniak (2011) in a multi-asset model. All these approaches take it for granted that the buyer can only exercise the option instantly, at an ordinary stopping time of his choosing.

By contrast, in the present paper we allow the buyer the flexibility to exercise an American option gradually, rather than all at a single time instance. Though it would be difficult in practice to exercise a fraction of an option contract and to hold on to the remainder to exercise it later, the holder of a large portfolio of options may well choose to exercise the individual contracts on different dates if that proves beneficial. Does this ability to exercise gradually affect the pricing bounds, hedging strategies and optimal stopping times for the buyer and/or seller? Perhaps surprisingly, the answer to this question is yes, it does in the presence of transaction costs.

Gradual exercise turns out to be linked to another feature, referred to as deferred solvency, which will also be studied here. If a temporary loss of liquidity occurs in the market, as reflected by unusually large bid-ask spreads, agents may become insolvent. Being allowed to defer closing their positions until liquidity is restored might enable them to become solvent once again. This gives more leeway when constructing hedging strategies than the usual requirement that agents should remain solvent at all times.

Tien (2012) was the first to explore the consequences of gradual exercise and deferred solvency using a model with a single risky asset as a testing ground. In the present paper these ideas are developed in a systematic manner and extended to the much more general setting of the multi-asset market model with transaction costs due to Kabanov (1999); see also Kabanov & Stricker (2001) and Schachermayer (2004).

Pricing and hedging for the seller of an American option under transaction costs is a convex optimisation problem irrespective of whether instant or gradual exercise is permitted. However, this is not so for the buyer. In this case one has to tackle a non-convex optimisation problem for options that can only be exercised instantly. A very interesting consequence of gradual exercise is that pricing and hedging becomes a convex optimisation problem also for the buyer of an American option, making it possible to deploy convex duality methods. The convexity of the problem also makes it much easier to implement the pricing and hedging algorithms numerically. We will make use of this new opportunity in this paper.

The paper is organised as follows. Section 2 recalls the general setting of Kabanov's multi-asset model with transaction costs. In Section 3 the hedging strategies for the buyer and seller and the corresponding option prices under gradual exercise are introduced and compared with the same notions under instant exercise. A toy example is set up to demonstrate that it is easier to hedge an option and that

the bid-ask spread of the option prices can be narrower under gradual exercise as compared to instant exercise. In Section 4 the seller's case is studied in detail. The notion of deferred solvency is first discussed and linked in Proposition 4.3 with the hedging problem for the seller of an American option with gradual exercise. The sets of seller's hedging portfolios are then constructed and related to the ask price of the option under gradual exercise and to a construction of a seller's hedging strategy realising the ask price; see Theorem 4.1. A dual representation of the seller's price is established in Theorem 4.2. The toy example is revisited to illustrate the various constructions and results for the seller. Section 5 is devoted to the buyer's case. Buyer's hedging portfolios and strategies are constructed and used to compute the bid price of the option; see Theorem 5.1. Finally, the dual representation for the buyer is explored in Theorem 5.2. Once again, the toy example serves to illustrate the results. A numerical example with three assets can be found in Section 6. Some conclusions and possible further developments and ramifications are touched upon in Section 7. Technical information and proofs are collected in the Appendix.

2. Multi-currency model with proportional transaction costs

Let $(\Omega, \mathcal{F}, \mathbb{P}; (\mathcal{F}_t)_{t=0}^T)$ be a filtered probability space. We assume that Ω is finite, $\mathcal{F}_0 = \{\emptyset, \Omega\}$, $\mathcal{F}_T = \mathcal{F} = 2^\Omega$ and $\mathbb{P}(\omega) > 0$ for all $\omega \in \Omega$. For each $t = 0, \dots, T$ let Ω_t be the collection of atoms of \mathcal{F}_t , called the *nodes* of the associated tree model. A node $\nu \in \Omega_{t+1}$ is said to be a *successor* of a node $\mu \in \Omega_t$ if $\nu \subseteq \mu$. For each $t = 0, \dots, T-1$ we denote the collection of successors of any given node $\mu \in \Omega_t$ by $\text{succ } \mu$.

For each $t = 0, \dots, T$ let $\mathcal{L}_t := \mathcal{L}^0(\mathbb{R}^d; \mathcal{F}_t)$ be the collection of \mathcal{F}_t -measurable \mathbb{R}^d -valued random variables. We identify elements of \mathcal{L}_t with functions on Ω_t whenever convenient.

We consider the discrete-time currency model introduced by Kabanov (1999) and developed further by Kabanov & Stricker (2001) and Schachermayer (2004) among others. The model contains d assets or currencies. At each trading date $t = 0, 1, \dots, T$ and for each $k, j = 1, \dots, d$ one unit of asset k can be obtained by exchanging $\pi_t^{jk} > 0$ units of asset j . We assume that the exchange rates π_t^{jk} are \mathcal{F}_t -measurable and $\pi_t^{jj} = 1$ for all t and j, k .

We say that a portfolio $x \in \mathcal{L}_t$ is can be *exchanged* into a portfolio $y \in \mathcal{L}_t$ at time t whenever there are \mathcal{F}_t -measurable random variables $\beta^{jk} \geq 0$, $j, k = 1, \dots, d$ such that for all $k = 1, \dots, d$

$$y^k = x^k + \sum_{j=1}^d \beta^{jk} - \sum_{j=1}^d \beta^{kj} \pi_t^{kj}, \quad (2.1)$$

where β^{jk} represents the number of units of asset k received as a result of exchanging some units of asset j .

The *solvency cone* $\mathcal{K}_t \subseteq \mathcal{L}_t$ is the set of portfolios that are *solvent* at time t , i.e. the portfolios at time t that can be exchanged into portfolios with non-negative

holdings in all d assets. It is straightforward to show that \mathcal{K}_t is the convex cone generated by the canonical basis e^1, \dots, e^d of \mathbb{R}^d and the vectors $\pi_t^{jk} e^j - e^k$ for $j, k = 1, \dots, d$, and so \mathcal{K}_t is a polyhedral cone, hence closed. Note that \mathcal{K}_t contains all the non-negative elements of \mathcal{L}_t .

A trading strategy $y = (y_t)_{t=0}^{T+1}$ is a predictable \mathbb{R}^d -valued process with final value $y_{T+1} = 0$ and initial endowment $y_0 \in \mathbb{R}^d$. For each $t > 0$ the portfolio $y_t \in \mathcal{L}_{t-1}$ is held from time $t-1$ to time t . Let Φ be the set of trading strategies. We say that $y \in \Phi$ is a *self-financing* strategy whenever $y_t - y_{t+1} \in \mathcal{K}_t$ for all $t = 0, \dots, T-1$. Note that no implicitly assumed self-financing condition is included in the definition of Φ .

A trading strategy $y \in \Phi$ is an *arbitrage opportunity* if it is self-financing, $y_0 = 0$ and there is a portfolio $x \in \mathcal{L}_T \setminus \{0\}$ with non-negative holdings in all d assets such that $y_T - x \in \mathcal{K}_T$. This notion of arbitrage was considered by Schachermayer (2004), and its absence is formally different but equivalent to the weak no-arbitrage condition introduced by Kabanov & Stricker (2001).

Theorem 2.1 (Kabanov & Stricker (2001), Schachermayer (2004)). *The model admits no arbitrage opportunity if and only if there exists a probability measure \mathbb{Q} equivalent to \mathbb{P} and an \mathbb{R}^d -valued \mathbb{Q} -martingale $S = (S_t)_{t=0}^T$ such that*

$$S_t \in \mathcal{K}_t^* \setminus \{0\} \text{ for all } t = 0, \dots, T, \quad (2.2)$$

where \mathcal{K}_t^* is the polar of $-\mathcal{K}_t$; see (A.1) in the Appendix.

We denote by \mathcal{P} the set of pairs (\mathbb{Q}, S) satisfying the conditions in Theorem 2.1, and by $\bar{\mathcal{P}}$ the set of pairs (\mathbb{Q}, S) satisfying the conditions in Theorem 2.1 but with \mathbb{Q} absolutely continuous with respect to (and not necessarily equivalent to) \mathbb{P} . We assume for the remainder of this paper that the model admits no arbitrage opportunities, i.e. $\mathcal{P} \neq \emptyset$.

Remark 2.1. In place of a pair $(\mathbb{Q}, S) \in \mathcal{P}$ one can equivalently use the so-called *consistent price process* $S_t \mathbb{E}_{\mathbb{P}}(\frac{d\mathbb{Q}}{d\mathbb{P}} | \mathcal{F}_t)$; see Schachermayer (2004).

We also define for any $j = 1, \dots, d$

$$\mathcal{P}_j := \{(\mathbb{Q}, S) \in \mathcal{P} \mid S^j = 1\}, \quad \bar{\mathcal{P}}_j := \{(\mathbb{Q}, S) \in \bar{\mathcal{P}} \mid S^j = 1\}.$$

In the absence of arbitrage \mathcal{K}_t^* is a non-empty compactly j -generated polyhedral cone for all t (Roux & Zastawniak 2011 Remark 2.2), which means that $\mathcal{P}_j \neq \emptyset$. (For the definition of a compactly j -generated cone, see Appendix A.2.)

3. Instant versus gradual exercise

The payoff of an American option in the model with d underlying currencies is, in general, an \mathbb{R}^d -valued adapted process $\xi = (\xi_t)_{t=0}^T$. The seller of the American option is obliged to deliver, and the buyer is entitled to receive the portfolio of currencies ξ_τ at a stopping time $\tau \in \mathcal{T}$ chosen by the buyer. Here \mathcal{T} denotes the

family of stopping times with values in $\{0, \dots, T\}$. This is the usual setup in which the option is exercised *instantly* at a stopping time $\tau \in \mathcal{T}$.

American options with the provision for instant exercise in the multi-currency model under proportional transaction costs have been studied by Bouchard & Temam (2005), who established a characterisation of the superhedging strategies for the option seller only, and by Roux & Zastawniak (2011), who provided iterative constructions of the ask and bid option prices and the superhedging strategies for both the option seller and buyer, as well as the first constructive proofs of the corresponding dual representations.

In the present paper we relax the requirement that the option needs to be exercised instantly at a stopping time $\tau \in \mathcal{T}$. Instead, we allow the buyer to exercise *gradually* at a mixed stopping time $\chi \in \mathcal{X}$. (For the definition of mixed stopping times, see Appendix A.3.)

If the buyer chooses to exercise the option gradually according to a mixed stopping time $\chi \in \mathcal{X}$, then the seller of the American option will be obliged to deliver, and the buyer will be entitled to receive the fraction χ_t of the portfolio of currencies ξ_t at each time $t = 0, \dots, T$.

The question then arises whether or not it would be more beneficial for the buyer to exercise the option gradually rather than instantly? What will be the optimal mixed stopping time $\chi \in \mathcal{X}$ for the buyer? How should the seller hedge against gradual exercise? Are the ask (seller's) and bid (buyer's) option prices and hedging strategies affected by gradual exercise as compared to instant exercise?

3.1. Instant exercise

In the case of instant exercise the seller of an American option ξ needs to hedge by means of a trading strategy $y \in \Phi$ against all ordinary stopping times $\tau \in \mathcal{T}$ chosen by the buyer. The trading strategy y needs to be self-financing up to time τ and to allow the seller to remain solvent on delivering the portfolio ξ_τ at time τ , for any $\tau \in \mathcal{T}$. Hence the family of seller's superhedging strategies is defined as

$$\Phi^a(\xi) := \{y \in \Phi \mid \forall \tau \in \mathcal{T} : y_t - y_{t+1} \in \mathcal{K}_t \text{ for } t = 0, \dots, \tau - 1, y_\tau - \xi_\tau \in \mathcal{K}_\tau\},$$

and the *ask price (seller's price)* of the option in currency $j = 1, \dots, d$ is

$$\pi_j^a(\xi) := \inf\{x \in \mathbb{R} \mid \exists y \in \Phi^a(\xi) : xe^j = y_0\}.$$

This is the smallest amount in currency j needed to superhedge a short position in ξ .

On the other hand, the buyer of an American option ξ can select both a stopping time $\tau \in \mathcal{T}$ and a trading strategy $y \in \Phi$. The trading strategy y needs to be self-financing up to time τ and to allow the buyer to remain solvent on receiving the portfolio ξ_τ at time τ . Thus, the family of buyer's superhedging strategies is defined as

$$\Phi^b(\xi) := \{(y, \tau) \in \Phi \times \mathcal{T} \mid y_t - y_{t+1} \in \mathcal{K}_t \text{ for } t = 1, \dots, \tau - 1, y_\tau + \xi_\tau \in \mathcal{K}_\tau\},$$

6 Alet Roux and Tomasz Zastawniak

and the *bid price* (*buyer's price*) of the option in currency $j = 1, \dots, d$ is

$$\pi_j^b(\xi) := \sup\{-x \in \mathbb{R} \mid \exists(y, \tau) \in \Phi^b(\xi) : xe^j = y_0\}.$$

This is the largest amount in currency j that the buyer can raise using the option ξ as surety.

For American options with instant exercise, iterative constructions of the ask and bid option prices $\pi_j^a(\xi)$ and $\pi_j^b(\xi)$ and the corresponding seller's and buyer's superhedging strategies from $\Phi^a(\xi)$ and $\Phi^b(\xi)$ were established by Roux & Zastawniak (2011).

3.2. Gradual exercise

When the buyer is allowed to exercise gradually, the seller needs to follow a suitable trading strategy to hedge his exposure. Since the seller can react to the buyer's actions, this strategy may in general depend on the mixed stopping time $\chi \in \mathcal{X}$ followed by the buyer, and will be denoted by $Y^\chi \in \Phi$. In other words, we consider a function $Y : \mathcal{X} \rightarrow \Phi$.

At each time t the seller will be holding a portfolio Y_t^χ and will be obliged to deliver a fraction χ_t of the payoff ξ_t . He can then rebalance the remaining portfolio $Y_t^\chi - \chi_t \xi_t$ into Y_{t+1}^χ in a self-financing manner, so that

$$Y_t^\chi - \chi_t \xi_t - Y_{t+1}^\chi \in \mathcal{K}_t \text{ for each } t = 0, \dots, T. \quad (3.1)$$

The self-financing and superhedging conditions have merged into one. We call (3.1) the *rebalancing* condition.

When creating the portfolio Y_t^χ at time $t-1$, the seller can only use information available at that time. This includes $\chi_0, \dots, \chi_{t-1}$, but the seller has no way of knowing the future values χ_t, \dots, χ_T that will be chosen by the buyer. The trading strategies $Y^\chi \in \Phi$ that can be adopted by the seller are therefore restricted to those satisfying the *non-anticipation* condition

$$\begin{aligned} & \forall \chi, \chi' \in \mathcal{X} \forall t = 0, \dots, T \forall \omega \in \Omega : \\ & \chi_s(\omega) = \chi'_s(\omega) \text{ for each } s = 0, \dots, t-1 \implies Y_t^\chi(\omega) = Y_t^{\chi'}(\omega). \end{aligned} \quad (3.2)$$

In particular, the initial endowment Y_0^χ of the trading strategy Y^χ is the same for all $\chi \in \mathcal{X}$. We denote this common value by Y_0 .

We define the family of seller's superhedging strategies against gradual exercise by

$$\Phi^{\text{ag}}(\xi) := \{Y : \mathcal{X} \rightarrow \Phi \mid Y \text{ satisfies (3.1) and (3.2)}\}$$

and the corresponding *ask price* (*seller's price*) of the option in currency $j = 1, \dots, d$ by

$$\pi_j^{\text{ag}}(\xi) := \inf\{x \in \mathbb{R} \mid \exists Y \in \Phi^{\text{ag}}(\xi) : xe^j = Y_0\}. \quad (3.3)$$

This is the smallest amount in currency j that the seller needs to superhedge a short position in the American option ξ when the buyer is allowed to exercise gradually.

On the other hand, the buyer is able to select both a mixed stopping time $\chi \in \mathcal{X}$ and a trading strategy $y \in \Phi$, and will be taking delivery of a fraction χ_t of the payoff ξ_t at each time t . Because the choice of the mixed stopping time χ is up to the buyer, the trading strategy y needs to be good just for the one chosen stopping time, and does not need to be considered as a function of χ , in contrast to the seller's case. The *rebalancing* condition

$$y_t + \chi_t \xi_t - y_{t+1} \in \mathcal{K}_t \text{ for each } t = 0, \dots, T \quad (3.4)$$

needs to be satisfied.

Hence, the family of superhedging strategies for the buyer of an American option ξ with gradual exercise is defined as

$$\Phi^{\text{bg}}(\xi) := \{(y, \chi) \in \Phi \times \mathcal{X} \mid (y, \chi) \text{ satisfies (3.4)}\},$$

and the corresponding *bid price* (*buyer's price*) of the option in currency $j = 1, \dots, d$ is

$$\pi_j^{\text{bg}}(\xi) := \sup\{-x \in \mathbb{R} \mid \exists (y, \chi) \in \Phi^{\text{bg}}(\xi) : x e^j = y_0\}. \quad (3.5)$$

This is the largest amount in currency j that can be raised using the option as surety by a buyer who is able to exercise gradually.

Example 3.1. We consider a toy example with two assets, a foreign currency (asset 1) and domestic currency (asset 2) in a two-step binomial tree model with the following bid/ask foreign currency prices $S_t^b \leq S_t^a$ in each of the four scenarios in $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4\}$:

	S_0^b	S_0^a	S_1^b	S_1^a	S_2^b	S_2^a
ω_1	5	5	3	9	4	8
ω_2	5	5	3	9	4	4
ω_3	5	5	2	2	3	3
ω_4	5	5	2	2	1	1

Note there are only two nodes with a non-trivial bid/ask spread, namely the 'up' node $U = \{\omega_1, \omega_2\}$ and the 'up-up' node $UU = \{\omega_1\}$. The corresponding exchange rates are

$$\begin{bmatrix} \pi_t^{11} & \pi_t^{12} \\ \pi_t^{21} & \pi_t^{22} \end{bmatrix} = \begin{bmatrix} 1 & 1/S_t^b \\ S_t^a & 1 \end{bmatrix}.$$

In this model we consider an American option with the following payoff process $\xi_t = (\xi_t^1, \xi_t^2)$:

	ξ_0	ξ_1	ξ_2
ω_1	(0, 0)	(0, 4)	(2, -8)
ω_2	(0, 0)	(0, 4)	(0, 0)
ω_3	(0, 0)	(0, 0)	(0, 0)
ω_4	(0, 0)	(0, 0)	(0, 0)

8 *Alet Roux and Tomasz Zastawniak*

In the case when the option can only be exercised instantly, using the algorithms of Roux & Zastawniak (2011) we can compute the bid and ask prices of the option in the domestic currency to be

$$\pi_2^b(\xi) = 2, \quad \pi_2^a(\xi) = \frac{28}{5}.$$

Now consider $Y : \mathcal{X} \rightarrow \Phi$ given by

	Y_0^χ	Y_1^χ	Y_2^χ
ω_1	(0, 5)	(1, 0)	$(1, -4\chi_1^{\omega_1})$
ω_2	(0, 5)	(1, 0)	$(1, -4\chi_1^{\omega_2})$
ω_3	(0, 5)	(1, 0)	(0, 0)
ω_4	(0, 5)	(1, 0)	(0, 0)

for any $\chi \in \mathcal{X}$. Also consider $y \in \Phi$ and $\chi \in \mathcal{X}$ such that

	y_0	y_1	y_2	χ_0	χ_1	χ_2
ω_1	(0, -3)	(-1, 2)	(-1, 4)	0	$\frac{1}{2}$	$\frac{1}{2}$
ω_2	(0, -3)	(-1, 2)	(-1, 4)	0	$\frac{1}{2}$	$\frac{1}{2}$
ω_3	(0, -3)	(-1, 2)	(0, 0)	0	0	1
ω_4	(0, -3)	(-1, 2)	(0, 0)	0	0	1

We can verify that $Y \in \Phi^{\text{ag}}(\xi)$ and $(y, \chi) \in \Phi^{\text{bg}}(\xi)$. The existence of these strategies means that

$$\pi_2^b(\xi) = 2 < 3 \leq \pi_2^{\text{bg}}(\xi), \quad \pi_2^{\text{ag}}(\xi) \leq 5 < \frac{28}{5} = \pi_2^a(\xi).$$

This example demonstrates that the seller's and buyer's prices $\pi_j^{\text{ag}}(\xi), \pi_j^{\text{bg}}(\xi)$ under gradual exercise may differ from their respective counterparts $\pi_j^a(\xi), \pi_j^b(\xi)$ under instant exercise. It demonstrates the need to revisit and investigate the pricing and superhedging results in the case when the instant exercise provision is relaxed and replaced by gradual exercise.

4. Pricing and superhedging for the seller under gradual exercise

We have seen in Example 3.1 that the seller's price $\pi_j^a(\xi)$ may be higher than $\pi_j^{\text{ag}}(\xi)$. The reason is that an option seller who follows a hedging strategy $y \in \Phi^a(\xi)$ is required to be instantly solvent upon delivering the payoff at the stopping time $\tau \in \mathcal{T}$ when the buyer has chosen to exercise the option. Meanwhile, a seller who follows a strategy $Y \in \Phi^{\text{ag}}(\xi)$ will be able to continue rebalancing the strategy up to the time horizon T as long as a solvent position can be reached eventually. Being able to defer solvency in this fashion allows more flexibility for the seller, resulting in a lower seller's price.

On the other hand, it might appear that a seller who hedges against gradual exercise (against mixed stopping times) would have a harder task to accomplish than someone who only needs to hedge against instant exercise (ordinary stopping times). However, this turns out not to be a factor affecting the seller's price, as we shall see in Proposition 4.3.

4.1. Deferred solvency

These considerations indicate that the notion of solvency needs to be relaxed.

We say that a portfolio $z \in \mathcal{L}_t$ satisfies the *deferred solvency* condition at time t if it can be exchanged into a solvent portfolio by time T without any additional investment, i.e. if there is a sequence y_{t+1}, \dots, y_{T+1} such that $y_s \in \mathcal{L}_{s-1}$ for all $s = t + 1, \dots, T$ and

$$z - y_{t+1} \in \mathcal{K}_t, \quad y_s - y_{s+1} \in \mathcal{K}_s \text{ for all } s = t + 1, \dots, T, \quad y_{T+1} = 0.$$

We call such a sequence y_{t+1}, \dots, y_{T+1} a *liquidation strategy* starting from z at time t .

The set of portfolios satisfying the deferred solvency condition at time t is a cone. We call it the *deferred solvency cone* and denote by \mathcal{Q}_t .

Example 4.1. In Example 3.1 the portfolio with 8 in the domestic currency and -1 in the foreign currency is insolvent at the 'up' node $U = \{\omega_1, \omega_2\}$ at time 1, that is, $(-1, 8) \notin \mathcal{K}_1^U$. It does, however, satisfy the deferred solvency condition at that node, i.e. $(-1, 8) \in \mathcal{Q}_1^U$. The large bid-ask spread $[S_1^{bU}, S_1^{aU}] = [3, 9]$ at node U indicates a temporary loss of liquidity. Although the portfolio is insolvent at that node, waiting until the market recovers from the loss of liquidity can restore solvency. The liquidation strategy is to hold the portfolio until time 2 and to buy the foreign currency then.

The following result shows that the deferred solvency cones \mathcal{Q}_t can be regarded as the sets of time t superhedging portfolios for the seller of a European option with expiry time T and zero payoff; see Roux & Zastawniak (2011).

Proposition 4.1. *The deferred solvency cones can be constructed by backward induction as follows:*

$$\mathcal{Q}_T = \mathcal{K}_T, \tag{4.1}$$

$$\mathcal{Q}_t = \mathcal{Q}_{t+1} \cap \mathcal{L}_t + \mathcal{K}_t \text{ for all } t = 0, \dots, T - 1. \tag{4.2}$$

The proof of Proposition 4.1 can be found in Appendix A.4.1.

From (4.2) we can see that for any $t = 0, \dots, T - 1$ and for any $\mu \in \Omega_t$

$$\mathcal{Q}_t^\mu = \bigcap_{\nu \in \text{succ } \mu} \mathcal{Q}_{t+1}^\nu + \mathcal{K}_t^\mu. \tag{4.3}$$

By backward induction, \mathcal{Q}_t^μ is given as an intersection and algebraic sum of a finite number of polyhedral cones, so it is a polyhedral cone. This also means the solvency cones can readily be computed using standard operations on polyhedral convex sets.

The next result shows that Theorem 2.1 can be formulated equivalently in terms of the deferred solvency cones \mathcal{Q}_t instead of the solvency cones \mathcal{K}_t .

Proposition 4.2. *If \mathbb{Q} is a probability measure and $S = (S_t)_{t=0}^T$ is an \mathbb{R}^d -valued \mathbb{Q} -martingale, then S satisfies (2.2) if and only if*

$$S_t \in \mathcal{Q}_t^* \setminus \{0\} \text{ for all } t = 0, \dots, T, \quad (4.4)$$

where \mathcal{Q}_t^* is the polar of $-\mathcal{Q}_t$.

The proof of Proposition 4.2 is in Appendix A.4.1.

4.2. Construction of seller's price and superhedging strategy

We extend the family $\Phi^a(\xi)$ of seller's superhedging strategies by allowing for deferred solvency:

$$\Phi^{\text{ad}}(\xi) := \{y \in \Phi \mid \forall \tau \in \mathcal{T} : y_t - y_{t+1} \in \mathcal{Q}_t \text{ for } t = 1, \dots, \tau - 1, y_\tau - \xi_\tau \in \mathcal{Q}_\tau\},$$

The following proposition shows that the set of initial endowments that allow the seller to hedge against gradual exercise is the same as that allowing to hedge against instant exercise with deferred solvency.

Proposition 4.3. *For any American option ξ*

$$\{x \in \mathbb{R}^d \mid \exists Y \in \Phi^{\text{ag}}(\xi) : x = Y_0\} = \{x \in \mathbb{R}^d \mid \exists y \in \Phi^{\text{ad}}(\xi) : x = y_0\}.$$

For the proof of Proposition 4.3, see Appendix A.4.2.

We now present an iterative construction of the set of initial endowments that allow superhedging for the seller under deferred solvency. By Proposition 4.3, this also gives the set of initial endowments that allow superhedging for the seller under gradual exercise.

Construction 4.1. Construct adapted sequences $\mathcal{U}_t^{\text{ad}}, \mathcal{V}_t^{\text{ad}}, \mathcal{W}_t^{\text{ad}}, \mathcal{Z}_t^{\text{ad}}$ for $t = 0, \dots, T$ by

$$\begin{aligned} \mathcal{U}_t^{\text{ad}} &:= \xi_t + \mathcal{Q}_t \text{ for all } t = 0, \dots, T, \\ \mathcal{Z}_T^{\text{ad}} &:= \mathcal{V}_T^{\text{ad}} := \mathcal{W}_T^{\text{ad}} := \mathcal{U}_T^{\text{ad}}, \end{aligned} \quad (4.5)$$

and by backward induction on all $t = 0, \dots, T - 1$

$$\mathcal{W}_t^{\text{ad}} := \mathcal{Z}_{t+1}^{\text{ad}} \cap \mathcal{L}_t, \quad (4.6)$$

$$\mathcal{V}_t^{\text{ad}} := \mathcal{W}_t^{\text{ad}} + \mathcal{Q}_t, \quad (4.7)$$

$$\mathcal{Z}_t^{\text{ad}} := \mathcal{U}_t^{\text{ad}} \cap \mathcal{V}_t^{\text{ad}}. \quad (4.8)$$

It follows by backward induction that the sets

$$\mathcal{W}_t^{\text{ad}\mu} = \bigcap_{\nu \in \text{succ } \mu} \mathcal{Z}_{t+1}^{\text{ad}\nu}, \quad \mathcal{V}_t^{\text{ad}\mu} = \mathcal{W}_t^{\text{ad}\mu} + \mathcal{Q}_t^\mu, \quad \mathcal{Z}_t^{\text{ad}\mu} = \mathcal{U}_t^{\text{ad}\mu} \cap \mathcal{V}_t^{\text{ad}\mu}$$

are convex and polyhedral for each $t = 0, \dots, T-1$ and $\mu \in \Omega_t$ because the algebraic sum and the intersection of a finite number of convex polyhedral sets are convex and polyhedral, and

$$\mathcal{Z}_T^{\text{ad}\mu} = \mathcal{V}_T^{\text{ad}\mu} = \mathcal{W}_T^{\text{ad}\mu} = \mathcal{U}_T^{\text{ad}\mu} = \xi_T^\mu + \mathcal{Q}_T^\mu$$

are convex polyhedral sets for each $\mu \in \Omega_T$. Moreover, $\mathcal{U}_t^{\text{ad}}$, $\mathcal{V}_t^{\text{ad}}$, $\mathcal{W}_t^{\text{ad}}$, $\mathcal{Z}_t^{\text{ad}}$ are non-empty for each $t = 0, \dots, T$ because the portfolio $(z, \dots, z) \in \mathbb{R}^d$ belongs to all of them when $z \in \mathbb{R}$ is large enough.

Theorem 4.1. *The set of initial endowments that superhedge the seller's position in the American option ξ under gradual exercise is equal to*

$$\mathcal{Z}_0^{\text{ad}} = \{x \in \mathbb{R}^d \mid \exists Y \in \Phi^{\text{ag}}(\xi) : x = Y_0\}, \quad (4.9)$$

and the ask (seller's) price of the option in currency $j = 1, \dots, d$ can be computed as

$$\pi_j^{\text{ag}}(\xi) = \min\{x \in \mathbb{R} \mid xe^j \in \mathcal{Z}_0^{\text{ad}}\}.$$

Moreover, a strategy $Y \in \Phi^{\text{ag}}(\xi)$ can be constructed such that

$$\pi_j^{\text{ag}}(\xi)e^j = Y_0.$$

The proof of Theorem 4.1 can be found in Appendix A.4.2.

We can conclude that the set of initial endowments $\mathcal{Z}_0^{\text{ad}}$ superhedging the seller's position, the option ask price $\pi_j^{\text{ag}}(\xi)$, and a superhedging strategy Y realising the ask price can be computed by means of standard operations on convex polyhedral sets.

Example 4.2. Working within the setting of Example 3.1, we can now apply the constructions described in the current section to compute the sets $\mathcal{Z}_t^{\text{ad}}$ of superhedging portfolios for the seller. These are sets of portfolios $(x^1, x^2) \in \mathbb{R}^2$ satisfying the inequalities

	$\mathcal{Z}_0^{\text{ad}}$	$\mathcal{Z}_1^{\text{ad}}$	$\mathcal{Z}_2^{\text{ad}}$
ω_1	$5x^1 + x^2 \geq 5$	$8x^1 + x^2 \geq 8$ $4x^1 + x^2 \geq 0$	$8x^1 + x^2 \geq 8$ $4x^1 + x^2 \geq 0$
ω_2	$5x^1 + x^2 \geq 5$	$8x^1 + x^2 \geq 8$ $4x^1 + x^2 \geq 0$	$4x^1 + x^2 \geq 0$
ω_3	$5x^1 + x^2 \geq 5$	$2x^1 + x^2 \geq 0$	$3x^1 + x^2 \geq 0$
ω_4	$5x^1 + x^2 \geq 5$	$2x^1 + x^2 \geq 0$	$x^1 + x^2 \geq 0$

From $\mathcal{Z}_0^{\text{ad}}$ we obtain the ask price

$$\pi_2^{\text{ag}}(\xi) = \min\{x \in \mathbb{R} \mid (0, x) \in \mathcal{Z}_0^{\text{ad}}\} = 5.$$

We can also construct a superhedging strategy $Y \in \Phi^{\text{ag}}(\xi)$ such that

$$(0, \pi_2^{\text{ag}}(\xi)) = Y_0 = (0, 5).$$

It is the strategy Y specified in Example 3.1.

4.3. Dual representation of seller's price

A dual representation of the seller's price $\pi_j^{\text{ag}}(\xi)$ can be obtained with the aid of the support function Z_0^{ad} of $-\mathcal{Z}_0^{\text{ad}}$. For the definition of the support function of a convex set, see Appendix A.1. More generally, let $U_t^{\text{ad}}, V_t^{\text{ad}}, W_t^{\text{ad}}, Z_t^{\text{ad}}$ be the support functions of the sets $-\mathcal{U}_t^{\text{ad}}, -\mathcal{V}_t^{\text{ad}}, -\mathcal{W}_t^{\text{ad}}, -\mathcal{Z}_t^{\text{ad}}$ of Construction 4.1. The functions $U_t^{\text{ad}}, V_t^{\text{ad}}, W_t^{\text{ad}}, Z_t^{\text{ad}}$ are polyhedral (Rockafellar 1996 Corollary 19.2.1), hence continuous. Proposition A.1 in Appendix A.4.2 lists a number of properties of support functions, which will prove useful in what follows.

Proposition 4.4. *The seller's price of an American option ξ with gradual exercise can be written as*

$$\pi_j^{\text{ag}}(\xi) = \max\{-Z_0^{\text{ad}}(s) \mid s \in \sigma_j(\mathbb{R}^d)\} = \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{\chi})$$

for some mixed stopping time $\chi \in \mathcal{X}$, a probability measure \mathbb{Q} and an \mathbb{R}^d -valued adapted process S such that

$$S_t \in \mathcal{Q}_t^* \setminus \{0\} \quad \text{and} \quad \mathbb{E}_{\mathbb{Q}}(S_{t+1}^{X^*} | \mathcal{F}_t) \in \mathcal{Q}_t^* \quad \text{for each } t = 0, \dots, T, \quad (4.10)$$

and $S_t^j = 1$ for all $t = 0, \dots, T$. Such χ , \mathbb{Q} and S can be constructed by a recursive procedure.

The notation $\sigma_j(\mathbb{R}^d)$, $(\xi \cdot S)_{\chi}$ and S^{X^*} used in Proposition 4.4 is defined by (A.2), (A.5) and (A.6). The proof is provided in Appendix A.4.2.

For any $\chi \in \mathcal{X}$ denote by $\bar{\mathcal{P}}^d(\chi)$ the set of pairs (\mathbb{Q}, S) such that \mathbb{Q} is a probability measure and S is an \mathbb{R}^d -valued adapted process satisfying (4.10). Also define for $j = 1, \dots, d$

$$\bar{\mathcal{P}}_j^d(\chi) := \{(\mathbb{Q}, S) \in \bar{\mathcal{P}}^d(\chi) \mid S_t^j = 1 \text{ for all } t = 0, \dots, T\}.$$

The lack of arbitrage opportunities and Proposition 4.2 ensure that

$$\emptyset \neq \mathcal{P} \subseteq \bar{\mathcal{P}}^d(\chi), \quad \emptyset \neq \mathcal{P}_j \subseteq \bar{\mathcal{P}}_j^d(\chi)$$

for all $\chi \in \mathcal{X}$.

Remark 4.1. The superscript d indicating deferred solvency distinguishes $\bar{\mathcal{P}}^d(\chi)$ and $\bar{\mathcal{P}}_j^d(\chi)$ from the collections $\bar{\mathcal{P}}(\chi)$ and $\bar{\mathcal{P}}_j(\chi)$ defined by Roux & Zastawniak (2011) in a similar way as above, but with the weaker condition

$$S_t \in \mathcal{K}_t^* \setminus \{0\} \quad \text{and} \quad \mathbb{E}_{\mathbb{Q}}(S_{t+1}^{X^*} | \mathcal{F}_t) \in \mathcal{K}_t^* \quad \text{for each } t = 0, \dots, T$$

in place of (4.10).

The following result provides a representation of $\pi_j^{\text{ag}}(\xi)$ dual to the representation (3.3) in terms of superhedging strategies.

Theorem 4.2. *The ask price in currency $j = 1, \dots, d$ of an American option ξ with gradual exercise can be written as*

$$\pi_j^{\text{ag}}(\xi) = \max_{\chi \in \mathcal{X}} \max_{(Q, S) \in \bar{\mathcal{P}}_j^d(\chi)} \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{\chi}).$$

Moreover, we can algorithmically construct $\hat{\chi} \in \mathcal{X}$, and $(\hat{Q}, \hat{S}) \in \bar{\mathcal{P}}_j^d(\hat{\chi})$ such that

$$\pi_j^{\text{ag}}(\xi) = \mathbb{E}_{\hat{\mathbb{Q}}}((\xi \cdot \hat{S})_{\hat{\chi}}).$$

This theorem is proved in Appendix A.4.2.

Example 4.3. We continue working in the setting of Example 3.1. The mixed stopping time $\hat{\chi} \in \mathcal{X}$ and a pair $(\hat{Q}, \hat{S}) \in \bar{\mathcal{P}}_2^d(\hat{\chi})$ such that

$$\pi_2^{\text{ag}}(\xi) = \mathbb{E}_{\hat{\mathbb{Q}}}((\xi \cdot \hat{S})_{\hat{\chi}}) = 5$$

are

	\hat{Q}	\hat{S}_0	\hat{S}_1	\hat{S}_2	$\hat{\chi}_0$	$\hat{\chi}_1$	$\hat{\chi}_2$
ω_1	1	(5, 1)	(4, 1)	(8, 1)	0	$\frac{3}{4}$	$\frac{1}{4}$
ω_2	0	(5, 1)	(4, 1)	(4, 1)	0	$\frac{3}{4}$	$\frac{1}{4}$
ω_3	0	(5, 1)	(2, 1)	(3, 1)	0	0	1
ω_4	0	(5, 1)	(2, 1)	(1, 1)	0	0	1

5. Pricing and superhedging for the buyer under gradual exercise

The buyer of an American option ξ is entitled to receive the payoff according to a mixed stopping time $\chi \in \mathcal{X}$ of his choosing. In other words, the buyer receives $\chi_t \xi_t$ at each time $t = 0, \dots, T$. The family $\Phi^{\text{bg}}(\xi)$ of superhedging strategies for the buyer and the bid price (buyer's price) $\pi_j^{\text{bg}}(\xi)$ under gradual exercise are defined in Section 3.2. We turn to the task of computing the bid price and an optimal superhedging strategy for the buyer.

5.1. Construction of buyer's price and superhedging strategy

We start by computing the set of initial endowments that allow superhedging for the buyer.

Construction 5.1. Construct adapted sequences $\mathcal{U}_t^{\text{bd}}, \mathcal{V}_t^{\text{bd}}, \mathcal{W}_t^{\text{bd}}, \mathcal{Z}_t^{\text{bd}}$ for $t = 0, \dots, T$ by

$$\begin{aligned} \mathcal{U}_t^{\text{bd}} &:= -\xi_t + Q_t \text{ for all } t = 0, \dots, T, \\ \mathcal{Z}_T^{\text{bd}} &:= \mathcal{V}_T^{\text{bd}} := \mathcal{W}_T^{\text{bd}} := \mathcal{U}_T^{\text{bd}}, \end{aligned} \tag{5.1}$$

and by backward induction on all $t < T$

$$\mathcal{W}_t^{\text{bd}} := \mathcal{Z}_{t+1}^{\text{bd}} \cap \mathcal{L}_t, \quad (5.2)$$

$$\mathcal{V}_t^{\text{bd}} := \mathcal{W}_t^{\text{bd}} + \mathcal{Q}_t, \quad (5.3)$$

$$\mathcal{Z}_t^{\text{bd}} := \text{conv} \{ \mathcal{U}_t^{\text{bd}}, \mathcal{V}_t^{\text{bd}} \}. \quad (5.4)$$

For each t the convex hull in (5.4) is taken on each atom of \mathcal{F}_t , i.e. for all $\mu \in \Omega_t$

$$\mathcal{Z}_t^{\text{bd}\mu} = \text{conv} \{ \mathcal{U}_t^{\text{bd}\mu}, \mathcal{V}_t^{\text{bd}\mu} \}.$$

The index d indicates that the deferred solvency cones \mathcal{Q}_t are used in this construction. The sets $\mathcal{U}_t^{\text{bd}}$, $\mathcal{V}_t^{\text{bd}}$, $\mathcal{W}_t^{\text{bd}}$, $\mathcal{Z}_t^{\text{bd}}$ are non-empty for each $t = 0, \dots, T$ because the portfolio $(z, \dots, z) \in \mathbb{R}^d$ belongs to all of them when $z \in \mathbb{R}$ is large enough.

In contrast with Construction 4.6 of Roux & Zastawniak (2011), which was used the case of instant exercise at an ordinary stopping time, we have the convex hull of $\mathcal{U}_t^{\text{bd}}$, $\mathcal{V}_t^{\text{bd}}$ in (5.4) rather than the union of sets. This means that $\mathcal{U}_t^{\text{bd}}$, $\mathcal{V}_t^{\text{bd}}$, $\mathcal{W}_t^{\text{bd}}$, $\mathcal{Z}_t^{\text{bd}}$ are convex sets, unlike their counterparts in Construction 4.6 of Roux & Zastawniak (2011). This is important because, once it is established in the next proposition that the $\mathcal{Z}_t^{\text{bd}}$ are polyhedral, it becomes possible to implement techniques from convex analysis to compute them.

Another consequence of having the convex hull in (5.4) is that there is no analogue of Proposition 4.3 in the buyer's case. Gradual exercise allows the option buyer more room for manoeuvre when hedging their position as compared to instant exercise with or without deferred solvency. This is down to the fact that the convex hull $\text{conv} \{ \mathcal{U}_t^{\text{bd}}, \mathcal{V}_t^{\text{bd}} \}$ in (5.4) is, in general, larger than the union of such sets.

Remark 5.1. It is important to understand why we need the convex hull $\text{conv} \{ \mathcal{U}_t^{\text{bd}}, \mathcal{V}_t^{\text{bd}} \}$ in (5.4) instead of the union of sets as in Construction 4.6 of Roux & Zastawniak (2011). In a nutshell, the convex hull appears because an option buyer who is allowed to exercise gradually has the choice not only between exercising the option fully or continuing to hold it at any given time, but can also decide to exercise a fraction of the option and continue to hold the remaining fraction. More precisely, suppose that the buyer of an option with payoff ξ_t is also holding a portfolio $z \in \text{conv} \{ \mathcal{U}_t^{\text{bd}}, \mathcal{V}_t^{\text{bd}} \}$ at time t . We have $z = \lambda u + (1 - \lambda)v$ for some $u \in \mathcal{U}_t^{\text{bd}}$, $v \in \mathcal{V}_t^{\text{bd}}$ and $\lambda \in [0, 1]$. Under gradual exercise the buyer can exercise a fraction λ of the option at time t , which will add a fraction $\lambda\xi_t$ of the payoff to the portfolio z . The resulting portfolio $z + \lambda\xi_t$ belongs to $(1 - \lambda)\mathcal{V}_t^{\text{bd}}$ since, by (5.3),

$$z + \lambda\xi_t = \lambda(u + \xi_t) + (1 - \lambda)v \in \lambda\mathcal{Q}_t + (1 - \lambda)\mathcal{V}_t^{\text{bd}} \subseteq (1 - \lambda)\mathcal{V}_t^{\text{bd}}.$$

But any portfolio in $(1 - \lambda)\mathcal{V}_t^{\text{bd}}$ is enough to hedge the remaining fraction $1 - \lambda$ of the option, which the buyer will continue to hold beyond time t . It means that portfolios in $\text{conv} \{ \mathcal{U}_t^{\text{bd}}, \mathcal{V}_t^{\text{bd}} \}$ are exactly those that can hedge the buyer's position in the option at time t under gradual exercise.

Proposition 5.1. *The set $\mathcal{Z}_t^{\text{bd}}$ in Construction 5.1 is polyhedral with recession cone \mathcal{Q}_t for each $t = 0, \dots, T$.*

The proof of Proposition 5.1 can be found in Appendix A.4.3.

The next result shows that Construction 5.1 produces the set of initial endowments that superhedges ξ for the buyer, which in turn makes it possible to compute the option bid price and also to construct a strategy that realises this price. This is similar to Theorem 4.1 for the seller.

Theorem 5.1. *The set of initial endowments that superhedge the buyer's position in the American option ξ with gradual exercise is equal to*

$$\mathcal{Z}_0^{\text{bd}} = \{x \in \mathbb{R}^d \mid \exists(y, \chi) \in \Phi^{\text{bg}}(\xi) : x = y_0\}, \quad (5.5)$$

and the bid (buyer's) price of the option in currency $j = 1, \dots, d$ can be computed as

$$\pi_j^{\text{bg}}(\xi) = \max \{-x \in \mathbb{R} \mid xe^j \in \mathcal{Z}_0^{\text{bd}}\}.$$

Moreover, a strategy $(y, \chi) \in \Phi^{\text{bg}}(\xi)$ can be constructed such that

$$\pi_j^{\text{bg}}(\xi)e^j = -y_0.$$

The proof of this theorem is also in Appendix A.4.3.

Example 5.1. Still within the setting of Example 3.1, we apply the constructions described in the current section to compute the sets $\mathcal{Z}_t^{\text{bd}}$ of superhedging portfolios for the buyer. These are sets of portfolios $(x^1, x^2) \in \mathbb{R}^2$ satisfying the inequalities

	$\mathcal{Z}_0^{\text{bd}}$	$\mathcal{Z}_1^{\text{bd}}$	$\mathcal{Z}_2^{\text{bd}}$
ω_1	$5x^1 + x^2 \geq -3$	$8x^1 + x^2 \geq -8$	$8x^1 + x^2 \geq -8$
ω_2		$6x^1 + x^2 \geq -4$	$4x^1 + x^2 \geq 0$
ω_3	$5x^1 + x^2 \geq -3$	$4x^1 + x^2 \geq -4$	$4x^1 + x^2 \geq 0$
ω_4		$8x^1 + x^2 \geq -8$	$3x^1 + x^2 \geq 0$
	$5x^1 + x^2 \geq -3$	$6x^1 + x^2 \geq -4$	$3x^1 + x^2 \geq 0$
		$4x^1 + x^2 \geq -4$	$x^1 + x^2 \geq 0$
	$5x^1 + x^2 \geq -3$	$2x^1 + x^2 \geq 0$	$2x^1 + x^2 \geq 0$
	$5x^1 + x^2 \geq -3$	$2x^1 + x^2 \geq 0$	$x^1 + x^2 \geq 0$

From $\mathcal{Z}_0^{\text{bd}}$ we obtain the ask price

$$\pi_2^{\text{bg}}(\xi) = \max\{-x \in \mathbb{R} \mid (0, x) \in \mathcal{Z}_0^{\text{bd}}\} = 3.$$

We can also construct a superhedging strategy $(y, \chi) \in \Phi^{\text{bg}}(\xi)$ such that

$$(0, \pi_2^{\text{bg}}(\xi)) = -y_0 = (0, 3).$$

It is the strategy (y, χ) specified in Example 3.1.

5.2. Dual representation of buyer's price

Since the $\mathcal{U}_t^{\text{bd}}$, $\mathcal{V}_t^{\text{bd}}$, $\mathcal{W}_t^{\text{bd}}$, $\mathcal{Z}_t^{\text{bd}}$ are convex, it becomes possible to apply convex duality methods not just in the seller's case but also in the buyer's case. (This was impossible to do in Roux & Zastawniak (2011) for American options with instant exercise because of the lack of convexity in the buyer's case.)

In particular, in a similar way as in the proof of Proposition 4.4, we can show that the bid price of an American option with payoff ξ under gradual exercise can be expressed as

$$\pi_j^{\text{bg}}(\xi) = \max \{ Z_0^{\text{bd}}(s) \mid s \in \sigma_j(\mathbb{R}^d) \}$$

in terms of the support function Z_0^{bd} of $-\mathcal{Z}_0^{\text{bd}}$.

However, we follow a different approach to obtain a representation of the bid price $\pi_j^{\text{bg}}(\xi)$ dual to the representation (3.5) of $\pi_j^{\text{bg}}(\xi)$ by means of superhedging strategies. In Theorem 5.1 a mixed stopping time $\chi \in \mathcal{X}$ has already been constructed as part of a superhedging strategy $(y, \chi) \in \Phi^{\text{bg}}(\xi)$ such that $\pi_j^{\text{bg}}(\xi) = y_0$. As a result, the bid price given by (3.5) can be written as

$$\pi_j^{\text{bg}}(\xi) = \max \{ -x \in \mathbb{R} \mid \exists y \in \Phi : (y, \chi) \in \Phi^{\text{bg}}(\xi), xe^j = y_0 \}$$

for this mixed stopping time χ . It turns out that the set on the right-hand side can be expressed by means of the family $\Psi^{\text{a}}(-\xi_\chi)$ of superhedging strategies for the seller of a European option with expiry time T and payoff $-\xi_\chi$ as described in Appendix A.5, where ξ_χ is defined by (A.6).

Proposition 5.2. *For any American option ξ and any mixed stopping time $\chi \in \mathcal{X}$ we have*

$$\begin{aligned} \{ -x \in \mathbb{R}^d \mid \exists y \in \Phi : (y, \chi) \in \Phi^{\text{bg}}(\xi), x = y_0 \} \\ = \{ -x \in \mathbb{R}^d \mid \exists z \in \Psi^{\text{a}}(-\xi_\chi) : x = z_0 \}. \end{aligned}$$

This proposition is proved in Appendix A.4.3.

We are now in a position to state a representation of the bid price dual to (3.5), and to prove it with the aid of Proposition 5.2.

Theorem 5.2. *The buyer's (bid) price of an American option ξ in currency $j = 1, \dots, d$ can be represented as*

$$\pi_j^{\text{bg}}(\xi) = \max_{\chi \in \mathcal{X}} \min_{(\mathbb{Q}, S) \in \mathcal{P}_j} \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_\chi), \quad (5.6)$$

where $(\xi \cdot S)_\chi$ is defined by (A.6). Moreover, we can algorithmically construct $\hat{\chi} \in \mathcal{X}$ and $(\hat{\mathbb{Q}}, \hat{S}) \in \mathcal{P}_j$ such that

$$\pi_j^{\text{bg}}(\xi) = \mathbb{E}_{\hat{\mathbb{Q}}}((\xi \cdot \hat{S})_{\hat{\chi}}) = \min_{(\mathbb{Q}, S) \in \mathcal{P}_j} \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{\hat{\chi}}).$$

The proof of Theorem 5.2 is in Appendix A.4.3.

Example 5.2. We revisit Example 3.1 one more time to construct a mixed stopping time $\hat{\chi} \in \mathcal{X}$ and a pair $(\hat{Q}, \hat{S}) \in \bar{\mathcal{P}}_2$ such that

$$\pi_2^{\text{bg}}(\xi) = \mathbb{E}_{\hat{Q}}((\xi \cdot \hat{S})_{\hat{\chi}}) = \min_{(Q,S) \in \bar{\mathcal{P}}_2} \mathbb{E}_Q((\xi \cdot S)_{\hat{\chi}}) = 3.$$

They are

	\hat{Q}	\hat{S}_0	\hat{S}_1	\hat{S}_2	$\hat{\chi}_0$	$\hat{\chi}_1$	$\hat{\chi}_2$
ω_1	1	(5, 1)	(5, 1)	(5, 1)	0	$\frac{1}{2}$	$\frac{1}{2}$
ω_2	0	(5, 1)	(5, 1)	(4, 1)	0	$\frac{1}{2}$	$\frac{1}{2}$
ω_3	0	(5, 1)	(2, 1)	(3, 1)	0	0	1
ω_4	0	(5, 1)	(2, 1)	(1, 1)	0	0	1

6. Numerical example

In this section we present a three-dimensional numerical example with a realistic flavour to illustrate Constructions 4.1 and 5.1. The numerical procedures below were implemented in *Maple* with the aid of the *Convex* package (Franz 2009).

Consider a model involving a domestic currency and two foreign currencies, with time horizon $\tau = 1$ and with $T = 10$ time steps. The friction-free nominal exchange rates $E_t = (E_t^1, E_t^2)$ between the domestic currency and the two foreign currencies follow the two-asset recombining Korn & Müller (2009) model with Cholesky decomposition. That is, there are $(t + 1)^2$ possibilities for the exchange rates at each time step $t = 0, \dots, T$, indexed by pairs (j_1, j_2) with $1 \leq j_1, j_2 \leq t + 1$, and each non-terminal node with exchange rates $E_t(j_1, j_2)$ has four successors, associated with exchange rates $E_{t+1}(j_1, j_2)$, $E_{t+1}(j_1 + 1, j_2)$, $E_{t+1}(j_1, j_2 + 1)$ and $E_{t+1}(j_1 + 1, j_2 + 1)$. With $\Delta = \frac{\tau}{T}$ defined for convenience, the exchange rates are given by

$$E_t^1(j_1, j_2) = E_0^1 e^{-\frac{1}{2}\sigma_1^2 t \Delta + (2j_1 - t - 2)\sigma_1 \sqrt{\Delta}},$$

$$E_t^2(j_1, j_2) = E_0^2 e^{-\frac{1}{2}\sigma_2^2 t \Delta + ((2j_1 - t - 2)\rho + (2j_2 - t - 2)\sqrt{1 - \rho^2})\sigma_2 \sqrt{\Delta}}$$

for $t = 0, \dots, T$ and $j_1, j_2 = 1, \dots, t + 1$, where $E_0^1 = 40$ and $E_0^2 = 50$ are the initial exchange rates, $\sigma_1 = 15\%$ and $\sigma_2 = 10\%$ are the volatilities and $\rho = 50\%$ is the correlation between the logarithmic growth of the exchange rates.

Assume that proportional transaction costs of 0.5% are payable on all currency exchanges, except at time step 1, when 10% is payable, modelling a temporary loss of liquidity. In other words, the matrix of exchange rates between each pair among the three currencies at each time step t is

$$\begin{pmatrix} \pi_t^{11} & \pi_t^{12} & \pi_t^{13} \\ \pi_t^{21} & \pi_t^{22} & \pi_t^{23} \\ \pi_t^{31} & \pi_t^{32} & \pi_t^{33} \end{pmatrix} = \begin{pmatrix} 1 & \frac{E_t^2}{E_t^1}(1 + k_t) & \frac{1}{E_t^1}(1 + k_t) \\ \frac{E_t^1}{E_t^2}(1 + k_t) & 1 & \frac{1}{E_t^2}(1 + k_t) \\ \frac{E_t^1}{E_t^1}(1 + k_t) & \frac{E_t^2}{E_t^2}(1 + k_t) & 1 \end{pmatrix},$$

18 *Alet Roux and Tomasz Zastawniak*

where

$$k_t = \begin{cases} 0.1 & \text{if } t = 1, \\ 0.005 & \text{otherwise.} \end{cases}$$

Consider an American put option with physical delivery and strike 90 on a basket containing one unit of each of the foreign currencies. It offers the payoff

$$\xi_t = (-1, -1, 90) \quad \text{for } t = 0, \dots, T.$$

We allow for the possibility that the option may never be exercised by adding an extra time step $T + 1$ to the model and setting the payoff to be $(0, 0, 0)$ at that time step. Constructions 4.1 and 5.1 give

$$\begin{aligned} \mathcal{Z}_0^{\text{ad}} &= \text{conv} \left\{ \begin{pmatrix} -0.749 \\ -0.218 \\ 47.587 \end{pmatrix}, \begin{pmatrix} -0.166 \\ -0.727 \\ 49.773 \end{pmatrix} \right\} + \mathcal{Q}_0, \\ \mathcal{Z}_0^{\text{bd}} &= \text{conv} \left\{ \begin{pmatrix} 0.631 \\ 0.783 \\ -65.310 \end{pmatrix}, \begin{pmatrix} 0.453 \\ 0.286 \\ -33.404 \end{pmatrix}, \begin{pmatrix} 0.419 \\ 0.782 \\ -56.798 \end{pmatrix}, \begin{pmatrix} 0.249 \\ 0.267 \\ -24.342 \end{pmatrix}, \right. \\ &\quad \left. \begin{pmatrix} 0.329 \\ 0.513 \\ -39.790 \end{pmatrix} \right\} + \mathcal{Q}_0, \end{aligned}$$

where \mathcal{Q}_0 is the convex cone generated by the vectors

$$\begin{pmatrix} 0.000 \\ 0.020 \\ -1.000 \end{pmatrix}, \begin{pmatrix} -1.000 \\ 0.804 \\ 0.000 \end{pmatrix}, \begin{pmatrix} 1.246 \\ -1.000 \\ 0.000 \end{pmatrix}, \begin{pmatrix} -1.000 \\ 0.000 \\ 40.200 \end{pmatrix}, \begin{pmatrix} 0.025 \\ 0.000 \\ -1.000 \end{pmatrix}, \begin{pmatrix} 0.000 \\ -1.000 \\ 50.250 \end{pmatrix}.$$

The sets $\mathcal{Z}_0^{\text{ad}}$ and $\mathcal{Z}_0^{\text{bd}}$, which appear in Figure 1, yield the ask and bid prices

$$\begin{aligned} \pi_1^{\text{ag}}(\xi) &= 0.174, & \pi_2^{\text{ag}}(\xi) &= 0.140, & \pi_3^{\text{ag}}(\xi) &= 6.941, \\ \pi_1^{\text{bg}}(\xi) &= 0.022, & \pi_2^{\text{bg}}(\xi) &= 0.017, & \pi_3^{\text{bg}}(\xi) &= 0.879 \end{aligned}$$

in each of the three currencies.

7. Conclusions and Outlook

In this paper we have explored American options with gradual exercise within Kabanov's model (Kabanov 1999) of many assets under transaction costs, along with the related notion of deferred solvency, which helps to deal with a temporary loss of liquidity (large bid-ask spreads) in the market. We have demonstrated that gradual exercise (at a mixed stopping time chosen by the buyer) can reduce the ask (seller's) price and increase the bid (buyer's) price of the option compared with the case when the option can only be exercised instantly (at an ordinary stopping time).

In this context we have constructed and implemented algorithms to compute the ask and bid option prices, the buyer's and seller's optimal hedging portfolios




Fig. 1. Boundaries of the sets of superhedging endowments for the seller and buyer

and strategies, and their optimal mixed stopping times. We have studied dual representations for both the buyer and the seller of an American option with gradual exercise. The results have been illustrated by numerical examples.

Compared to options with instant exercise, a novel feature is that pricing and hedging an American option is a convex optimisation problem not just for the seller but also for the buyer of the option, making it possible to use convex duality in both cases. Ramifications to be explored further may include an extension of Bouchard and Temam’s representation of the strategies hedging the seller’s (short) position (Bouchard & Temam 2005) to the case of hedging the buyer’s (long) position in the option.

The computational efficiency of the algorithms under gradual exercise is similar to the seller’s algorithm in the case of instant exercise, explored numerically by Zhang & Roux & Zastawniak (2012). These calculations took no more than a couple of seconds using a modest parallel computer with 8 processors to solve the pricing problem for the seller under instant exercise in a realistic setting on a recombining binomial tree with up to 1500 time steps. There reasons for this high efficiency are two-fold: the convexity of the optimisation problem, and the fact that the procedure follows the principles of dynamic programming, and therefore shows merely polynomial growth for options with path-independent payoffs (such as puts and calls) in a recombining tree model, and additionally lends itself to parallelisation. The computational overhead (due to precomputing the deferred solvency cones) needed to implement the case of gradual exercise will have only a minor effect on computational efficiency. Moreover, since pricing and hedging becomes a convex optimisation problem for the buyer when gradual exercise is allowed, similarly efficient calculations can also be performed for the buyer and not just for the

20 *Alet Roux and Tomasz Zastawniak*

seller of the option.

We also conjecture that it should be possible to adapt the constructions presented here so that linear vector optimisation methods can be used to price and hedge both the seller's and buyer's positions in an American option with gradual exercise, along similar lines as was done by Löhne & Rudloff (2014) for European options under transaction costs, possibly resulting in further efficiency gains.

A. Appendix

A.1. *Some notation and facts from convex analysis*

For any non-empty convex cone $A \subseteq \mathbb{R}^d$, denote by A^* the *polar* of $-A$, i.e.

$$A^* := \{y \in \mathbb{R}^d \mid y \cdot x \geq 0 \text{ for all } x \in A\}. \quad (\text{A.1})$$

For any set $A \subseteq \mathbb{R}^d$ define the *cone generated by* A as

$$\text{cone } A := \{\lambda x \mid \lambda \geq 0, x \in A\}.$$

The *recession cone* of a non-empty convex set $A \subseteq \mathbb{R}^d$ is defined as

$$0^+ A := \{y \in \mathbb{R}^d \mid A + \lambda y \subseteq A \text{ for all } \lambda \geq 0\}.$$

It is a convex cone containing the origin (Rockafellar 1996 Theorem 8.1). If A is a polyhedral cone, then $0^+ A = A$ (Rockafellar 1996 Corollary 8.3.2).

The *convex hull* of sets A_1, \dots, A_n in \mathbb{R}^d is the smallest convex set in \mathbb{R}^d that contains A_1, \dots, A_n , and is denoted by $\text{conv}\{A_1, \dots, A_n\}$. The *convex hull* of convex functions $f_1, \dots, f_n : \mathbb{R} \rightarrow \mathbb{R} \cup \{\infty\}$ is the function $f := \text{conv}\{f_1, \dots, f_n\}$ defined by

$$f(x) := \inf \left\{ \sum_{i=1}^n \lambda_i f_i(x_i) \mid 0 \leq \lambda_1, \dots, \lambda_n \leq 1, \sum_{i=1}^n \lambda_i = 1, \sum_{i=1}^n \lambda_i x_i = x \right\}.$$

The *effective domain* of a convex function $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\infty\}$ is defined as

$$\text{dom } f := \{y \in \mathbb{R}^d \mid f(y) < \infty\}.$$

The *support function* δ_A^* of a convex set $A \subseteq \mathbb{R}^d$ is defined as

$$\delta_A^*(x) := \sup\{x \cdot y \mid y \in A\}.$$

A.2. *Compactly generated cones*

For any set $A \subseteq \mathbb{R}^d$ and $j = 1, \dots, d$ define

$$\sigma_j(A) := \{x = (x^1, \dots, x^d) \in A \mid x^j = 1\}. \quad (\text{A.2})$$

We say that a cone $A \subset \mathbb{R}^d$ is *compactly j -generated* if $\sigma_j(A)$ is compact, non-empty and A is generated by $\sigma_j(A)$.

Lemma A.1. *If two cones $A \subset \mathbb{R}^d$ and $B \subset \mathbb{R}^d$ are compactly j -generated and $A \cap B \setminus \{0\} \neq \emptyset$, then $A \cap B$ is compactly j -generated and*

$$\sigma_j(A \cap B) = \sigma_j(A) \cap \sigma_j(B). \quad (\text{A.3})$$

Proof. Equality (A.3) follows directly from (A.2). A vector $x = (x^1, \dots, x^d)$ is an element of $\sigma_j(A \cap B)$ if and only if $x \in A \cap B$ and $x^j = 1$, if and only if $x \in A$ and $x \in B$ and $x^j = 1$, if and only if $x \in \sigma_j(A)$ and $x \in \sigma_j(B)$, if and only if $x \in \sigma_j(A) \cap \sigma_j(B)$.

The set $\sigma_j(A \cap B)$ is compact since it is the intersection of two compact sets $\sigma_j(A)$ and $\sigma_j(B)$. It remains to show that $\sigma_j(A \cap B)$ is non-empty and generates $A \cap B$. To this end, fix any $x \in A \cap B \setminus \{0\}$. As A and B are generated, respectively, by $\sigma_j(A)$ and $\sigma_j(B)$, there exist $\lambda_A \geq 0$, $\lambda_B \geq 0$, $x_A \in \sigma_j(A)$ and $x_B \in \sigma_j(B)$ such that

$$\lambda_A x_A = x = \lambda_B x_B.$$

As $x \neq 0$, we must have $\lambda_A > 0$ and $\lambda_B > 0$. Moreover, since $x_A^j = x_B^j = 1$, we have

$$\lambda_A = \lambda_A x_A^j = x^j = \lambda_B x_B^j = \lambda_B,$$

which implies $x_A = x_B \in \sigma_j(A) \cap \sigma_j(B) = \sigma_j(A \cap B)$, completing the proof. \square

In this paper we also make use of the following result by Roux & Zastawniak (2011 Lemma A.1).

Lemma A.2. *Fix any $j = 1, \dots, d$, and suppose that A_1, \dots, A_n are non-empty closed convex sets in \mathbb{R}^d such that $A := \bigcap_{i=1}^n A_i \neq \emptyset$ and $\text{dom } \delta_{A_i}^*$ is compactly j -generated for all i . Then*

$$\delta_A^* = \text{conv}\{\delta_{A_1}^*, \dots, \delta_{A_n}^*\},$$

the cone $\text{dom } \delta_A^*$ is compactly j -generated and

$$\text{dom } \delta_A^* = \text{conv}\{\text{dom } \delta_{A_1}^*, \dots, \text{dom } \delta_{A_n}^*\},$$

and for each $x \in \sigma_j(\text{dom } \delta_A^*)$ there exist $\alpha_1, \dots, \alpha_n \geq 0$ and x_1, \dots, x_n with $x_i \in \sigma_j(\text{dom } \delta_{A_i}^*)$ for all $i = 1, \dots, n$ such that

$$\delta_A^*(x) = \sum_{i=1}^n \alpha_i \delta_{A_i}^*(x_i), \quad x = \sum_{i=1}^n \alpha_i x_i, \quad 1 = \sum_{i=1}^n \alpha_i.$$

A.3. Mixed stopping times

A *mixed* (or *randomised*) *stopping time* is a non-negative adapted process $\chi = (\chi_t)_{t=0}^T$ with values in $[0, 1]$ such that

$$\sum_{t=0}^T \chi_t = 1.$$

22 Alet Roux and Tomasz Zastawniak

The family of mixed stopping times will be denoted by \mathcal{X} .

For any $\chi \in \mathcal{X}$ we put

$$\chi_t^* := \sum_{s=t}^T \chi_s \text{ for } t = 0, \dots, T, \quad \chi_{T+1}^* := 0. \quad (\text{A.4})$$

Moreover, for any adapted process X and for any $\chi \in \mathcal{X}$ we put

$$X_t^{\chi^*} := \sum_{s=t}^T \chi_s X_s \text{ for } t = 0, \dots, T, \quad X_{T+1}^{\chi^*} := 0. \quad (\text{A.5})$$

We also define X evaluated at χ by

$$X_\chi := \sum_{s=0}^T \chi_s X_s. \quad (\text{A.6})$$

With each ordinary stopping time $\tau \in \mathcal{T}$ we associate the mixed stopping time $\chi^\tau \in \mathcal{X}$ defined as

$$\chi_t^\tau := \begin{cases} 1 & \text{on } \{\tau = t\} \\ 0 & \text{on } \{\tau \neq t\} \end{cases} \text{ for each } t = 0, \dots, T. \quad (\text{A.7})$$

A.4. Proofs and technical results

A.4.1. Deferred solvency

Proof of Proposition 4.1. Equality (4.1) is obvious. By the definition of the deferred solvency cones, for any $t = 0, \dots, T-1$ the following conditions are equivalent: $z \in \mathcal{Q}_{t+1} \cap \mathcal{L}_t + \mathcal{K}_t$ if and only if there is a $y_{t+1} \in \mathcal{L}_t$ such that

$$y_{t+1} \in \mathcal{Q}_{t+1}, \quad z - y_{t+1} \in \mathcal{K}_t,$$

if and only if there is a sequence y_{t+1}, \dots, y_{T+1} such that $y_s \in \mathcal{L}_{s-1}$ for each $s = t+1, \dots, T$ and

$$z - y_{t+1} \in \mathcal{K}_t, \quad y_s - y_{s+1} \in \mathcal{K}_s \text{ for all } s = t+1, \dots, T, \quad y_{T+1} = 0,$$

if and only if $z \in \mathcal{Q}_t$. This proves (4.2). \square

Proof of Proposition 4.2. In view of (4.2), we have $\mathcal{K}_t \subseteq \mathcal{Q}_t$, so $\mathcal{Q}_t^* \subseteq \mathcal{K}_t^*$, and (4.4) implies (2.2).

Conversely, suppose that S is an \mathbb{R}^d -valued \mathbb{Q} -martingale that satisfies (2.2). To show that it satisfies (4.4) we proceed by backward induction. By (4.1), we have $S_T \in \mathcal{K}_T^* \setminus \{0\} = \mathcal{Q}_T^* \setminus \{0\}$. For any $t = 0, \dots, T-1$ suppose that $S_{t+1} \in \mathcal{Q}_{t+1}^* \setminus \{0\}$. As S is a \mathbb{Q} -martingale, we have for all $\mu \in \Omega_t$ that

$$S_t^\mu = \mathbb{E}_{\mathbb{Q}}(S_{t+1} | \mu) \in \text{conv}\{\mathcal{Q}_{t+1}^{\nu^*} | \nu \in \text{succ } \mu\}.$$

For every $t = 0, \dots, T-1$ and $\mu \in \Omega_t$, observe from (4.3) that

$$\mathcal{Q}_t^{\mu^*} = \left[\bigcap_{\nu \in \text{succ } \mu} \mathcal{Q}_{t+1}^{\nu} + \mathcal{K}_t^\mu \right]^*.$$

Successive application of Corollaries 16.4.2 and 16.5.2 in Rockafellar (1996) then gives

$$\mathcal{Q}_t^{\mu^*} = \left[\bigcap_{\nu \in \text{succ } \mu} \mathcal{Q}_{t+1}^\nu \right]^* \cap \mathcal{K}_t^{\mu^*} = \text{conv}\{\mathcal{Q}_{t+1}^{\nu^*} \mid \nu \in \text{succ } \mu\} \cap \mathcal{K}_t^{\mu^*}. \quad (\text{A.8})$$

Since $S_t^\mu \in \mathcal{K}_t^{\mu^*} \setminus \{0\}$ by (2.2), it follows that $S_t^\mu \in \mathcal{Q}_t^{\mu^*} \setminus \{0\}$, which concludes the inductive step. \square

A.4.2. Seller's case

Proof of Proposition 4.3. We show first that for any $y \in \Phi^{\text{ad}}(\xi)$ there exists $Y \in \Phi^{\text{ag}}(\xi)$ such that $Y_0 = y_0$. If $y \in \Phi^{\text{ad}}(\xi)$, then for each $t = 0, \dots, T-1$ we have $y_t - y_{t+1} \in \mathcal{Q}_t$, i.e. there exists a liquidation strategy $z_{t+1}^t, \dots, z_{T+1}^t$ starting from $y_t - y_{t+1}$ at time t . We also put $z_{T+1}^T := 0$ for notational convenience. Moreover, for each $t = 0, \dots, T$ we have $y_t - \xi_t \in \mathcal{Q}_t$, i.e. there exists a liquidation strategy $x_{t+1}^t, \dots, x_{T+1}^t$ starting from $y_t - \xi_t$ at time t . For each $\chi \in \mathcal{X}$ define

$$Y_0^\chi := y_0, \\ Y_t^\chi := \chi_t^* y_t + \sum_{s=0}^{t-1} \chi_{s+1}^* z_t^s + \sum_{s=0}^{t-1} \chi_s x_t^s \text{ for } t = 1, \dots, T+1,$$

where χ^* is defined by (A.4). The process Y^χ belongs to Φ and satisfies the non-anticipation condition (3.2). Moreover, for each $t = 0, \dots, T$

$$\begin{aligned} Y_t^\chi - \chi_t \xi_t - Y_{t+1}^\chi &= \chi_t^* y_t + \sum_{s=0}^{t-1} \chi_{s+1}^* z_t^s + \sum_{s=0}^{t-1} \chi_s x_t^s - \chi_t \xi_t \\ &\quad - \chi_{t+1}^* y_{t+1} - \sum_{s=0}^t \chi_{s+1}^* z_{t+1}^s - \sum_{s=0}^t \chi_s x_{t+1}^s \\ &= \chi_{t+1}^* (y_t - y_{t+1} - z_{t+1}^t) + \chi_t (y_t - \xi_t - x_{t+1}^t) \\ &\quad + \sum_{s=0}^{t-1} \chi_{s+1}^* (z_t^s - z_{t+1}^s) + \sum_{s=0}^{t-1} \chi_s (x_t^s - x_{t+1}^s) \\ &\in \chi_{t+1}^* \mathcal{K}_t + \chi_t \mathcal{K}_t + \sum_{s=0}^{t-1} \chi_{s+1}^* \mathcal{K}_t + \sum_{s=0}^{t-1} \chi_s \mathcal{K}_t \subseteq \mathcal{K}_t \end{aligned}$$

because $\chi_t^* = \chi_t + \chi_{t+1}^*$ and \mathcal{K}_t is a convex cone. Hence Y satisfies (3.1) in addition to (3.2), and so $Y \in \Phi^{\text{ag}}(\xi)$.

Conversely, fix any $Y \in \Phi^{\text{ag}}(\xi)$ and put $y := Y^{\chi^T} \in \Phi$, where χ^T is defined by (A.7). Then for all $s = 0, \dots, T-1$ we have $\chi_s^T = 0$ and

$$y_s - y_{s+1} = Y_s^{\chi^T} - Y_{s+1}^{\chi^T} = Y_s^{\chi^T} - \chi_s^T \xi_s - Y_{s+1}^{\chi^T} \in \mathcal{K}_s \subseteq \mathcal{Q}_s.$$

Fix any $t = 0, \dots, T$. Then $\chi_s^T = \chi_s^t = 0$ for each $s = 0, \dots, t-1$, and the non-anticipation property (3.2) of Y gives $y_t = Y_t^{\chi^T} = Y_t^{\chi^t}$. Since $\chi_t^t = 1$, it means

24 Alet Roux and Tomasz Zastawniak

that

$$y_t - \xi_t - Y_{t+1}^{\chi^t} = Y_t^{\chi^t} - \chi_t^t \xi_t - Y_{t+1}^{\chi^t} \in \mathcal{K}_t \subseteq \mathcal{Q}_t. \quad (\text{A.9})$$

Moreover, for each $s = t + 1, \dots, T$ we have $\chi_s^t = 0$, and so

$$Y_s^{\chi^t} - Y_{s+1}^{\chi^t} = Y_s^{\chi^t} - \chi_s^t \xi_s - Y_{s+1}^{\chi^t} \in \mathcal{K}_s \subseteq \mathcal{Q}_s. \quad (\text{A.10})$$

We verify by backward induction that $Y_{s+1}^{\chi^t} \in \mathcal{Q}_s$ for each $s = t, \dots, T$. Clearly, $Y_{T+1}^{\chi^t} = 0 \in \mathcal{Q}_T$. Now suppose that $Y_{s+1}^{\chi^t} \in \mathcal{Q}_s$ for some $s = t + 1, \dots, T$. From (A.10) we can see that $Y_s^{\chi^t} = (Y_s^{\chi^t} - Y_{s+1}^{\chi^t}) + Y_{s+1}^{\chi^t} \in \mathcal{Q}_s + \mathcal{Q}_s = \mathcal{Q}_s$. Because Y^{χ^t} is predictable, we have $Y_s^{\chi^t} \in \mathcal{Q}_s \cap \mathcal{L}_{s-1} \subseteq \mathcal{Q}_{s-1}$ by (4.2), completing the backward induction argument. In particular, this means that $Y_{t+1}^{\chi^t} \in \mathcal{Q}_t$. Together with (A.9) it gives $y_t - \xi_t \in \mathcal{Q}_t$ for any $t = 0, \dots, T$. As a result, we have constructed $y \in \Phi^{\text{ad}}(\xi)$ such that $y_0 = Y_0$. \square

Proof of Theorem 4.1. Suppose that $x \in \mathcal{Z}_0^{\text{ad}}$. We construct a sequence $y = (y_t)_{t=0}^{T+1}$ of random variables by induction. First take $y_0 = x$. Now suppose that we have already constructed $y_t \in \mathcal{Z}_t^{\text{ad}}$ such that $y_t \in \mathcal{L}_{(t-1) \vee 0}$ for some $t = 0, \dots, T-1$. From (4.8) we obtain $y_t \in \mathcal{U}_t^{\text{ad}}$, whence

$$y_t - \xi_t \in \mathcal{Q}_t$$

by (4.5). We also obtain $y_t \in \mathcal{V}_t^{\text{ad}}$, and by (4.7) there exists a random variable $y_{t+1} \in \mathcal{W}_t^{\text{ad}}$ such that $y_t - y_{t+1} \in \mathcal{Q}_t$. From (4.6) we have $y_{t+1} \in \mathcal{Z}_{t+1}^{\text{ad}} \cap \mathcal{L}_t$, which concludes the inductive step. Finally, we put $y_{T+1} := 0$. It follows that $y \in \Phi^{\text{ad}}(\xi)$ with $y_0 = x$. By Proposition 4.3, a strategy $Y \in \Phi^{\text{ag}}(\xi)$ can be constructed such that $Y_0 = y_0 = x$.

Suppose now that $Y \in \Phi^{\text{ag}}(\xi)$. By Proposition 4.3, there is a $y \in \Phi^{\text{ad}}(\xi)$ such that $Y_0 = y_0$. Clearly,

$$y_t \in \xi_t + \mathcal{Q}_t = \mathcal{U}_t^{\text{ad}} \quad (\text{A.11})$$

for all $t = 0, \dots, T$, and in particular $y_T \in \mathcal{Z}_T^{\text{ad}} = \mathcal{U}_T^{\text{ad}}$. We now show by backward induction that $y_t \in \mathcal{Z}_t^{\text{ad}}$ for all $t = 0, \dots, T$. Suppose that $y_{t+1} \in \mathcal{Z}_{t+1}^{\text{ad}}$ for some $t = 0, \dots, T-1$. Since $y_{t+1} \in \mathcal{L}_t$, this means by (4.6) that $y_{t+1} \in \mathcal{W}_t^{\text{ad}}$. The condition $y_t - y_{t+1} \in \mathcal{Q}_t$ implies that $y_t \in \mathcal{W}_t^{\text{ad}} + \mathcal{Q}_t = \mathcal{V}_t^{\text{ad}}$. Property (A.11) then gives $y_t \in \mathcal{Z}_t^{\text{ad}}$ by (4.8), which completes the inductive step. We conclude that $Y_0 = y_0 \in \mathcal{Z}_0^{\text{ad}}$.

We have proved (4.9). It follows that

$$\begin{aligned} \pi_j^{\text{ag}}(\xi) &= \inf\{x \in \mathbb{R} \mid \exists Y \in \Phi^{\text{ag}}(\xi) : xe^j = Y_0\} \\ &= \inf\{x \in \mathbb{R} \mid xe^j \in \mathcal{Z}_0^{\text{ad}}\}. \end{aligned}$$

We know that $\mathcal{Z}_0^{\text{ad}}$ is polyhedral, hence closed, so $\{x \in \mathbb{R} \mid xe^j \in \mathcal{Z}_0^{\text{ad}}\}$ is also a closed set. It is non-empty and bounded below because $xe^j \in \mathcal{Z}_0^{\text{ad}}$ for any $x \in \mathbb{R}$ large enough, and $xe^j \notin \mathcal{Z}_0^{\text{ad}}$ for any $x \in \mathbb{R}$ small enough. As a result, the infimum

is attained. It means, in particular, that $\pi_j^{\text{ag}}(\xi)e^j \in \mathcal{Z}_0^{\text{ad}}$, so we know that a strategy $Y \in \Phi^{\text{ag}}(\xi)$ can be constructed such that $\pi_j^{\text{ag}}(\xi)e^j = Y_0$. \square

The following result is similar to Lemma 5.5 in (Roux & Zastawniak 2011), but with the solvency cones \mathcal{K}_t replaced by the deferred solvency cones \mathcal{Q}_t .

Proposition A.1.

- (1) For each $t = 0, \dots, T$ the set \mathcal{Q}_t^* is compactly j -generated.
- (2) For all $t = 0, \dots, T$ and $y \in \mathcal{L}_t$ we have

$$U_t^{\text{ad}}(y) = \begin{cases} -y \cdot \xi_t & \text{on } \{y \in \mathcal{Q}_t^*\}, \\ \infty & \text{on } \{y \notin \mathcal{Q}_t^*\}, \end{cases} \quad (\text{A.12})$$

$$V_t^{\text{ad}}(y) = \begin{cases} W_t^{\text{ad}}(y) & \text{on } \{y \in \mathcal{Q}_t^*\}, \\ \infty & \text{on } \{y \notin \mathcal{Q}_t^*\}. \end{cases} \quad (\text{A.13})$$

- (3) We have $\text{dom } Z_t^{\text{ad}} = \mathcal{Q}_t^*$ for all $t = 0, \dots, T$. For all $t = 0, \dots, T-1$ and $\mu \in \Omega_t$ we have

$$Z_t^{\text{ad}\mu} = \text{conv}\{U_t^{\text{ad}\mu}, V_t^{\text{ad}\mu}\}, \quad (\text{A.14})$$

and for each $y \in \sigma_j(\mathcal{Q}_t^{\mu*})$ there exist $\lambda \in [0, 1]$, $x \in \sigma_j(\text{dom } V_t^{\text{ad}\mu})$ and $s \in \sigma_j(\mathcal{Q}_t^{\mu*})$ such that

$$Z_t^{\text{ad}\mu}(y) = \lambda U_t^{\text{ad}\mu}(s) + (1 - \lambda)V_t^{\text{ad}\mu}(x), \quad y = \lambda s + (1 - \lambda)x.$$

- (4) For every $t = 0, \dots, T-1$ and $\mu \in \Omega_t$ we have

$$W_t^{\text{ad}\mu} = \text{conv}\{Z_{t+1}^{\text{ad}\nu} \mid \nu \in \text{succ } \mu\},$$

and for each $x \in \sigma_j(\text{dom } W_t^{\text{ad}\mu})$ there exist $q^\nu \in [0, 1]$ and $y^\nu \in \sigma_j(\mathcal{Q}_{t+1}^{\nu*})$ for all $\nu \in \text{succ } \mu$ such that

$$W_t^{\text{ad}\mu}(x) = \sum_{\nu \in \text{succ } \mu} q^\nu Z_{t+1}^{\text{ad}\nu}(y^\nu), \quad x = \sum_{\nu \in \text{succ } \mu} q^\nu y^\nu, \quad 1 = \sum_{\nu \in \text{succ } \mu} q^\nu.$$

Proof. We first consider claim (2). As \mathcal{Q}_t is a cone,

$$\delta_{-\mathcal{Q}_t}^*(x) = \begin{cases} 0 & \text{if } x \in \mathcal{Q}_t^*, \\ \infty & \text{otherwise.} \end{cases}$$

Note in particular that $\text{dom } \delta_{-\mathcal{Q}_t}^* = \mathcal{Q}_t^*$. For all t and $y \in \mathcal{L}_t$ we have (Rockafellar 1996 p. 113)

$$U_t^{\text{ad}}(y) = \delta_{-\xi_t - \mathcal{Q}_t}^*(y) = \delta_{\{-\xi_t\}}^*(y) + \delta_{-\mathcal{Q}_t}^*(y),$$

which leads to (A.12). Equation (A.13) follows similarly from

$$V_t^{\text{ad}} = \delta_{-\mathcal{W}_t^{\text{ad}} - \mathcal{Q}_t}^* = \delta_{-\mathcal{W}_t^{\text{ad}}}^* + \delta_{-\mathcal{Q}_t}^* = W_t^{\text{ad}} + \delta_{-\mathcal{Q}_t}^*.$$

26 Alet Roux and Tomasz Zastawniak

Claims (1), (3) and (4) can be obtained by backward induction. We clearly have

$$\text{dom } Z_T^{\text{ad}} = \text{dom } U_T^{\text{ad}} = \text{dom } \delta_{-\mathcal{Q}_T}^* = \mathcal{Q}_T^*,$$

and this set is compactly j -generated because $\mathcal{Q}_T^* = \mathcal{K}_T^*$.

Now fix any $t = 0, \dots, T-1$ and $\mu \in \Omega_t$, and suppose that $\text{dom } Z_{t+1}^{\text{ad}} = \mathcal{Q}_{t+1}^*$ and that this set is compactly j -generated. Since

$$\bigcap_{\nu \in \text{succ } \mu} Z_{t+1}^{\text{ad}\nu} = W_t^{\text{ad}\mu} \neq \emptyset,$$

Lemma A.2 can be applied to the sets $-Z_{t+1}^{\text{ad}\nu}$ for all $\nu \in \text{succ } \mu$. This justifies claim (4) for this t and also that

$$\text{dom } W_t^{\text{ad}\mu} = \text{conv} \{ \text{dom } Z_{t+1}^{\text{ad}\nu} \mid \nu \in \text{succ } \mu \} = \text{conv} \{ \mathcal{Q}_{t+1}^{\nu*} \mid \nu \in \text{succ } \mu \}$$

is compactly j -generated.

By Theorem 2.1 and Proposition 4.2, the lack of arbitrage opportunities implies that there is a pair $(\mathbb{Q}, S) \in \mathcal{P}$ such that $S_t^\mu \in \mathcal{Q}_t^{\mu*} \setminus \{0\} \subseteq \mathcal{K}_t^{\mu*} \setminus \{0\}$ and $S_{t+1}^\nu \in \mathcal{Q}_{t+1}^{\nu*}$ for each $\nu \in \text{succ } \mu$. Since S is a martingale under \mathbb{Q} , it follows that

$$S_t^\mu = \mathbb{E}_{\mathbb{Q}}(S_{t+1} \mid \mu) \in \text{conv} \{ \mathcal{Q}_{t+1}^{\nu*} \mid \nu \in \text{succ } \mu \},$$

and so $S_t^\mu \in \text{conv} \{ \mathcal{Q}_{t+1}^{\nu*} \mid \nu \in \text{succ } \mu \} \cap \mathcal{K}_t^{\mu*} \setminus \{0\} \neq \emptyset$. As $\text{conv} \{ \mathcal{Q}_{t+1}^{\nu*} \mid \nu \in \text{succ } \mu \}$ and $\mathcal{K}_t^{\mu*}$ are compactly j -generated, it follows from Lemma A.1 and (A.8) that $\mathcal{Q}_t^{\mu*} = \text{conv} \{ \mathcal{Q}_{t+1}^{\nu*} \mid \nu \in \text{succ } \mu \} \cap \mathcal{K}_t^{\mu*}$ is compactly j -generated, which justifies claim (1) for this value of t .

In view of (A.12) and (A.13), Lemma A.1 consequently shows that

$$\text{dom } V_t^{\text{ad}\mu} = \text{dom } W_t^{\text{ad}\mu} \cap \text{dom } U_t^{\text{ad}\mu} = \text{dom } W_t^{\text{ad}\mu} \cap \mathcal{Q}_t^{\mu*}$$

is also compactly j -generated. We may apply Lemma A.2 to the sets $-\mathcal{U}_t^{\text{ad}\mu}$ and $-\mathcal{V}_t^{\text{ad}\mu}$ since

$$\mathcal{U}_t^{\text{ad}\mu} \cap \mathcal{V}_t^{\text{ad}\mu} = \mathcal{Z}_t^{\text{ad}\mu} \neq \emptyset.$$

Claim (3) for this value of t follows upon observing that

$$\text{dom } Z_t^{\text{ad}\mu} = \text{conv} \{ \text{dom } U_t^{\text{ad}\mu}, \text{dom } V_t^{\text{ad}\mu} \} = \text{conv} \{ \mathcal{Q}_t^{\mu*}, \text{dom } W_t^{\text{ad}\mu} \cap \mathcal{Q}_t^{\mu*} \} = \mathcal{Q}_t^{\mu*}.$$

This concludes the inductive step. \square

Proof of Proposition 4.4. By Proposition A.1, $\text{dom } Z_0^{\text{ad}} = \mathcal{Q}_0^*$ is compactly j -generated. Since $-Z_0^{\text{ad}}$ is polyhedral, it is continuous on its effective domain and

therefore attains a maximum on the non-empty compact set $\sigma_j(\mathcal{Q}_0^*)$. From Theorem 4.1 it follows (Rockafellar 1996 Theorem 13.1) that

$$\begin{aligned}\pi_j^{\text{ag}}(\xi) &= \min\{x \in \mathbb{R} \mid xe^j \in \mathcal{Z}_0^{\text{ad}}\} \\ &= \min\{x \in \mathbb{R} \mid xe^j \cdot s \geq -Z_0^{\text{ad}}(s) \text{ for all } s \in \mathbb{R}^d\} \\ &= \min\{x \in \mathbb{R} \mid xe^j \cdot s \geq -Z_0^{\text{ad}}(s) \text{ for all } s \in \mathcal{Q}_0^*\} \\ &= \min\{x \in \mathbb{R} \mid x \geq -Z_0^{\text{ad}}(s) \text{ for all } s \in \sigma_j(\mathcal{Q}_0^*)\} \\ &= \max\{-Z_0^{\text{ad}}(s) \mid s \in \sigma_j(\mathcal{Q}_0^*)\} \\ &= \max\{-Z_0^{\text{ad}}(s) \mid s \in \sigma_j(\mathbb{R}^d)\}.\end{aligned}$$

The following construction produces adapted processes y_t , λ_t , x_t and S_t for $t = 0, \dots, T$, and q_t for $t = 1, \dots, T$. We already know that the maximum of $-Z_0^{\text{ad}}$ over the set $\sigma_j(\mathbb{R}^d)$ is attained, i.e. there exists some $y_0 \in \sigma_j(\mathcal{Q}_0^*)$ such that

$$\pi_j^{\text{ag}}(\xi) = -Z_0^{\text{ad}}(y_0) = \max\{-Z_0^{\text{ad}}(s) \mid s \in \sigma_j(\mathbb{R}^d)\}. \quad (\text{A.15})$$

For any $t = 0, \dots, T-1$, suppose that $y_t \in \sigma_j(\mathcal{Q}_t^*)$ is given, and fix any $\mu \in \Omega_t$. Then by Proposition A.1(3), there exist $\lambda_t^\mu \in [0, 1]$, $x_t^\mu \in \sigma_j(\text{dom } V_t^{\text{ad}\mu})$ and $S_t^\mu \in \sigma_j(\mathcal{Q}_t^{\mu*})$ such that

$$Z_t^{\text{ad}\mu}(y_t^\mu) = \lambda_t^\mu U_t^{\text{ad}\mu}(S_t^\mu) + (1 - \lambda_t^\mu) V_t^{\text{ad}\mu}(x_t^\mu), \quad (\text{A.16})$$

$$y_t^\mu = \lambda_t^\mu S_t^\mu + (1 - \lambda_t^\mu) x_t^\mu. \quad (\text{A.17})$$

By (A.13) and Proposition A.1(4), there exist $q_{t+1}^\nu \in [0, 1]$ and $y_{t+1}^\nu \in \sigma_j(\mathcal{Q}_{t+1}^{\nu*})$ for all $\nu \in \text{succ } \mu$ such that

$$V_t^{\text{ad}\mu}(x_t^\mu) = W_t^{\text{ad}\mu}(x_t^\mu) = \sum_{\nu \in \text{succ } \mu} q_{t+1}^\nu Z_{t+1}^{\text{ad}\nu}(y_{t+1}^\nu), \quad (\text{A.18})$$

$$x_t^\mu = \sum_{\nu \in \text{succ } \mu} q_{t+1}^\nu y_{t+1}^\nu, \quad (\text{A.19})$$

$$1 = \sum_{\nu \in \text{succ } \mu} q_{t+1}^\nu.$$

This completes the inductive step. Also define for all $\mu \in \Omega_T$

$$\lambda_T^\mu := 1, \quad x_T^\mu := S_T^\mu := y_T^\mu.$$

Then (A.16), (A.17) are also satisfied when $t = T$.

Define the probability measure \mathbb{Q} on $\mathcal{F}_T = 2^\Omega$ as

$$\mathbb{Q}(\{\omega\}) := \prod_{t=1}^T q_t^{\omega \uparrow t} \text{ for all } \omega \in \Omega,$$

where $\omega \uparrow t$ denotes the element of Ω_t that contains ω . It then follows from (A.18), (A.19) that for all $t = 0, \dots, T-1$

$$V_t^{\text{ad}}(x_t) = \mathbb{E}_{\mathbb{Q}}(Z_{t+1}^{\text{ad}}(y_{t+1}) \mid \mathcal{F}_t), \quad (\text{A.20})$$

$$x_t = \mathbb{E}_{\mathbb{Q}}(y_{t+1} \mid \mathcal{F}_t). \quad (\text{A.21})$$

28 *Alet Roux and Tomasz Zastawniak*

The mixed stopping time χ is defined by setting $\chi_0 := \lambda_0$ and

$$\chi_t := \lambda_t \left(1 - \sum_{s=0}^{t-1} \chi_s \right) \text{ for all } t = 1, \dots, T.$$

It is straightforward to show by induction that $\chi_t \geq 0$ for all t . Moreover, since $\lambda_T = 1$, we have

$$\sum_{t=0}^T \chi_t = 1.$$

Observe also that

$$\lambda_t \chi_t^* = \chi_t, \quad (1 - \lambda_t) \chi_t^* = \chi_t^* - \chi_t = \chi_{t+1}^*$$

for all $t = 0, \dots, T$, where χ^* is defined by (A.4). It then follows from (A.16), (A.17) and (A.12) that for all $t = 0, \dots, T$

$$\chi_t^* Z_t^{\text{ad}}(y_t) = -\chi_t \xi_t \cdot S_t + \chi_{t+1}^* V_t^{\text{ad}}(x_t), \quad (\text{A.22})$$

$$\chi_t^* y_t = \chi_t S_t + \chi_{t+1}^* x_t. \quad (\text{A.23})$$

We now show by backward induction that for all $t = 0, \dots, T$

$$\mathbb{E}_{\mathbb{Q}}(S_{t+1}^{\chi^*} | \mathcal{F}_t) = \chi_{t+1}^* x_t. \quad (\text{A.24})$$

At time T the result is trivial because $S_{T+1}^{\chi^*} = \chi_{T+1}^* = 0$. Suppose now that (A.24) holds for some $t = 1, \dots, T$. Then, by the tower property of conditional expectation,

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}(S_t^{\chi^*} | \mathcal{F}_{t-1}) &= \mathbb{E}_{\mathbb{Q}}(\chi_t S_t + S_{t+1}^{\chi^*} | \mathcal{F}_{t-1}) \\ &= \mathbb{E}_{\mathbb{Q}}(\chi_t S_t + \mathbb{E}_{\mathbb{Q}}(S_{t+1}^{\chi^*} | \mathcal{F}_t) | \mathcal{F}_{t-1}) \\ &= \mathbb{E}_{\mathbb{Q}}(\chi_t S_t + \chi_{t+1}^* x_t | \mathcal{F}_{t-1}) \end{aligned}$$

and, by (A.23), the predictability of χ^* , and (A.21),

$$\mathbb{E}_{\mathbb{Q}}(S_t^{\chi^*} | \mathcal{F}_{t-1}) = \mathbb{E}_{\mathbb{Q}}(\chi_t^* y_t | \mathcal{F}_{t-1}) = \chi_t^* \mathbb{E}_{\mathbb{Q}}(y_t | \mathcal{F}_{t-1}) = \chi_t^* x_{t-1}.$$

This concludes the inductive step.

We also show by backward induction that for all $t = 0, \dots, T$

$$\chi_t^* Z_t^{\text{ad}}(y_t) = -\mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_t^{\chi^*} | \mathcal{F}_t). \quad (\text{A.25})$$

At time T

$$\chi_T^* Z_T^{\text{ad}}(y_T) = \chi_T^* U_T^{\text{ad}}(y_T) = -\chi_T \xi_T \cdot S_T = -(\xi \cdot S)_T^{\chi^*} = -\mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_T^{\chi^*} | \mathcal{F}_T).$$

Suppose now that (A.25) holds for some $t = 1, \dots, T$. Then by (A.22), (A.20) and the tower property of conditional expectation, we have

$$\begin{aligned} \chi_{t-1}^* Z_{t-1}^{\text{ad}}(y_{t-1}) &= -\chi_{t-1} \xi_{t-1} \cdot S_{t-1} + \chi_t^* V_{t-1}^{\text{ad}}(x_{t-1}) \\ &= -\chi_{t-1} \xi_{t-1} \cdot S_{t-1} + \chi_t^* \mathbb{E}_{\mathbb{Q}}(Z_t^{\text{ad}}(y_t) | \mathcal{F}_{t-1}) \\ &= -\chi_{t-1} \xi_{t-1} \cdot S_{t-1} + \mathbb{E}_{\mathbb{Q}}(\chi_t^* Z_t^{\text{ad}}(y_t) | \mathcal{F}_{t-1}) \\ &= -\chi_{t-1} \xi_{t-1} \cdot S_{t-1} - \mathbb{E}_{\mathbb{Q}}(\mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_t^{\chi^*} | \mathcal{F}_t) | \mathcal{F}_{t-1}) \\ &= -\chi_{t-1} \xi_{t-1} \cdot S_{t-1} - \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_t^{\chi^*} | \mathcal{F}_{t-1}) \\ &= -\mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{t-1}^{\chi^*} | \mathcal{F}_{t-1}). \end{aligned}$$

This concludes the inductive step.

Since $x_t \in \text{dom } V_t^{\text{ad}} \subseteq \mathcal{Q}_t^*$ for all t , property (A.24) implies that $\mathbb{E}_{\mathbb{Q}}(S_{t+1}^{\chi^*} | \mathcal{F}_t) \in \mathcal{Q}_t^*$ for all $t = 0, \dots, T$. Moreover, by (A.25),

$$-Z_0^{\text{ad}}(y_0) = -\chi_0^* Z_0^{\text{ad}}(y_0) = \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_0^{\chi^*} | \mathcal{F}_0) = \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_\chi).$$

From (A.15) we therefore have $\pi_j^{\text{ag}}(\xi) = \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_\chi)$, as required. \square

Proof of Theorem 4.2. By Proposition 4.4, a stopping time $\hat{\chi} \in \mathcal{X}$ and a pair $(\hat{\mathbb{Q}}, \hat{S}) \in \bar{\mathcal{P}}_j^{\text{d}}(\hat{\chi})$ can be constructed such that

$$\pi_j^{\text{ag}}(\xi) = \mathbb{E}_{\hat{\mathbb{Q}}}((\xi \cdot \hat{S})_{\hat{\chi}}) \leq \max_{\chi \in \mathcal{X}} \max_{(\mathbb{Q}, S) \in \bar{\mathcal{P}}_j^{\text{d}}(\chi)} \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_\chi).$$

To establish the reverse inequality we prove by backward induction that for any $y \in \Phi^{\text{ad}}(\xi)$, $\chi \in \mathcal{X}$ and $(\mathbb{Q}, S) \in \bar{\mathcal{P}}^{\text{d}}(\chi)$

$$y_t \cdot \mathbb{E}_{\mathbb{Q}}(S_t^{\chi^*} | \mathcal{F}_t) \geq \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_t^{\chi^*} | \mathcal{F}_t) \text{ for all } t = 0, \dots, T. \quad (\text{A.26})$$

When $t = T$,

$$y_T \cdot \mathbb{E}_{\mathbb{Q}}(S_T^{\chi^*} | \mathcal{F}_T) = \chi_T y_T \cdot S_T \geq \chi_T \xi_T \cdot S_T = \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_T^{\chi^*} | \mathcal{F}_T)$$

since $y_T - \xi_T \in \mathcal{Q}_T$ and $S_T \in \mathcal{Q}_T^*$. Now fix any $t = 0, \dots, T-1$, and suppose that

$$y_{t+1} \cdot \mathbb{E}_{\mathbb{Q}}(S_{t+1}^{\chi^*} | \mathcal{F}_{t+1}) \geq \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{t+1}^{\chi^*} | \mathcal{F}_{t+1}).$$

Then, by the tower property of conditional expectation, and since $y_t - y_{t+1} \in \mathcal{Q}_t$ and $\mathbb{E}_{\mathbb{Q}}(S_t^{\chi^*} | \mathcal{F}_t) \in \mathcal{Q}_t^*$, it follows that

$$\begin{aligned} y_t \cdot \mathbb{E}_{\mathbb{Q}}(S_t^{\chi^*} | \mathcal{F}_t) &= \chi_t y_t \cdot S_t + y_t \cdot \mathbb{E}_{\mathbb{Q}}(S_{t+1}^{\chi^*} | \mathcal{F}_t) \\ &\geq \chi_t \xi_t \cdot S_t + y_{t+1} \cdot \mathbb{E}_{\mathbb{Q}}(S_{t+1}^{\chi^*} | \mathcal{F}_t) \\ &= \chi_t \xi_t \cdot S_t + \mathbb{E}_{\mathbb{Q}}(y_{t+1} \cdot \mathbb{E}_{\mathbb{Q}}(S_{t+1}^{\chi^*} | \mathcal{F}_{t+1}) | \mathcal{F}_t) \\ &\geq \chi_t \xi_t \cdot S_t + \mathbb{E}_{\mathbb{Q}}(\mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{t+1}^{\chi^*} | \mathcal{F}_{t+1}) | \mathcal{F}_t) \\ &= \chi_t \xi_t \cdot S_t + \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{t+1}^{\chi^*} | \mathcal{F}_t) = \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_t^{\chi^*} | \mathcal{F}_t), \end{aligned}$$

which proves (A.26). The construction in the proof of Theorem 4.1 with initial portfolio $\hat{y}_0 = \pi_j^{\text{ag}}(\xi) e^j \in \mathcal{Z}_0^{\text{ad}}$ yields a strategy $\hat{y} \in \Phi^{\text{ad}}(\xi)$. For any $\chi \in \mathcal{X}$ and $(\mathbb{Q}, S) \in \bar{\mathcal{P}}_j^{\text{d}}(\chi)$ we have $\mathbb{E}_{\mathbb{Q}}(S_\chi^j) = 1$, and therefore (A.26) with $t = 0$ yields

$$\pi_j^{\text{ag}}(\xi) = \hat{y}_0 \cdot \mathbb{E}_{\mathbb{Q}}(S_\chi) \geq \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_\chi).$$

30 Alet Roux and Tomasz Zastawniak

It follows that

$$\pi_j^{\text{ag}}(\xi) \geq \max_{\chi \in \mathcal{X}} \max_{(Q,S) \in \mathcal{P}_j^{\text{d}}(\chi)} \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{\chi}). \quad \square$$

A.4.3. Buyer's case

Proof of Proposition 5.1. As \mathcal{Q}_t and \mathcal{L}_t are polyhedral cones, they are closed and convex. We have $0^+ \mathcal{Q}_t = \mathcal{Q}_t$ and $0^+ \mathcal{L}_t = \mathcal{L}_t$. It follows that $0^+ \mathcal{U}_t^{\text{bd}} = \mathcal{Q}_t$ for all $t = 0, \dots, T$ (Rockafellar 1996 Corollary 9.1.2).

The set $\mathcal{Z}_T^{\text{bd}} = \mathcal{U}_T^{\text{bd}} = -\xi_T + \mathcal{Q}_T$ is clearly polyhedral with recession cone \mathcal{Q}_T . For $t = 0, \dots, T-1$ we proceed by induction. Suppose that $\mathcal{Z}_{t+1}^{\text{bd}}$ is polyhedral and its recession cone is \mathcal{Q}_{t+1} . Then $\mathcal{W}_t^{\text{bd}} = \mathcal{Z}_{t+1}^{\text{bd}} \cap \mathcal{L}_t$ is polyhedral and its recession cone is $\mathcal{Q}_{t+1} \cap \mathcal{L}_t$ (Rockafellar 1996 Corollary 8.3.3). Being polyhedral, $\mathcal{W}_t^{\text{bd}}$ is the convex hull of a finite set of points and directions, and its recession cone $\mathcal{Q}_{t+1} \cap \mathcal{L}_t$ is the convex hull of the origin and the directions in $\mathcal{W}_t^{\text{bd}}$.

The set $\mathcal{V}_t^{\text{bd}} = \mathcal{W}_t^{\text{bd}} + \mathcal{Q}_t$ is polyhedral (Rockafellar 1996 Corollary 19.3.2) and hence it is the convex hull of a finite set of points and directions. Since the cone \mathcal{Q}_t can be written as the convex hull of the origin and a finite number of directions, it is possible to write $\mathcal{V}_t^{\text{bd}}$ as the convex hull of a finite set of points, all in $\mathcal{W}_t^{\text{bd}}$, and a finite set of directions. These directions are exactly the directions in $\mathcal{W}_t^{\text{bd}}$ and \mathcal{Q}_t , i.e. the directions in $\mathcal{Q}_{t+1} \cap \mathcal{L}_t$ and \mathcal{Q}_t . Thus the recession cone of $\mathcal{V}_t^{\text{bd}}$ is

$$0^+ \mathcal{V}_t^{\text{bd}} = \text{conv}\{\mathcal{Q}_t, \mathcal{Q}_{t+1} \cap \mathcal{L}_t\} = \mathcal{Q}_t$$

since $\mathcal{Q}_{t+1} \cap \mathcal{L}_t \subseteq \mathcal{Q}_t$ by (4.2). This means that the set $\mathcal{Z}_t^{\text{bd}} = \text{conv}\{\mathcal{V}_t^{\text{bd}}, \mathcal{U}_t^{\text{bd}}\}$ is closed and its recession cone is \mathcal{Q}_t (Rockafellar 1996 Corollary 9.8.1). Moreover, since $\mathcal{V}_t^{\text{bd}}$ and $\mathcal{U}_t^{\text{bd}}$ are polyhedral, it follows that $\text{cl} \mathcal{Z}_t^{\text{bd}} = \text{cl}(\text{conv}\{\mathcal{V}_t^{\text{bd}}, \mathcal{U}_t^{\text{bd}}\})$ is polyhedral (Rockafellar 1996 Theorem 19.6), which means that $\mathcal{Z}_t^{\text{bd}} = \text{cl} \mathcal{Z}_t^{\text{bd}}$ is polyhedral, concluding the inductive step. \square

Proposition A.2. *If $(y, \chi) \in \Phi^{\text{bg}}(\xi)$, then for all $t = 0, \dots, T$*

$$y_t \in \begin{cases} \chi_t^* \mathcal{Z}_t^{\text{bd}} & \text{on } \{\chi_t^* > 0\}, \\ \mathcal{Q}_t & \text{on } \{\chi_t^* = 0\}. \end{cases}$$

Proof. The proof is by backward induction. Since $y_{T+1} = 0$, from (3.4) we have

$$y_T \in -\chi_T \xi_T + \mathcal{K}_T \subseteq -\chi_T \xi_T + \mathcal{Q}_T.$$

It immediately follows that $y_T \in \mathcal{Q}_T$ on the set $\{\chi_T^* = 0\} = \{\chi_T = 0\}$. On the set $\{\chi_T^* > 0\} = \{\chi_T > 0\}$ we have $\mathcal{Q}_T = \chi_T \mathcal{Q}_T$ because \mathcal{Q}_T is a cone, and therefore

$$y_T \in \chi_T(-\xi_T + \mathcal{Q}_T) = \chi_T \mathcal{U}_T^{\text{bd}} = \chi_T^* \mathcal{Z}_T^{\text{bd}}.$$

Suppose now for some $t = 0, \dots, T-1$ that

$$y_{t+1} \in \begin{cases} \chi_{t+1}^* \mathcal{Z}_{t+1}^{\text{bd}} & \text{on } \{\chi_{t+1}^* > 0\}, \\ \mathcal{Q}_{t+1} & \text{on } \{\chi_{t+1}^* = 0\}. \end{cases}$$

Because $y_{t+1}, \chi_{t+1}^* \in \mathcal{L}_t$, this means that

$$y_{t+1} \in \begin{cases} \chi_{t+1}^* \mathcal{W}_t^{\text{bd}} & \text{on } \{\chi_{t+1}^* > 0\}, \\ \mathcal{Q}_{t+1} \cap \mathcal{L}_t & \text{on } \{\chi_{t+1}^* = 0\}. \end{cases}$$

Since $y_t \in -\chi_t \xi_t + y_{t+1} + \mathcal{K}_t \subseteq -\chi_t \xi_t + y_{t+1} + \mathcal{Q}_t$ by (3.4), it follows that

$$y_t \in \begin{cases} -\chi_t \xi_t + \chi_{t+1}^* \mathcal{W}_t^{\text{bd}} + \mathcal{Q}_t & \text{on } \{\chi_{t+1}^* > 0\}, \\ -\chi_t \xi_t + \mathcal{Q}_{t+1} \cap \mathcal{L}_t + \mathcal{Q}_t & \text{on } \{\chi_{t+1}^* = 0\}. \end{cases}$$

We consider the two possibilities separately.

- On the set $\{\chi_{t+1}^* > 0\}$ we have $\chi_t^* > 0$ and therefore

$$\mathcal{Q}_t = \chi_t^* \mathcal{Q}_t \subseteq \chi_t \mathcal{Q}_t + \chi_{t+1}^* \mathcal{Q}_t,$$

so that

$$\begin{aligned} y_t &\in -\chi_t \xi_t + \chi_{t+1}^* \mathcal{W}_t^{\text{bd}} + \chi_t \mathcal{Q}_t + \chi_{t+1}^* \mathcal{Q}_t \\ &= \chi_t (-\xi_t + \mathcal{Q}_t) + \chi_{t+1}^* (\mathcal{W}_t^{\text{bd}} + \mathcal{Q}_t) = \chi_t \mathcal{U}_t^{\text{bd}} + \chi_{t+1}^* \mathcal{V}_t^{\text{bd}}. \end{aligned}$$

Since

$$\chi_t \mathcal{U}_t^{\text{bd}} + \chi_{t+1}^* \mathcal{V}_t^{\text{bd}} = \chi_t^* \left(\frac{\chi_t}{\chi_t^*} \mathcal{U}_t^{\text{bd}} + \frac{\chi_{t+1}^*}{\chi_t^*} \mathcal{V}_t^{\text{bd}} \right) \subseteq \chi_t^* \mathcal{Z}_t^{\text{bd}},$$

it follows that $y_t \in \chi_t^* \mathcal{Z}_t^{\text{bd}}$ on $\{\chi_{t+1}^* > 0\}$.

- On the set $\{\chi_{t+1}^* = 0\}$ we have

$$y_t \in -\chi_t \xi_t + \mathcal{Q}_{t+1} \cap \mathcal{L}_t + \mathcal{Q}_t \subseteq -\chi_t \xi_t + \mathcal{Q}_t$$

because $\mathcal{Q}_{t+1} \cap \mathcal{L}_t \subseteq \mathcal{Q}_t$ by (4.2). There are two further possibilities.

- On $\{\chi_t^* > 0\} \cap \{\chi_{t+1}^* = 0\}$ we have $\chi_t = \chi_t^* > 0$ and therefore

$$y_t \in -\chi_t \xi_t + \mathcal{Q}_t = \chi_t (-\xi_t + \mathcal{Q}_t) = \chi_t \mathcal{U}_t^{\text{bd}} \subseteq \chi_t \mathcal{Z}_t^{\text{bd}} = \chi_t^* \mathcal{Z}_t^{\text{bd}}.$$

- On $\{\chi_t^* = 0\} \subseteq \{\chi_{t+1}^* = 0\}$ we have $\chi_t = 0$ and therefore $y_t \in \mathcal{Q}_t$ as claimed. \square

Proof of Theorem 5.1. In view of Proposition A.2, to verify (5.5) it is sufficient to show that for every $x \in \mathcal{Z}_0^{\text{bd}}$ there exists a pair $(y, \chi) \in \Phi^{\text{bg}}(\xi)$ such that $y_0 = x$. To this end, define $p_0 := 1$ and $z_0 := x \in p_0 \mathcal{Z}_0^{\text{bd}}$. Suppose by induction that for some $t = 0, \dots, T-1$ we have constructed predictable sequences z_0, \dots, z_t and p_0, \dots, p_t such that $1 = p_0 \geq \dots \geq p_t \geq 0$ and

$$\begin{aligned} z_s + (p_s - p_{s+1}) \xi_s - z_{s+1} &\in \mathcal{Q}_s && \text{for all } s = 0, \dots, t-1, \\ z_s &\in p_s \mathcal{Z}_s^{\text{bd}} && \text{for all } s = 0, \dots, t. \end{aligned}$$

Because of (5.4), there exists an \mathcal{F}_t -measurable random variable λ_t such that $0 \leq \lambda_t \leq 1$ and

$$z_t \in (1 - \lambda_t) p_t \mathcal{U}_t^{\text{bd}} + \lambda_t p_t \mathcal{V}_t^{\text{bd}}.$$

32 Alet Roux and Tomasz Zastawniak

Equations (5.1) and (5.3) then give

$$z_t \in (1 - \lambda_t)p_t(-\xi_t + \mathcal{Q}_t) + \lambda_t p_t(\mathcal{W}_t^{\text{bd}} + \mathcal{Q}_t) = -(1 - \lambda_t)p_t\xi_t + p_t\mathcal{Q}_t + \lambda_t p_t\mathcal{W}_t^{\text{bd}},$$

where

$$(1 - \lambda_t)p_t\mathcal{Q}_t + \lambda_t p_t\mathcal{Q}_t = p_t\mathcal{Q}_t$$

follows from the fact that \mathcal{Q}_t is a convex cone. This means there exists a random variable

$$z_{t+1} \in \lambda_t p_t \mathcal{W}_t^{\text{bd}} = \lambda_t p_t (\mathcal{Z}_{t+1}^{\text{bd}} \cap \mathcal{L}_t) = (\lambda_t p_t \mathcal{Z}_{t+1}^{\text{bd}}) \cap \mathcal{L}_t$$

such that

$$z_t + (1 - \lambda_t)p_t\xi_t - z_{t+1} \in p_t\mathcal{Q}_t \subseteq \mathcal{Q}_t.$$

Put $p_{t+1} := \lambda_t p_t$. Then $z_{t+1} \in p_{t+1}\mathcal{Z}_{t+1}^{\text{bd}}$, which concludes the inductive step. Now define the mixed stopping time $\chi = (\chi_t)$ by

$$\chi_t := \begin{cases} p_T & \text{if } t = T, \\ p_t - p_{t+1} & \text{if } t = 0, \dots, T-1. \end{cases}$$

We also put $z_{T+1} := 0$. We have constructed $z \in \Phi$ and $\chi \in \mathcal{X}$ such that $z_0 = x$ and

$$z_t + \chi_t\xi_t - z_{t+1} \in \mathcal{Q}_t \text{ for each } t = 0, \dots, T.$$

Finally, we construct $y \in \Phi$ such that $(y, \chi) \in \Phi^{\text{bg}}(\xi)$ and $y_0 = x$. By the definition of the deferred solvency cones \mathcal{Q}_t , for each $t = 0, \dots, T$ there is a liquidation strategy $v_{t+1}^t, \dots, v_{T+1}^t$ starting from $z_t + \chi_t\xi_t - z_{t+1}$ at time t . We put

$$y_t := z_t + \sum_{r=0}^{t-1} \sum_{s=t}^T (v_s^r - v_{s+1}^r),$$

which means that

$$\begin{aligned} y_t + \chi_t\xi_t - y_{t+1} &= z_t + \sum_{r=0}^{t-1} \sum_{s=t}^T (v_s^r - v_{s+1}^r) + \chi_t\xi_t - z_{t+1} - \sum_{r=0}^t \sum_{s=t+1}^T (v_s^r - v_{s+1}^r) \\ &= z_t + \chi_t\xi_t - z_{t+1} - v_{t+1}^t + \sum_{r=0}^{t-1} (v_t^r - v_{t+1}^r) \in \mathcal{K}_t \end{aligned}$$

for each $t = 0, \dots, T$, with $y_0 = z_0 = x$, completing the proof of (5.5).

Next, it follows from (5.5) that

$$\begin{aligned} \pi_j^{\text{bg}}(\xi) &= \sup \{ -x \in \mathbb{R} \mid \exists (y, \chi) \in \Phi^{\text{bg}}(\xi) : xe^j = y_0 \} \\ &= \sup \{ -x \in \mathbb{R} \mid xe^j \in \mathcal{Z}_0^{\text{bd}} \}. \end{aligned}$$

By Proposition 5.1, $\mathcal{Z}_0^{\text{bd}}$ is polyhedral, hence closed. As a result, the set $\{ -x \in \mathbb{R} \mid xe^j \in \mathcal{Z}_0^{\text{bd}} \}$ is also closed. It is non-empty and bounded above because $xe^j \in \mathcal{Z}_0^{\text{bd}}$ for any $x \in \mathbb{R}$ large enough, and $xe^j \notin \mathcal{Z}_0^{\text{bd}}$ for any $x \in \mathbb{R}$ small enough.

This means that the supremum is attained. It follows that $-\pi_j^{\text{bg}}(\xi)e^j \in \mathcal{Z}_0^{\text{bd}}$, so a strategy $(y, \chi) \in \Phi^{\text{bg}}(\xi)$ can be constructed such that $\pi_j^{\text{bg}}(\xi)e^j = -y_0$. \square

Proof of Proposition 5.2. For any $y \in \Phi$ such that $(y, \chi) \in \Phi^{\text{bg}}(\xi)$, put

$$z_t := y_t - \sum_{s=0}^{t-1} \chi_s \xi_s$$

for each $t = 0, \dots, T$, and $z_{T+1} := 0$. Then

$$\begin{aligned} z_t - z_{t+1} &= y_t - \sum_{s=0}^{t-1} \chi_s \xi_s - y_{t+1} + \sum_{s=0}^t \chi_s \xi_s \\ &= y_t + \chi_t \xi_t - y_{t+1} \in \mathcal{K}_t \end{aligned}$$

for each $t = 0, \dots, T-1$, and

$$\begin{aligned} z_T + \xi_\chi &= y_T - \sum_{s=0}^{T-1} \chi_s \xi_s + \sum_{s=0}^T \chi_s \xi_s \\ &= y_T + \chi_T \xi_T - y_{T+1} \in \mathcal{K}_T \end{aligned}$$

since $y_{T+1} = 0$, so $z \in \Psi^{\text{a}}(-\xi_\chi)$ with $z_0 = y_0$.

Conversely, for any $z \in \Psi^{\text{a}}(-\xi_\chi)$ we put

$$y_t := z_t + \sum_{s=0}^{t-1} \chi_s \xi_s$$

for each $t = 0, \dots, T$, and $y_{T+1} := 0$. Then

$$\begin{aligned} y_t + \chi_t \xi_t - y_{t+1} &= z_t + \sum_{s=0}^{t-1} \chi_s \xi_s + \chi_t \xi_t - z_{t+1} - \sum_{s=0}^t \chi_s \xi_s \\ &= z_t - z_{t+1} \in \mathcal{K}_t \end{aligned}$$

for each $t = 0, \dots, T-1$, and

$$\begin{aligned} y_T + \chi_T \xi_T - y_{T+1} &= z_T + \sum_{s=0}^{T-1} \chi_s \xi_s + \chi_T \xi_T \\ &= z_T + \sum_{s=0}^T \chi_s \xi_s \in \mathcal{K}_T. \end{aligned}$$

It follows that $(y, \chi) \in \Phi^{\text{bg}}(\xi)$ and $y_0 = z_0$, completing the proof. \square

Proof of Theorem 5.2. Theorem 5.1 gives

$$\pi_j^{\text{bg}}(\xi) = \max\{-x \in \mathbb{R} \mid xe^j \in \mathcal{Z}_0^{\text{bd}}\}.$$

The maximum is attained, so $-\pi_j^{\text{bg}}(\xi)e^j \in \mathcal{Z}_0^{\text{bd}}$. The strategy $(\hat{y}, \hat{\chi}) \in \Phi^{\text{bg}}(\xi)$ constructed by the method in the proof of Theorem 5.1 from the initial portfolio

34 *Alet Roux and Tomasz Zastawniak*

$\hat{y}_0 = -\pi_j^{\text{bg}}(\xi)e^j$ therefore realises the supremum in (3.5). We write this supremum as a maximum,

$$\begin{aligned}\pi_j^{\text{bg}}(\xi) &= \max\{-x \in \mathbb{R} \mid \exists(y, \chi) \in \Phi^{\text{bg}}(\xi) : xe^j = y_0\} \\ &= \max_{\chi \in \mathcal{X}} \left[\max\{-x \in \mathbb{R} \mid \exists y \in \Phi : (y, \chi) \in \Phi^{\text{bg}}(\xi), xe^j = y_0\} \right],\end{aligned}$$

and apply Proposition 5.2, which gives

$$\begin{aligned}\pi_j^{\text{bg}}(\xi) &= \max_{\chi \in \mathcal{X}} \left[\max\{-x \in \mathbb{R} \mid \exists y \in \Psi^{\text{a}}(-\xi_\chi) : xe^j = y_0, \} \right] \\ &= \max_{\chi \in \mathcal{X}} \left[-p_j^{\text{a}}(-\xi_\chi) \right],\end{aligned}$$

where $p_j^{\text{a}}(-\xi_\chi)$ is the ask (seller's) price in currency j of a European option with expiry time T and payoff $-\xi_\chi \in \mathcal{L}_T$ as defined in Appendix A.5. We can now apply Lemma A.3 to write

$$-p_j^{\text{a}}(-\xi_\chi) = - \max_{(\mathbb{Q}, S) \in \bar{\mathcal{P}}_j} \mathbb{E}_{\mathbb{Q}}(-\xi_\chi \cdot S_T) = \min_{(\mathbb{Q}, S) \in \bar{\mathcal{P}}_j} \mathbb{E}_{\mathbb{Q}}(\xi_\chi \cdot S_T).$$

For any $(\mathbb{Q}, S) \in \bar{\mathcal{P}}_j$, since S is a martingale under \mathbb{Q} , we have

$$\mathbb{E}_{\mathbb{Q}}(\xi_\chi \cdot S_T) = \mathbb{E}_{\mathbb{Q}}\left(\sum_{t=0}^T \chi_t \xi_t \cdot S_T\right) = \mathbb{E}_{\mathbb{Q}}\left(\sum_{t=0}^T \chi_t \xi_t \cdot S_t\right) = \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_\chi).$$

This means that

$$\pi_j^{\text{bg}}(\xi) = \max_{\chi \in \mathcal{X}} \min_{(\mathbb{Q}, S) \in \bar{\mathcal{P}}_j} \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_\chi),$$

proving (5.6). We know that $(\hat{y}, \hat{\chi})$ realises the supremum in (3.5), and therefore the above maxima over $\chi \in \mathcal{X}$ are attained at $\hat{\chi}$. A pair $(\hat{\mathbb{Q}}, \hat{S}) \in \bar{\mathcal{P}}_j$ such that

$$\pi_j^{\text{bg}}(\xi) = -p_j^{\text{a}}(-\xi_{\hat{\chi}}) = \mathbb{E}_{\hat{\mathbb{Q}}}((\xi \cdot \hat{S})_{\hat{\chi}}) = \min_{(\mathbb{Q}, S) \in \bar{\mathcal{P}}_j} \mathbb{E}_{\mathbb{Q}}((\xi \cdot S)_{\hat{\chi}})$$

can be constructed by the method of Roux & Zastawniak (2011 Proposition 5.3) for the European option with payoff $-\xi_{\hat{\chi}}$, completing the proof. \square

A.5. European options

We recall a result for European options in the market model with d assets under transaction costs. This is needed in the proof of the dual representation for the bid price of an American option.

A European option obliges the seller (writer) to deliver a portfolio $\zeta \in \mathcal{L}_T$ at time T . The set of strategies superhedging the seller's position is given as

$$\Psi^{\text{a}}(\zeta) := \{y \in \Phi \mid y_t - y_{t+1} \in \mathcal{K}_t \text{ for } t = 0, \dots, T-1, y_T - \zeta \in \mathcal{K}_T\}$$

and the *ask price (seller's price)* of such an option in currency $j = 1, \dots, d$ is

$$p_j^{\text{a}}(\zeta) := \inf \{x \in \mathbb{R} \mid \exists y \in \Psi^{\text{a}}(\zeta) : xe^j = y_0\}.$$

The following result can be found in Roux & Zastawniak (2011 Section 4.3.1).

Lemma A.3. *The ask price in currency $j = 1, \dots, d$ of a European option ζ can be represented as*

$$p_j^a(\zeta) = \max_{(\mathbb{Q}, S) \in \bar{\mathcal{P}}_j} \mathbb{E}_{\mathbb{Q}}(\zeta \cdot S_T).$$

Moreover, a pair $(\hat{\mathbb{Q}}, \hat{S}) \in \bar{\mathcal{P}}_j$ such that $p_j^a(\zeta) = \mathbb{E}_{\hat{\mathbb{Q}}}(\zeta \cdot \hat{S}_T)$ can be constructed algorithmically.

References

- B. Bouchard & J.-F. Chassagneux (2009) Representation of continuous linear forms on the set of l adlag processes and the hedging of American claims under proportional costs, *Electronic Journal of Probability* **14**, 612–632.
- B. Bouchard & E. Temam (2005) On the hedging of American options in discrete time markets with proportional transaction costs, *Electronic Journal of Probability* **10**, 746–760.
- P. Chalasani & S. Jha (2001) Randomized stopping times and American option pricing with transaction costs, *Mathematical Finance* **11**(1), 33–77.
- D. De Valli ere, E. Denis & Y. Kabanov (2009) Hedging of American options under transaction costs, *Finance and Stochastics* **13**, 105–119.
- M. Franz (2009) *Convex – a Maple package for convex geometry*.
URL: <http://www.math.uwo.ca/~mfranz/convex/>
- Y. M. Kabanov (1999) Hedging and liquidation under transaction costs in currency markets, *Finance and Stochastics* **3**, 237–248.
- Y. M. Kabanov & C. Stricker (2001) The Harrison-Pliska arbitrage pricing theorem under transaction costs, *Journal of Mathematical Economics* **35**, 185–196.
- R. Korn & S. M uller (2009) The decoupling approach to binomial pricing of multi-asset options, *Journal of Computational Finance* **12**, 1–30.
- A. L ohne & B. Rudloff (2014) An algorithm for calculating the set of superhedging portfolios in markets with transaction costs, *International Journal of Theoretical and Applied Finance* **17** (2), 1450012-1–1450012-33.
- R. T. Rockafellar (1996) *Convex Analysis*, Princeton Landmarks in Mathematics and Physics, Princeton University Press.
- A. Roux & T. Zastawniak (2009) American options under proportional transaction costs: Pricing, hedging and stopping algorithms for long and short positions, *Acta Applicandae Mathematicae* **106**, 199–228.
- A. Roux & T. Zastawniak (2011) American and Bermudan options in currency markets under proportional transaction costs. arXiv:1108.1910v3.
- W. Schachermayer (2004) The fundamental theorem of asset pricing under proportional transaction costs in finite discrete time, *Mathematical Finance* **14** (1), 19–48.
- C.-Y. Tien (2012) *Mixed Stopping Times and American Options under Transaction Costs*, PhD Thesis, University of York.
- N. Zhang, A. Roux & T. Zastawniak (2012) Parallel binomial American option pricing under proportional transaction costs, *Applied Mathematics* **3**, 1795–1810.