

This is a repository copy of *On interpreting Patterson - : Sullivan measures of geometrically finite groups as Hausdorff and packing measures.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/91009/>

Version: Submitted Version

Article:

Simmons, David orcid.org/0000-0002-9136-6635 (2016) On interpreting Patterson - : Sullivan measures of geometrically finite groups as Hausdorff and packing measures. *Ergodic Theory and Dynamical Systems*. pp. 2675-2686. ISSN 1469-4417

<https://doi.org/10.1017/etds.2015.27>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

ON INTERPRETING PATTERSON–SULLIVAN MEASURES OF GEOMETRICALLY FINITE GROUPS AS HAUSDORFF AND PACKING MEASURES

DAVID SIMMONS

ABSTRACT. We provide a new proof of a theorem whose proof was sketched by Sullivan ('82), namely that if the Poincaré exponent of a geometrically finite Kleinian group G is strictly between its minimal and maximal cusp ranks, then the Patterson–Sullivan measure of G is not proportional to the Hausdorff or packing measure of any gauge function. This disproves a conjecture of Stratmann ('97, '06).

1. INTRODUCTION

Let G be a nonelementary geometrically finite Kleinian group with at least one cusp.¹ Let k_{\min} and k_{\max} denote the smallest and largest cusp ranks, respectively, and let δ denote the Poincaré exponent of G . Let Λ be the limit set of G , and let μ be the Patterson–Sullivan measure of G , i.e. the unique probability measure on Λ which is δ -conformal with respect to G . It is known [15, Theorem 2] that

$$\begin{aligned}\mu \propto \mathcal{H}^\delta \upharpoonright \Lambda &\Leftrightarrow \delta \geq k_{\max} \text{ and} \\ \mu \propto \mathcal{P}^\delta \upharpoonright \Lambda &\Leftrightarrow \delta \leq k_{\min}.\end{aligned}$$

Here \mathcal{H}^δ and \mathcal{P}^δ denote the Hausdorff and packing measures, respectively, in dimension δ , and \propto is the proportionality symbol. The above conditions can be thought of as giving a “geometric interpretation” of the Patterson–Sullivan measure μ in the cases $\delta \geq k_{\max}$ and $\delta \leq k_{\min}$, since the measures $\mathcal{H}^\delta \upharpoonright \Lambda$ and $\mathcal{P}^\delta \upharpoonright \Lambda$ are defined using only the metric structure of Λ , without reference to the group G .

If $k_{\min} < \delta < k_{\max}$, then the above geometric interpretations fail, but it is natural to ask whether μ is proportional to $\mathcal{H}^\psi \upharpoonright \Lambda$ or $\mathcal{P}^\psi \upharpoonright \Lambda$ for some Hausdorff gauge function ψ (cf. §2.1). In fact, this question has a confusing history: Sullivan originally asked it in 1982 [13, p.71] and then sketched a proof of a negative answer later that year [14, second Corollary on p.235] (see also [15, Remark (2) on p.261]). Over a decade later, and without referencing Sullivan’s sketch, Stratmann stated that the question was open on two separate occasions [9, p.68], [10, p.235], even conjecturing that the answer is positive for the gauge function

$$(1.1) \quad \psi(r) = r^\delta \exp\left(\frac{k_{\max} - \delta}{2\delta - k_{\max}} \left(\log \log \frac{1}{r} + \log \log \log \log \frac{1}{r}\right)\right)^2.$$

(Cf. [2], where a similar result is proven for analytically finite but geometrically infinite groups.) In 2011 Ala-Mattila, a doctoral student of Tukia, noticed these inconsistencies in the literature in his dissertation [1, p.105], and stated that he was not able to follow Sullivan’s proof. (Nevertheless, his thesis answers a different but related question; namely, it provides a dynamics-independent construction of the Patterson–Sullivan measure, and thus a “geometric interpretation” of the Patterson–Sullivan measure in the sense described above.)

The author of the present paper confesses that he is also not able to follow Sullivan’s proof, so based on the above, it would seem that the question is still open. The purpose of this note is to prove using different methods that Sullivan’s answer is correct. Specifically, we prove the following theorem:

Key words and phrases. Patterson–Sullivan measures, geometrically finite Kleinian groups, Hausdorff and packing measures.

¹In this paper, a *Kleinian group* is a discrete subgroup of $\text{Isom}(\mathbb{B}^d)$ for any $d \geq 2$, where \mathbb{B}^d denotes d -dimensional hyperbolic space.

²Since Stratmann was working in \mathbb{B}^3 only, in his formula, $k_{\min} = 1$ and $k_{\max} = 2$.

Theorem 1.1. *Let G be a nonelementary geometrically finite Kleinian group with at least one cusp, let ψ be a Hausdorff gauge function, and let $\Psi : \mathbb{R} \rightarrow \mathbb{R}$ be the unique function satisfying*

$$(1.2) \quad \psi(r) = r^\delta \exp\left(\Psi\left(\log\frac{1}{r}\right)\right),$$

where δ is the Poincaré exponent. Assume that

$$(1.3) \quad \Psi \text{ is eventually differentiable and monotonic, and the limit } \lim_{t \rightarrow \infty} \Psi'(t) \text{ exists.}$$

Let k_{\min} and k_{\max} denote the smallest and largest cusp ranks, respectively. Then

(i) *If $\delta < k_{\max}$, then $\mathcal{H}^\psi(\mu) = 0$ or ∞ , according to whether the series*

$$(1.4) \quad \sum_{t=1}^{\infty} \exp\left(-\frac{2\delta - k_{\max}}{k_{\max} - \delta} \Psi(t)\right)$$

diverges or converges, respectively.

(ii) *If $\delta > k_{\min}$, then $\mathcal{P}^\psi(\mu) = 0$ or ∞ , according to whether the series*

$$(1.5) \quad \sum_{t=1}^{\infty} \exp\left(\frac{2\delta - k_{\min}}{\delta - k_{\min}} \Psi(t)\right)$$

converges or diverges, respectively.

In particular, if $k_{\min} < \delta < k_{\max}$ then neither $\mathcal{H}^\psi(\mu)$ nor $\mathcal{P}^\psi(\mu)$ is positive and finite for any Hausdorff gauge function ψ satisfying (1.3), so the Patterson–Sullivan measure cannot be interpreted geometrically as a Hausdorff or packing measure of such a gauge function. (See also Corollary 1.5 below.)

Example 1.2. For the value of ψ given in (1.1), we have

$$\Psi(t) = \frac{k_{\max} - \delta}{2\delta - k_{\max}} (\log(t) + \log \log \log(t)),$$

so the series (1.4) reduces to

$$\sum_{t=1}^{\infty} \frac{1}{t \log \log(t)},$$

which diverges. Therefore $\mathcal{H}^\psi(\mu) = 0$ for this ψ .

Remark 1.3. The relations in Theorem 1.1 can be a little disorienting at first – divergence of the series (1.4) guarantees $\mathcal{H}^\psi(\mu) = 0$, while divergence of the series (1.5) guarantees $\mathcal{P}^\psi(\mu) = \infty$. The unifying insight is that higher values of Ψ correspond to higher (decreasing more slowly as $r \searrow 0$) values of ψ , which in turn correspond to higher values of $\mathcal{H}^\psi(\mu)$ and $\mathcal{P}^\psi(\mu)$. Since the series (1.4) is decreasing with respect to Ψ while (1.5) is increasing with respect to Ψ , divergence of (1.4) indicates low (not close to $+\infty$) values of Ψ while divergence of (1.5) indicates high (not close to $-\infty$) values of Ψ .

Remark 1.4. Theorem 1.1 can be generalized to the setting of pinched Hadamard manifolds,³ under some additional assumptions regarding the group G . Indeed, the three main results used in the proof of Theorem 1.1, namely the Rogers–Taylor–Tricot density theorem, the Global Measure Formula, and Stratmann–Velani’s Khinchin-type Theorem for Geometrically Finite Groups, are generalized to this setting in [6, §8],⁴ [8, Théorème 7.2], and [5, Theorem 3], respectively, and the deduction of Theorem 1.1 from these theorems does not make use of constant curvature in any essential way. However, for simplicity of exposition we stick to the case of standard hyperbolic space.

Corollary 1.5. *Let G and ψ be as in Theorem 1.1.*

(i) *If $\delta < k_{\max}$, then $\mathcal{H}^\psi(\Lambda) = 0$ or ∞ .*

³In this setting, the cusp rank k_ξ of a parabolic point ξ is interpreted to be the number $2\delta(G_\xi)$, where G_ξ is the stabilizer of ξ in G and $\delta(G_\xi)$ is the Poincaré exponent of G_ξ ; cf. [8, Théorème 3.2].

⁴The “arbitrary metric space” of [6, §8] should be interpreted to be the Gromov boundary of the pinched Hadamard manifold in question, endowed with a visual metric.

(ii) If $\delta > k_{\min}$, then $\mathcal{P}^\psi(\Lambda) = 0$ or ∞ .

Proof. Without loss of generality we may assume that

$$(1.6) \quad \lim_{t \rightarrow \infty} \Psi'(t) = 0.$$

Indeed, since the Hausdorff and packing dimensions of Λ are both equal to δ [15, 11], if (1.6) fails then the corollary follows from comparing the measures \mathcal{H}^ψ and \mathcal{P}^ψ with the measures \mathcal{H}^s and \mathcal{P}^s , where s lies between δ and $\delta + \lim_{t \rightarrow \infty} \Psi'(t)$.

Suppose $\delta < k_{\max}$ but $0 < \mathcal{H}^\psi(\Lambda) < \infty$. Then (1.6) implies that the measure $\nu = \mathcal{H}^\psi \upharpoonright \Lambda$ is δ -conformal with respect to G (Lemma 4.3). Next, the uniqueness of δ -conformal measures for geometrically finite groups (e.g. [15, Theorem 1]) implies that $\nu = \alpha\mu$ for some $\alpha > 0$. But then $\mathcal{H}^\psi(\mu) = \mathcal{H}^\psi(\nu) = \mathcal{H}^\psi(\Lambda) \in (0, \infty)$, contradicting Theorem 1.1. This demonstrates (i), and (ii) is proven similarly. \square

Note however that this argument does not tell us that the value of $\mathcal{H}^\psi(\Lambda)$ is determined by the convergence or divergence of the series (1.4); *a priori* it may be the case that $\mathcal{H}^\psi(\Lambda) = \infty$ while $\mathcal{H}^\psi(\mu) = 0$. An interesting question is whether or not this is possible.

Convention. The symbols \lesssim , \gtrsim , and \asymp will denote coarse asymptotics; a subscript of $+$ indicates that the asymptotic is additive, and a subscript of \times indicates that it is multiplicative. For example, $A \lesssim_{\times} B$ means that there exists a constant $C > 0$ (the *implied constant*) such that $A \leq CB$. It is understood that the implied constant C is only allowed to depend on certain “universal” parameters, to be understood from context.

Acknowledgements. The author thanks Tushar Das for calling his attention to Ala-Mattila’s thesis and to the fact that the question addressed in this paper is still considered open, as well as for helpful comments on an earlier version of this paper. The author also thanks the referee for valuable comments.

2. PRELIMINARIES

Our theorem is essentially a combination of three known theorems: the Rogers–Taylor–Tricot density theorem, the Global Measure Formula, and Stratmann–Velani’s Khinchin-type Theorem for Geometrically Finite Groups. In this section we recall these theorems and the definitions used in them.

2.1. The Rogers–Taylor–Tricot density theorem. We first recall the basics of generalized Hausdorff and packing measures. Let $\psi : [0, \infty) \rightarrow [0, \infty)$ be a *Hausdorff gauge function*, i.e. a positive increasing continuous function satisfying $\psi(0) = 0$. The ψ -dimensional Hausdorff measure of a set $A \subseteq \mathbb{R}^d$ is

$$\mathcal{H}^\psi(A) = \liminf_{\varepsilon \searrow 0} \left\{ \sum_{i=1}^{\infty} \psi(\text{diam}(U_i)) : (U_i)_{i=1}^{\infty} \text{ is a countable cover of } A \text{ with } \text{diam}(U_i) \leq \varepsilon \ \forall i \right\}$$

and the ψ -dimensional packing measure of A is defined by the formulas

$$\tilde{\mathcal{P}}^\psi(A) = \limsup_{\varepsilon \searrow 0} \left\{ \sum_{j=1}^{\infty} \psi(\text{diam}(B_j)) : (B_j)_{j=1}^{\infty} \text{ is a countable disjoint collection of balls with centers in } A \text{ and with } \text{diam}(B_j) \leq \varepsilon \ \forall j \right\}$$

$$\mathcal{P}^\psi(A) = \inf \left\{ \sum_{i=1}^{\infty} \tilde{\mathcal{P}}^\psi(A_i) : A \subseteq \bigcup_{i=1}^{\infty} A_i \right\}.$$

A special case is when $\psi(r) = r^s$ for some $s > 0$, in which case the shorthands $\mathcal{H}^\psi = \mathcal{H}^s$ and $\mathcal{P}^\psi = \mathcal{P}^s$ are used.

If μ is a Borel measure on \mathbb{R}^d , then we let

$$\begin{aligned} \mathcal{H}^\psi(\mu) &= \inf \{ \mathcal{H}^\psi(A) : \mu(\mathbb{R}^d \setminus A) = 0 \}, \\ \mathcal{P}^\psi(\mu) &= \inf \{ \mathcal{P}^\psi(A) : \mu(\mathbb{R}^d \setminus A) = 0 \}. \end{aligned}$$

Moreover, for each point $x \in \mathbb{R}^d$ let

$$\begin{aligned}\overline{D}_\mu^\psi(x) &= \limsup_{r \searrow 0} \frac{\mu(B(x, r))}{\psi(r)}, \\ \underline{D}_\mu^\psi(x) &= \liminf_{r \searrow 0} \frac{\mu(B(x, r))}{\psi(r)}.\end{aligned}$$

Theorem 2.1 (Rogers–Taylor–Tricot density theorem). *Fix $d \in \mathbb{N}$, let μ be a Borel measure on \mathbb{R}^d , and let ψ be a Hausdorff gauge function. Then for every Borel set $A \subseteq \mathbb{R}^d$,*

$$(2.1) \quad \mu(A) \inf_{x \in A} \frac{1}{\overline{D}_\mu^\psi(x)} \lesssim_\times \mathcal{H}^\psi(A) \lesssim_\times \mu(\mathbb{R}^d) \sup_{x \in A} \frac{1}{\overline{D}_\mu^\psi(x)}$$

$$(2.2) \quad \mu(A) \inf_{x \in A} \frac{1}{\underline{D}_\mu^\psi(x)} \lesssim_\times \mathcal{P}^\psi(A) \lesssim_\times \mu(\mathbb{R}^d) \sup_{x \in A} \frac{1}{\underline{D}_\mu^\psi(x)}.$$

Formula (2.1) was proven by Rogers and Taylor in 1959 [7], while formula (2.2) was proven by Taylor and Tricot in 1985 [16, Theorem 2.1]. A short proof of (2.1) can be found in [3, Proposition 3.4 of Chapter 4]. A generalization to arbitrary metric spaces can be found in [6, §8].

Corollary 2.2. *With d , μ , and ψ as above,*

$$(2.3) \quad \mu(\mathbb{R}^d) \operatorname{ess\,inf}_{x \in \mathbb{R}^d} \frac{1}{\overline{D}_\mu^\psi(x)} \lesssim_\times \mathcal{H}^\psi(\mu) \lesssim_\times \mu(\mathbb{R}^d) \operatorname{ess\,sup}_{x \in \mathbb{R}^d} \frac{1}{\overline{D}_\mu^\psi(x)}$$

$$(2.4) \quad \mu(\mathbb{R}^d) \operatorname{ess\,inf}_{x \in \mathbb{R}^d} \frac{1}{\underline{D}_\mu^\psi(x)} \lesssim_\times \mathcal{P}^\psi(\mu) \lesssim_\times \mu(\mathbb{R}^d) \operatorname{ess\,sup}_{x \in \mathbb{R}^d} \frac{1}{\underline{D}_\mu^\psi(x)}.$$

Proof. Let

$$A_0 = \left\{ x \in \mathbb{R}^d : \operatorname{ess\,inf} \overline{D}_\mu^\psi \leq \overline{D}_\mu^\psi(x) \leq \operatorname{ess\,sup} \overline{D}_\mu^\psi \right\}.$$

Since $\mu(\mathbb{R}^d \setminus A_0) = 0$, we have

$$\mathcal{H}^\psi(\mu) \leq \mathcal{H}^\psi(A_0) \lesssim_\times \mu(\mathbb{R}^d) \operatorname{ess\,sup}_{x \in \mathbb{R}^d} \frac{1}{\overline{D}_\mu^\psi(x)}.$$

On the other hand, given any set $B \subseteq \mathbb{R}^d$ such that $\mu(\mathbb{R}^d \setminus B) = 0$, applying (2.1) with $A = A_0 \cap B$ gives

$$\mathcal{H}^\psi(B) \gtrsim_\times \mu(A_0 \cap B) \operatorname{ess\,inf}_{x \in \mathbb{R}^d} \frac{1}{\overline{D}_\mu^\psi(x)} = \mu(\mathbb{R}^d) \operatorname{ess\,inf}_{x \in \mathbb{R}^d} \frac{1}{\overline{D}_\mu^\psi(x)},$$

and taking the infimum over all such B s demonstrates (2.3). The proof of (2.4) proceeds in the same manner. \square

2.2. Geometrically finite groups. Fix $d \geq 2$, let \mathbb{B}^d denote the d -dimensional Poincaré disk, and let G be a nonelementary geometrically finite Kleinian group acting on \mathbb{B}^d with at least one cusp. Recall that this means that there exists a disjoint G -invariant collection of horoballs \mathcal{H} with the property that the quotient $(\mathcal{C}_G \setminus \bigcup(\mathcal{H}))/G$ is compact, where \mathcal{C}_G is the convex hull of the limit set of G (e.g. [4, Definition (GF1)]). The elements of \mathcal{H} are centered at the parabolic fixed points of G . For each parabolic point ξ , let H_ξ denote the unique element of \mathcal{H} centered at ξ , and let k_ξ denote the rank of ξ .

Let δ denote the *Poincaré exponent* of G , i.e.

$$\delta = \inf \left\{ s \geq 0 : \sum_{g \in G} e^{-sd(0, g(0))} < \infty \right\},$$

where d denotes the hyperbolic distance on \mathbb{B}^d . Let Λ denote the *limit set* of G , i.e. the set

$$\Lambda = \left\{ \xi \in \partial \mathbb{B}^d : \xi = \lim_{n \rightarrow \infty} g_n(0) \text{ for some sequence } (g_n)_1^\infty \text{ in } G \right\}.$$

Finally, let μ denote the *Patterson–Sullivan measure* of G , i.e. the unique Borel probability measure on Λ obeying the transformation rule

$$\mu(g(A)) = \int_A |g'(\xi)|^\delta d\mu(\xi) \quad \forall A \subseteq \partial\mathbb{B}^d \quad \forall g \in G.$$

2.2.1. *The global measure formula.* Let the functions $k, b : \mathbb{B}^d \rightarrow [0, \infty)$ be defined as follows:

- If $x \in \mathbb{B}^d \setminus \bigcup(\mathcal{H})$, then $k(x) = b(x) = 0$.
- If $x \in H_\xi$ for some $H_\xi \in \mathcal{H}$, then $k(x) = k_\xi$ and $b(x) = d(x, \partial H_\xi)$.

Given a point $\eta \in \Lambda$, let η_t denote the unique point on the geodesic ray connecting 0 and η whose hyperbolic distance from 0 is equal to t .

Theorem 2.3 (Global measure formula, [12, Theorem 2]). *For any $\eta \in \Lambda$ and for any $t > 0$,*

$$\mu(B(\eta, e^{-t})) \asymp_\times e^{-\delta t} e^{b(\eta_t)(k(\eta_t) - \delta)}.$$

2.2.2. *Khinchin-type theorem for geometrically finite groups.* For each $\xi \in \partial\mathbb{B}^d$ and $r > 0$, let $H(\xi, r)$ be the unique horoball centered at ξ with Euclidean radius r , i.e.

$$H(\xi, r) := B((1-r)\xi, r).$$

If ξ is a parabolic point, then we let r_ξ denote the unique value such that $H(\xi, r_\xi) \in \mathcal{H}$, so that $H_\xi = H(\xi, r_\xi)$.

Let $\phi : (0, \infty) \rightarrow (0, \infty)$ be a *Khinchin function*, i.e. a positive monotonic doubling function. Here, by saying that ϕ is *doubling*, we mean that there exist constants $C_1, C_2 > 1$ such that

$$\frac{1}{C_1} \leq \frac{y}{x} \leq C_1 \quad \Rightarrow \quad \frac{1}{C_2} \leq \frac{\phi(y)}{\phi(x)} \leq C_2 \quad \forall x, y > 0.$$

(For ϕ to be doubling, it is sufficient but not necessary that the function $\log \circ \phi \circ \exp$ is uniformly continuous.) Now fix a parabolic point p , and consider the set

$$\Omega_p(\phi) := \{\eta \in \Lambda : \|\xi - \eta\| \leq \phi(r_\xi)r_\xi \text{ for infinitely many } \xi \in G(p)\}.$$

Theorem 2.4 (Khinchin-type theorem for geometrically finite Kleinian groups, [12, Theorem 4]). $\mu(\Omega_p(\phi)) = 0$ or 1 according to whether the series

$$\sum_{n=1}^{\infty} \phi(\lambda^n)^{2\delta - k_p}$$

converges or diverges, respectively. Here $\lambda \in (0, 1)$ is a constant depending only on G .

3. PROOF OF THEOREM 1.1

As in Theorem 1.1, let G be a nonelementary geometrically finite Kleinian group with at least one cusp, let ψ be a Hausdorff gauge function, let Ψ be given by (1.2), and assume that (1.3) holds. As in the proof of Corollary 1.5, we may without loss of generality assume that

$$(3.1) \quad \lim_{t \rightarrow \infty} \Psi'(t) = 0.$$

Let P be a complete set of inequivalent parabolic points of G , and let

$$P_{>\delta} = \{p \in P : k_p > \delta\}, \quad P_{<\delta} = \{p \in P : k_p < \delta\}.$$

Lemma 3.1. *For each $p \in P_{<\delta} \cup P_{>\delta}$ and $\alpha > 0$ let $\psi_p, \theta_p, \phi_{p,\alpha} : (0, \infty) \rightarrow (0, \infty)$ be defined by*

$$\psi_p(r) = \exp\left(-\frac{\Psi(\log(1/r))}{k_p - \delta}\right), \quad \theta_p(r) = \frac{r}{\psi_p(r)}, \quad \phi_{p,\alpha}(r) = \frac{\theta_p^{-1}(r/\alpha)}{r}.$$

(i) *If $\delta < k_{\max}$ and $\lim_{t \rightarrow \infty} \Psi(t) = +\infty$, then for μ -a.e. $\eta \in \Lambda$,*

$$\overline{D}_\mu^\psi(\eta) \asymp_\times \max_{p \in P_{>\delta}} \left(\sup\{\alpha > 0 : \eta \in \Omega_p(\phi_{p,\alpha})\} \right)^{k_p - \delta}.$$

(ii) If $\delta > k_{\min}$ and $\lim_{t \rightarrow \infty} \Psi(t) = -\infty$, then for μ -a.e. $\eta \in \Lambda$,

$$\underline{D}_\mu^\psi(\eta) \asymp_\times \min_{p \in P_{<\delta}} \left(\sup\{\alpha > 0 : \eta \in \Omega_p(\phi_{p,\alpha})\} \right)^{k_p - \delta}.$$

Remark 3.2. From the assumptions (1.3) and (3.1), we can see that

- (i) ψ_p is monotonic in a neighborhood of 0 (since Ψ is eventually monotonic),
- (ii) θ_p is increasing in a neighborhood of 0 (since $\lim_{t \rightarrow -\infty} [\log \circ \theta_p \circ \exp]'(t) = 1$),
- (iii) $\phi_{p,\alpha}$ is monotonic in a neighborhood of 0 (since $\phi_{p,\alpha}(r) = \frac{1}{\alpha} \psi_p(\theta_p^{-1}(r/\alpha))$), and
- (iv) $\phi_{p,\alpha}$ is doubling

$$\left(\text{since } \lim_{t \rightarrow -\infty} [\log \circ \phi_{p,\alpha} \circ \exp]'(t) = \frac{\lim_{t \rightarrow -\infty} [\log \circ \psi_p \circ \exp]'(t)}{\lim_{t \rightarrow -\infty} [\log \circ \theta_p \circ \exp]'(t)} = \frac{0}{1} = 0 \right).$$

In particular, $\phi_{p,\alpha}$ is a Khinchin function.

Proof of Lemma 3.1. We prove only (ii); the proof of (i) is similar but easier. By Theorem 2.3,

$$\underline{D}_\mu^\psi(\eta) = \liminf_{t \rightarrow \infty} \frac{\mu(B(\eta, e^{-t}))}{\psi(e^{-t})} \asymp_\times \liminf_{t \rightarrow \infty} \frac{e^{-\delta t} e^{b(\eta_t)(k(\eta_t) - \delta)}}{e^{-\delta t} e^{\Psi(t)}} = \exp \liminf_{t \rightarrow \infty} [b(\eta_t)(k(\eta_t) - \delta) - \Psi(t)].$$

Since by assumption $\lim_{t \rightarrow \infty} \Psi(t) = -\infty$, values of t for which $b(\eta_t)(k(\eta_t) - \delta) \geq 0$ will not affect the lim-inf. On the other hand, since $\delta > k_{\min}$, for μ -a.e. $\eta \in \Lambda$ we have $b(\eta_t)(k(\eta_t) - \delta) < 0$ for a sequence of t tending to infinity. Thus

$$(3.2) \quad \log \underline{D}_\mu^\psi(\eta) \asymp_+ \liminf_{\substack{t \rightarrow \infty \\ b(\eta_t)(k(\eta_t) - \delta) < 0}} [b(\eta_t)(k(\eta_t) - \delta) - \Psi(t)] = \liminf_{\xi \in G(P_{<\delta})} \inf_{\substack{t > 0 \\ \eta_t \in H_\xi}} [b(\eta_t)(k_\xi - \delta) - \Psi(t)].$$

Fix $\xi \in G(P_{<\delta})$. To estimate the infimum on the right hand side, we use the following estimate for $b(\eta_t)$ which is valid for $\eta_t \in H_\xi$ (cf. Lemma 4.2 below):

$$(3.3) \quad b(\eta_t) \asymp_+ f_\xi(t) := \min \left(t - \log \frac{1}{r_\xi}, 2 \log \frac{1}{\|\xi - \eta\|} - \log \frac{1}{r_\xi} - t \right).$$

Let t_ξ be the unique point at which f_ξ is not differentiable, i.e. $t_\xi = \log \frac{1}{\|\xi - \eta\|}$. Then $f'_\xi(t) = 1$ for $t < t_\xi$ and $f'_\xi(t) = -1$ for $t > t_\xi$. Letting

$$h_\xi(t) = f_\xi(t)(k_\xi - \delta) - \Psi(t),$$

the assumption (3.1) guarantees that $h'_\xi(t) < 0$ for $t < t_\xi$ and $h'_\xi(t) > 0$ for $t > t_\xi$, assuming t is sufficiently large. It follows that for all but finitely many $\xi \in G(P_{<\delta})$,

$$\inf_{\substack{t > 0 \\ \eta_t \in H_\xi}} h_\xi(t) = \inf_{t > 0} h_\xi(t) = h_\xi(t_\xi) = \left[\left(t_\xi - \log \frac{1}{r_\xi} \right) (k_\xi - \delta) - \Psi(t_\xi) \right]$$

and thus

$$\log \underline{D}_\mu^\psi(\eta) \asymp_+ \liminf_{\xi \in G(P_{<\delta})} \left[\left(\log \frac{1}{\|\xi - \eta\|} - \log \frac{1}{r_\xi} \right) (k_\xi - \delta) - \Psi \left(\log \frac{1}{\|\xi - \eta\|} \right) \right].$$

Exponentiating gives

$$\underline{D}_\mu^\psi(\eta) \asymp_\times \min_{p \in P_{<\delta}} \liminf_{\xi \in G(p)} \left(\frac{1/\|\xi - \eta\|}{1/r_\xi} \psi_p(\|\xi - \eta\|) \right)^{k_p - \delta} = \min_{p \in P_{<\delta}} \left(\limsup_{\xi \in G(p)} \frac{r_\xi}{\theta_p(\|\xi - \eta\|)} \right)^{k_p - \delta}.$$

The calculation

$$\begin{aligned} \limsup_{\xi \in G(p)} \frac{r_\xi}{\theta_p(\|\xi - \eta\|)} &= \sup \left\{ \alpha > 0 : \frac{r_\xi}{\theta_p(\|\xi - \eta\|)} \geq \alpha \text{ for infinitely many } \xi \in G(p) \right\} \\ &= \sup \left\{ \alpha > 0 : \|\xi - \eta\| \leq \theta_p^{-1}(r_\xi/\alpha) \text{ for infinitely many } \xi \in G(p) \right\} \\ &= \sup \left\{ \alpha > 0 : \eta \in \Omega_p(\phi_{p,\alpha}) \right\} \end{aligned}$$

finishes the proof. \square

Fix $p \in P_{<\delta} \cup P_{>\delta}$ and $\alpha > 0$, and let us determine whether the series

$$\Sigma_{p,\alpha} = \sum_{n=1}^{\infty} \phi_{p,\alpha}(\lambda^n)^{2\delta-k_p}$$

occurring in Theorem 2.4 converges or diverges. For convenience write $\Delta_p = 2\delta - k_p$. We have⁵

$$\begin{aligned} \Sigma_{p,\alpha} &\asymp_+ \int_{\clubsuit}^{\infty} \phi_{p,\alpha}(\lambda^t)^{\Delta_p} dt = \int_{\clubsuit}^{\infty} \frac{\theta_p^{-1}(\lambda^t/\alpha)^{\Delta_p}}{\lambda^{\Delta_p t}} dt \\ &\asymp_{\times} \int_{\clubsuit}^{\infty} \theta_p^{-1}(x^{-1/\Delta_p})^{\Delta_p} dx && \text{(letting } x = \alpha^{\Delta_p} \lambda^{-\Delta_p t}\text{)} \\ &\asymp_+ \int_0^{\clubsuit} \theta_p(x^{1/\Delta_p})^{-\Delta_p} dx && \text{(by Lemma 4.1)} \\ &= \int_0^{\clubsuit} \psi_p(x^{1/\Delta_p})^{\Delta_p} \frac{dx}{x} \\ &\asymp_{\times} \int_{\clubsuit}^{\infty} \psi_p(e^{-t})^{\Delta_p} dt && \text{(letting } t = -\log(x^{1/\Delta_p})\text{)} \\ &= \int_{\clubsuit}^{\infty} \exp\left(-\frac{2\delta - k_p}{k_p - \delta} \Psi(t)\right) dt \\ &\asymp_+ \Sigma_p := \sum_{t=1}^{\infty} \exp\left(-\frac{2\delta - k_p}{k_p - \delta} \Psi(t)\right). \end{aligned}$$

By Theorem 2.4, for μ -a.e. $\eta \in \Lambda$, for every $p \in P_{<\delta} \cup P_{>\delta}$, and for every rational $\alpha > 0$, we have $\eta \in \Omega_p(\phi_{p,\alpha})$ if and only if the series $\Sigma_{p,\alpha} \asymp_{+, \times} \Sigma_p$ diverges. Thus by Lemma 3.1, for μ -a.e. $\eta \in \Lambda$ we have

$$(3.4) \quad \overline{D}_{\mu}^{\psi}(\eta) \asymp_{\times} \max_{p \in P_{>\delta}} \left(\begin{cases} \infty & \Sigma_p = \infty \\ 0 & \Sigma_p < \infty \end{cases} \right)^{k_p - \delta} = \begin{cases} \infty & \exists p \in P_{>\delta} \ \Sigma_p = \infty \\ 0 & \forall p \in P_{>\delta} \ \Sigma_p < \infty \end{cases} = \begin{cases} \infty & \text{(1.4) diverges} \\ 0 & \text{(1.4) converges} \end{cases}$$

(if $\delta < k_{\max}$ and $\Psi \rightarrow +\infty$)

$$(3.5) \quad \underline{D}_{\mu}^{\psi}(\eta) \asymp_{\times} \min_{p \in P_{<\delta}} \left(\begin{cases} \infty & \Sigma_p = \infty \\ 0 & \Sigma_p < \infty \end{cases} \right)^{k_p - \delta} = \begin{cases} 0 & \exists p \in P_{<\delta} \ \Sigma_p = \infty \\ \infty & \forall p \in P_{<\delta} \ \Sigma_p < \infty \end{cases} = \begin{cases} 0 & \text{(1.5) diverges} \\ \infty & \text{(1.5) converges} \end{cases}$$

(if $\delta > k_{\min}$ and $\Psi \rightarrow -\infty$)

Combining with Corollary 2.2, we see that Theorem 1.1 holds if the hypothesis $\Psi \rightarrow +\infty$ is added to case (i) and the hypothesis $\Psi \rightarrow -\infty$ is added to case (ii). But these extra hypotheses can be added without loss of generality, since if they fail, Theorem 1.1 can be deduced by comparing the measures \mathcal{H}^{ψ} and \mathcal{P}^{ψ} to the measures \mathcal{H}^{δ} and \mathcal{P}^{δ} , respectively.

4. PROOFS OF AUXILIARY FACTS

The following lemmas are probably well-known to experts, so we have separated them from the main part of the proof for ease of exposition.

Lemma 4.1. *Let $f : (0, \infty) \rightarrow (0, \infty)$ be a decreasing homeomorphism. Then*

$$\int_0^{\infty} f(x) dx = \int_0^{\infty} f^{-1}(x) dx.$$

Proof. These integrals are respectively equal to the Lebesgue measures of the sets

$$\begin{aligned} S_f &= \{(x, y) \in (0, \infty)^2 : y < f(x)\} \\ S_{f^{-1}} &= \{(x, y) \in (0, \infty)^2 : y < f^{-1}(x)\}, \end{aligned}$$

⁵In the calculation below, an integration bound of \clubsuit means that the precise integration bound is irrelevant.

which are conjugate to each other via the measure-preserving isometry $(x, y) \mapsto (y, x)$. \square

Lemma 4.2. *Let $H_\xi = H(\xi, r_\xi)$ be a horoball not containing the origin, and fix $\eta \in \partial\mathbb{B}^d \setminus \{\xi\}$ and $t > 0$ such that $\eta_t \in H_\xi$. Then*

$$b(\eta_t) \asymp_+ \min \left(t - \log \frac{1}{r_\xi}, 2 \log \frac{1}{\|\xi - \eta\|} - \log \frac{1}{r_\xi} - t \right).$$

Proof. Let \mathcal{B}_ξ denote the Busemann function of ξ , i.e.

$$\mathcal{B}_\xi(y, z) = \lim_{x \rightarrow \xi} [d(x, y) - d(x, z)],$$

where d denotes the hyperbolic distance in \mathbb{B}^d . Then

$$(4.1) \quad b(\eta_t) = \mathcal{B}_\xi(0, \eta_t) - d(0, H_\xi) \asymp_+ \mathcal{B}_\xi(0, \eta_t) - \log \frac{1}{r_\xi}.$$

To estimate $\mathcal{B}_\xi(0, \eta_t)$, we use the well-known fact that the incircle radius of a (possibly ideal) hyperbolic triangle is uniformly bounded. Specifically, when we consider the triangle $\Delta = \Delta(0, \eta, \xi)$ and let $a \in [0, \eta]$, $b \in [0, \xi]$, $c \in [\xi, \eta]$ be the points of Δ tangent to the incircle, then $d(a, b) \asymp_+ d(a, c) \asymp_+ 0$. Here $[p, q]$ denotes the geodesic (or geodesic ray) connecting p and q . Write $s = d(0, a)$, and for each $u > 0$ let $[c, \eta]_u$ denote the unique point on the geodesic $[c, \eta]$ such that $d(c, [c, \eta]_u) = u$. Then

$$\begin{cases} d(\eta_t, \xi_t) \asymp_+ 0 & \text{if } t \leq s \\ d(\eta_t, [c, \eta]_{t-s}) \asymp_+ 0 & \text{if } t \geq s \end{cases}$$

and thus

$$(4.2) \quad \mathcal{B}_\xi(0, \eta_t) \begin{cases} \asymp_+ \mathcal{B}_\xi(0, \xi_t) = t & \text{if } t \leq s \\ \asymp_+ \mathcal{B}_\xi(0, [c, \eta]_{t-s}) \asymp_+ 2\langle \xi | \eta \rangle_0 - t & \text{if } t \geq s \end{cases}.$$

Here $\langle \xi | \eta \rangle_0$ denotes the *Gromov product*

$$\langle \xi | \eta \rangle_0 = \lim_{\substack{x \rightarrow \xi \\ y \rightarrow \eta}} \frac{1}{2} [d(0, x) + d(0, y) - d(x, y)].$$

Plugging $t = s$ into (4.2) gives $s \asymp_+ \langle \xi | \eta \rangle_0$, and thus

$$\mathcal{B}_\xi(0, \eta_t) \asymp_+ \min(t, 2\langle \xi | \eta \rangle_0 - t).$$

Combining with (4.1) together with the well-known asymptotic

$$\langle \xi | \eta \rangle_0 \asymp_+ \log \frac{1}{\|\xi - \eta\|}$$

completes the proof. \square

Lemma 4.3. *Let G be a Kleinian group, and let ψ be a Hausdorff gauge function satisfying (1.6). Then the measures $\nu_1 = \mathcal{H}^\psi \upharpoonright \Lambda$ and $\nu_2 = \mathcal{P}^\psi \upharpoonright \Lambda$ are δ -conformal with respect to G .*

Proof. Fix $\lambda > 0$, and note that (1.6) guarantees that

$$(4.3) \quad \lim_{r \searrow 0} \frac{\psi(\lambda r)}{\psi(r)} = \lambda^\delta.$$

Now let $A \subseteq \Lambda$ be a Borel set on which $g \in G$ is λ -Lipschitz continuous. If $(U_i)_1^\infty$ is a countable cover of A satisfying $\text{diam}(U_i) \leq \varepsilon \forall i$, then $(\tilde{U}_i := g(U_i \cap A))_1^\infty$ is a cover of $g(A)$ satisfying $\text{diam}(\tilde{U}_i) \leq \lambda \varepsilon \forall i$ and

$$\sum_{i=1}^{\infty} \psi(\text{diam}(\tilde{U}_i)) \leq \left(\sup_{r \leq \varepsilon} \frac{\psi(\lambda r)}{\psi(r)} \right) \sum_{i=1}^{\infty} \psi(\text{diam}(U_i)).$$

Taking the infimum, letting $\varepsilon \rightarrow 0$, and using (4.3) shows that $\nu_1(g(A)) \leq \lambda^\delta \nu_1(A)$. Similarly, if $(B_i = B(x_i, r_i))_1^\infty$ is a countable disjoint collection of balls with centers in $g(A)$ satisfying $\text{diam}(B_i) \leq \varepsilon \forall i$, then

$(\tilde{B}_i := B(g^{-1}(x_i), r_i/\lambda))_1^\infty$ is a disjoint⁶ collection of balls with centers in A satisfying $\text{diam}(\tilde{B}_i) \leq \varepsilon/\lambda \ \forall i$ and

$$\sum_{i=1}^\infty \psi(\text{diam}(B_i)) \leq \left(\sup_{r \leq \varepsilon/\lambda} \frac{\psi(\lambda r)}{\psi(r)} \right) \sum_{i=1}^\infty \psi(\text{diam}(\tilde{B}_i)).$$

Taking the supremum, letting $\varepsilon \rightarrow 0$, and using (4.3) shows that $\nu_2(g(A)) \leq \lambda^\delta \nu_2(A)$. So

$$\nu_i(g(A)) \leq \lambda^\delta \nu_i(A) \quad (i = 1, 2).$$

Setting $\lambda = \sup_A |g'|$ and using the geometric mean value theorem gives

$$(4.4) \quad \nu_i(g(A)) \leq \sup_A |g'|^\delta \nu_i(A).$$

Now let $B \subseteq \Lambda$ be an arbitrary Borel set, and fix $g \in G$. Fix $\varepsilon > 0$, and let \mathcal{A} be a partition of Λ such that $\sup_A |g'|/\inf_A |g'| \leq 1 + \varepsilon$ for all $A \in \mathcal{A}$. We have

$$\begin{aligned} \nu_i(g(B)) &= \sum_{A \in \mathcal{A}} \nu_i(g(B \cap A)) \leq \sum_{A \in \mathcal{A}} \sup_A |g'|^\delta \nu_i(B \cap A) && \text{(by (4.4))} \\ &\leq (1 + \varepsilon) \sum_{A \in \mathcal{A}} \inf_A |g'|^\delta \nu_i(B \cap A) \leq (1 + \varepsilon) \int_B |g'|^\delta d\nu_i. \end{aligned}$$

Letting ε tend to zero shows that $\nu_i(g(B)) \leq \int_B |g'|^\delta d\nu_i$. The reverse inequality is proved similarly. □

REFERENCES

1. V. Ala-Mattila, *Geometric characterizations for Patterson-Sullivan measures of geometrically finite Kleinian groups*, Ph.D. thesis, University of Helsinki, 2011.
2. C. J. Bishop and P. W. Jones, *The law of the iterated logarithm for Kleinian groups*, Lipa's legacy (New York, 1995), Contemp. Math., 211, Amer. Math. Soc., Providence, RI, 1997, pp. 17–50.
3. C. J. Bishop and Y. Peres, *Fractal sets in probability and analysis*, <http://www.math.washington.edu/~solomyak/TEACH/582/12/BishopPeres.pdf> preprint.
4. B. H. Bowditch, *Geometrical finiteness for hyperbolic groups*, J. Funct. Anal. **113** (1993), no. 2, 245–317.
5. S. D. Hersonsky and F. Paulin, *Counting orbit points in coverings of negatively curved manifolds and Hausdorff dimension of cusp excursions*, Ergodic Theory Dynam. Systems **24** (2004), no. 3, 803–824.
6. R. D. Mauldin, T. Szarek, and M. Urbański, *Graph directed Markov systems on Hilbert spaces*, Math. Proc. Cambridge Philos. Soc. **147** (2009), 455–488.
7. C. A. Rogers and S. J. Taylor, *The analysis of additive set functions in Euclidean space*, Acta Math. **101** (1959), 273–302.
8. B. Schapira, *Lemme de l'ombre et non divergence des horosphères d'une variété géométriquement finie. (the shadow lemma and nondivergence of the horospheres of a geometrically finite manifold)*, Ann. Inst. Fourier (Grenoble) **54** (2004), no. 4, 939–987 (French).
9. B. O. Stratmann, *Multiple fractal aspects of conformal measures; a survey*, Workshop on fractals and dynamics (M. Denker, S.-M. Heinemann, B. Stratmann, eds.), Math. Gottingensis, vol. 5, Springer-Verlag, Berlin, 1997, pp. 65–71.
10. ———, *Fractal geometry on hyperbolic manifolds*, Non-Euclidean geometries, Math. Appl. (N.Y.), 581, Springer, New York, 2006, pp. 227–247.
11. B. O. Stratmann and M. Urbański, *The box-counting dimension for geometrically finite Kleinian groups*, Fund. Math. **149** (1996), no. 1, 83–93.
12. B. O. Stratmann and S. L. Velani, *The Patterson measure for geometrically finite groups with parabolic elements, new and old*, Proc. London Math. Soc. (3) **71** (1995), no. 1, 197–220.
13. D. P. Sullivan, *Discrete conformal groups and measurable dynamics*, Bull. Amer. Math. Soc. **6** (1982), no. 1, 57–73.
14. ———, *Disjoint spheres, approximation by imaginary quadratic numbers, and the logarithm law for geodesics*, Acta Math. **149** (1982), no. 3–4, 215–237.
15. ———, *Entropy, Hausdorff measures old and new, and limit sets of geometrically finite Kleinian groups*, Acta Math. **153** (1984), no. 3–4, 259–277.
16. S. J. Taylor and C. Tricot, *Packing measure, and its evaluation for a Brownian path*, Trans. Amer. Math. Soc. **288** (1985), no. 2, 679–699.

OHIO STATE UNIVERSITY, DEPARTMENT OF MATHEMATICS, 231 W. 18TH AVENUE, COLUMBUS, OH 43210-1174, USA
 E-mail address: simmons.465@osu.edu
 URL: <https://sites.google.com/site/davidsimmonsmath/>

⁶Here we use the fact that for all $x_1, x_2 \in \mathbb{R}^d$ and $r_1, r_2 > 0$, $B(x_1, r_1)$ is disjoint from $B(x_2, r_2)$ if and only if $d(x_1, x_2) \geq r_1 + r_2$.