



UNIVERSITY OF LEEDS

This is a repository copy of *A Facile Palladium Catalysed 3-Component Cascade Route to Functionalised Isoquinolinones and Isoquinolines*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/91357/>

Version: Accepted Version

Article:

Grigg, R, Elboray, EE, Akkarasamiyo, S et al. (7 more authors) (2016) A Facile Palladium Catalysed 3-Component Cascade Route to Functionalised Isoquinolinones and Isoquinolines. *Chemical Communications*, 52 (1). pp. 164-166. ISSN 1359-7345

<https://doi.org/10.1039/c5cc07526b>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

A Facile Palladium Catalysed 3-Component Cascade Route to Functionalised Isoquinolinones and Isoquinolines

Ronald Grigg,^{*a} Elghareeb E. Elboray,^{a,b} Sunisa Akkarasamiyo,^{a,c} Nutthawat Chuanopparat,^{a,c} H. Ali Dondas^d, Hussien H. Abbas-Temirek,^b Colin W.G. Fishwick,^{*a} Moustafa F. Aly,^b Boonsong Kongkathip^c and Ngampong Kongkathip^c

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

Palladium catalysed three component cascade process, involving coupling of 2-iodobenzoates, -benzaldehydes, or acetophenones with substituted allenes and ammonium tartrate as an ammonium surrogate, provides a novel and facile route to substituted functionalised isoquinolinones and isoquinolines in good yields.

Isoquinolinone and isoquinoline derivatives are important constituents of a diverse range of molecules of biological and pharmaceutical relevance as well as a common structural component within many alkaloids.^{1,2,3} However, little attention has been given to approaches to these systems which utilise metal-catalysed aryl-allene couplings, despite the obvious potential of this route to produce such systems with high degrees of chemo- and regioselectivity. Nonetheless, the potential of this approach has been underlined previously. Thus, Larock reported that annulation of a substituted allene with N-tosyl-2-iodobenzylamine under Pd(II) catalysis afforded isoquinolines as a mixture of three regio- and stereo-isomers.⁴ Additionally, 2-iodobenzaldehyde imines have also been used with Pd(0) catalysis to annulate substituted allenes giving isoquinoline derivatives.⁵ Ni(0)/chiral phosphine ligand mediated regio- and enantioselective synthesis of isoquinoline-1(2H)-one derivatives has been reported via denitrogenation or decarbonylation of N-aryl-1,2,3-benzotriazin-4(3H)-ones or N-substituted phthalimide, respectively, followed by intermolecular annulation with substituted allenes.⁶ Recently, Glorius et al., employed Rh(III) to catalyse C-H activation of N-(pivaloyloxy)benzamide involving

intermolecular annulation with substituted allenes to furnish isoquinoline-1(2H)-ones.⁷

As part of our program of research into the development and application of palladium catalysed allene insertion cascades, we have previously reported a number of examples of three-component cascades for the synthesis of N-substituted 4-methylene-3,4-dihydro-1(2H)-isoquinolinones. A feature of these reactions is that, following an initial Pd-mediated intramolecular allene insertion, both intra- and intermolecular nucleophilic addition then occurs to give tetra-fused ring systems containing an isoquinolinone ring.^{8,9a} These include, following the initial Pd catalysed intramolecular allene insertion, intermolecular nucleophilic addition involving N-allenyl-2-iodobenzamide,^{9b,c} and nitrogen-tethered 1,6-enynes^{9d} respectively. Additionally, we have also reported two types of cascade reactions that can furnish isoquinolines; (i) intermolecular allene insertion into the C-I bond of an aryl iodide linked N-nucleophile followed by intramolecular N-addition to the generated π -allyl,¹⁰ and (ii) intermolecular allene insertion to an aryl iodide carrying a dipolarophile/Michael acceptor followed by intermolecular N-addition of an azide/amine and finally intramolecular 1,3-dipolar cycloaddition/Michael addition.¹¹

In the present study, we report a new approach utilising our “ammonium surrogate” technology¹² as a novel ammonia source to furnish substituted functionalised isoquinolinone and isoquinoline derivatives. Thus, methyl 2-iodobenzoate derivatives **1** were reacted with a range of substituted allenes **2** in the presence of ammonium tartrate (6 equiv.), Pd₂(dba)₃ (2.5 mol%), TFP (10 mol%), and K₂CO₃ (3 equiv.), to afford isoquinolinones **4** via intramolecular cyclisation of the intermediate **3** in 51-78% yield (Table 1). Z-configuration of the exocyclic double bonds were established using NOE data (see the Experimental Section) and in the absence of ammonium tartrate, no reaction occurred, and only starting materials were observed. This appears to be consistent with a mechanism involving the addition of ammonia to the π -allyl intermediate forming amine **3** which subsequently cyclises to give **4**. Thus, the cyclisation step in **3**→**4** is faster than further allylation of the allyl-NH₂ group. In the case of methyl 5-bromo-2-iodobenzoate (**1**, R¹ = 5-Br), the reaction is chemoselective for oxidative addition at the C-I bond leaving the

^a School of Chemistry, University of Leeds. Leeds, LS6 9JT, UK. Fax: +44-0113 343 6501; Tel: +44-0113 343 6510; E-mail: r.grigg@leeds.ac.uk; c.w.g.fishwick@leeds.ac.uk.

^b Department of Chemistry, Faculty of Science at Qena, South Valley University, Qena, Egypt


^c Natural Products and Organic Synthesis Research Unit (NPOS), Department of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of Science, Kasetsart University, Chatuchak, Bangkok 10900, Thailand

^d Department of Chemistry, Faculty of Pharmacy, Mersin University, 33342, Mersin, Turkey

† Electronic Supplementary Information (ESI) available: [Full characterisation of all new compounds]. See DOI: 10.1039/b000000x/

C-Br bond intact. It is also noteworthy that the ester moieties in **4k-m** were unchanged under the reaction conditions.


Table 1. Pd(0) catalysed annulation of allenes with methyl 2-iodobenzoates.



Reactions carried out at 100 °C in 1,4-dioxane/DMF (5:1) for 16-31h and employed substituted allene **2** (1 equiv.), **1** (1.2 equiv.), ammonium tartrate (6 equiv.), Pd₂(dba)₃ (2.5 mol%), TFP (10 mol%), and K₂CO₃ (3 equiv.). Figures in brackets indicate reaction time and isolated yields.

In order to briefly explore the potential of adducts **4** for further synthetic manipulation, compounds **4f** and **4k**, selected as representative examples, were converted into the corresponding 1-chloroisoquinolines **5a** and **5b** in the presence of POCl₃, (Scheme 1). The assignment of the structures of the chlorination


products to chloropyridines **5a** and **5b** followed from analysis of the ¹H-NMR data for these compounds. This revealed the absence of allyl signals (typically a triplet at ~6-6.5 ppm and doublet at ~4.5-5 ppm respectively), and instead, comprised an AA'BB' nmr pattern for the two methylene groups at 3-3.5 and 4-4.5 ppm respectively, consistent with the assigned structures. (see Supporting Information).



Scheme 1. Conversion of isoquinolinone to isoquinoline.

To further probe the scope of this process, the reaction of 2-iodobenzaldehydes/2'-iodoacetophenone **6** with substituted allenes **2** was explored. This reaction presumably goes via intermediate **7** which undergoes a 1,3-hydrogen rearrangement generating isoquinolines **8**, Table 2. Analogously to isoquinoline **5a** and **5b**, ¹H-NMR data (see Supporting Information) showed no indication of allyl signals but instead included an AA'BB' pattern for the two methylene groups present in **8**. The somewhat low yields of products from this reaction may reflect the thermal instability of the substrates or the products. This hypothesis appears to be supported by the isolation of theobromine (in the case of **8a,f** and **g**), 2',3',5'-tri-O-acetyluridine (in case of **8d**), 3',5'-tri-O-acetylthymidine (in case of **8e**), quinazolin-4-one (in case of **8b**) and chloroquinazolin-4-one (in case of **8c**) as by-products. It is noteworthy that thermal degradation of products was not observed in the preparation of isoquinolinones **4a-m**.

Table 2. Pd(0) catalysed preparation of isoquinolines **8**.



Reaction carried out at 100 °C in 1,4-dioxane/DMF (5:1) for 12-26h and employed substituted allene **2** (1 equiv.), **6** (1.2 equiv.), ammonium tartrate (6 equiv.), Pd₂(dba)₃ (2.5 mol%), TFP (10 mol%), and K₂CO₃ (2-3 equiv.).

In summary, a novel and powerful cascade approach has been applied to the synthesis of substituted functionalised isoquinolinone and isoquinoline derivatives via 3-component palladium catalysed cascade chemistry. The utility of ammonium tartrate as a novel ammonia source is underlined in this simple one-pot cascade protocol.

We acknowledge support from the University of Leeds, the Egyptian Government, South Valley University, Thailand Research Fund (TRF), Royal Golden Jubilee Program, the Center of Excellence for Innovation in Chemistry (PERCH-CIC), Mersin University and The Scientific and Technological Research Council of Turkey (TUBİTAK), Commission on Higher Education, Ministry of Education and Kasetsart University Research and Development Institute.

Notes and references

- See for example: (a) J. D. Scott and R. M. Williams, *Chem. Rev.* 2002, 102, 1669-1730; (b) S.-S. Lee, Y.-C. Lai, C.-K. Chen, L.-H. Tseng and C.-Y. Wang, *J. Nat. Prod.* 2007, 70, 637-642; (c) A. Padwa and H. Zhang, *J. Org. Chem.* 2007, 72, 2570-2582; (d) M. Matveenko, O. J. Kokas, M. G. Banwell and A. C. Willis, *Org. Lett.* 2007, 9, 3683-3685; (e) C. T. Goralski, D. L. Hasha, D. R. Henton, R. C. Krauss, C. D. Pfeiffer and B. M. Williams, *Org. Process Res. Dev.* 1997, 1, 273-279; (f) C.-J. Chou, L.-C. Lin, K.-T. Chen and C.-F. Chen, *J. Nat. Prod.* 1994, 57, 689-694. (g) K. Iwasa, M. Moriyasu, Y. Tachibana, H. S. Kim, Y. Watanya, W. Wiegrebe, K. F. Bastow, M. Cosentino, M. Kozuka, K. H. Lee. *Bioorg. Med. Chem.* 2001, 9, 2781-2884
- See for example: (a) Y. Asano, S. Kitamura, T. Ohra, F. Itoh, M. Kajino, T. Tamura, M. Kaneko, S. Ikeda, H. Igata, T. Kawamoto, S. Sogabe, S.-I. Matsumoto, T. Tanaka, M. Yamaguchi, H. Kimurab and S. Fukumoto, *Bioorg. Med. Chem.* 2008, 16, 4699-4714; (b) R. Pellicciari, E. Camaioni, G. Costantino, L. Formentini, P. Sabbatini, F. Venturoni, G. Eren, D. Bellocchi, A. Chiarugi and F. Moroni, *ChemMedChem.* 2008, 3, 914-923; (c) W.-H. Chueh and J.-Y. Lin, *J. Agric. Food Chem.* 2011, 59, 8021-8027; (d) H.-P. Kuo, T.-C. Chuang, S.-C. Tsai, H.-H. Tseng, S.-C. Hsu, Y.-C. Chen, C.-L. Kuo, Y.-H. Kuo, J.-Y. Liu and M.-C. Kao, *J. Agric. Food Chem.* 2012, 60, 9649-9658; (e) M. Jayaraman, B. M. Fox, M. Hollingshead, G. Kohlhausen, Y. Pommier and M. Cushman, *J. Med. Chem.* 2002, 45, 242-249; (f) L. Ingrassia, F. Lefranc, J. Dewelle, L. Pottier, V. Mathieu, S. Spiegl-Kreinecker, S. Sauvage, M. El Yazidi, M. Dehoux, W. Berger, E. Van Quaquebeke and R. Kiss, *J. Med. Chem.* 2009, 52, 1100-1114; (g) L. Chen, M. Conda-Sheridan, P. V. N. Reddy, A. Morrell, E.-J. Park, T. P. Kondratyuk, J. M. Pezzuto, R. d B. van Breemen and M. Cushman, *J. Med. Chem.* 2012, 55, 5965-5981; (h) A. Chiarugi, E. Meli, M. Calvani, R. Picca, R. Baronti, E. Camaioni, G. Costantino, M. Marinozzi, D. E. Pellegrini-Giampietro, R. Pellicciari and F. Moroni, *J. Pharmacol Exp Ther.* 2003, 305, 943-949; (i) X. Xiao and M. Cushman, *J. Org. Chem.* 2005, 70, 6496-6498.
- See for example: (a) M. Álvarez and J. A. Joule, in *Science of Synthesis*, ed. D. S. Black, Thieme, Stuttgart, 2004, vol 15, pp. 661-836; (b) M. Álvarez and J. A. Joule, in *Science of Synthesis*, ed. D. S. Black, Thieme, Stuttgart, 2004, vol 15, pp. 839-906; (c) M. Chrzanowska and M. D. Rozwadowska, *Chem. Rev.* 2004, 104, 3341-3370; (d) G. Zeni and R. C. Larock, *Chem. Rev.* 2006, 106, 4644-4680; (e) J. Lu and H. Fu, *J. Org. Chem.* 2011, 76, 4600-4605; (f) L. Ackermann and S. Fenner, *Org. Lett.* 2011, 13, 6548-6551; (g) N. Guimond, C. Gouliaras and K. Fagnou, *J. Am. Chem. Soc.* 2010, 132, 6908-6909; (h) K. Fujiwara, T. Kurahashi and S. Matsubara, *Org. Lett.* 2010, 12, 4548-4551; (i) P. C. Too and S. Chiba, *Chem. Commun.* 2012, 48, 7634-7636.
- (a) T. Lechel, F. Pfrengle, H.-U. Reissig and R. Zimmer, *ChemCatChem.* 2013, 5, 2100-2130. (b) R. C. Larock, N. G. Berrios-Peiiia and C. A. Fried, *J. Org. Chem.* 1991, 56, 2615-2617.
- J. J. H. Diederer, R. W. Sinkeldam, H.-W. Frühauf, H. Hiemstra and K. Vrieze, *Tetrahedron Lett.* 1999, 40, 4255-4258.
- (a) M. Yamauchi, M. Morimoto, T. Miura and M. Murakami, *J. Am. Chem. Soc.* 2010, 132, 54-55; (b) Y. Ochi, T. Kurahashi and S. Matsubara, *Org. Lett.* 2011, 13, 1374-1377.
- H. Wang and F. Glorius, *Angew. Chem. Int. Ed.* 2012, 51, 7318-7322.
- R. Grigg, T. Khamnaen, S. Rajviroongit and V. Sridharan, *Tetrahedron Lett.* 2002, 43, 2601-2603
- (a) R. Grigg, I. Köppen, M. Rasparini and V. Sridharan, *Chem. Commun.* 2001, 964-965; (b) M. Gardiner, R. Grigg, M. Kordes, V. Sridharan and N. Vicker, *Tetrahedron* 2001, 57, 7729-7735; (c) R. Grigg, V. Sridharan and L.-H. Xu, *J. Chem. Soc. Chem. Commun.* 1995, 1903-1904. ; (d) N. M. Groome, E. E. Elboray, M.W. Inman, H. A. Dondas, R. M. Phillips, C. Kilner, R. Grigg, *Chem. Eur. J.* 2013, 19, 2180 - 2184
- (a) R. Grigg, V. Sridharan and A. Thayaparan, *Tetrahedron Lett.* 2003, 44, 9017-9019; (b) H. A. Dondas, C. W. G. Fishwick, X. Gai, R. Grigg, C. Kilner, N. Dumrongchai, B. Kongathip, N. Kongathip, C. Polysuk and V. Sridharan, *Angew. Chem. Int. Ed.* 2005, 44, 7570-7574.
- (a) R. Grigg, M. Inman, C. Kliner, I. Köppen, J. Marchbank, P. Selby and V. Sridharan, *Tetrahedron* 2007, 63, 6152-6169; (b) X. Gai, R. Grigg, I. Köppen, J. Marchbank and V. Sridharan, *Tetrahedron Lett.* 2003, 44, 7445-7448; (c) X. Gai, R. Grigg, S. Rajviroongit, S. Songarsa and V. Sridharan, *Tetrahedron Lett.* 2005, 46, 5899-5902.
- R. Grigg, S. Akkarasamiyo, N. Chuanopparat, E. E. Elboray, M. F. Aly, H. H. Abbas-Temirek, B. Kongkathip and N. Kongkathip, *Chem. Commun.* 2013, 49, 2007-2009.