



This is a repository copy of *An exploratory randomised trial of a simple, brief psychological intervention to reduce subsequent suicidal ideation and behaviour in patients admitted to hospital for self-harm..*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/95403/>

Version: Accepted Version

Article:

Armitage, C.J., Abdul Rahim, W., Rowe, R. et al. (1 more author) (2016) An exploratory randomised trial of a simple, brief psychological intervention to reduce subsequent suicidal ideation and behaviour in patients admitted to hospital for self-harm. *British Journal of Psychiatry*. ISSN 0007-1250

<https://doi.org/10.1192/bjp.bp.114.162495>

This is an author-produced electronic version of an article accepted for publication in the *British Journal of Psychiatry*. The definitive publisher-authenticated version is available online at <http://bjp.rcpsych.org>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

An Exploratory Randomized Trial of a Simple, Brief Psychological Intervention to Reduce Subsequent Suicidal Ideation and Behaviour in Patients Hospitalized for Self-Harm

Christopher J. Armitage

University of Manchester

Wirda Abdul Rahim and Richard Rowe

University of Sheffield

Rory C. O'Connor

University of Glasgow

Correspondence to:

Chris Armitage

Manchester Centre for Health Psychology, School of Psychological Sciences

Manchester Academic Health Science Centre, University of Manchester

Coupland Street, Oxford Road

Manchester, M13 9PL, UK

E-mail: chris.armitage@manchester.ac.uk

Tel: +44 (0) 161 275 2556

Fax: +44 (0) 161 275 2623

Date: 18 December 2014

Words: 3906 (excluding references)

Running head: BRIEF INTERVENTION TO REDUCE SUICIDAL IDEATION AND BEHAVIOUR

The authors declare no conflict of interest and thank the Malaysian Ministry of Health for their financial support of this project.

An Exploratory Randomized Trial of a Simple, Brief Psychological Intervention to Reduce Subsequent Suicidal Ideation and Behaviour in Patients Hospitalized for Self-Harm

Background. “Implementation intentions” (IIs) link triggers for self-harm with coping skills and appear to create an automatic tendency to invoke coping responses when faced with a triggering situation.

Methods. 226 self-harm patients were randomized to: (a) form IIs with a “volitional help sheet”; (b) self-generate IIs without help; or (c) think about triggers and coping, but not form IIs. Self-reported suicidal ideation and behaviour, threats of suicide, and likelihood of future suicide attempt were measured at baseline and 3-month follow-up.

Results. All suicide-related outcome measures were significantly lower at follow-up among patients forming IIs compared with those in the control condition ($d_s > 0.35$). The volitional help sheet caused fewer suicide threats ($d = 0.59$) and lower likelihood of future suicide attempts ($d = 0.29$) compared with patients who self-generated IIs.

Conclusions. IIs-based interventions, particularly when supported by a volitional help sheet, show promise in reducing future suicidal ideation and behaviour.

Key words: self-harm, suicide, implementation intention, brief intervention, volitional help sheet, Malaysia

An Exploratory Randomized Trial of a Simple, Brief Psychological Intervention to Reduce Subsequent Suicidal Ideation and Behaviour in Patients Hospitalized for Self-Harm

Suicidal and non-suicidal self-harm¹ extorts significant social and economic costs. People admitted to hospital following an episode of self-harm are thirty-times more likely to die by suicide than those in the general population (1) and even modest reductions in self-harm would bring considerable savings to health care services (2). Recent research attention has therefore focused on testing brief interventions that can be delivered to patients before they are discharged from hospital (3). These brief interventions have shown promise in reducing suicidal ideation and behaviour, but they may be limited by typically not being based explicitly on psychological theories of behaviour change and in focusing on heterogeneous patient groups (3). The aim of the present research was to test a brief theory-based psychological intervention to reduce suicidal ideation and behaviour among patients admitted following an episode of self-harm.

People engage in self-harm for many different reasons (4), but common among these reasons are triggers or critical situations (e.g., defeat, entrapment) in which people feel compelled to self-harm (5), and the implication is that providing people with the means to respond effectively to these critical situations might lessen the likelihood of an act of self-harm. Implementation intentions – tools based on Gollwitzer’s (6) model of action phases – might be helpful in this regard because they work by automatizing appropriate responses to critical situations.

Implementation intentions are “if-then” plans that work by linking in memory a critical situation (“if”) with an appropriate response (“then”). For the purposes of the present research, participants will be asked to identify “critical situations” in which they may be tempted to self-harm and to link them with “appropriate responses” such as consciousness

raising and stimulus control (7) that will help overcome those situations. The principal idea behind implementation intentions is that specifying the circumstances in which one will act (e.g., “if I am tempted to self-harm when I feel trapped”) ensures that the appropriate response (e.g., “then I will do something else instead of self-harming”) will be triggered at the appropriate time and place in the future. One key feature of implementation intentions is that they seem to operate beyond conscious awareness by enhancing the salience of the critical situation and automatizing the appropriate response, mechanisms that are supported by meta-analysis (8). It is also clear that the way in which these plans are formed is critical: For example, Armitage (9) showed that asking people simply to form plans had no effect on subsequent alcohol consumption whereas implementation intentions significantly reduced alcohol intake.

There is a large body of research attesting to the efficacy of implementation intention-based interventions for changing behaviour in field settings. Gollwitzer and Sheeran (8) identified 94 independent tests of implementation intentions (including laboratory tests) that yielded an average effect size of $d = 0.65$. Of these 94 studies, however, none was concerned with self-harm, and no studies of the potential impact of implementation intentions on suicidal ideation and behaviour have yet been published.

Thus the main aim of the present research was, for the first time, to test the ability of implementation intentions to reduce suicidal ideation and behaviour in a high risk group. In addition, we sought to address two further issues that have arisen in the broader literature in relation to implementation intentions. First, the effects of implementation intentions on behaviour change have typically been tested among students (8) and have not yet been tested among clinical populations.

Second, in field studies, people are asked to self-generate implementation intentions, i.e., participants are provided with instructions that ask them to produce both critical

situations and appropriate responses and then create their own implementation intentions.

Although this approach has been shown to be successful in bringing about behaviour change, it is not clear whether this method will be effective in a sample of people who have been admitted to hospital following an episode of self-harm. Thus, in addition to asking people to self-generate implementation intentions to reduce suicidal ideation and behaviour, we will test a tool that is designed to assist implementation intention formation, namely, a volitional help sheet (10).

Volitional help sheets are designed to provide a standard means by which people can form their own implementation intentions and so overcome the need for participants to self-generate implementation intentions (10). The volitional help sheet for self-harm provides participants with the critical situations they may encounter and the responses they might use to ensure they avoid self-harming. The content of the volitional help sheet draws on theories of suicidal behaviour (3), the self-harm motives literature (11) and the transtheoretical model of change (7) and provides a theoretically-driven framework on which participants can build their own implementation intentions. To date, volitional help sheets have successfully reduced cigarette smoking (10), alcohol consumption (12) and weight (13), but – consistent with the broader implementation intention literature – the volitional help sheet has not yet been tested in the domain of self-harm.

Based upon the research reviewed above, there are two rationales underpinning the present research. First, there is a need to reduce suicidal ideation and behaviour in a cost-effective manner. Second, although implementation intentions have been shown consistently to change behaviour, no studies have yet tested the ability of implementation intentions to improve treatment outcomes in relation to self-harm. It is hypothesized that: (a) implementation intentions will significantly reduce suicidal ideation and behaviour, and (b) using a tool to support the formation of implementation intentions (a “volitional help sheet”,

10), as opposed to asking people to form their own (“self-generated”) implementation intentions, will maximize reductions in suicidal ideation and behaviour.

Method

Participants

Both Kuala Lumpur Hospital ethics committee and the University of Sheffield ethics committee gave approval to conduct the research. Participants were assured of confidentiality and anonymity (personal codes were used to identify individuals to preserve confidentiality and facilitate blinding), and were made aware of their right to withdraw from the study or have their data removed at any point with no adverse consequences.

Two hundred and seventy eight patients who had been admitted to Kuala Lumpur Hospital following an episode of self-harm (ICD-10 cause codes X60-X84, [14]) were approached between 1 March 2010 and 28 February 2011. Potential participants were asked to read a patient information sheet and provide signed informed consent before taking part in the study (Figure 1). Although no incentive was offered for participation, 226 (81%) people initially agreed to take part in the study.

Design

A mixed measures design was employed with one between-participants factor and one within-participants factor. *Condition* (control versus self-generated implementation intention versus volitional help sheet implementation intention) was the between-participants factor, and *time* (baseline versus 3-month follow-up) was the within-participants variable. The main outcome measure was suicidal ideation and behaviour. The three-month follow-up period was chosen because this is the modal time to self-harm repetition (15)².

Procedure

Once informed consent was received, participants were given a baseline questionnaire to complete alone, which was collected subsequently by a site investigator who was

independent of the research team. The interventions were placed at the end of the identical-looking questionnaires, which had previously been sorted into random order using a web-based randomizer. This meant that, as far as was feasible, the site investigator was blind to condition: Research staff with knowledge of treatment allocation had no interactions with patients while they were in hospital or at discharge. Because Kuala Lumpur Hospital does not have an anonymized central database for self-harm events, participants were asked to provide contact details if they were willing to complete follow-up measures. One hundred and seven participants (47%) were successfully contacted again and completed 3-month follow-up questionnaires (Figure 1). Baseline and follow-up questionnaires were matched using personal codes: Contact details were kept separate from the data. The data were analyzed according to intention to treat, with the last observation carried forward.

Interventions

Participants in all three conditions were presented with a brief statement designed to encourage them to plan not to self-harm (“We want you to plan not to self-harm. Research shows that you are much more likely to be successful in your intention not to self-harm if you can identify critical situations and appropriate responses”). Following this statement, participants randomized to the *self-generated* implementation intention condition were given standard (9) implementation intention instructions: “You are free to choose how you will do this, but we want you to formulate your plans in as much detail as possible. Please pay particular attention to the situations in which you will implement these plans” and participants were left space in which to write their implementation intentions.

Participants in the *volitional help sheet* implementation intention condition had a volitional help sheet appended to their questionnaires following the brief statement encouraging them to plan to stop self-harming. The volitional help sheet was similar to those used to support implementation intention formation and successful health behaviour change

in previous research (10). It consisted of a table with two columns each containing lists of eleven critical situations and eleven appropriate responses (see Appendix). The eleven critical situations were derived from items used to measure self-harm triggers (4, 11, 16, 17) and the eleven appropriate responses were derived from items used to measure the processes of change from Prochaska and DiClemente's (7) transtheoretical model. The critical situations tap the range of motives that typically underpin self-harm (including suicidal and non-suicidal motives). The temptation items were translated into "if" statements, for example: "If I am tempted to self-harm when I want to get relief from a terrible state of mind;" the processes of change items were translated into "then" statements, for example, "then I will think about the impact of my self-harming on the people around me." There was one item for each of the eleven processes of change. Participants in the *volitional help sheet* condition were then asked to draw links between as many critical situations and appropriate responses as they wanted and thereby form implementation intentions.

Participants in the *control condition* were also given a volitional help sheet, but were not instructed to form implementation intentions. Instead they were simply asked to identify critical situations and appropriate responses that might be useful to them.

Measures

Suicidal ideation and behaviour. Suicidal ideation and behaviour was measured using the revised suicide behaviours questionnaire (18). The four items assess suicidal ideation and behaviour ("Have you ever thought about or attempted to kill yourself?"), frequency of suicidal thoughts ("How often have you thought about killing yourself in the past year?"), threats to commit suicide ("Have you ever told someone that you were going to commit suicide, or that you might do it?"), and self-reported likelihood of suicide attempts ("How likely is it that you will attempt suicide someday?"). The suicidal ideation and behaviour items ask about suicidal thoughts, plans and attempts with and without a wish to

die. At follow-up, we framed these questions in relation to suicidal ideation and behaviour in the preceding 3-months.

Depression. Depression was measured using the Beck Depression Inventory - II (19), which had good internal reliability at baseline ($\alpha = .75$) and at 3-month follow-up ($\alpha = .75$).

Motivation. Motivation to avoid self-harming in the future was assessed by adapting standard measures of behavioural intention and self-efficacy assessed on 7-point (+1 to +7) scales (10). Behavioural intention was measured using three items (e.g., “I intend to avoid deliberately harming myself *definitely do not-definitely do*”). Internal reliability at both baseline ($\alpha = .41$) and 3-month follow-up ($\alpha = .40$) was low. Self-efficacy was measured using three items (e.g., “My avoiding deliberately harming self is *difficult-easy*”). Internal reliability at baseline was $\alpha = .71$, at 3-month follow-up it was $\alpha = .69$.

Data Analysis

Randomization was tested using MANOVA; the effects of the interventions were tested using repeated measures ANOVAs as well as ANCOVAs that controlled for baseline values. Because this was an exploratory trial, we chose not to specify a main outcome measure *a priori*, nor to apply Bonferroni’s correction to the statistical comparisons.

Results

Representativeness Check

Consistent with studies of self-harm prevalence around the world, significantly more women than men were admitted for self-harm. Consistent with Malaysian data (20, 21), the Indian subgroup accounted for a significantly higher proportion of admissions than would be expected by chance (Table 1).

Randomization Check

Randomization was checked using MANOVA. The independent variable was *condition* with three levels: Control versus self-generated implementation intention versus volitional help sheet implementation intention. The dependent variables were age, gender, suicidal ideation and behaviour, frequency of suicidal thoughts, threats to commit suicide, likelihood of attempting suicide again, depression, behavioural intention, and self-efficacy at baseline. The multivariate test was nonsignificant, $F(18, 430) = 1.23, p = .23, \eta_p^2 = .05, d = 0.46$, as were all the univariate tests, indicating success in the randomization procedure.

Effects of the Interventions

The effect of the interventions was tested initially using mixed ANOVAs. *Condition* was the between-participants factor and *time* (baseline versus 3-month follow-up) was the within-persons factor. The dependent variables were: Suicidal ideation and behaviour, frequency of suicidal thoughts, threats to commit suicide, likelihood of attempting suicide again, depression, behavioural intention, and self-efficacy (Table 3).

There were nonsignificant condition \times time interactions for: Frequency of suicidal thoughts, $F(2, 223) = 2.66, p = .07, \eta_p^2 = .02, d = 0.29$ (Figure 2), depression, $F(2, 223) = 1.48, p = .23, \eta_p^2 = .01, d = 0.20$, behavioural intention, $F(2, 223) = 1.42, p = .24, \eta_p^2 = .01, d = 0.20$, and self-efficacy, $F(2, 223) = 2.83, p = .06, \eta_p^2 = .02, d = 0.29$ (Table 3). However, there were significant interactions between time and condition for: Suicidal ideation and behaviour, $F(2, 223) = 4.95, p < .01, \eta_p^2 = .04, d = 0.41$, threats to commit suicide, $F(2, 223) = 6.20, p < .01, \eta_p^2 = .05, d = 0.46$, and likelihood of future suicide attempt, $F(2, 223) = 4.78, p < .01, \eta_p^2 = .04, d = 0.41$. The subsequent analyses decompose these significant interactions (see also Figure 2).

Suicidal ideation and behaviour. Repeated measures ANOVAs, run separately for each condition, revealed significant decreases in suicidal ideation and behaviour between baseline and follow-up, $F_s(1, 72-77) = 9.99-40.58, p_s < .01, \eta_p^2_s > .12, d_s > 0.74$; the largest

decrease in suicidal ideation and behaviour was associated with the volitional help sheet condition, $F(2, 74) = 40.58, p < .01, \eta_p^2 = .35, d = 1.47$. Between-participants ANCOVAs controlling for suicidal ideation and behaviour at baseline showed significant differences between conditions at follow-up, $F(2, 222) = 8.70, p < .01, \eta_p^2 = .07, d = 0.55$. Planned simple contrasts revealed significant differences ($ps < .01$) between the control and both implementation intention formation conditions, but the self-generated and volitional help sheet conditions did not differ significantly from one another ($p = .13$).

Threats to commit suicide. Repeated measures ANOVAs, run separately for each condition, revealed significant decreases in threats to commit suicide between baseline and three months in the volitional help sheet condition, $F(1, 74) = 13.77, p < .01, \eta_p^2 = .16, d = 0.87$, but not in the control or self-generated conditions, $F_s(2, 72, 77) = 1.00, ps > .30, \eta_p^2s < .02, ds < 0.29$. Between-participants ANCOVAs controlling for threats to commit suicide at baseline showed significant differences between conditions at follow-up, $F(2, 222) = 8.96, p < .01, \eta_p^2 = .07, d = 0.55$. Planned simple contrasts revealed a significant difference ($p < .01$) between the control and volitional help sheet conditions, but no significant difference ($p = .70$) between the control and self-generated conditions. The volitional help sheet condition and self-generated conditions differed significantly from one another ($p < .01, d = 0.59$), meaning participants in the volitional help sheet condition made significantly fewer threats to commit suicide at follow-up.

Likelihood of future suicide attempt. Repeated measures ANOVAs, run separately for each condition, revealed significant decreases in likelihood of future suicide attempt between baseline and follow-up across all conditions, $F_s(1, 72-77) = 4.25-23.52, ps < .05, \eta_p^2s > .05, ds > 0.46$, although the largest decrease in likelihood of future suicide attempt was associated with the volitional help sheet condition, $F(1, 74) = 23.52, p < .01, \eta_p^2 = .24, d = 1.12$. Between-participants ANCOVAs controlling for likelihood of future suicide attempt at

baseline showed significant differences between conditions at follow-up, $F(2, 222) = 3.81, p = .02, \eta_p^2 = .03, d = 0.35$. Planned simple contrasts revealed a significant difference ($p < .01$) between the control and the volitional help sheet conditions, but not between the control and self-generated conditions ($p = .56$). The self-generated and volitional help sheet conditions also differed significantly from one another at follow-up ($p = .04, d = 0.29$) with people in the volitional help sheet condition reporting significant lower likelihood of their future suicide attempt controlling for baseline.

Discussion

This is the first study to use implementation intentions to try to reduce suicidal ideation and behaviour among patients admitted to hospital following an episode of self-harm. The principal finding was that implementation intention formation was associated with lower levels of suicidal ideation and behaviour, fewer threats to commit suicide, and lower reported likelihood of future suicide attempt at follow-up. The effects were more pronounced when implementation intention formation was structured by use of a supporting tool, the volitional help sheet, which significantly augmented the effects of implementation intention formation on threats to commit suicide, and reported likelihood of future suicide attempt. Moreover, because we employed an “active” control group, we were able to control for people’s exposure to the critical situations and appropriate responses described in the volitional help sheet. This means that explicit implementation intention formation is necessary for behaviour change to occur. The following discussion focuses on the theoretical and practical implications of the findings.

To date, implementation intention research has been characterized by student samples and short follow-up periods (8) and the present research extends the evidence base to a clinical sample with a three-month follow-up. This is important because one implication of the present research is that some samples may need support in implementation intention

formation and that the volitional help sheet represents one way in which this might be achieved.

Consistent with a large body of research on the impact of implementation intention-based interventions on behaviour change, the present study showed that implementation intentions were effective in reducing suicidal ideation and behaviour in a clinical setting without potentially costly health professional time, tailoring or targeting. Together, these findings suggest that the volitional help sheet potentially represents a unique, non-invasive, low-cost tool that can be used to prevent repetition of suicidal ideation and behaviour. However, the question arises as to whether the effects could be enhanced with further input from a health professional. For example, Luszczynska, Sobczyk, and Abraham (22) showed that, in a study of women enrolled in a commercial weight loss programme, participants who were asked to form implementation intentions with the help of feedback lost significantly more weight than women in the control group. Thus, the beneficial effects of implementation intention formation may be augmented by interaction with a health professional and could boost the effects observed in the present study. Indeed, as implementation intention-based interventions are not focused on the causes of an individual's distress, such interventions are likely to be of most use when used as an adjunct to other forms of psychosocial or pharmacological treatments.

Consistent with previous research (10), the impact of implementation intentions on suicidal ideation and behaviour was not mediated by behavioural intention or self-efficacy, which provides further support for the claim that implementation intentions represent a case of strategic automaticity whereby they operate immediately, efficiently and beyond conscious awareness (6). It would be valuable to develop measures that could tap these constructs in field settings so that the mechanism by which implementation intentions operate in the field

could be verified. Nevertheless, there is considerable laboratory research showing the hypothesized mechanisms (8).

Although the present research makes contributions both to the literature on suicidal ideation and behaviour and the implementation intention literature, it is important to highlight some potential limitations. First, the sample was heterogeneous, including all self-harm presentations irrespective of whether the participants presented with suicidal or non-suicidal self-harm. This is not problematic for the administration of the help sheet per se as the volitional help sheet was designed for use in all cases of self-harm, irrespective of motivation. However, as a consequence of the heterogeneity, it is not clear whether the effects are equally applicable to individuals who present to hospital with suicidal and non-suicidal behaviour or with people with different suicidal histories. Second, the outcome measures, although selected for their brevity and established psychometric properties, did not distinguish clearly between suicidal thoughts and behaviours: For example, one of the items was “Have you ever thought about or attempted to kill yourself?”.

A third limitation is that, for practical reasons, the follow-up was restricted to mailed materials, which meant that attrition was inevitable and that the outcome measures had to be self-reported. Although attrition was handled using standard intention-to-treat analyses and made no difference to the principal findings, it would be valuable in future to devote more resources to ensuring participants complete the study. Fourth, it would be valuable to have a more objective outcome measure such as future hospital admissions, although this is not currently possible in the Malaysian context. Fifth, as far as was feasible, the person collecting the data was blind to condition, but in future research it would be valuable explicitly to probe the success or failure of such concealment procedures. Sixth, because this was an exploratory trial, we chose not to specify a main outcome measure a priori, nor to apply Bonferroni’s correction to the statistical comparisons. In future definitive trials, we

will be able to generate more precise estimates of effect size and hence specify the main outcome measure in advance.

In conclusion, the volitional help sheet shows promise as a brief, cost-effective tool to reduce suicidal ideation and behaviour, threats to commit suicide, and reported likelihood of future suicide attempt. Further research is required to replicate the findings with a more complete dataset and objective outcome measures.

Footnotes

¹ Despite the inclusion of non-suicidal self-injury and suicidal behaviour disorder in section 3 of DSM-5 (for disorders requiring further research), there is still no international consensus on which terms best describe the wide range of self-injurious behaviours (see Kapur et al., 2013). As self-injurious behaviour is rarely driven by a sole motive, in this paper, self-harm is used to refer to all forms of self-injurious behaviour irrespective of motive(s).

² Note that we attempted to conduct an additional follow-up at six months. However, substantial attrition (>80% from baseline, $n = 35$ at 6-month follow-up) means that we lack sufficient confidence in the reliability/validity of the findings to present them in the main text. We attribute the substantial rate of attrition to three main factors that were due to the administrative arrangements in Malaysian healthcare and/or a condition of gaining ethical approval: (a) postal follow-up, (b) lack of incentives, and (c) lack of reminders.

Nevertheless, using identical analyses to those reported in the body of the text (i.e., standard intention to treat, last observation carried forward), the pattern of findings at six-month follow-up is identical (i.e., statistically significant differences in favour of the volitional help sheet condition) to the pattern of findings at the three-month follow-up.

References

1. Cooper, J., Kapur, N., Webb, R., Lawlor, M., Guthrie, E., Mackway-Jones, K., & Appleby, L. (2005). Suicide after deliberate self-harm: A 4-year cohort study. *American Journal of Psychiatry, 162*, 297-303. doi: 10.1176/appi.ajp.162.2.297
2. Sinclair, J. M. A., Gray, A., Rivero-Arias, O., Saunders, K. E. A., & Hawton, K. (2011). Healthcare and social services resource use and costs of self-harm patients. *Social Psychiatry and Psychiatric Epidemiology, 46*, 263-271. doi: 10.1007/s00127-010-0183-5
3. O'Connor, R. C., Rasmussen, S., & Beautrais, A. (2011). Recognition of suicide risk, crisis helplines, and psychosocial interventions: A selective review. In R. C. O'Connor, S. Platt, and J. Gordon (Eds.), *International handbook of suicide prevention: Research policy and practice* (pp. 435-456). Chichester, UK: Wiley-Blackwell.
4. Hjelmeland, H., Hawton, K., Nordvik, H., Bille-Brahe, U., De Leo, D., Fekete, S., Grad, O., Haring, C., Kerkhof, A. J. F. M., Lonnqvist, J., Michel, K., Renberg, E. S., Schmidtke, A., Van Heeringen, K., & Wasserman, D. (2002). Why people engage in parasuicide: A cross-cultural study of intentions. *Suicide and Life-Threatening Behavior, 32*, 380-393. doi: 10.1521/suli.32.4.380.22336
5. O'Connor, R. C., Smyth, R., Ferguson, E., Ryan, C., & Williams, J. M. G. (2013). Psychological processes and repeat suicidal behavior: A four year prospective study. *Journal of Consulting and Clinical Psychology, 81*, 1137-1143. doi: 10.1037/a0033751
6. Gollwitzer, P. M. (1999). Implementation intentions: Strong effects of simple plans. *American Psychologist, 54*, 493-503.
7. Prochaska, J. O., & DiClemente, C. C. (1983). Stages and processes of self-change of smoking – toward an integrative model of change. *Journal of Consulting and Clinical Psychology, 51*, 390-395. doi: 10.1037/0022-006X.51.3.390

8. Gollwitzer, P. M., & Sheeran, P. (2006). Implementation intentions and goal achievement: A meta-analysis of effects and processes. *Advances in Experimental Social Psychology*, 38, 69–119.
9. Armitage, C. J. (2009). Effectiveness of experimenter-provided and self-generated implementation intentions to reduce alcohol consumption in a sample of the general population: A randomized exploratory trial. *Health Psychology*, 28, 545-553. doi: 10.1037/a0015984
10. Armitage, C. J. (2008). A volitional help sheet to encourage smoking cessation: A randomized exploratory trial. *Health Psychology*, 27, 557–566. doi: 10.1037/0278-6133.27.5.557
11. Madge, N., Hewitt, A., Hawton, K., de Wilde, E. J., Corcoran, P., Fekete, S., van Heeringen, K., De Leo, D., & Ystgaard, M. (2008). Deliberate self-harm within an international community sample of young people: Comparative findings from the Child and Adolescent Self-harm in Europe (CASE) study. *Journal of Child Psychology and Psychiatry*, 49, 667-677. doi: 10.1111/j.1469-7610.2008.01879.x
12. Armitage, C. J., & Arden, M. A. (2012). A volitional help sheet to reduce alcohol consumption in the general population: A field experiment. *Prevention Science*, 13, 635-643. doi: 10.1007/s11121-012-0291-4
13. Armitage, C. J., Norman, P., Noor, M., Alganem, S., & Arden, M. A. (2014). Evidence that a very brief psychological intervention boosts weight loss in a weight loss program. *Behavior Therapy*, 45, 700-707. doi: 10.1016/j.beth.2014.04.001
14. World Health Organization (WHO). (2011). Retrieved 30 October 2011 from <http://apps.who.int/classifications/icd10/browse/2010/en#/X64>
15. Kapur, N., & Gask, L. (2009). Introduction to suicide and self-harm. *Psychiatry*, 8, 233-236. doi: 10.1016/j.mppsy.2009.04.008

16. O'Connor, R. C. (2011). The integrated motivational-volitional model of suicidal behavior. *Crisis, 32*, 295-298. doi: 10.1027/0227-5910/a000120
17. Williams, J. M. G. (2001). *Suicide and attempted suicide: Understanding the cry of pain*. London: Penguin Psychology.
18. Osman, A., Bagge, C. L., Gutierrez, P. M., Konick, L. C., Kopper, B. A., & Barrios, F. X. (2001). The Suicidal Behaviors Questionnaire-Revised (SBQ-R): Validation with clinical and nonclinical samples. *Assessment, 8*, 443-454. doi: 10.1177/107319110100800409
19. Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory II*. San Antonio, TX: Psychological Corporation.
20. Armitage, C. J., Abdul Rahim, W., Rowe, R., & O'Connor, R. C. (in press). Trends in self-harm in Kuala Lumpur, 2005-2011. *Archives of Suicide Research*.
21. Armitage, C. J., Panagioti, M., Abdul Rahim, W., Rowe, R., & O'Connor, R. C. (in press). Completed suicides and self-harm in Malaysia: A systematic review. *General Hospital Psychiatry*.
22. Luszczynska, A., Sobczyk, A., & Abraham, C. (2007). Planning to lose weight: Randomized controlled trial of an implementation intention prompt to enhance weight reduction among overweight and obese women. *Health Psychology, 26*, 507-512.

Table 1

Baseline Characteristics of the Sample Compared With Population

| Variable | Sample (<i>n</i> = 226) | Population ^a (<i>N</i> = 1,674,621) | χ^2 for difference between sample and population |
|----------------------------|-----------------------------|--|--|
| Gender (%) | | | 39.16 (<i>p</i> < .01) |
| Male | 30.1 | 50.9 | |
| Female | 69.9 | 49.1 | |
| Age (%) | | | |
| 0-14 years | 0.0 | 22.1 | 64.11 (<i>p</i> < .01) |
| 15-64 years | 96.0 | 73.2 | 59.98 (<i>p</i> < .01) |
| 65 years and older | 4.0 | 4.7 | 0.26 (<i>p</i> = .61) |
| Ethnicity (%) ^b | | | |
| Malay | 30.1 | 45.9 | 24.17 (<i>p</i> < .01) |
| Chinese | 10.6 | 43.2 | 97.76 (<i>p</i> < .01) |
| Indian | 50.0 | 10.3 | 385.30 (<i>p</i> < .01) |
| Others | 1.3 | 0.6 | 2.00 (<i>p</i> = .16) |

Note. ^aDepartment of Statistics Malaysia (2011). ^bEighteen (8.0%) people chose not to report their ethnicity.

Table 2

Baseline Characteristics of the Sample by Randomized Intervention Group

| Baseline Variables | Control, <i>n</i> = 73 | | Self-Generated Intervention, <i>n</i> = 78 | | Volitional Help Sheet Intervention, <i>n</i> = 75 | | <i>p</i> ^a | <i>d</i> ^a |
|---|------------------------|-----------|---|-----------|--|-----------|-----------------------|-----------------------|
| | <i>M</i> | <i>SD</i> | <i>M</i> | <i>SD</i> | <i>M</i> | <i>SD</i> | | |
| Age (years) | 28.25 | 11.44 | 30.26 | 12.34 | 31.57 | 16.07 | .32 | 0.20 |
| Gender (Men = 1, Women = 2) | 1.71 | 0.46 | 1.63 | 0.49 | 1.76 | 0.43 | .20 | 0.20 |
| Suicidal Ideation and Behaviour | 2.79 | 1.30 | 2.47 | 1.39 | 2.83 | 1.51 | .23 | 0.20 |
| Frequency of Suicidal Thoughts | 1.99 | 0.72 | 1.74 | 0.75 | 1.92 | 0.78 | .12 | 0.29 |
| Threats to Commit Suicide | 1.30 | 0.62 | 1.36 | 0.60 | 1.29 | 0.65 | .78 | 0.11 |
| Likelihood of Attempting Suicide Again | 2.42 | 2.05 | 2.13 | 1.92 | 2.69 | 2.01 | .22 | 0.20 |
| Depression | 20.19 | 7.52 | 18.74 | 6.84 | 18.09 | 5.58 | .15 | 0.29 |
| Behavioral Intention | 3.44 | 0.52 | 3.42 | 0.65 | 3.51 | 0.54 | .63 | 0.13 |
| Self-Efficacy | 3.32 | 0.55 | 3.21 | 0.78 | 3.25 | 0.56 | .54 | 0.14 |

Note. The reported means are “raw” and not adjusted for baseline values.

^a*p* values and *d* values associated with the univariate *F*s testing for differences in baseline values between intervention conditions and control condition. All comparisons are nonsignificant. The omnibus test was also nonsignificant.

Table 3


Effects of the Interventions on Self-Harm, Depression and Motivation

| Variables | Baseline | | Follow-Up | |
|---|----------|-----------|-----------|-----------|
| | <i>M</i> | <i>SD</i> | <i>M</i> | <i>SD</i> |
| <i>Suicidal Ideation and Behaviour</i> | | | | |
| Control, <i>n</i> = 73 | 2.79 | 1.30 | 2.44 | 1.01 |
| Self-Generated, <i>n</i> = 78 | 2.47 | 1.39 | 1.96 | 0.99 |
| Volitional Help Sheet, <i>n</i> = 75 | 2.83 | 1.51 | 1.95 | 1.02 |
| <i>Frequency of Suicidal Thoughts</i> | | | | |
| Control, <i>n</i> = 73 | 1.99 | 0.72 | 2.00 | 0.73 |
| Self-Generated, <i>n</i> = 78 | 1.74 | 0.75 | 1.77 | 0.77 |
| Volitional Help Sheet, <i>n</i> = 75 | 1.92 | 0.78 | 1.79 | 0.81 |
| <i>Threats to Commit Suicide</i> | | | | |
| Control, <i>n</i> = 73 | 1.30 | 0.62 | 1.26 | 0.53 |
| Self-Generated, <i>n</i> = 78 | 1.36 | 0.60 | 1.32 | 0.52 |
| Volitional Help Sheet, <i>n</i> = 75 | 1.29 | 0.65 | 1.03 | 0.66 |
| <i>Likelihood of Future Suicide Attempt</i> | | | | |
| Control, <i>n</i> = 73 | 2.42 | 2.05 | 2.16 | 1.82 |
| Self-Generated, <i>n</i> = 78 | 2.13 | 1.92 | 1.87 | 1.60 |
| Volitional Help Sheet, <i>n</i> = 75 | 2.69 | 2.01 | 1.92 | 1.57 |
| <i>Depression</i> | | | | |
| Control, <i>n</i> = 73 | 20.19 | 7.52 | 17.78 | 6.51 |
| Self-Generated, <i>n</i> = 78 | 18.74 | 6.84 | 16.20 | 6.56 |
| Volitional Help Sheet, <i>n</i> = 75 | 18.09 | 5.58 | 16.65 | 5.92 |
| <i>Behavioural Intention</i> | | | | |

| | | | | |
|---------------------------------|------|------|------|------|
| Control, $n = 73$ | 3.44 | 0.52 | 3.56 | 0.54 |
| Self-Generated, $n = 78$ | 3.42 | 0.65 | 3.58 | 0.68 |
| Volitional Help Sheet, $n = 75$ | 3.51 | 0.54 | 3.51 | 0.62 |
| <i>Self-Efficacy</i> | | | | |
| Control, $n = 73$ | 3.32 | 0.55 | 3.60 | 0.55 |
| Self-Generated, $n = 78$ | 3.21 | 0.78 | 3.57 | 0.58 |
| Volitional Help Sheet, $n = 75$ | 3.25 | 0.56 | 3.44 | 0.55 |

Note. The reported means are “raw” and not adjusted for baseline values.

Figure 1

Flow of Participants Through the Phases of the Field Experiment


Note. Data were analyzed according to intention-to-treat with the last observation carried forward.

Figure 2


Effects of the Interventions on the Main Suicide-Related Outcomes at 3-Month Follow-up

(Adjusted for Baseline)


Panel A: Suicidal Ideation and Behaviour




Panel B: Frequency of Suicidal Thoughts



Panel C: Threats to Commit Suicide



Panel D: Likelihood of Future Suicide



Appendix

Self-Harm Volitional Help Sheet (Intervention Instructions)

We want you to plan to avoid self-harming. Research shows that if people can spot situations in which they will be tempted to self-harm and then link them with a way to overcome those situations, they are much more likely to be successful in avoiding self-harming.

On the left hand side of the page below is a list of common situations in which people feel tempted to self-harm; on the right hand side of the page is a list of possible solutions.

For each situation that applies to you personally (left hand side), **please draw a line linking it to a solution** (right hand side) that you think might work for you. Please draw a line linking one situation to one solution at a time, **but make as many (or as few) situation-solution links as you like.**

| SITUATIONS | SOLUTIONS |
|---|---|
| <input type="checkbox"/> If I am tempted to self-harm when I want to get relief from a terrible state of mind | <input type="checkbox"/> Then I will do something else instead of self-harming |
| <input type="checkbox"/> If I am tempted to self-harm when I want to punish myself | <input type="checkbox"/> Then I will tell myself that I can stop self-harming if I want to |
| <input type="checkbox"/> If I am tempted to self-harm when I want to die | <input type="checkbox"/> Then I will recall information people have given me about the benefits of stopping self-harming |
| <input type="checkbox"/> If I am tempted to self-harm when I want to show how desperate I am feeling | <input type="checkbox"/> Then I will tell myself that Society is changing in ways that make it easier for people to stop self-harming |
| <input type="checkbox"/> If I am tempted to self-harm when I want to find out whether someone really loves me | <input type="checkbox"/> Then I will make sure I am rewarded by others if I don't self-harm |
| <input type="checkbox"/> If I am tempted to self-harm when I want to get some attention | <input type="checkbox"/> Then I will think about the impact of my self-harming on the people around me |
| <input type="checkbox"/> If I am tempted to self-harm when I want to frighten someone | <input type="checkbox"/> Then I will remember that I react emotionally to warnings about my self-harming |
| <input type="checkbox"/> If I am tempted to self-harm when I want to get my own back on someone | <input type="checkbox"/> Then I will remember that I get upset when I think about my self-harming |
| <input type="checkbox"/> If I am tempted to self-harm when I feel defeated | <input type="checkbox"/> Then I will put things around my home or place of work that remind me not to self-harm |
| <input type="checkbox"/> If I am tempted to self-harm when I feel trapped | <input type="checkbox"/> Then I will seek out someone who listens when I need to talk about self-harm |

| | |
|---|--|
| <input type="checkbox"/> If I am tempted to self-harm when I feel hopeless | <input type="checkbox"/> Then I will take medication |
|---|--|