



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/96029/>

Version: Accepted Version

Article:

Lyons, I, Barber, N, Raynor, DK et al. (2015) The Medicines Advice Service Evaluation (MASE): An RCT of an Intervention to Improve Medication Adherence in a Mail-order Pharmacy Population. *Pharmacoepidemiology and Drug Safety*, 24 (S1-Special Issue: Ab). p. 209. ISSN: 1053-8569

<https://doi.org/10.1002/pds.3838>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The Medicines Advice Service Evaluation (MASE): An RCT Of An Intervention To Improve Medication Adherence In A Mail-Order Pharmacy Population

Imogen Lyons, BA MSc¹, Nick Barber, BPharm PhD², D.K. Raynor, BPharm PhD³ and Li Wei, MBChB MSc MPH PhD¹.

¹Practice and Policy, School of Pharmacy, University College London, London, United Kingdom

²The Health Foundation, London, United Kingdom

³University of Leeds, Leeds, United Kingdom.

Background

Non-adherence to medicines for long-term conditions is a complex, prevalent phenomenon, with significant clinical and economic consequences for patients and health providers worldwide.

Objectives

To test the effectiveness of a pharmacist-led intervention to improve adherence, in the context of mail-order pharmacy.

Methods

A parallel-group RCT was conducted. 677 patients prescribed at least one oral medication for type 2 diabetes and/or lipid regulation were recruited from a UK mail-order pharmacy between Nov 2012-Sept 2013, and randomised (340 intervention, 337 control). The intervention was patient-centred, comprising information and advice by phone and written information by post, delivered by a pharmacist. All elements of the intervention were tailored to the individuals' needs. The primary outcome was self-reported adherence to medication at 6-month follow-up, measured using the Diagnostic Adherence to Medication Scale. Generalised estimating equations analyses were conducted according to the intention-to-treat principle. Secondary outcomes included prescription refill adherence defined as a medication possession ratio and lipid and glycemic control.

Results

Patients who received the intervention had 54% increased odds of being adherent (defined as $\geq 90\%$ of medication taken in the past 7 days), compared with the control group (OR 1.54, 95%CI 1.11-2.15, $p=0.01$). Analyses of dispensing data also showed that the odds of being classified as adherent ($\geq 90\%$) were 60% greater for the intervention group compared with the control group (OR 1.60, 95%CI 1.14-2.24, $p<0.01$). For patients who provided a blood sample at 6-month follow up, 67% vs 31% (16 intervention, 5 control, $p=0.06$) and 65% vs 55% (64 intervention, 38 control, $p=0.24$) achieved guideline targets for glycemic and lipid control, respectively.

Conclusions

Intervention, led by a pharmacist and tailored to the individuals' needs, can significantly improve medication adherence in patients with long-term conditions. The findings provide further support for the enhanced role of

pharmacists in supporting and advising patients with their medicines, and improving outcomes.