

This is a repository copy of *Nurse led care*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/964/>

Article:

Cullum, N, Spilsbury, K orcid.org/0000-0002-6908-0032 and Richardson, G orcid.org/0000-0002-2360-4566 (2005) *Nurse led care*. *British medical journal*. pp. 682-683. ISSN 1756-1833

<https://doi.org/10.1136/bmj.330.7493.682>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

- 1 Sabin CA, Hill T, Lampe F, Matthias R, Baghani S, Gilson R, et al. Treatment exhaustion of highly active antiretroviral therapy (HAART) among individuals infected with HIV in the United Kingdom: multicentre cohort study. *BMJ* 2005;330:695-8.
- 2 Bangsberg DR, Porco TC, Kagay C, Charlebois ED, Deeks SG, Guzman D, et al. Modeling the HIV protease inhibitor adherence-resistance curve by use of empirically derived estimates. *J Infect Dis* 2004;190:162-5.
- 3 D'Arminio Monforte A, Cozzi Lepri A, Rezza G, Pezzotti P, Antinori A, Phillips AN et al. Insights into the reasons for discontinuation of the first highly active antiretroviral therapy (HAART) regimen in a cohort of antiretroviral naive patients. I.CO.N.A. Study Group. *AIDS* 2000;14:499-507.
- 4 Ammassari A, Murri R, Pezzotti P, Trotta MP, Ravasio L, De Longis P, et al. Self-reported symptoms and medication side effects influence adherence to highly active antiretroviral therapy in persons with HIV infection. *J Acquir Immune Defic Syndr* 2001;28:445-9.
- 5 Justice AC, Rabeneck L, Hays RD, Wu AW, Bozzette SA. Sensitivity, specificity, reliability, and clinical validity of provider-reported symptoms: a comparison with self-reported symptoms. *J Acquir Immune Defic Syndr* 1999;21:126-33.
- 6 Guidelines for the use of antiretroviral agents in HIV-1 infected adults and adolescents. 29 October 2004. http://aidsinfo.nih.gov/guidelines/adult/AH_102904.pdf (accessed 10 March 2005).
- 7 Garcia F, De Lazzari E, Plana M, Castro P, Mestre G, Nomdedu M, et al. Long-term CD4+ T-cell response to highly active antiretroviral therapy according to baseline CD4+ T-cell count. *J Acquir Immune Defic Syndr* 2004;36:702-13.
- 8 Hunt PW, Deeks SG, Rodriguez B, Valdez H, Shade SB, Abrams DI, et al. Continued CD4 cell count increases in HIV-infected adults experiencing 4 years of viral suppression on antiretroviral therapy. *AIDS* 2003;17:1907-15.
- 9 Centers for Diseases Control. Report of the NIH panel to define principles of therapy of HIV infection and guidelines for the use of antiretroviral agents in HIV-infected adults and adolescents. *Morb Mortal Wkly Rep MMWR* 1998;47(RR-5):1-41.
- 10 Badri SM, Adeyemi OM, Max BE, Zagorski BM, Barker DE. How does expert advice impact genotypic resistance testing in clinical practice? *Clin Infect Dis* 2003;37:708-13.
- 11 Raffanti SP, Fusco JS, Sherrill BH, Hansen NI, Justice AC, D'Aquila R, et al. Effect of persistent moderate viremia on disease progression during HIV therapy. *J Acquir Immune Defic Syndr* 2004;37:1147-54.
- 12 The PLATO Collaboration. Predictors of trend in CD4-positive T-cell count and mortality among HIV-1-infected individuals with virological failure to all three antiretroviral-drug classes. *Lancet* 2004;364:51-62.

Nurse led care

Determining long term effects is harder than measuring short term costs

What's the difference between medical and nursing care? The answer is not straightforward, but shortages in the medical workforce mean that nurses are increasingly called on to undertake work that was previously done by doctors (such as undertaking surgery,¹ prescribing drugs, performing triage in emergency departments), whereas shortages in the nursing workforce mean that healthcare assistants now do many tasks that nurses are trained to do. This fluidity in professional roles and competencies enables the health workforce to respond to need, but are outcomes for patients being improved? Do these benefits come at an additional cost, and if so, are they worth paying for?

Over the past decade, research has increasingly compared nurse led care with usual care for aspects of health care previously delivered by doctors. However, nurse led care does not have one meaning. Nurse led care can be usefully viewed as a continuum with, at one end, nurses undertaking highly protocol driven, focused tasks (cardioversion,² colposcopy, smoking cessation) and, at the other end, responding to far more diverse challenges in terms of clinical decision making, such as first contact care and rehabilitation. The extent to which doctors' work can be delegated effectively is likely to be influenced, in part, by the type and complexity of the associated decision tasks. This issue of the *BMJ* presents two economic evaluations of nurse led care—each occupying a different place on this continuum. The paper by Raftery et al (p 707) is an evaluation of nurse led secondary prevention of coronary heart disease and has several strengths, including its basis in a randomised controlled trial with four years' follow up and a cost effectiveness analysis.³ The authors conclude that primary care based, nurse led secondary prevention of coronary heart disease is highly cost effective, since the cost per patient was only £136 (\$260; €195) greater in the intervention group, but the benefits (fewer deaths and improvements in medical care and patient lifestyle) make this highly worth while, with a cost per quality adjusted life year (QALY) of £1097.

The second study, an evaluation of nurse led intermediate care in an acute setting (p 699), represents a more complex nursing role that demands multifarious clinical decisions (although patients reaching intermediate care have been "filtered" through medical diagnosis and initial treatments).⁴ These authors undertook a cost minimisation analysis—they viewed the clinical outcomes in the intermediate care and standard hospital care arms as equivalent and merely totalled up and compared the costs. Walsh et al found, as have others,⁵ that nurse led intermediate care in acute settings is more expensive than standard hospital based care for the inpatient phase, but the longer term costs and benefits are more uncertain.

Close inspection of the clinical outcomes in the trial by Walsh et al⁶ reveals that patients who received nurse led intermediate care had better functional outcomes at discharge, although this did not reach significance. However, this lack of statistical significance is not the same as "no difference" in functional outcomes. A meta-analysis of 10 studies of nurse led intermediate care⁷ (which includes the Walsh trial⁶) identified a statistically significant benefit of nurse led intermediate care on functional status at discharge, as well as reductions in the proportion of patients discharged to institutional care and in readmissions. This indicates that the increase in functional status may be clinically (and potentially economically) important and warrants further study.

In an editorial in the *BMJ* Briggs counselled against cost minimisation analysis in favour of cost effectiveness analysis since studies are rarely powered to confidently identify clinical equivalence.⁸ Hence, the lack of a statistically significant difference in effectiveness should not be used as a justification for a cost minimisation analysis. While the higher costs of nurse led intermediate care are due to an increased length of stay, existing analyses have failed to determine whether these costs are offset by lower costs (of health care and particularly social care) and health benefits gained in the longer term.

Papers p 699 and
Primary care p 707

The ways in which nursing teams in the nurse led units make decisions about discharge also need to be explored. Nurses may, rightly or wrongly, be more conservative in discharging patients. They may err on the side of caution, but the benefits of these conservative decisions can only be judged with longer term follow up.

Do these two new studies help us understand the differences between medical and nursing care? We think they usefully remind us that nursing care is not necessarily less costly and that the extra costs may be worth the benefits but that health outcomes need to be measured carefully in studies of sufficient power. It should not be assumed that the outcomes of nursing and medical care are equivalent.

The skills of healthcare professionals and their assistants are much in demand and constitute a limited

resource that needs to be deployed in the most cost effective way. Although UK health policy supports the development of nursing roles, as nurses take on more duties and responsibilities we must also question what, if anything, is being lost from nursing, to whom and does it matter?

Nicky Cullum *professor*

(nac2@york.ac.uk)

Karen Spilsbury *research fellow*

Department of Health Sciences, University of York, York YO10 5DD

Gerry Richardson *research fellow*

Centre for Health Economics, University of York, York YO10 5DD

Competing interests: KS and GR have conducted and published evaluations of nurse led intermediate care.

- 1 BBC News Online. Training nurses to do surgery. <http://news.bbc.co.uk/2/hi/health/3580453.stm> (accessed 10 Mar 2005).
- 2 Currie MP, Karwatowski SP, Perera J, Langford EJ. Introduction of nurse led DC cardioversion service in day surgery unit: prospective audit. *BMJ* 2004;329:892-4.
- 3 Raftery JP, Yao GL, Murchie P, Campbell NC, Ritchie LD. The cost effectiveness of nurse led secondary prevention clinics for coronary heart disease in primary care: follow up of a randomised trial. *BMJ* 2005;330:707-10.
- 4 Walsh B, Steiner A, Pickering RM, Ward-Basu J. Economic evaluation of nurse led intermediate care versus standard acute care for post-acute medical patients: cost minimisation analysis of data from a randomised controlled trial. *BMJ* 2005;330:699-702.
- 5 Griffiths P, Harris R, Richardson G, Hallett N, Heard S, Wilson-Barnett J. Substitution of a nursing-led inpatient unit for acute services: randomized controlled trial of outcomes and cost of nursing-led intermediate care. *Age Ageing* 2001;30:483-8.
- 6 Steiner A, Walsh B, Pickering RM, Wiles R, Ward J, Brooking JI. Therapeutic nursing or unblocking beds? A randomised controlled trial of a post-acute intermediate care service. *BMJ* 2001;322:453-60.
- 7 Griffiths PD, Edwards MH, Forbes A, Harris RL, Ritchie G. Effectiveness of intermediate care in nursing-led in-patient units. *Cochrane Database Syst Rev* 2004;(4):CD002214.
- 8 Briggs A. Economic evaluation and clinical trials: size matters. *BMJ* 2000;321:1362-3.

Large scale food retail interventions and diet

Improving retail provision alone may not have a substantial impact on diet

Ensuring communities have good access to healthy affordable food is one of the government's joined up strategies to improve public health and reduce health inequalities.^{1,2} Policy solutions for deprived communities without good access—food deserts—have focused on improving provision of food retail as part of a wider suite of recommendations for population dietary change focused around awareness, affordability, and acceptability.³ However, the evidence for the widespread existence of food deserts and their impact on population health has been contested.^{4,5} This has meant that although retail based policy recommendations to reduce diet related health inequalities now exist,^{1,2} the evidence to inform how, when, and where to reduce these inequalities is only now emerging.

Recently completed projects in Newcastle, Leeds, and Glasgow have started to provide us with this evidence.⁶⁻⁸ The Newcastle study concludes that food deserts exist only for a minority of people who do not or cannot shop outside their immediate locality and for whom the locality suffers from poor retail provision of foods that compose a healthy diet. Key predictors of healthy eating were found to be dietary knowledge, relative affluence, and healthy lifestyle—retail provision was not independently associated with diet.

The Leeds and Glasgow studies were both prospective evaluations of the impact of large scale food retailing. Utilising an uncontrolled before-after design the Leeds study concluded that access to food improved notably after the intervention. The average

distance travelled to the main food store fell to under 1 km, and the percentage of people walking to the main food store tripled to over 30%. Substantial increases in consumption of fruit and vegetables of between 0.25 and 0.5 portions per day were also reported, particularly for respondents who switched to the new provision. In contrast the Glasgow study, a controlled quasi-experimental study, found little evidence for an overall effect of the intervention for fruit and vegetable consumption in portions per day. For those consumers who switched their main food shopping to the new store an improvement in consumption of around 0.35 portions per day was seen though the evidence for this was very weak. A substantial positive improvement in one measure of psychological health (GHQ-12) and a weak positive effect on self reported health was seen in switchers.

How should this evidence be interpreted? Firstly, the term food desert, although a striking metaphor, has unintentionally led to such polarisation of views by researchers, policy makers, and other interest groups so as to be of limited further use. The authors of the Newcastle study propose that the focus should be on food equity instead.⁶

Secondly, ambiguity remains over whether large scale food retail interventions work. Despite the reporting of positive changes in fruit and vegetable consumption in the Leeds study, pre-intervention and post-intervention designs alone rarely provide compelling evidence that an intervention has been successful. Changes in the prevalence of risk factors and