

This is a repository copy of *The molecular basis of polysaccharide cleavage by lytic polysaccharide monoxygenases*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/97868/>

Version: Accepted Version

Article:

Frandsen, Kristian E H, Simmons, Thomas J, Dupree, Paul et al. (18 more authors) (2016) The molecular basis of polysaccharide cleavage by lytic polysaccharide monoxygenases. NATURE CHEMICAL BIOLOGY. 298–303. ISSN 1552-4450

<https://doi.org/10.1038/nchembio.2029>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The molecular basis of polysaccharide cleavage by *lytic polysaccharide monoxygenases*

Kristian E. H. Frandsen,¹ Thomas J. Simmons,² Paul Dupree,² Jens-Christian N. Poulsen,¹ Glyn R. Hemsworth,³ Luisa Ciano,³ Esther M. Johnston,³ Morten Tovborg,⁴ Katja S. Johansen,^{4,5} Pernille von Freiesleben,⁴ Laurence Marmuse,⁶ Sébastien Fort,⁶ Sylvain Cottaz,⁶ Hugues Driguez,⁶ Bernard Henrissat,^{7,8,9} Nicolas Lenfant,^{7,8} Floriana Tuna,¹⁰ Amgalanbaatar Baldansuren,¹⁰ Gideon J. Davies,³ Leila Lo Leggio^{1,*} and Paul H. Walton^{3,*}

1. Department of Chemistry, University of Copenhagen, Universitetsparken 5, 2100, Copenhagen Ø, Denmark.
2. Department of Biochemistry, University of Cambridge, Tennis Court Road, Cambridge CB2 1QW, U.K.
3. Department of Chemistry, University of York, York YO10 5DD, U.K.
4. Novozymes A/S, Krogshoejvej 36, 2880 Bagsvaerd, Denmark.
5. Current address: Division of Industrial Biotechnology, Chalmers University of Technology, Kemivägen 10, SE-412 96 Göteborg, Sweden.
6. Université de Grenoble Alpes, CERMAV, F-38000 Grenoble, France. CNRS, CERMAV, F-38000 Grenoble, France.
7. Architecture et Fonction des Macromolécules Biologiques, CNRS, Aix-Marseille Université, 13288 Marseille, France.
8. INRA, USC 1408 AFMB, 13288 Marseille, France.
9. Department of Biological Sciences, King Abdulaziz University, Jeddah, Saudi Arabia.
10. EPSRC National EPR Facility, School of Chemistry and Photon Science Institute, University of Manchester, Oxford Road, Manchester M13 9PL, U.K.

***Lytic polysaccharide monooxygenases* (LPMOs) are copper-containing enzymes which oxidatively break down recalcitrant polysaccharides such as cellulose and chitin. Since their discovery, LPMOs have become integral factors in the industrial utilization of biomass, especially in the sustainable generation of cellulosic bioethanol. As such LPMOs are pivotal in addressing world energy needs. We report here the first structural determination of an LPMO–oligosaccharide complex, yielding detailed new insights into the mechanism of action of LPMOs. Using a combination of structure and electron paramagnetic resonance spectroscopy, we reveal the means by which these enzymes interact with saccharide substrates. We further uncover electronic and structural features of the enzyme active site that show how LPMOs orchestrate the reaction of oxygen with polysaccharide chains.**

The sustainable production of energy and materials from biomass is of global importance, particularly in the need for carbon-neutral sources as contributors to balanced energy portfolios.^{1,2} The capacity of biofuels to make a significant contribution, however, depends upon finding a sustainable means of overcoming the recalcitrance of some of the most abundant and widespread polysaccharides, most notably cellulose. Indeed, a viable solution to this problem has been elusive, with recent decades having seen a worldwide research effort directed at providing a means of sustainably deconstructing cellulose or cellulose-related materials. Progress has been slow, but recent advances in the use of cocktails of glycosidic hydrolases identified from the secretomes of saprophytic organisms are promising.³⁻⁵ For instance, several newly-opened cellulosic-bioethanol

biorefineries which employ enzyme cocktails appear, for the first time, to be close to both commercial and environmental viability. The success of these biorefineries hinges on the efficiency of the enzymatic saccharification step in which lignocellulosic material is decomposed into soluble sugars. The efficacy of this step further rests on advances in the bespoke content of the cocktails, of which perhaps the most conspicuous is the recent inclusion of *lytic polysaccharide monooxygenases* (LPMOs).

LPMOs are metalloenzymes classified in the Carbohydrate-Active Enzymes database (CAZy) as Auxiliary Activity families AA9–AA11 and AA13.⁶⁻⁹ The ability of LPMOs to boost biomass degradation was reported in the late 2000s^{10,11} followed in 2010 by the seminal discovery of their oxidative mode of action,¹² and then in 2011 by further activity studies¹³ and identification of copper at the active site.^{14,15} The unusual activity and active site of LPMOs mean that they are challenging precepts not only in the mechanisms by which biological degradation of biomass occurs, but also in copper bioinorganic chemistry.^{16,17} LPMOs catalyze the scission of a polysaccharide chain by harnessing the oxidative capacity of molecular oxygen to cleave a glycosidic C–H bond, the strength of which is estimated to be at least 95 kcal mol⁻¹ (**Fig. 1a**).¹⁸⁻²¹ To break these bonds LPMOs activate oxygen, in a reducing-agent dependent manner, at a copper-containing active site known as the histidine brace (**Fig. 1b**).^{14,16,22} In doing so, LPMOs boost the action of canonical glycoside hydrolases by up to two orders of magnitude,¹¹ thereby greatly bringing down the financial and environmental penalties normally associated with using recalcitrant polysaccharides as a feedstock, and moreover, placing LPMOs at the center of considerable research interest.⁴ This research interest is focused on improving

the efficiencies of LPMOs, with particular attention on determining the detailed mechanism of action of the enzymes, an endeavor which is hindered by the lack of detailed structural information of an LPMO in direct contact with a polysaccharide. It is in this context that we report here the first crystal structure of an LPMO–oligosaccharide complex. This much-anticipated structure, in combination with its spectroscopy, now reveals the essential molecular features of LPMO interaction with substrates, paving the way forward for improvements in catalytic efficiencies.




Figure 1 | LMPO activity and active site. **a**, Oxidative cleavage of a polysaccharide chain carried out by LPMOs, showing different routes of oxidation carried out by different sub-groups of LPMOs.²³ **b**, The histidine brace active site and proposed first step in the reaction with oxygen.¹⁸

Isolation and activity

An LPMO (CAZy classification AA9) from *Lentinus similis*, naturally devoid of a carbohydrate-binding-module, was expressed in *Aspergillus oryzae* (hereafter

Ls(AA9)A). Using PACE gel product analysis, *Ls*(AA9)A was found to be able to cleave phosphoric acid swollen cellulose (PASC) in the presence of a reducing agent, yielding a range of cello-oligosaccharides dominated by cellobiose and cellotriose (**Fig. 2a**, **Supplementary Fig. 1**). This product profile is notably different from that of most other AA9s which yield longer oligosaccharides; for example, **Figure 2a** shows the action of an AA9 from *Thermoascus aurantiacus* (*Ta*(AA9)A).¹⁴ Additionally, unlike *Ta*(AA9)A, *Ls*(AA9)A also shows cleavage activity on soluble oligosaccharides (cellotriose to cellohexaose, G3–G6, **Supplementary Fig. 1c,d**), a feature which has been reported previously for a handful of other AA9s and which may be an important factor in the effectiveness of these particular LPMOs to degrade polysaccharides.²⁴⁻²⁷ *Ls*(AA9)A products were analyzed by MALDI-TOF-MS and HPLC, showing that C4 was the sole site of oxidation (**Fig. 1a**, **Supplementary Fig. 2** and **Supplementary Note 1**).^{23,25,27}

Taking advantage of the reactivity of *Ls*(AA9)A with oligosaccharides, we monitored the kinetics of oxidative cleavage of a cellotetraose labelled with fluorescent groups (**Fig. 2b**, **Supplementary Note 2**, **Supplementary Fig. 3**). This Förster-resonance-energy-transfer (FRET) substrate is fluorescently silent when the molecule is intact, but fluoresces upon cleavage in dilute solution. Before analysis LPMO samples were freshly prepared with careful removal of any excess copper, and only initial rates were used to avoid any complications due to enzyme inactivation.²⁶ The resultant kinetics could be interpreted using classic Michaelis-Menten parameters, with $K_m = 43 \pm 9 \mu\text{M}$ and $k_{\text{cat}} = 0.11 \pm 0.01 \text{ s}^{-1}$ at pH 7.0 and 37 °C, comparable to previously reported turnover numbers.^{12,25,28}




Figure 2 | *Ls*(AA9)A and *Ta*(AA9)A activities on insoluble and soluble cellulose. a, Comparison of *Ls*(AA9)A and *Ta*(AA9)A activities on PASC. Reaction products were analyzed by PACE. +, incubation with ascorbate; -, incubation without ascorbate. **b,** Cleavage of FRET substrate and subsequent Michaelis-Menten plot, error bars represent random error of triplicate measurements.

Structures of *Ls*(AA9)A and *Ls*(AA9)A–oligosaccharide complexes

The crystallographic structure of *Ls*(AA9)A was determined by molecular replacement using PDB deposition 4EIR as a model.²³ The resulting structure displays a known AA9 topology with an immunoglobulin-type fold (**Fig. 3a**). Initial structures yielded copper coordination geometries consistent with photoreduction of the copper site to Cu(I) by incident X-rays, as has been seen in previously reported structures of other LPMOs (**Fig. 3a,c, Supplementary Fig. 4a**). Accordingly, further datasets were collected at reduced X-ray doses (**Fig. 3b,d, Supplementary Note 3**).^{22,29} In the low X-ray dose structure the copper ion has a coordination sphere expected for a Cu(II) oxidation state with three

nitrogen atoms of the histidine brace and a water molecule in a near square planar geometry (Cu–N(His1) 1.9 Å, Cu–N(His78) 2.1 Å, Cu–NH₂ 2.2 Å, Cu–OH₂ 2.2 Å, remote distances Cu...OH₂ (axial) 2.8 Å and Cu...OTyr164 2.7 Å) (**Fig. 3b**). The topography of the putative polysaccharide binding surface of *Ls*(AA9)A is distinctive when compared to the structures of most other AA9s insofar as the active site is located within a contoured binding surface (**Supplementary Fig. 4d, Supplementary Note 3**), which may be a factor in the ability of *Ls*(AA9)A to react with both oligosaccharides and polysaccharides. As observed in the structures of other LPMOs, the surface in *Ls*(AA9)A further harbors an aromatic residue (Tyr203) which is conserved in all but one of the structurally characterized AA9 LPMOs to date and is approximately 15 Å away from the copper active site.




Figure 3 | Structural views of *Ls(AA9)A* with the copper ion depicted as an orange sphere, Cu(I), or a blue sphere, Cu(II). a, Ribbon view of *Ls(AA9)A*. **b,** Electron density of active site obtained with low X-ray dose. **c,** Principal protein contacts between G3 and the binding surface of *Ls(AA9)A*. **d,** Electron density of *Ls(AA9)A*–G3 with low X-ray dose.

Guided by the activity studies described above, soaking crystals of *Ls(AA9)A* with substrates cellotriose (G3) or cellohexaose (G6) at pH 5.5 afforded new structures which showed clear and unambiguous density associated with the oligosaccharide in contact with the binding surface of *Ls(AA9)A* (**Fig. 3d**, **Supplementary Fig. 4b**, **Supplementary Note 3**), confirming the long-held proposals that this surface is indeed the site of interaction with substrates.^{12,14} Using a sugar subsite nomenclature, analogous to that used for glycoside hydrolases, G6 binds to *Ls(AA9)A* within a shallow groove from subsites –4 to +2 (**Supplementary Fig. 4b,c**), and G3 from –1 to +2 (**Fig. 3c**). In both structures, the terminal glucosyl unit at subsite +2 is held within a network of hydrogen bonds with protein residues (Asn28, His66 and Asn67) anchoring this unit into a fixed position (**Supplementary Table 3**). As illustrated for *Ls(AA9)A*–G3 in **Figure 3c** and **Figure 4**, the glucosyl unit at subsite +1 stacks directly on top of His1 *via* a striking lone-pair...aromatic interaction (ring O5... His1 centroid, 3.4 Å). The glycosidic bond between the units at subsites +1 and –1 arches directly over the putative O₂ binding site at the copper ion (**Fig. 3d**), bringing the C4–H bond in subsite +1 and the C1–H bond in subsite –1 into close proximity to the exogenous ligand on the copper ion. The glucosyl units in subsites –3 and –4 in *Ls(AA9)A*–G6 are less well defined (see

Supplementary Note 3), although the unit in subsite -3 forms close contacts with the aromatic ring of the surface-exposed and highly conserved Tyr203 (C...C distances of 3.7 Å), presumably via CH... π interactions (estimated to give ~2 kcal mol⁻¹ binding energy), again confirming long-held proposals that aromatic groups are effective at binding poly/oligosaccharides in LPMOs (**Supplementary Fig. 4b,c**).^{11,23,30,31} Through a combination of hydrogen bonds (**Supplementary Table 3**), the lone-pair-aromatic interaction with His1, and (for longer oligosaccharides) a CH... π interaction with Tyr203, the orientation and position of the oligosaccharide is held in a fixed aspect relative to the copper active site. In this regard, it is notable that the array of interactions between G3 substrate and enzyme does not drive any significant distortion of torsion angles in the oligosaccharide away from their canonical geometries (**Supplementary Table 2**).³² This lack of distortion reinforces the prevailing idea that LPMO catalysis is largely mediated by the exceptional chemical environment of the copper histidine brace active site,^{16,33} rather than aided by the enzyme-mediated distortion of the substrate.

In a comparison of the structures of *Ls*(AA9)A before and after substrate binding there are clear differences within error (>0.1 $\delta\text{\AA}$) in the copper coordination spheres. Most notably, in the presence of the oligosaccharide the exogenous ligand on the copper is best modeled by a fully occupied chloride ion (**Fig. 3d**, Cu-Cl, 2.3 Å, **Supplementary Note 3**), rather than a water molecule at 2.2 Å as in the non-substrate structure (chloride is present in both soaking conditions at ca. 3.9 M, **Supplementary Table 4**), and there is a shortening of the Cu...OTyr164 contact from 2.7 to 2.5 Å. Beyond the immediate copper coordination sphere, the C6 hydroxymethyl group of the unit at subsite +1 is placed close

to the copper ion (C...Cu, 3.8 Å) such that the group occludes the axial binding site, displacing the water molecule that is seen in the non-substrate structure (*cf.* **Fig. 3b,d**). In the non-substrate structure the main chain carbonyl of His1 forms a hydrogen bond to a water molecule (O...O, 2.8 Å) which is held within a small pocket circumscribed by the side chain of His66 and the main chain peptides of His1 and Ala75. Upon substrate binding, the C6 –CH₂OH group of the glucosyl unit at subsite +1 forms hydrogen bonds with the C3–OH group of adjacent unit at subsite +2 and with the ‘pocket’ water molecule (O...O, 2.8 Å). The hydrogen bonding pattern around the ‘pocket’ water further includes the NH₂ terminus (N...O, 2.9 Å), thereby connecting the substrate to the amino terminus and the copper (Fig. 4).




Figure 4 | Structure of *Ls*(AA9)A around active site before and after binding of G3.

The ‘pocket’ water molecule is shown in red. The extent of hydrogen bonding between the ‘pocket’ water molecule and the amino terminus is informed by EPR spectroscopy, discussed below.

Electron paramagnetic resonance spectroscopy of *Ls*(AA9)A

To probe the solution phase interaction between *Ls*(AA9)A and oligosaccharide substrates, continuous wave (cw) X-band EPR spectra of the *met*-forms of *Ls*(AA9)A and *Ls*(AA9)A–G6 were obtained under both high and low chloride conditions (**Fig. 5**).




Figure 5 | EPR spectra of *Ls*(AA9)A. * indicates signals from copper site where water is the exogenous ligand, other signals are from species with chloride as exogenous ligand. **a**, X-band cw EPR spectra in low chloride conditions, *Ls*(AA9)A–G6 (top) and *Ls*(AA9)A with no oligosaccharide (bottom). **b**, X-band cw EPR spectra in high chloride conditions, *Ls*(AA9)A–G6 (top) and *Ls*(AA9)A with no oligosaccharide (bottom). **c**, Contour presentation of the ^1H -HYSCORE spectrum of *Ls*(AA9)A–G6 in the presence of chloride ions (magnetic field 343.6 mT, time between first and second pulse $\tau = 136$ ns, microwave frequency 9.7 GHz). The anti-diagonal cross-peaks are assigned to anisotropic protons, labelled 1_H and 2_H .

The spectrum of *met-Ls*(AA9)A in the absence of substrate is shown in **Figure 5a**. As with previous EPR spectra of AA9 enzymes, the spin Hamiltonian parameters are diagnostic of a type 2 copper site albeit with some rhombicity evident in the xy direction,

$g_z = 2.28$, $g_y = 2.09$, $g_x = 2.04$, $|A_z| = 458$ MHz. The spectrum was sensitive to the presence of chloride in solution and under saturating conditions (~200 mM chloride) the presence of a second, somewhat rhombic minor species (~20%) could be determined $g_z = 2.25$, $g_y = 2.09$, $g_x = 2.01$, $|A_z| = 455$ MHz, (**Fig. 5b**), although there is some uncertainty in the perpendicular g values.

Addition of G6 to *Ls(AA9)A* under low chloride conditions results in a substrate-induced perturbation of the copper active site (**Fig. 5a**), shown by a small increase in $|A_z|$ to 515 MHz, a decrease in g_y and g_z ($g_z = 2.27$, $g_y = 2.07$, $g_x = 2.04$), and the appearance of well-resolved superhyperfine (SHF) coupling due to an increase in the number of magnetically coupled nitrogen nuclei from two to three (34, 37 MHz in the absence of G6 and 19, 30, 36 MHz with G6, simulations in **Supplementary Fig. 6a,b,d** and **Supplementary Table 6**). These observations suggest that the addition of substrate drives a larger contribution of the amino terminus to the singly-occupied-molecular-orbital (SOMO) on the copper ion.³⁴ Under high chloride conditions, additional SHF features appear due to the presence of a chloride nucleus (**Fig. 5b**) and there is a significant decrease in g_z to 2.23, showing that when chloride is available addition of substrate is accompanied by the coordination of chloride to the copper site, consistent with crystal structures. The synergy of chloride and substrate binding is confirmed by isothermal calorimetry experiments in which the dissociation constant of G6 binding to *Ls(AA9)A* in the presence of 200 mM chloride was found to be 3.7 ± 0.1 μ M, significantly lower than the dissociation constant in solution at the same ionic strength but in the absence of chloride, $K_d \sim 1$ mM (**Supplementary Fig. 8**). Moreover, after G6 and chloride binding, anisotropic cross-peaks arising from

magnetic coupling with the protons of the NH₂ ligand appear in the ¹H HYSCORE spectrum (**Fig. 5c**, **Supplementary Fig. 7**, and **Supplementary Note 4**), in which the two individual protons exhibit large dipolar coupling constants and are chemically distinct. This chemical asymmetry likely arises from the formation of a hydrogen bond between one of the protons of the amino terminus and the pocket water molecule upon G6 addition (**Fig. 4**), which connects the amino terminus to a hydrogen bonding network involving G6 and the remote residues His66 and Asp72. This hydrogen bonding network provides a proton transfer channel for potential deprotonation of the amino terminus to form a strongly electron-donating amide ligand, which has been proposed from small model complex studies to stabilize a high valent reactive copper-oxygen species.^{33,35}

To compare the interactions between the copper site of *Ls*(AA9)A and oligosaccharide substrates to those with insoluble polysaccharide substrates, the X-band cw EPR spectrum of *Ls*(AA9)A was collected in the presence of excess PASC (**Supplementary Fig. 6c**). The spectrum of *Ls*(AA9)A–PASC is virtually identical to that of *Ls*(AA9)A with G6, showing that the binding interactions of *Ls*(AA9)A to polysaccharide substrates like PASC are the same as those observed for oligosaccharide substrates.

Discussion

Exploiting the fact that *Ls*(AA9)A is able to act on soluble cello-oligosaccharides, we have been able for the first time to characterize the detailed interaction of an LPMO with a substrate. The oligosaccharide chain is shown to bind to the enzyme in a fixed orientation relative to the copper active site. A comparison of the *Ls*(AA9)A structures

with those of other AA9 LMPOs not able to act on oligosaccharides, exemplified by *Ta*(AA9)A (**Supplementary Fig. 4** and **5** and **Supplementary Note 3**), shows that the latter lack many of the residues that mediate hydrogen bond interactions with the oligosaccharide substrate(s).

The orientation of substrate binding to *Ls*(AA9)A brings the C1–H bond of the sugar moiety at subsite –1 and the C4–H bond of the sugar moiety at subsite +1 into close proximity to the equatorially-bound ligand at the copper ion (**Fig. 3d**, **Fig. 4**). The exclusive oxidation of the C4–H bond by *Ls*(AA9)A is commensurate with the closer distance of the exogenous ligand to this bond compared to the C1–H bond (3.7 Å vs 3.9 Å, respectively), although slight structural changes would shift the attack to the C1–H bond, highlighting that small differences between different LPMOs can lead to different patterns of reactivity. The proximity of the C4–H bond to the exogenous ligand atom directly bound to the copper is consistent with the possibility of a copper(III)-hydroxide or copper(II)-oxyl being the reactive intermediate (both of which can be formed by further reduction of a copper(II)-superoxide).^{19,36-38}

Further examination of the *Ls*(AA9)A structures shows that oligosaccharide binding leads to the occlusion of the axial binding site on the copper ion by the C6 CH₂OH group of subsite +1, displacing an axial water molecule. This leads to a movement of the copper ion towards Tyr164 (**Fig. 3d**), thereby linking the copper to the potentially redox-active tyrosine. In addition, the occlusion of the axial binding site prevents the potential loss of

a coordinated superoxide anion through the axial site. Such a mechanism of potential deactivation *via* associative displacement of superoxide was proposed for *Ta*(AA9)A.¹⁸

Our studies also reveal an interaction between the ring oxygen of the glucosyl unit occupying subsite +1 and the face of the imidazole side chain of His1.³⁹ Characterized contacts like this are rare in biological chemistry with only a handful of examples, including a ribose sugar ring-oxygen lone pair interacting with the aromatic face of a guanine group.⁴⁰ Despite its scarcity in biology this type of interaction is prevalent in supramolecular chemistry where it is often labeled as a lone-pair to π^* overlap, but is now believed to be principally electrostatic in character.⁴¹ The electrostatic nature of the interaction is likely to be enhanced by the presence of alkyl groups on aromatic ring nitrogen atoms, thereby providing some rationale for the previously puzzling δ -N methylation of His1 observed in fungal LPMOs.

Our studies further point to an interesting interdependence of binding of substrate and chloride on the copper ion. As chloride and superoxide have similar size and polarizability, the greatly enhanced affinity for chloride after oligosaccharide binding raises the possibility that binding of dioxygen and the polysaccharide substrate to the LPMOs is similarly synergistic, thereby affording LPMOs a mechanism by which the production of reactive oxygen species such as superoxide is controlled by the presence of the substrate.⁴² Insight into the origin of this synergistic binding can be found from the crystal structures, where the binding of substrate results in the formation of a cavity at the active site that is of the correct shape to accommodate a chloride or superoxide ion

(**Supplementary Fig. 9**). The change in binding affinity of the exogenous ligand may also arise from changes in the coordinating power of the amino terminus, which is made more basic by the presence of a hydrogen bond to the pocket water molecule upon substrate binding, thus raising the energy of $d(x^2-y^2)$ orbital and stabilizing the Cu(II) oxidation state (**Fig. 4**).

Finally, the observation that the EPR spectra of *Ls*(AA9)A binding to PASC and to the oligosaccharide G6 are close to identical indicates that oligosaccharide substrates faithfully mimic the enzyme's binding to extended polysaccharides. This observation connects our detailed studies of LPMO interactions with an oligosaccharide with the *in vivo* activity of these enzymes, which are expected to interact with extended polysaccharides. Our studies therefore give detailed insight into the means by which LPMOs react with insoluble polysaccharide substrates and pave the way forward for improving the efficacies of these industrially-important enzymes.

Supplementary Information is linked to the online version of the paper. Raw data for EPR, ITC and kinetics experiments is available through the University of York Library, DOI: 10.15124/92868bbb-f21a-41e9-a27f-3078b40d144e

Acknowledgments This work was supported by the UK Biotechnology and Biological Sciences Research Council (BB/L000423 and BB/L021633/1), Agence Française de l'Environnement et de la Maîtrise de l'Energie (1201C102), Danish Council for Strategic Research (12-134923, 12-134922). L.L.L. acknowledges the experimental assistance of

Kathrine Rasmussen and Rasmus M. Borup and thanks MAXLAB, Sweden and the ESRF, France, for provision of synchrotron beamtime and assistance. The Danish Ministry of Higher Education and Science through the Instrument Center DANSCATT and the European Community's Seventh Framework Programme (FP7/2007-2013) under BioStruct-X (grant agreement N°283570) funded travel to synchrotrons. L.M., S.F., S.C. and H.D. acknowledge support from ICMG FR 2607, LabEx ARCANE (ANR-11-LABX-0003-01), PolyNat Carnot Institute, French Agence Nationale de la Recherche (PNRB2005-11).

Author Contributions K.F. crystallized protein, collected and analysed crystallographic data, solved crystal structures and made structural figures and tables; P.D. and T.J.S. conceived, and T.J.S. performed, the activity, oxidation state and MS experiments; J.C.P. crystallized protein and collected crystallographic data; G.R.H. designed and performed the FRET kinetics experiments; L.C. performed EPR experiments and simulations; E.J. performed EPR experiments; M.T. purified the recombinant enzymes; L.M., S.C., S.F. and H.D. conceived and performed the FRET substrate synthesis; B.H. and N.L. performed bioinformatics analyses and alignments; F.T. collected pulsed EPR data; A.B. collected and simulated pulsed EPR spectra; G.J.D. conceived the FRET kinetics study; L.L.L. conceived the crystallographic study, collected and analyzed crystallographic data, solved crystal structures and co-wrote the paper; P.H.W. conceived the EPR study and co-wrote the paper.

Additional Information Atomic coordinates and structure factors for the reported crystal structures have been deposited with the Protein Data Bank under accession codes 5ACH (*Ls*(AA9)A high resolution), 5ACG (*Ls*(AA9)A low X-ray dose), 5ACI (*Ls*(AA9)A with G6), 5ACJ (*Ls*(AA9)A with G3) and 5ACF (*Ls*(AA9)A with G3 and low X-ray dose). The authors declare competing financial interests: M.T. and P.vF. are employees of Novozymes, a producer of enzymes for industrial use. Reprints and permissions information is available at <http://www.nature.com/reprints/index.html>. Correspondence and requests for materials should be addressed to P.H.W. (paul.walton@york.ac.uk) or L.L.L. (leila@chem.ku.dk).

Figure Legends

Figure 1 | LMPO activity and active site. **a**, Oxidative cleavage of a polysaccharide chain carried out by LPMOs, showing different routes of oxidation carried out by different sub-groups of LPMOs.²³ **b**, The histidine brace active site and proposed first step in the reaction with oxygen.¹⁸

Figure 2 | *Ls*(AA9)A and *Ta*(AA9)A activities on insoluble and soluble cellulose. **a**, Comparison of *Ls*(AA9)A and *Ta*(AA9)A activities on PASC. Reaction products were analyzed by PACE. +, incubation with ascorbate; –, incubation without ascorbate. **b**, Cleavage of FRET substrate and subsequent Michaelis-Menten plot, error bars represent random error of triplicate measurements.

Figure 3 | Structural views of *Ls(AA9)A* with the copper ion depicted as an orange sphere, Cu(I), or a blue sphere, Cu(II). a, Ribbon view of *Ls(AA9)A*. **b,** Electron density of active site obtained with low X-ray dose. **c,** Principal protein contacts between G3 and the binding surface of *Ls(AA9)A*. **d,** Electron density of *Ls(AA9)A*–G3 with low X-ray dose.

Figure 4 | Structure of *Ls(AA9)A* around active site before and after binding of G3. The ‘pocket’ water molecule is shown in red. The extent of hydrogen bonding between the ‘pocket’ water molecule and the amino terminus is informed by EPR spectroscopy, discussed below.

Figure 5 | EPR spectra of *Ls(AA9)A*. * indicates signals from copper site where water is the exogenous ligand, other signals are from species with chloride as exogenous ligand. **a,** X-band cw EPR spectra in low chloride conditions, *Ls(AA9)A*–G6 (top) and *Ls(AA9)A* with no oligosaccharide (bottom). **b,** X-band cw EPR spectra in high chloride conditions, *Ls(AA9)A*–G6 (top) and *Ls(AA9)A* with no oligosaccharide (bottom). **c,** Contour presentation of the ¹H-HYSCORE spectrum of *Ls(AA9)A*-G6 in the presence of chloride ions (magnetic field 343.6 mT, time between first and second pulse $\tau = 136$ ns, microwave frequency 9.7 GHz). The anti-diagonal cross-peaks are assigned to anisotropic protons, labelled 1_H and 2_H.

Online Methods

Cloning, expression and purification

The gene encoding *Ls(AA9)A* was PCR-amplified from genomic DNA of *Lentinus similis* and cloned in *E. coli* as described in ⁴³ using the following forward primer F-P247JK: 5'-ACACA ACTGGGGATCCACCATGAAGTACTCCATCCTCGGGCT-3' and reverse primer R-P247JK: 5'-CCCTCTAGATCTCGAGCCTTGTCGAGCGACTCTATCCA-3', containing insertion sites for the vector pDau109 used for cloning. The sequence was deposited as KT368674 (Genbank). The fragments were then cloned into Bam HI and Xho I digested pDau109 using an IN-FUSION™ Cloning Kit. Cloning of the genes into Bam HI-Xho I digested pDau109 resulted in transcription of the *Ls(AA9)A* gene under the control of a NA2-tpi double promoter. The treated plasmids and inserts were transformed into One Shot TOP10F Chemically Competent *E. coli* cells (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol, spread onto LB plates supplemented with 0.1 mg of ampicillin per mL and incubated at 37 °C overnight. Colonies of each transformation were cultivated in LB medium supplemented with 0.1 mg of ampicillin per mL and plasmids were isolated using a QIAPREP Spin Miniprep Kit (QIAGEN Inc., Valencia, CA, USA).

Ls(AA9)A was expressed in *Aspergillus oryzae* MT3568 as also described in ⁴³. A transformant producing the recombinant *Ls(AA9)A* was inoculated in 2 L of Dap-4C medium and incubated at 30 °C for 4 days. Mycelia were removed by filtration and the broth collected for purification. Ammonium sulfate was added to the sterile filtered broth to a conductivity of 200 mSi/cm and the pH adjusted to 7.5. The broth was applied to a 50/15 Butyl Toyopearl column (Tosoh Biosciences, Stuttgart, Germany) equilibrated

with 25 mM Tris, 1.5 M ammonium sulfate, pH 7.5. The column was washed in the same buffer and eluted with a gradient to 25 mM Tris pH 7.5. Fractions containing *Ls(AA9)A* were combined and washed with milliQ water by ultrafiltration (10 KDa MWCO, PES filter, Sartorius, Goettingen, Germany) to a conductivity of 1.2 mSi/cm. The pH was adjusted to 8.0 and applied to a 50/40 Q Sepharose FF column (GE Healthcare, Pittsburgh, USA) equilibrated with 20 mM Tris pH 8.0. The column was washed in the same buffer and the enzyme eluted with a gradient from 0 to 0.5 M sodium chloride. Fractions containing *Ls(AA9)A* were combined and concentrated by ultrafiltration using VIVASPIN 20 (10 KDa MWCO) spin concentrators.

Activity measurements by polysaccharide analysis by carbohydrate gel electrophoresis (PACE) and matrix assisted laser-desorption ionisation (MALDI) mass spectrometry

Ls(AA9)A, at 125 μ M, was pre-incubated for 1 h at 5 °C in 0.9 stoichiometric Cu(II)NO₃ before enzyme reactions. Enzyme reactions on cello-oligosaccharides were in 10 μ L containing 5 nmol cello-oligosaccharide, 100 mM ammonium formate pH 6, \pm 4 mM ascorbate, \pm 5 pmol *Ls(AA9)A/Ta(AA9)A*, and were incubated at 20 °C for 4 h. Reactions on phosphoric acid-swollen cellulose (PASC) were in 100 μ L containing 0.5% (w/v) PASC, 100 mM ammonium formate pH 6, \pm 4 mM ascorbate, \pm 63 pmol *Ls(AA9)A/Ta(AA9)A*, and were incubated at 20 °C for 16 h. PASC was prepared beforehand by making a slurry of 1 g of Avicel cellulose (Sigma-Aldrich) with 3 mL H₂O, before adding 30 mL ice-cold phosphoric acid and incubating at 0 °C for 1 h. The

cellulose was then washed numerous times with water until the flow-through had a neutral pH before use in reactions.

For PACE, reaction products and oligosaccharide standards (Megazyme) were reductively aminated with 8-aminonaphthalene-1,3,6-trisulfonic acid (ANTS; Invitrogen, www.lifetechnologies.com), and separated by acrylamide gel electrophoresis. PACE was performed using a 192 mM glycine, 25 mM tris, pH 8.5 running buffer. Two gels contained: resolving gels, 37.5 mL 40% (w/v) acrylamide, 12.5 mL 375 mM Tris-HCl buffer pH 8.8, 100 μ L 10% ammonium persulphate, 50 μ L tetramethylethylenediamine (TEMED); stacking gels, 2 mL 40% (w/v) acrylamide, 2.5 mL 375 mM Tris-HCl buffer pH 8.8, 5.4 mL water, 100 μ L 10% ammonium persulphate, 10 μ L TEMED. Electrophoresis was carried out at 100 V for 30 min, 500 V for 30 min and 1000 V for 180–210 min, and were then visualised with a G-box (Syngene) equipped with a short pass detection filter (500–600 nm) and long-wave UV tubes (365 nm emission). Representative gels from replicate experiments are shown.

MALDI mass spectrometry was performed as described previously.⁷ Trifluoroacetic acid (TFA) hydrolysis was performed by incubating reaction products in 2 M TFA for 1 h at 120 °C before drying *in vacuo*. High-performance liquid chromatography (HPLC) was performed on a CarboPac PA1 column (high-performance anion-exchange chromatography, HPAEC; Dionex, Camberley, UK) with elution at 0.4 mL min⁻¹ and injections of 20 μ L. The elution profile was: 0–3 min, 10 mM NaOH (isocratic); 3–6, 10→1 mM NaOH (linear gradient); 6–19 min, 1 mM NaOH (isocratic); 19–37 min, 45

mM NaOH, 225 mM Na:Acetate (isocratic). A pulsed amperometric detector (PAD) with a gold electrode was used. PAD response was calibrated using markers (500 pmol).

Synthesis of Förster-resonance-energy-transfer (FRET) substrate

Chemicals were purchased from Sigma-Aldrich Chimie (Saint Quentin-Fallavier, France). Reactions were monitored by thin layer chromatography (TLC) using Silica Gel 60 F254 pre-coated plates (E. Merck, Darmstadt). Detection of carbohydrates was achieved by charring with the sulfuric acid/methanol/H₂O (1:1:0.5 v/v). See **Supplementary Note 2** and **Supplementary Figure 3** for details on analysis of individual compounds.

2,3,4-Tri-*O*-acetyl-6-*O*-levulinyl- β -D-galactopyranosyl-(1 \rightarrow 4)-2,3,6-tri-*O*-acetyl- α -D-glucopyranosyl fluoride (**2**): Hepta-acetyl α -lactosyl fluoride **1**⁴⁴ (955 mg, 1.05 mmol) was suspended in 2-methyl-2-butanol (36 mL) in presence of *Candida antarctica* lipase (Novozyme 435) (1g) and trifluoroethyl levulinate⁴⁵ (3.1 mL) was added. The reaction was placed on a rotative shaker for 3 days at 45~50 °C. The reaction was filtered, the filtrate evaporated and the residue was purified by flash column chromatography (dichloromethane/methanol, 10:0 \rightarrow 9:1 v/v). The expected product was acetylated (acetic anhydride/pyridine, 1:10 v/v, 9 mL) in the presence of catalytic amount of 4-dimethylaminopyridine (DMAP). After 3 h of stirring at room temperature, the reaction mixture was evaporated *in vacuo* and co-evaporated with toluene. The crude product was purified by flash column chromatography (toluene/ethyl acetate, 1:1 \rightarrow 2:3 v/v) to give fluoride **2** as a white solid (350 mg, 48 %).

2,3,4-Tri-*O*-acetyl-6-*O*-methanesulfonyl- β -D-galactopyranosyl-(1 \rightarrow 4)-2,3,6-tri-*O*-acetyl- α -D-glucopyranosyl fluoride (**3**): Fluoride **2** (500 mg, 0.72 mmol) was suspended in ethanol (20 mL) and hydrazine acetate (140 mg, 1.55 mmol) was added. The reaction was stirred at room temperature for 2 h and neutralized with Et₃N. The reaction was concentrated *in vacuo* and the residue was diluted with CH₂Cl₂ and washed with a sat. aq NaCl solution and H₂O. The crude product was dissolved in pyridine (12 mL) in presence of catalytic amount of DMAP. Methanesulfonyl chloride (240 μ L, 1.44 mmol) was added the mixture was stirred for 2 h at room temperature. The solution was evaporated *in vacuo* and co-evaporated twice with toluene. The residue was purified by flash column chromatography (toluene/ethyl acetate, 7:3 \rightarrow 3:2 v/v) to give mesylate **3** as white foam (440 mg, 90%).

2,3,4-Tri-*O*-acetyl-6-azido-6-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)-2,3,6-tri-*O*-acetyl- α -D-glucopyranosyl fluoride (**4**): Compound **3** (380 mg, 0.56 mmol) was dissolved in dimethylformamide (DMF) (8 mL), NaN₃ (610 mg, 7.84 mmol) and 18-Crown-6 (65 mg) were added. The reaction was heated at 80 °C for 4 days and concentrated *in vacuo*. The residue was diluted with CH₂Cl₂ and washed with a sat. aq. NaCl solution then H₂O. The crude product was purified by flash column chromatography (toluene/ethyl acetate, 1:1 \rightarrow 2:3 v/v) to give compound **4** as white foam (206 mg, 59 %).

6-Azido-6-deoxy- β -D-galactopyranosyl-(1 \rightarrow 4)- α -D-glucopyranosyl fluoride (**5**): The peracetylated fluoride **4** (183 mg, 0.29 mmol) was treated with NaOMe (300 μ L, 1M) in

MeOH (7 mL) at 0 °C. The reaction mixture was stirred for 2 h and neutralized with Amberlite IR 120 H⁺. After filtration of the resin and evaporation under reduced pressure the residue was freeze-dried to give the fluoride **5** (103 mg, 96%). This compound was used immediately without any further characterization.

Sodium *N*-[2-*N*[(S-(6-azido-6-deoxy-β-D-galactopyranosyl-(1→4)-β-D-glucopyranosyl-(1→4)-β-D-glucopyranosyl)-2-thioacetyl]aminoethyl]-1-naphthylamine-5-sulfonate (**7**): Cel7B Glu197Ala glycosynthase ⁴⁶ (0.62 mg) was added to a solution of fluoride **5** (31 mg, 0.083 mmol) and *N*-[2-*N*[(S-(β-D-glucopyranosyl)-(1→4)-β-D-glucopyranosyl)-2-thioacetyl]aminoethyl]-1-naphthylamine-5-sulfonate ⁴⁷ **6** (51 mg, 0.076 mmol) in sodium phosphate buffer (1.5 mL, 0.1M, pH 7). The solution was placed in a rotative shaker for 4 h at 37°C. Then the reaction was evaporated and the crude product was purified on octadecyl reversed phase silica cartridge (H₂O/MeOH, 1:0→19:1 v/v) to give tetrasaccharide **7** as an amorphous white solid (61 mg, 79 %).

Sodium *N*-[2-*N*[(S-(6-deoxy-6-dimethylaminophenylazophenylthioureido-β-D-galactopyranosyl-(1→4)-β-D-glucopyranosyl-(1→4)-β-D-glucopyranosyl)-2-thioacetyl]aminoethyl]-1-naphthylamine-5-sulfonate (**8**) (F*-G4-F): A solution of azido tetrasaccharide **7** (50 mg, 49 μmol) in pyridine and H₂O (1:1 v/v, 10 mL) was saturated with H₂S. The reaction was stirred at room temperature overnight. After evaporation under reduced pressure, the residue was dissolved in DMF (10 mL) and aq sodium hydrogenocarbonate solution (6 mL, 0.35M). Then 4-(4-isothiocyanatophenylazo)-*N,N*-dimethylaniline (DABITC) (22 mg, 78 μmol) was added.

The reaction was stirred for 12 h at 40°C. The solution was evaporated under reduced pressure and the residue was purified on octadecyl reversed phase silica cartridge (H₂O/MeOH, 1:0→19:1 v/v). Appropriate fractions were pooled, concentrated *in vacuo* and freeze dried to give the title compound (F*-G4-F) as amorphous orange solid (45 mg, 69 %).

Substrate preparation and kinetics measurements

F*-G4-F was dissolved in DMSO to a final stock concentration of 10 mM. This was then diluted to 1.33x the working concentration with 20 mM Bis-Tris pH 7 and 133 nM *Ls*(AA9)A ensuring the DMSO concentration was the same in all samples. 15 µL aliquots were then spread into a 96 well plate and fluorescence was monitored in an Agilent MX3000P QPCR machine using an excitation wavelength of 330 nm and emission wavelength of 492 nm. The increase in fluorescence in triplicate samples was monitored following the addition of 5 µL of ascorbate (prepared in 20 mM Bis-Tris pH 7) giving final concentrations of 100 nM protein and 5 mM ascorbate. Fluorescence was quantified by plotting a standard curve of fluorescence against an equimolar mix of fluorophore (EDANS – Santa Cruz Biosciences) and quencher (DABCYL - Sigma-Aldrich) from 2 to 10 µM concentrations. Inner filter effects were corrected by measuring standard curves in the presence and absence of substrate at each concentration used and calculating correction factors appropriately.⁴⁸ Data were analysed using SigmaPlot.

Crystallization, Data Collection and Structure determination

All crystals were obtained by vapor diffusion technique set up in sitting drop MRC-plates, with a reservoir volume of 100 μ L and at room temperature using an Oryx-8 robot (Douglas Instrument). Crystallization and post-crystallization experimental details are shown in **Supplementary Table 4**. All data were collected at cryogenic temperatures (100 K) after cryocooling the crystals in liquid nitrogen. Cellotriose (G3) for soaking was purchased from Megazyme, while the cellohexaose (G6) preparation (which contained about 20% cellopentaose as contaminant) was provided by Novozymes/AS.

Intergrown plates of glycosylated *Ls*(AA9)A (6.12 mg/mL in 20mM MES pH 6) were initially obtained in a Morpheus screen. Single plates could be separated and mounted, and a data set extending to a maximum resolution of 2.8 Å was collected at beamline I911-3, Maxlab,Lund, Sweden. A dataset was collected and processed in space group C222₁ (details not shown, this and subsequent data processing carried out with XDS⁴⁹). A partial structure could be determined by Molecular Replacement with MOLREP⁵⁰ using modified coordinates of PDB 4EIR as search model. Despite poor statistics (starting R-factors and R-free around 50%), solutions with 2 mols/asu could be obtained that reproduced well the features seen in a self-rotation function. By alternated manual building in density modified maps in COOT⁵¹ and rigid body/restrained refinement in REFMAC5⁵² a partial model (with rather poor geometry and not containing the active site copper) was obtained with an R-factor of 39.2% and R-free of 37.7% (details not shown), and was used as starting model for further molecular replacement.

For subsequent crystallization trials *Ls*(AA9)A was deglycosylated in 20 mM MES pH 6.0, 125 mM NaCl by incubation with 0.05-0.06 units/mg *Ls*(AA9)A of Endoglycosidase

H from Roch Diagnostics GmbH (REF: 11 643 053 001), and then buffer exchanged to 20 mM sodium acetate buffer pH5.5. Crystallization trials also included pre-incubation with Cu(II)acetate (see details in **Supplementary Table 4**).

A relatively large single crystal (maximum dimension 180 μm) was first obtained in an IndexHT screen (Molecular Dimensions) after two weeks of set up, with precipitant containing 0.1 M Citric Acid pH 3.5 and 3.0 M NaCl. The crystals were later found to be reproducible and grew after few days in conditions ranging from 3.0 - 4.7 M NaCl and pH 3.0 – 4.0. Crystals could be transferred to buffers at higher pH, to provide conditions closer to the pH of activity and spectroscopy studies, and maintaining high resolution diffraction, in general better than 2.0 \AA . The crystals were the basis for determination of a series of structures of which a selection is presented in this manuscript.

The preliminary model obtained for the glycosylated protein was used as search model for the newly obtained crystal form in space group $P4_132$, for which the first structure could be refined using COOT/REFMAC5. All subsequently determined structures were isomorphous to this.

The highest resolution structure obtained (data set denoted $Ls(\text{AA9})\text{A_highres}$) was at 1.28 \AA resolution, for which data was collected at $\lambda = 0.965\text{\AA}$ and processed automatically at ID30A-1 of the ESRF using the MASSIF-1 protocol (statistics in **Supplementary Table 5**). This structure was refined anisotropically for all atoms and had 94.44% of residues in the Ramachandran most favorable regions. Preliminary

experiments at ID30A-1 of the ESRF also showed that complexes with celooligosaccharides could be obtained by soaking.

By soaking crystals in pH 5.5 solutions containing celooligosaccharides G3 (the minimal substrate for *Ls(AA9)A*) and G6, two complexes were obtained (*Ls(AA9)A_G3* and *Ls(AA9)A_G6*, details in **Supplementary Table 4** and **Supplementary Table 5**) at Maxlab in Lund, Sweden, at beamline I911-2 and I911-3, respectively (with $\lambda = 1.03841\text{\AA}$ and 1.06883 respectively). The final structures had 93.81% and 93.36% of residues in the most favorable regions of the Ramachandran plot, respectively, and the bound ligands were evident in the electron density (**Fig. 3d** and **Supplementary Fig. 4b**). In all the structures described so far, the density suggests that Cu is in a mixture of oxidation states, with significant disorder of the non-protein ligands. In order to obtain optimal structures with Cu(II) for interpretation of the EPR data, data was collected (at $\lambda = 1.0000\text{\AA}$ at I911-3) with short exposure times, beam attenuation (3% transmission) and the minimal number of X-ray images to ensure an acceptable completeness (*Ls(AA9)A_lowdose* and *Ls(AA9)A_G3_lowdose*, see **Supplementary Tables 4** and **5** for details). The strategy succeeded in obtaining structures where the geometry strongly supports a Cu(II) state (see **Supplementary Note 3**), although at the cost of somewhat lower quality crystallographic statistics. For the *Ls(AA9)A_lowdose* and *Ls(AA9)A_G3_lowdose* structures 92.14% and 92.27% of residues were in the most favorable regions of the Ramachandran plot.

EPR spectroscopy

Continuous wave X-band frozen solution EPR spectra of single sample of 0.2 to 0.6 mM solutions of Cu(II)-*Ls*(AA9)A and with 2-fold excess of G6 oligosaccharide or 25 mg/mL PASC (in 10% *v/v* glycerol) at pH 6.0 (20 mM MES buffer, 200 mM NaCl or 20 mM phosphate buffer) and 150 K were acquired on a Bruker EMX spectrometer operating at ~9.30 GHz, with a modulation amplitude of 4 G and microwave power of 10.02 mW. Chloride containing samples were prepared by the addition of an equimolar amount of CuCl₂ prepared in MilliQ water to the protein. The sample was then buffer exchanged on a HiLoad 16/600 Superdex 75 column (GE Healthcare) into 20 mM MES pH 6, 200 mM NaCl to remove any excess copper. For chloride free samples *Ls*(AA9)A was copper loaded with an equimolar amount of CuSO₄ before being buffer exchanged into 20 mM sodium phosphate buffer pH 6 on a HiLoad 16/600 Superdex 75 column (GE Healthcare). Spectral simulations were carried out using EasySpin 5.0.0 integrated into MATLAB R2014a software on a desktop PC. Simulation parameters are given in **Supplementary Table 6**. g_z and $|A_z|$ values were determined accurately from the three absorptions at low field. It was assumed that g and A tensors were axially coincident. Accurate determination of the g_x , g_y , $|A_x|$ and $|A_y|$ was not possible due to the second order nature of the perpendicular region, although it was noted that satisfactory simulation could only be achieved with one particular set of g values. The simulation of *Ls*(AA9)A-G6 spectrum at low chloride conditions was performed after subtraction of the normalized *Ls*(AA9)A spectrum recorded in the presence of 200 mM NaCl (~15% of the chloride-bound species was present). Spectra and simulations are shown in **Supplementary Figure 6**.

¹H HYSCORE

Hyperfine sublevel correlation (HYSCORE) spectra were collected on *Ls*(AA9)A and *Ls*(AA9)A-G6 samples at 1.2 and 3.2 mM protein concentration, respectively, (the high concentration sample showed a pale blue color) with 2-fold excess G6 oligosaccharide (in 10% v/v glycerol) at pH 6.0 (20 mM MES buffer, 200 mM NaCl) on a Bruker ElexSys E580 spectrometer. *Ls*(AA9)A was copper loaded as described in the EPR section. Standard dielectric ring Bruker EPR cavities (ER 4118X-MD5 and EN 4118X-MD4) equipped with an Oxford CF 935 helium flow cryostat were used. The ¹H HYSCORE spectra were recorded employing the sequence $\pi/2 - \tau - \pi/2 - t_1 - \pi - t_2 - \pi/2 - \tau - \text{echo}$ with microwave pulses of length $t_{\pi/2} = 16$ ns and $t_{\pi} = 32$ ns, $\tau = 136$ ns, starting times $t_{1,2} = 100$ ns and time increments $\Delta t_{1,2} = 20 - 32$ ns at 15 K. The intensity of the inverted echo following the fourth pulse is measured with t_1 and t_2 varied and constant τ . Unwanted features from the experimental echo envelopes were removed by using a four-step phase cycle.⁵³ 256 data points were collected in both dimensions. The relaxation decay was subtracted by baseline corrections (fitting by polynomials of 3-6 degree) in both time domains, subsequently applying apodization (Hamming window) and zero-filling to 1024 data points in both dimensions. After 2D fast Fourier transformation, absolute value spectra were simulated using EasySpin 5.0.0 integrated into MATLAB R2014a software. (**Supplementary Note 4** and **Supplementary Fig. 7**)

Isothermal Titration Calorimetry

Isothermal titration calorimetry was performed using an Auto-ITC microcalorimeter (Malvern). Typically protein was present in the cell at 50 μ M with a ten-fold more

concentrated solution of G6 (Seikagaku Biobusiness) in the syringe. Titrations were performed at 25 °C with an initial 0.5 µL injection which is discarded in the data analysis, followed by 18 2 µL injections with a 2 minute interval between each injection. The chloride containing buffer used was 20 mM MES pH 6.0, 200 mM NaCl. Copper loading of Ls(AA9)A was carried out as for the EPR spectroscopy sample. G6 for the syringe was prepared by dissolving solid G6 in the exact same buffer as used for size exclusion chromatography in the last step of protein preparation for experiments in both the presence and absence of chloride. All data were analyzed using the Origin 7 software package (MicroCal). Heats of dilution were subtracted from the data for the titration in the presence of chloride. The analysis typically returned a stoichiometry of between 0.7 and 0.9 for titration in the presence of chloride, probably due to inaccuracies in the measurement of small masses of G6. The G6 concentration was therefore adjusted in the software assuming a 1:1 stoichiometry of G6 to protein.

References

1. Himmel, M.E. *et al.* Biomass recalcitrance: engineering plants and enzymes for biofuels production. *Science* **315**, 804-807 (2007).
2. Sheldon, R.A. Green and sustainable manufacture of chemicals from biomass: state of the art. *Green Chem.* **16**, 950-963 (2014).
3. Payne, C.M. *et al.* Fungal cellulases. *Chem. Rev.* **115**, 1308-1448 (2015).
4. Beeson, W.T. *et al.* Cellulose degradation by polysaccharide monooxygenases. *Annu. Rev. of Biochem.* **84**, 923-946 (2015).
5. Horn, S.J., Vaaje-Kolstad, G., Westereng, B. & Eijsink, V.G.H. Novel enzymes for the degradation of cellulose. *Biotechnol. Biofuels* **5**, 45 (2012).
6. Hemsworth, G.R., Henrissat, B., Davies, G.J. & Walton, P.H. Discovery of a new family of lytic polysaccharide mono-oxygenases. *Nature Chem. Biol.* **10**, 122-126 (2014).
7. Lo Leggio, L. *et al.* Structure and boosting activity of a starch-degrading lytic polysaccharide monooxygenase. *Nature Commun.* **6**, 5961 (2015).
8. Cantarel, B.L. *et al.* The carbohydrate-active enzymes database (CAZy): an expert resource for glycogenomics. *Nucleic Acids Res.* **37**, D233-D238 (2009).
9. Lombard, V. *et al.* The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res.* **42**, D490-495 (2014).
10. Merino, S. & Cherry, J. in *Biofuels* Vol. 108 *Advances in Biochemical Engineering/Biotechnology* (ed Lisbeth Olsson) Ch. 66, 95-120 (Springer Berlin Heidelberg, 2007).

11. Harris, P.V. *et al.* Stimulation of lignocellulosic biomass hydrolysis by proteins of glycoside hydrolase family 61: structure and function of a large, enigmatic family. *Biochemistry* **49**, 3305-3316 (2010).
12. Vaaje-Kolstad, G. *et al.* An oxidative enzyme boosting the enzymatic conversion of recalcitrant polysaccharides. *Science* **330**, 219-222 (2010).
13. Forsberg, Z. *et al.* Cleavage of cellulose by a CBM33 protein. *Protein Sci.* **20**, 1479-1483 (2011).
14. Quinlan, R.J. *et al.* Insights into the oxidative degradation of cellulose by a copper metalloenzyme that exploits biomass components. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 15079-15084 (2011).
15. Phillips, C.M., Beeson, W.T., Cate, J.H. & Marletta, M.A. Cellobiose Dehydrogenase and a Copper-Dependent Polysaccharide Monooxygenase Potentiate Cellulose Degradation by *Neurospora crassa*. *ACS Chemical Biology* **6**, 1399-1406 (2011).
16. Hemsworth, G.R., Davies, G.J. & Walton, P.H. Recent insights into copper-containing lytic polysaccharide mono-oxygenases. *Curr. Opin. Struct. Biol.* **23**, 660-668 (2013).
17. Aachmann, F.L. *et al.* in *Encyclopedia of Inorganic and Bioinorganic Chemistry* (Wiley, 2015).
18. Kjaergaard, C.H. *et al.* Spectroscopic and computational insight into the activation of O₂ by the mononuclear Cu center in polysaccharide monooxygenases. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 8797-8802 (2014).
19. Kim, S. *et al.* Quantum mechanical calculations suggest that lytic polysaccharide monooxygenases use a copper-oxy, oxygen-rebound mechanism. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 149-154 (2014).
20. Gao, J., Thomas, D.A., Sohn, C.H. & Beauchamp, J.L. Biomimetic reagents for the selective free radical and acid-base chemistry of glycans: application to glycan structure determination by mass spectrometry. *J. Am. Chem. Soc.* **135**, 10684-10692 (2013).
21. Eriksson, K.-E., Pettersson, B. & Westermark, U. Oxidation: an important enzyme reaction in fungal degradation of cellulose. *FEBS Lett.* **49**, 282-285 (1974).
22. Hemsworth, G.R. *et al.* The copper active site of CBM33 polysaccharide oxygenases. *J. Am. Chem. Soc.* **135**, 6069-6077 (2013).
23. Li, X. *et al.* Structural basis for substrate targeting and catalysis by fungal polysaccharide monooxygenases. *Structure* **20**, 1051-1061 (2012).
24. Agger, J.W. *et al.* Discovery of LPMO activity on hemicelluloses shows the importance of oxidative processes in plant cell wall degradation. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 6287-6292 (2014).
25. Borisova, A.S. *et al.* Structural and functional characterization of a lytic polysaccharide monooxygenase with broad substrate specificity. *J. Biol. Chem.* **290**, 22955-22969 (2015).
26. Isaksen, T. *et al.* A C4-oxidizing lytic polysaccharide monooxygenase cleaving both cellulose and cello-oligosaccharides. *J. Biol. Chem.* **289**, 2632-2642 (2014).
27. Bennati-Granier, C. *et al.* Substrate specificity and regioselectivity of fungal AA9 lytic polysaccharide monooxygenases secreted by *Podospora anserina*. *Biotechnol. Biofuels* **8**, 90 (2015).
28. Loose, J.S.M. *et al.* A rapid quantitative activity assay shows that the *Vibrio cholerae* colonization factor GbpA is an active lytic polysaccharide monooxygenase. *FEBS Lett.* **588**, 3435-3440 (2014).
29. Gudmundsson, M. *et al.* Structural and electronic snapshots during the transition from a Cu(II) to Cu(I) metal center of a lytic polysaccharide monooxygenase by X-ray photoreduction. *J. Biol. Chem.* **289**, 18782-18792 (2014).
30. Asensio, J.L., Ardá, A., Cañada, F.J. & Jiménez-Barbero, J. Carbohydrate-aromatic interactions. *Acc. Chem. Res.* (2012).
31. Nishio, M. The CH/π hydrogen bond in chemistry. Conformation, supramolecules, optical resolution and interactions involving carbohydrates. *Phys. Chem. Chem. Phys.* **13**, 13873-13900 (2011).
32. Davies, G.J., Planas, A. & Rovira, C. Conformational analyses of the reaction coordinate of glycosidases. *Acc. Chem. Res.* **45**, 308-316 (2012).
33. Lee, J.Y. & Karlin, K.D. Elaboration of copper-oxygen mediated C-H activation chemistry in consideration of future fuel and feedstock generation. *Curr. Opin. Chem. Biol.* **25**, 184-193 (2015).

34. Iwaizumi, M., Kudo, T. & Kita, S. Correlation between the hyperfine coupling constants of donor nitrogens and the structures of the first coordination sphere in copper complexes as studied by nitrogen-14 ENDOR spectroscopy. *Inorg. Chem.* **25**, 1546-1550 (1986).
35. Dhar, D. & Tolman, W.B. Hydrogen atom abstraction from hydrocarbons by a copper(III)-hydroxide complex. *J. Am. Chem. Soc.* **137**, 1322-1329 (2015).
36. Karlin, K.D. & Itoh, S. *Copper-oxygen chemistry*. (Wiley, 2011).
37. Donoghue, P.J. *et al.* Rapid C–H bond activation by a monocopper(III)–hydroxide complex. *J. Am. Chem. Soc.* **133**, 17602-17605 (2011).
38. Wu, M. *et al.* Crystal structure and computational characterization of the lytic polysaccharide monoxygenase GH61D from the basidiomycota fungus *Phanerochaete chrysosporium*. *J. Biol. Chem.* **288**, 12828-12839 (2013).
39. Singh, S.K. & Das, A. The n - π^* interaction: a rapidly emerging non-covalent interaction. *Phys. Chem. Chem. Phys.* **17**, 9596-9612 (2015).
40. Egli, M. & Gessner, R.V. Stereoelectronic effects of deoxyribose O4' on DNA conformation. *Proc. Natl. Acad. Sci. U.S.A.* **92**, 180-184 (1995).
41. Pavlakos, I. *et al.* Noncovalent lone pair...($\text{No-}\pi!$)-heteroarene interactions: the Janus-faced hydroxy group. *Angew. Chem. Int. Ed.* **54**, 8169-8174 (2015).
42. Roeser, D. *et al.* A general binding mechanism for all human sulfatases by the formylglycine-generating enzyme. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 81-86 (2006).
43. Spodsberg, N. *et al.* Polypeptides having cellulolytic enhancing activity and polynucleotides encoding same. Novozymes A/S patent Patent WO2014066141 A2 (2014).
44. Jünnemann, J., Thiem, J. & Pedersen, C. Facile synthesis of acetylated glycosyl fluorides derived from di- and tri-saccharides. *Carbohydr. Res.* **249**, 91-94 (1993).
45. Rencurosi, A., Poletti, L., Panza, L. & Lay, L. Improvement on lipase catalysed regioselective O-acylation of lactose: a convenient route to 2'-O-fucosyllactose. *J. Carbohydr. Chem.* **20**, 761-765 (2001).
46. Fort, S. *et al.* Highly Efficient Synthesis of $\beta(1 \rightarrow 4)$ -Oligo- and -Polysaccharides Using a Mutant Cellulase. *J. Am. Chem. Soc.* **122**, 5429-5437 (2000).
47. Boyer, V. *et al.* Chemoenzymatic synthesis of a bifunctionalized cellohexaoside as a specific substrate for the sensitive assay of cellulase by fluorescence quenching. *Chem. Eur. J.* **8**, 1389-1394 (2002).
48. Liu, Y. *et al.* Use of a fluorescence plate reader for measuring kinetic parameters with inner filter effect correction. *Anal. Biochem.* **267**, 331-335 (1999).
49. Kabsch, W. XDS. *Acta Crystallogr.* **D66**, 125-132 (2010).
50. Vagin, A. & Teplyakov, A. Molecular replacement with MOLREP. *Acta Crystallogr.* **D66**, 22-25 (2010).
51. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr.* **D60**, 2126-2132 (2004).
52. Vagin, A.A. *et al.* REFMAC5 dictionary: organization of prior chemical knowledge and guidelines for its use. *Acta Crystallogr.* **D60**, 2184-2195 (2004).
53. Gemperle, C., Aebli, G., Schweiger, A. & Ernst, R.R. Phase cycling in pulse EPR. *J. Magn. Reson.* (1969) **88**, 241-256 (1990).