



UNIVERSITY OF LEEDS

This is a repository copy of *International cancer seminars: a focus on kidney cancer*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/98916/>

Version: Accepted Version

Article:

Scelo, G, Hofman, JN, Banks, RE et al. (26 more authors) (2016) International cancer seminars: a focus on kidney cancer. *Annals of Oncology*, 27 (8). pp. 1365-1366. ISSN 0923-7534

<https://doi.org/10.1093/annonc/mdw186>

© The Author 2016. Published by Oxford University Press on behalf of the European Society for Medical Oncology. All rights reserved. This is a pre-copyedited, author-produced PDF of an article accepted for publication in *Annals of Oncology* following peer review. The version of record: Scelo, G, Hofman, JN, Banks, RE, Bigot, P, Bhatt, RS, Cancel-Tassin, G, Chew, SK, Creighton, CJ, Cussenot, O, Davis, IJ, Escudier, B, Frayling, TM, Häggström, C, Hildebrandt, MAT, Holcatova, I, Johansson, M, Linehan, M, McDermott, DF, Nathanson, KL, Ogawa, S, Perlman, EJ, Purdue, MP, Stattin, P, Swanton, C, Vasudev, NS, Wu, X, Znaor, A, Brennan, P and Chanock, SJ (2016) International cancer seminars: a focus on kidney cancer. *Annals of Oncology*. ISSN 0923-7534 is available online at: <http://dx.doi.org/10.1093/annonc/mdw186>. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



International cancer seminars – A focus on kidney cancer

| | |
|-------------------------------|--|
| Journal: | <i>Annals of Oncology</i> |
| Manuscript ID | Draft |
| Manuscript Type: | Original Article |
| Date Submitted by the Author: | n/a |
| Complete List of Authors: | <p>Scélo, Ghislaine; International Agency for Research on Cancer, Genetic Epidemiology Group</p> <p>Hofmann, Jonathan; National Cancer Institute, National Institutes of Health, Division of Cancer Epidemiology and Genetics</p> <p>Banks, R.; Leeds Institute for Cancer Studies and Pathology, St James's University Hospital, Clinical and Biomedical Proteomics Group, Cancer Research UK Centre</p> <p>Bigot, Pierre; Centre Hospitalier Universitaire d'Angers, Department of Urology</p> <p>Bhatt, Rupal; Harvard Medical School, Department of Medicine</p> <p>Cancel-Tassin, Géraldine; Centre de Recherche sur les Pathologies Prostatiques et Urologiques (CeRePP), Groupe de Recherche GRC - UPMC n°5</p> <p>Chew, Su Kit; UCL Cancer Institute, University College London, Translational Cancer Therapeutics Laboratory</p> <p>Creighton, Chad; Baylor College of Medicine, Duncan Cancer Center-Biostatistics</p> <p>Cussenot, Olivier; Centre de Recherche sur les Pathologies Prostatiques et Urologiques (CeRePP), Groupe de Recherche GRC - UPMC n°5</p> <p>Davis, Ian; UNC School of Medicine, Department of Genetics</p> <p>Escudier, Bernard; Institut Gustave Roussy, Department of Medical Oncology</p> <p>Frayling, Timothy; University of Exeter Medical School, N/A</p> <p>Häggström, Christel; Umeå University, Department of Surgical and Perioperative Sciences, Urology and Andrology, and Department of Biobank Research</p> <p>Hildebrandt, Michelle; MD Anderson Cancer Center, Department of Epidemiology, Division of OVP, Cancer Prevention and Population Sciences</p> <p>Holcatova, I.; Charles University in Prague, Institute of Hygiene and Epidemiology, 1st Faculty of Medicine</p> <p>Johansson, Mattias; International Agency for Research on Cancer (IARC), Section of Genetics</p> <p>Linehan, Marston; Center for Cancer Research, National Cancer Institute,</p> |

| | |
|-----------|--|
| | <p>Urologic Oncology Branch McDermott, David; Beth Israel Deaconess Medical Center, Hematology/Oncology Nathanson, Katherine; University of Pennsylvania, Department of Medicine Ogawa, Seishi; University of Tokyo, Graduate School of Medicine Perlman, Elizabeth; Northwestern University Feinberg School of Medicine, Robert H. Lurie Comprehensive Cancer Center, Department of Pathology Purdue, Mark; National Cancer Institute, National Institutes of Health, Division of Cancer Epidemiology and Genetics Stattin, Pär; Umeå University, Department of Surgical and Perioperative Sciences Swanton, Charles; University College London Hospitals and Cancer Institute, N/A Vasudev, Naveen; Leeds Institute for Cancer Studies and Pathology, St James's University Hospital, Clinical and Biomedical Proteomics Group, Cancer Research UK Centre Wu, Xifeng; UT MD Anderson Cancer Center, Epidemiology Znaor, Ariana; International Agency for Research on Cancer (IARC), Section of Cancer Surveillance Brennan, Paul; International Agency for Research on Cancer (IARC), Section of Genetics Chanock, Stephen; National Cancer Institute, National Institutes of Health, Division of Cancer Epidemiology and Genetics</p> |
| Keywords: | kidney cancer, genomic, population, clinical research |
| Abstract: | <p>Recent years have seen important advances in our understanding of the etiology, biology and genetics of kidney cancer, some of which have been accompanied by impressive clinical advances. While these have occurred at a time when the incidence of kidney cancer among adults continues to increase in North America and most parts of Europe, elsewhere globally, rates remain stable. In order to summarize important achievements and identify prominent research questions that remain for kidney cancer, a workshop was organized by the International Agency for Research on Cancer (IARC) and the US National Cancer Institute (NCI) in the Spring of 2015. Based on a review of major themes in population, genomic and clinical research, a series of 'difficult questions' were formulated, which should be given future priority within each of these areas.</p> |
| | |

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article type: Special Article

International cancer seminars – A focus on kidney cancer

Ghislaine Scelo¹, Jonathan N. Hofmann², Rosamonde E. Banks³, Pierre Bigot⁴, Rupal S. Bhatt⁵, Géraldine Cancel-Tassin⁶, Su Kit Chew⁷, Chad J. Creighton⁸, Olivier Cussenot⁶, Ian J. Davis⁹, Bernard Escudier¹⁰, Timothy M. Frayling¹¹, Christel Häggström^{12,13}, Michelle A.T. Hildebrandt¹⁴, Ivana Holcatova¹⁵, Mattias Johansson¹, W. Marston Linehan¹⁶, David F. McDermott¹⁷, Katherine L. Nathanson¹⁸, Seishi Ogawa¹⁹, Elizabeth J. Perlman²⁰, Mark P. Purdue², Pär Stattin¹², Charles Swanton²¹, Naveen S. Vasudev³, Xifeng Wu²², Ariana Znaor¹, Paul Brennan^{1,*}, Stephen J. Chanock^{2,*}

¹ International Agency for Research on Cancer, Lyon, France

² Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Department of Health and Human Services, Bethesda, MD, USA

³ Clinical and Biomedical Proteomics Group, Cancer Research UK Centre, Leeds Institute for Cancer Studies and Pathology, St James's University Hospital, Leeds, United Kingdom

⁴ Department of Urology, Centre Hospitalier Universitaire d'Angers, Angers, France

⁵ Department of Medicine, Harvard Medical School, Boston, MA, USA

⁶ Groupe de Recherche GRC - UPMC n°5, Centre de Recherche sur les Pathologies Prostatiques et Urologiques (CeRePP), Paris, France

⁷ Translational Cancer Therapeutics Laboratory, UCL Cancer Institute, University College London, United Kingdom

⁸ Duncan Cancer Center-Biostatistics, Baylor College of Medicine, Houston, TX, USA

⁹ Department of Genetics, UNC School of Medicine, Chapel Hill, NC, USA

¹⁰ Department of Medical Oncology, Institut Gustave Roussy, Villejuif, France

¹¹ University of Exeter Medical School, Exeter, United Kingdom

¹² Department of Surgical and Perioperative Sciences, Urology and Andrology, Umeå University, Umeå, Sweden

¹³ Department of Biobank Research, Umeå University, Umeå, Sweden

¹⁴ Department of Epidemiology, Division of OVP, Cancer Prevention and Population Sciences, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

¹⁵ Institute of Public Health and Preventive Medicine, Charles University, 2nd Faculty of Medicine, Prague, Czech Republic

¹⁶ Urologic Oncology Branch, Center for Cancer Research, National Cancer Institute, Bethesda, MD, USA

¹⁷ Division of Hematology/Oncology, Beth Israel Deaconess Medical Center, Kidney Cancer Program, Dana-Farber/Harvard Cancer Center, Boston, MA, USA

¹⁸ Department of Medicine, University of Pennsylvania, Philadelphia, PA, USA

¹⁹ Graduate School of Medicine, University of Tokyo, Tokyo, Japan

²⁰ Department of Pathology, Northwestern University Feinberg School of Medicine, Robert H. Lurie Comprehensive Cancer Center, Chicago, IL, USA

²¹ University College London Hospitals and Cancer Institute, Huntley Street, London, United Kingdom

²² Department of Epidemiology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

*Correspondence to:

Dr Paul Brennan, International Agency for Research on Cancer, 150 cours Albert Thomas, 69372 Lyon cedex 08, France. Tel: +33 4 7273 8391, E-mail: brennanp@iarc.fr;

or

Stephen J. Chanock, Division of Cancer Epidemiology and Genetics, National Cancer Institute, 9609 Medical Center Dr. Rm. 7E412, MSC 9776, Bethesda, MD 20892-9776 (US Postal Service), Rockville, MD 20850 (courier services). Tel: +1 (240) 276-7150. E-mail: chanocks@mail.nih.gov;

Introduction (131)

Recent years have seen important advances in our understanding of the etiology, biology and genetics of kidney cancer, some of which have been accompanied by impressive clinical advances. While these have occurred at a time when the incidence of kidney cancer among adults continues to increase in North America and most parts of Europe, elsewhere globally, rates remain stable. In order to summarize important achievements and identify prominent research questions that remain for kidney cancer, a workshop was organized by the International Agency for Research on Cancer (IARC) and the US National Cancer Institute (NCI) in the Spring of 2015. Based on a review of major themes in population, genomic and clinical research, a series of 'difficult questions' were formulated, which should be given future priority within each of these areas.

Overview (1055):

Worldwide, it is estimated that there are over 300,000 new cases of kidney cancer per year. Rates are generally lower in most parts of Asia and Latin America when compared to Europe and North America.¹ In the United States, kidney cancer incidence varies across distinct populations, with rates highest among African Americans and lowest among Asian Americans. In Europe, a particularly notable feature is the strikingly high rate observed in the Czech Republic, with elevated rates observed in surrounding regions including Slovakia, the Baltic countries, eastern Germany and Northern Italy,² yet there is no evident explanation to account for this geographic pattern. There is also a consistent male to female excess of kidney cancer observed in both low and high incidence regions that is similarly not explained. Understanding these basic epidemiologic patterns of kidney cancer across the world could lead to insights into the biological mechanisms that contribute to this malignancy. Genetics and differences in detection of asymptomatic cases from diagnostic imaging could partially explain these geographic and ethnic discrepancies, although it is also likely that exposures and lifestyle choices contribute. The main established risk factors of kidney cancer are elevated body-mass index (BMI), hypertension, and tobacco use,³ which have moderate estimated effects.

Kidney cancer includes a spectrum of pathologies that can vary substantively, both with respect to rare inherited and sporadic forms, although clear cell renal cell carcinoma (ccRCC) is the most common. Family and population-based studies have successfully identified key components of the underlying genetic architecture of susceptibility to kidney cancer. Family studies have uncovered a series of rare but highly penetrant mutations, such as in the *VHL* gene, which not only have proven to be useful for screening and counseling in families laden with kidney cancer across generations but also for highlighting the importance of the VHL pathway in sporadic ccRCC.⁴ Other rare syndromes have revealed important insights into additional key genes for different kidney cancer subtypes and together have emerged as a panel for testing in kidney cancer family screening.⁵ Using genome-wide association studies, it has been possible to detect a fraction of the common variants conferring susceptibility to kidney cancer; these possess smaller estimated effects and tend to fall in regulatory regions that have an impact on redundant, key pathways related to kidney cancer development.⁶⁻⁹ Building on the knowledge of rare mutations and common variants, it is now possible to begin to build a polygenic risk model that will predict the familial risk for family members of a newly diagnosed case.

The application of new genomic technologies to investigate the spectrum of somatic alterations in kidney cancer has led to new insights into a series of events, which could lead to new

1
2
3 therapeutics and possibly earlier detection of kidney cancer. Large-scale sequence analyses of
4 sets of adult and pediatric kidney cancers have provided a view of the landscape of the more
5 common types of events across the genome.¹⁰⁻²² Adult kidney cancers harbor large numbers of
6 somatic mutations of all classes, from point mutations to structural alterations, whereas in
7 pediatric kidney cancers (e.g., Wilms tumor and at least half a dozen rarer types), there are
8 fewer somatic events and in fact, few instances in which a driver mutation has been identified.
9 Loss of the 3p chromosomal arm, which harbors the *VHL* gene, is commonly observed in
10 ccRCC, underscoring the importance of the VHL network and its effect on key metabolic
11 pathways in kidney carcinogenesis. VHL alteration also is the likely reason that VEGF pathway
12 directed anti-angiogenic strategies have shown some success. New drivers of kidney cancer
13 have been identified, partly on the basis of the frequency of mutations in key genes, and partly
14 on the basis of corroborative laboratory work.²³
15
16

17
18 Characterization of rarer types of kidney cancer has led to new insights, in some instances
19 providing opportunities to understand how germline mutations inform an understanding of the
20 somatic alterations. Similarly, initial studies have revealed the importance of the dynamic
21 genome, known as the epigenome in both adult and in particular, pediatric subtypes of kidney
22 cancer.²⁴ In this regard, the microenvironment has emerged as a critical focus for ongoing
23 research in order to understand how and in what way the ongoing interaction between host
24 factors and developing tumors either accelerates or inhibits tumor formation.
25

26
27 An important characteristic of kidney carcinogenesis is the role of alterations in metabolic
28 pathways, beginning with the VHL network²⁵. Germline genetic susceptibility (e.g., both common
29 variants and highly penetrant familial mutations) have pointed towards *VHL* and associated
30 genes, such as *EPAS1*. Metabolic disruption can directly contribute to kidney cancer
31 development, both in genomic studies and in laboratory investigations of models, tissue culture
32 and tumor tissue studies. In parallel with these seminal discoveries has been a strong interest in
33 characterizing the role of an elevated BMI, perhaps through a range of metabolic syndromes or
34 disruptions.
35

36
37 The discovery of stable and reproducible biomarkers for risk or earlier diagnosis of kidney
38 cancer using germline genetics or serum/urinary biomarkers remains a daunting challenge.
39 Many small studies have provided preliminary observations but in larger data sets, the utility of
40 candidate or pathway analyses remains elusive. Such molecular epidemiologic studies could
41 provide insight into mechanisms of kidney cancer and improve prediction models. Our capacity
42 to achieve these goals will be based on increasing the scope of our understanding of mutations
43 and biomarkers applied to substantively larger investigations, preferably using prospectively
44 collected biospecimens in order to identify markers of risk.
45

46
47 Therapeutic approaches to kidney cancer have blossomed in the last decade, fueled by new
48 targeted therapies (e.g., designer drugs that attack specific proteins or pathways critical for
49 angiogenesis and immune regulation). There are more than half a dozen new targeted agents in
50 use for metastatic cases, many showing impressive initial responses, but resistance is invariably
51 observed which limits the ability to impact long-term survival.²⁶ Proteomic and circulating
52 biomarker studies have provided promising leads for understanding determinants of tumor
53 progression and metastases, a major problem in kidney cancer. Immunotherapy for kidney
54 cancer has progressed from cytokine based therapies to PD-1 and PDL-1 blockade. Many
55 exciting reports have detailed successful applications of immunotherapy, with a recent phase 3
56 trial showing longer survival in patients receiving this new treatment.²⁷ More studies will be
57 needed to advance this exciting component of precision medicine, particularly as it relates to
58 selection of patients with a high probability of response to therapy.
59
60

Future directions (929):

The exciting trends in molecular characterization and targeted therapy of kidney cancer should be matched with a new commitment to understanding the etiology and pathogenesis of this malignancy. How and why there are major differences by sex, race, geography and exposure histories underscores the complexity of kidney cancer etiology. Global assessment of incidence but also outcomes of therapeutic approaches could reveal important observations that, in turn could inform both the underlying biology and identify new targets for early detection or therapy. The shift towards molecular characterization of adult and pediatric cancers has uncovered a series of important questions related to the key pathways underlying the spectrum of kidney cancer. Heterogeneity in somatic changes has emerged as a significant challenge in kidney cancer, revealing that distinct cancers within an individual may have to be targeted in parallel, unless there is an Achilles' heel that can be targeted upstream.²⁸ Moreover, the analysis of somatic signatures could reveal important clues to environmental and lifestyle factors that contribute to kidney cancer.^{16,29}

Over the past two decades, the detection of germline susceptibility alleles, both common and rare, has accelerated our understanding of the underlying genetic architecture of how kidney cancer develops. The new tools of genome-wide association and next generation sequencing studies should continue to be applied to progressively larger and better-annotated data sets to further add to the comprehensive catalog of susceptibility alleles. The utility of screening subsets of newly diagnosed cases of kidney cancers has already emerged as a compelling argument and merits further investigation, particularly in families or outliers (such as younger cases of ccRCC or the spectrum of rare subtypes). Similar commitment to the genetics of pediatric forms of kidney cancer could be useful for future screening approaches, especially in high-risk settings. Special attention should be given to conducting these studies in populations with different genetic histories and exposures. The ability to examine gene-environment interactions either directly or through the approach of Mendelian randomization should be a high priority for discovery of new relationships, some of which could be harnessed for detection or prevention.³⁰

Further characterization of the genomic landscape of kidney cancers, including larger numbers of rarer cancers, should provide an important foundation for identifying the catalog of driver events. While The Cancer Genome Atlas (TCGA) and International Cancer Genome Consortium (ICGC) have provided an impetus to conduct large scale characterization projects, the international community should develop a shared data resource to combine academic, clinical and research resources to rapidly accelerate the bioinformatics analyses of the spectrum of kidney cancer. So far, the number of samples studied has led to the identification of genes that are commonly mutated. As kidney cancer has emerged as one of the pivotal examples of tumor heterogeneity, together with its spectrum of rare and common subtypes, many more tumor/normal pairs will require characterization in order to capture key events, both for targets and prognostic factors.³¹ The next generation of genomic characterization efforts should be conducted in studies that have collected both epidemiologic risk factors, including pre-diagnostic biomarkers when possible, and extensive annotation of the clinical parameters including long-term outcome. The investigation of somatic alterations with clinical outcomes should be a high priority, including detailed analyses of tumor heterogeneity, which might require regional sampling guided by imaging. The challenge of heterogeneity has important implications for therapeutic decisions, and perhaps in the future therapeutic monitoring, especially if circulating tumor DNA or tumor cells can be adapted as an efficient and widespread technique. Current approaches to develop models for the evolution of kidney cancer represent an important parallel approach, one that should be iteratively coordinated with clinical characterization of the cancer

1
2
3 genome and its epigenomic changes.²⁸ Serial and regional biopsies in well-characterized
4 biorepositories should be available and include detailed follow-up on outcomes, therapeutic
5 interventions and new imaging techniques. A critical question of the future will be how and in
6 what way imaging-based techniques can be harmonized with genomic technologies to identify
7 targets, but equally new target identification will need to be combined with the parallel
8 development of predictive genomic and proteomic biomarkers to enable patient stratification.
9

10
11 The rapid acceleration of immune-based therapies should remain a major focus of kidney
12 cancer research, providing new insights into how and in what way the immune system could
13 better conduct surveillance against emerging cancers and to treat kidney cancer. The collective
14 assessment of the different strategies could shed light on prognostic factors, such as the HLA
15 haplotypes and circulating biomarkers. Eventually, the use of immunotherapy may be a central
16 component in the armamentarium of precision medicine tools. Determinants of immune-based
17 therapeutic success should provide new insights into stratification of individual patients, perhaps
18 based on emerging immune profiles of both measured biomarkers and genetic predisposition.
19

20
21 In conclusion, the tools of genomic characterization and targeted therapy have begun to
22 accelerate the investigation of the basic biology of kidney cancer, thus providing novel
23 approaches towards improving early detection, intervention and monitoring of kidney cancer.
24 The development of a more refined molecular taxonomy of the spectrum of kidney cancer will
25 advance most efficiently if the community continues to develop more robust approaches to data
26 sharing. Interoperability between data resources should remain a central goal both for resources
27 generated by “-omic” technologies and more precise phenotyping, based on sound
28 epidemiologic and clinical practice. By integrating data sets, the opportunity to determine shared
29 and unique features should benefit the diagnosis, treatment and monitoring of rare and common
30 kidney cancers alike. In the distant future, it may emerge that precision prevention strategies are
31 possible, but they will be based on further research in all corners of the globe.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

One dozen difficult questions that should be prioritized:

- 1 What is the explanation for the geographical, racial/ethnic and sex differences in the incidence of kidney cancer?
 - 2 What underlying biologic pathways drive the association between obesity/hypertension, and kidney cancer risk?
 - 3 Can a greater understanding of germline variation of kidney cancer inform us about the unknown lifestyle and environmental causes?
 - 4 Can a better understanding of the somatic signatures (both genomic and proteomic) of kidney cancer and its subtypes provide clues to etiologic risk factors and prognosis?
 - 5 How is response to targeted therapy and immunotherapy influenced by epigenetic variation?
 - 6 With more than a half a dozen approved drugs targeting the VHL/HIF pathway in clear cell renal cell carcinomas, what are the next steps towards improving therapy and survival in both early and advanced cases?
 - 7 What are the barriers for rapid data sharing of sequencing of kidney cancers, which could accelerate identification and characterization of drivers of oncogenesis?
 - 8 How do we select kidney cancer patients for constitutional genetic testing, beyond those fitting a 'classic' kidney cancer susceptibility syndrome?
 - 9 Why do more than half of pediatric Wilms tumors lack identifiable driver mutations and what does this suggest about epigenetic regulation?
 - 10 What new data are critical to develop more accurate models for understanding genomic and epigenetic changes across the spectrum of renal cancers?
 - 11 Are current pathological subtype classifications still clinically relevant?
 - 12 What factors can reliably predict durable response to immunotherapy?
-

Disclosure:

Organization of production of this seminar was funded jointly by the International Agency for Research on Cancer (IARC) and the Division of Cancer Epidemiology and Genetics (DCEG) of the US National Cancer Institute. We disclose no conflicts of interest.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References:

1. Znaor A, Lortet-Tieulent J, Laversanne M, Jemal A, Bray F. International variations and trends in renal cell carcinoma incidence and mortality. *Eur Urol* 2015; **67**(3): 519-30.
2. Li P, Znaor A, Holcatova I, et al. Regional Geographic Variations in Kidney Cancer Incidence Rates in European Countries. *Eur Urol* 2015; **67**(6): 1134-41.
3. Chow WH, Dong LM, Devesa SS. Epidemiology and risk factors for kidney cancer. *Nat Rev Urol* 2010; **7**(5): 245-57.
4. Haas NB, Nathanson KL. Hereditary kidney cancer syndromes. *Adv Chronic Kidney Dis* 2014; **21**(1): 81-90.
5. Linehan WM. Genetic basis of kidney cancer: role of genomics for the development of disease-based therapeutics. *Genome Res* 2012; **22**(11): 2089-100.
6. Purdue MP, Johansson M, Zelenika D, et al. Genome-wide association study of renal cell carcinoma identifies two susceptibility loci on 2p21 and 11q13.3. *Nat Genet* 2011; **43**(1): 60-5.
7. Wu X, Scelo G, Purdue MP, et al. A genome-wide association study identifies a novel susceptibility locus for renal cell carcinoma on 12p11.23. *Hum Mol Genet* 2012; **21**(2): 456-62.
8. Henrion M, Frampton M, Scelo G, et al. Common variation at 2q22.3 (ZEB2) influences the risk of renal cancer. *Hum Mol Genet* 2013; **22**(4): 825-31.
9. Henrion MY, Purdue MP, Scelo G, et al. Common variation at 1q24.1 (ALDH9A1) is a potential risk factor for renal cancer. *PLoS One* 2015; **10**(3): e0122589.
10. TCGA. Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature* 2013; **499**(7456): 43-9.
11. Dalgliesh GL, Furge K, Greenman C, et al. Systematic sequencing of renal carcinoma reveals inactivation of histone modifying genes. *Nature* 2010; **463**(7279): 360-3.
12. Davis CF, Ricketts CJ, Wang M, et al. The somatic genomic landscape of chromophobe renal cell carcinoma. *Cancer Cell* 2014; **26**(3): 319-30.
13. Network TCGAR. Comprehensive Molecular Characterization of Papillary Renal-Cell Carcinoma. *N Engl J Med* 2016; **374**(2): 135-45.
14. Pena-Llopis S, Vega-Rubin-de-Celis S, Liao A, et al. BAP1 loss defines a new class of renal cell carcinoma. *Nat Genet* 2012; **44**(7): 751-9.
15. Sato Y, Yoshizato T, Shiraishi Y, et al. Integrated molecular analysis of clear-cell renal cell carcinoma. *Nat Genet* 2013; **45**(8): 860-7.
16. Scelo G, Riazalhosseini Y, Greger L, et al. Variation in genomic landscape of clear cell renal cell carcinoma across Europe. *Nat Commun* 2014; **5**: 5135.

17. Varela I, Tarpey P, Raine K, et al. Exome sequencing identifies frequent mutation of the SWI/SNF complex gene PBRM1 in renal carcinoma. *Nature* 2011; **469**(7331): 539-42.
18. Lovvorn HN, 3rd, Pierce J, Libes J, et al. Genetic and chromosomal alterations in Kenyan Wilms Tumor. *Genes Chromosomes Cancer* 2015; **54**(11): 702-15.
19. Perlman EJ, Gadd S, Arold ST, et al. MLLT1 YEATS domain mutations in clinically distinctive Favourable Histology Wilms tumours. *Nat Commun* 2015; **6**: 10013.
20. Roy A, Kumar V, Zorman B, et al. Recurrent internal tandem duplications of BCOR in clear cell sarcoma of the kidney. *Nat Commun* 2015; **6**: 8891.
21. Walz AL, Ooms A, Gadd S, et al. Recurrent DGCR8, DROSHA, and SIX homeodomain mutations in favorable histology Wilms tumors. *Cancer Cell* 2015; **27**(2): 286-97.
22. Lawrence MS, Stojanov P, Polak P, et al. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* 2013; **499**(7457): 214-8.
23. Turajlic S, Larkin J, Swanton C. SnapShot: Renal Cell Carcinoma. *Cell* 2015; **163**(6): 1556- e1.
24. la Rosa AH, Acker M, Swain S, Manoharan M. The role of epigenetics in kidney malignancies. *Cent European J Urol* 2015; **68**(2): 157-64.
25. Linehan WM, Srinivasan R, Schmidt LS. The genetic basis of kidney cancer: a metabolic disease. *Nat Rev Urol* 2010; **7**(5): 277-85.
26. Powles T, Staehler M, Ljungberg B, et al. Updated EAU Guidelines for Clear Cell Renal Cancer Patients Who Fail VEGF Targeted Therapy. *Eur Urol* 2016; **69**(1): 4-6.
27. Motzer RJ, Escudier B, McDermott DF, et al. Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma. *N Engl J Med* 2015; **373**(19): 1803-13.
28. Gerlinger M, Rowan AJ, Horswell S, et al. Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N Engl J Med* 2012; **366**(10): 883-92.
29. Alexandrov LB, Nik-Zainal S, Wedge DC, et al. Signatures of mutational processes in human cancer. *Nature* 2013; **500**(7463): 415-21.
30. Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Hum Mol Genet* 2014; **23**(R1): R89-98.
31. Lawrence MS, Stojanov P, Mermel CH, et al. Discovery and saturation analysis of cancer genes across 21 tumour types. *Nature* 2014; **505**(7484): 495-501.