

This is a repository copy of *Proton imaging of an electrostatic field structure formed in laser-produced counter-streaming plasmas*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/99968/>

Version: Published Version

Article:

Morita, T., Kugland, N. L., Wan, W. et al. (27 more authors) (2016) Proton imaging of an electrostatic field structure formed in laser-produced counter-streaming plasmas. *Journal of Physics: Conference Series*. 012071. ISSN 1742-6596

<https://doi.org/10.1088/1742-6596/688/1/012071>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Proton imaging of an electrostatic field structure formed in laser-produced counter-streaming plasmas

This content has been downloaded from IOPscience. Please scroll down to see the full text.

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 144.32.225.118

This content was downloaded on 20/05/2016 at 09:40

Please note that [terms and conditions apply](#).

Proton imaging of an electrostatic field structure formed in laser-produced counter-streaming plasmas

T. Morita¹, N. L. Kugland², W. Wan³, R. Crowston⁴, R. P. Drake³, F. Fiuza², G. Gregori⁵, C. Huntington², T. Ishikawa⁶, M. Koenig⁷, C. Kuranz³, M. C. Levy^{2,8}, D. Martinez², J. Meinecke⁵, F. Miniati⁹, C. D. Murphy¹⁰, A. Pelka⁷, C. Plechaty², R. Presura¹¹, N. Quirós¹¹, B. A. Remington², B. Reville¹², J. S. Ross², D. D. Ryutov², Y. Sakawa¹³, L. Steele¹⁴, H. Takabe¹³, Y. Yamaura⁶, N. Woolsey⁴, H.-S. Park²

1. Department of Energy Engineering Science, Faculty of Engineering Sciences, Kyushu University, 6-1 Kasuga-koen, Kasuga, Fukuoka 816-8580, Japan

2. Lawrence Livermore National Laboratory, P.O. Box 808, Livermore, California 94550, USA

3. Department of Atmospheric, Oceanic, and Space Sciences, University of Michigan, Ann Arbor, MI 48109, USA

4. University of York, York, North Yorkshire YO10 5DD, United Kingdom

5. Department of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, UK

6. Graduate School of Science, Osaka University, 1-1 Machikane-yama, Toyonaka, Osaka 560-0043, Japan

7. Laboratoire pour l'Utilisation des Lasers Intenses (LULI), École Polytechnique-Univ, Paris VI, 91128 Palaiseau, France

8. Rice University, Houston, Texas 77251, USA

9. Physics Department, Wolfgang-Pauli-Strasse 27, ETH-Zürich, CH-8093 Zürich, Switzerland

10. University of Edinburgh, Edinburgh, EH8 9YL, United Kingdom

11. University of Nevada, Reno, Nevada 89557, USA

12. Centre for Plasma Physics, Queen's University Belfast, University Rd, Belfast BT7 1NN, UK

13. Institute of Laser Engineering, Osaka University, 2-6 Yamada-oka, Suita, Osaka 565-0871, Japan

14. University of California, Davis, California 95616, USA

E-mail: morita@aees.kyushu-u.ac.jp

Abstract. We report the measurements of electrostatic field structures associated with an electrostatic shock formed in laser-produced counter-streaming plasmas with proton imaging. The thickness of the electrostatic structure is estimated from proton images with different proton kinetic energies from 4.7 MeV to 10.7 MeV. The width of the transition region is characterized by electron scale length in the laser-produced plasma, suggesting that the field structure is formed due to a collisionless electrostatic shock.

1. Introduction

Shock waves are commonly observed in the universe, space plasmas, and laboratory plasmas. In particular, collisionless shocks play significant roles in particle acceleration, for example, in Earth's bow shock and supernova remnant shocks. In collisionless shocks, collisions between particles do not account for the formation mechanism and particle-field interactions are essential.






Figure 1. The target design and geometry of diagnostics for proton imaging. Two Mylar (CHO) disks were located in parallel separated by 3 mm, and one side of the target was irradiated by the drive laser. The laser-produced plasmas were diagnosed by a proton beam generated by focusing a short-pulse laser on an Au disk placed at 14.5 mm away from the plasma. The protons were detected by the stack of radiochromic films (RCFs) located at 127 mm from the proton source.

Therefore, the shock thickness is much smaller than ion-ion mean-free-path and a large electromagnetic field exists at the shock front. These shocks have been studied by observing emissions from astrophysical objects[1], and by in-situ measurements using satellites in space plasmas[2].

Laboratory experiments with high-power laser systems can be an alternative to observations or in-situ measurements by satellites[3]. Collisionless shocks have been produced and investigated in counter-streaming laser-produced plasmas[4]. They have been measured by optical diagnostics such as interferometry, shadowgraphy, optical pyrometry, and Thomson scattering to obtain the fundamental plasma parameters: density, temperature, charge state, and flow velocity[5]. However, the measurement of an electromagnetic field is indispensable for collisionless shock experiments.

Proton imaging is one of the methods to measure the field structures[6]; it is widely used for measuring the electric and magnetic fields in inertial confinement fusion experiments[7] and the field in laser-produced plasmas[8].

In this paper, we present the measurement of electric field structures formed at a shock with a proton beam generated by irradiating a thin solid foil by a short-pulse laser beam.

2. Experiment

The experiment was performed with the Jupiter laser facility at Lawrence Livermore National Laboratory. Double-disk Mylar (CHO) targets are used to produce counter-streaming plasmas as shown in Fig. 1. Each disk measures 2×2 mm and 0.5 mm in thickness, and two disks are separated by 3 mm. One of the disks (drive disk) was illuminated by a laser beam (drive laser), delivering 120 J on average in a 1.5 ns square pulse at a wavelength of 527 nm, with a focal spot of $500 \mu\text{m}$ in diameter. The other disk (second disk) was ablated by the radiation from the laser-produced plasma early in time, and by the plasma arriving later in time as described in the Refs. [4].

A laser pulse (160 J in 0.7 ps, wavelength of 1057 nm) was focused on an Au disk with a thickness of $25 \mu\text{m}$ to produce high-energy protons for radiography. The distance between the disk and the object plane, where the target was located, was $l = 14.5$ mm. The proton beam images electric and magnetic fields formed in the laser-produced plasmas on radiochromic films located at an imaging plane with a distance $L + l = 127$ mm from the Au disk, where L is the distance between the object and the radiochromic films. The radiochromic films were stacked to detect protons in different energy ranges: 4.7, 7.0, 8.8, and 10.7 MeV with 10% energy width for each film.




Figure 2. (a) The proton image taken at $t = 20$ ns at the proton energy of 4.7 MeV. (b) The line-out at $x \sim 15$ mm [square in (a)].

3. Result

Figure 2(a) shows the proton image at $t = 20$ ns at the proton energy of 4.7 MeV. A caustic structure is observed at $x \sim 15$ mm in front of the second disk. In general, this sudden change in proton intensity indicates a large electromagnetic field[6]. The line-out of the caustic for Fig. 2(a) is shown in Fig. 2(b). The widths of the caustics depend on the proton energy, i.e., high-energy protons make a thinner structure as shown in Fig. 3.

4. Discussion

The observed caustic structure corresponds to a shock front as discussed in previous works[4], and the left- and right-hand sides of this structure are upstream and downstream of a shock, respectively. Here, we assume that the protons are deflected by an electric field not by a magnetic field. If the shock is formed by a self-generated magnetic field due to electromagnetic Weibel instability, the shock thickness should be much larger ($c/\omega_{pi} > 10$ mm [9]) than experimental observation. Also, a spherically symmetric flat-top potential is assumed in the downstream region at the vicinity of a shock.

The position x on the imaging plane is expressed using the position x_0 on the object plane and the deflection angle $\alpha(x_0)$ as $x = L(x_0/l + \alpha(x_0))$ [6]. The deflection angle α of the proton beam by a spherically symmetric flat-top potential is expressed as $\alpha(x_0) = (e\phi_0 x_0 / 2W\sqrt{a\delta})F(\tilde{\xi}) = (e\phi_0(\tilde{\xi}\delta + a) / 2W\sqrt{a\delta})F(\tilde{\xi})$ [6], where $\tilde{\xi} = (x_0 - a)/\delta$, δ is the transition thickness, a is the curvature radius of the potential structure, W is the proton energy, $e\phi$ is the potential energy, and $F(\tilde{\xi}) = (2/\pi) \int_{-\infty}^{\infty} d\eta / ((\eta^2 + \tilde{\xi}^2) + 1)$. In case of caustic formation on the imaging plane, $\partial x / \partial x_0 = 0$ is satisfied. Therefore, $1/l + (e\phi_0 / 2W\delta\sqrt{a\delta})[\delta F(\tilde{\xi}) + (\tilde{\xi}\delta + a)F'(\tilde{\xi})] = 0$. Generally, this equation has two solutions ($\tilde{\xi}_1$ and $\tilde{\xi}_2$), however, the caustic positions $x(\tilde{\xi}_1)$ and $x(\tilde{\xi}_2)$ can not be resolved in this experiment because of small separation between them, and these two caustics are observed as a single peak in the proton intensity. Therefore, the thickness of an electric field at the shock front (δ) is comparable to

$$\delta \approx |x(\tilde{\xi}_2) - x(\tilde{\xi}_1)|. \quad (1)$$

The thicknesses of the caustics in different energy bins are shown in Fig. 3 obtained as full widths at half maximums of the caustics for the energies of 4.7, 7.0, 8.8, and 10.7 MeV. The experimental data are fitted with Eq. (1) for various thicknesses. If this caustic is formed due to an electrostatic shock, the thickness is characterized by electron inertial length $c/\omega_{pe} \sim 2 \mu\text{m}$ [$n_e \sim 5 \times 10^{18} \text{ cm}^{-3}$ evaluated from the interferometry (not shown)][9]. The solid and dotted lines show the fitted results with $\delta = 0.01 \mu\text{m} \sim \lambda_D$, where λ_D is the Debye length, and $0.2 \mu\text{m} \sim c/\omega_{pe}$, respectively. The fitted result with larger thickness of $\delta = 10 \mu\text{m}$ (dotted line) shows




Figure 3. Caustic thicknesses as a function of proton energy. The thickness is evaluated as a full width at half maximum of each line-out as shown in Fig. 2. The best-fit results using various thickness δ are shown with solid ($\delta = 10 \text{ nm} \sim \lambda_D$), dashed ($\delta = 2.0 \text{ } \mu\text{m} \sim c/\omega_{pe}$), and dotted ($\delta = 10 \text{ } \mu\text{m}$) lines.

large deviation from the experimental data, while the results from small thickness $\delta < c/\omega_{pe}$ shows good agreement with the experimental data, indicating that the structure is defined in an electron scale. Note that the ion inertial length is over hundreds of micron and is much larger than the thickness estimated here. Moreover, the ion-ion mean-free-path of counter-streaming plasmas is $\sim 200 \text{ } \mu\text{m}$ [4] using the velocity of $v = 3 \text{ mm}/20 \text{ ns} = 150 \text{ km/s}$. This small thickness of the electric field structure suggests that the caustic is formed by an electrostatic shock in collisionless counter-streaming plasmas.

5. Summary

We measured an electrostatic field structure formed in counter-streaming laser-produced plasmas with proton imaging technique. The thickness of the electrostatic structure is estimated from proton images with different proton kinetic energies from 4.7 MeV to 10.7 MeV. The width of the transition region is comparable to or less than the electron inertial length of laser-produced plasmas, suggesting that the field structure is formed due to a collisionless electrostatic shock.

Acknowledgments

The authors acknowledge the dedicated technical support by the staff at Jupiter laser facility for the laser operation and diagnostics. This work was performed for the U.S. Department of Energy by the Lawrence Livermore National Laboratory, under Contract No. DE-AC52-07NA27344. Support was provided by LLNL LDRD Grant No. 11-ERD-054 and Japan Society for the Promotion of Science (JSPS).

References

- [1] Bamba A, Yamazaki R, Ueno M and Koyama K 2003 *The Astrophysical Journal* **589** 827–837; Aharonian F A, *et al* 2004 *Nature* **432** 75–7; Uchiyama Y, Aharonian F A, Tanaka T, Takahashi T and Maeda Y 2007 *Nature* **449** 576–8;
- [2] Wenzel K, Reinhard R, Sanderson T and Sarris E 1985 *Journal of Geophysical Research* **90** 12–18; Sarris E T, Anagnostopoulos G C and Krimigis S M 1987 *Journal of Geophysical Research* **92** 12083; Anagnostopoulos G C 1994 *Physica Scripta* **T52** 142–151;
- [3] Remington B, Drake R and Ryutov D 2006 *Reviews of Modern Physics* **78** 755–807
- [4] Morita T *et al* *Physics of Plasmas* **17** 122702; Kuramitsu Y *et al* 2011 *Physical Review Letters* **106** 175002
- [5] Ross J S *et al* 2006 *Review of Scientific Instruments* **77** 10E520; Ross J S *et al* 2012 *Physics of Plasmas* **19** 056501
- [6] Kugland N L *et al* 2012 *The Review of scientific instruments* **83** 101301
- [7] Mackinnon A *et al* 2006 *Physical Review Letters* **97** 045001; Rygg J R *et al* 2008 *Science (New York, N.Y.)* **319** 1223–5; Li C K *et al* 2012 *Physical Review Letters* **108** 025001
- [8] Borghesi M *et al* 2002 *Physics of Plasmas* **9** 2214; Nilson P *et al* 2006 *Physical Review Letters* **97** 255001; Romagnani L *et al* 2008 *Physical Review Letters* **101** 025004; Kugland N L *et al* 2012 *Nature Physics* **8** 809–812
- [9] Kato T N and Takabe H 2010 *Physics of Plasmas* **17** 032114; Kato T N *et al* , 2008 *The Astrophysical Journal Letters* **681** 93–96