

This is a repository copy of *Scales of analysis:evidence of fish and fish processing at Star Carr*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/100106/>

Version: Accepted Version

---

**Article:**

Robson, Harry Kenneth [orcid.org/0000-0002-4850-692X](https://orcid.org/0000-0002-4850-692X), Little, Aimee Patrice [orcid.org/0000-0003-4713-4260](https://orcid.org/0000-0003-4713-4260), Jones, Andrew K. G. et al. (12 more authors) (2018) *Scales of analysis:evidence of fish and fish processing at Star Carr*. *Journal of Archaeological Science Reports*. ISSN 2352-409X

<https://doi.org/10.1016/j.jasrep.2016.02.009>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## 1 Scales of analysis: evidence of fish and fish processing at Star Carr

2  
3 Robson, Harry Kenneth<sup>a\*</sup>, Little, Aimée<sup>a</sup>, Jones, Andrew Kenneth George<sup>a</sup>, Blockley,  
4 Simon<sup>b</sup>, Candy, Ian<sup>b</sup>, Matthews, Ian<sup>b</sup>, Palmer, Adrian<sup>b</sup>, Schreve, Danielle<sup>b</sup>, Tong, Emma<sup>a</sup>,  
5 Pomstra, Diederik<sup>c</sup>, Fletcher, Lucie<sup>a</sup>, Hausmann, Niklas<sup>a</sup>, Taylor, Barry<sup>d</sup>, Conneller, Chantal<sup>e</sup>,  
6 and Milner, Nicky<sup>a</sup>

7  
8 <sup>a</sup>Department of Archaeology, The King's Manor, University of York, YO1 7EP, UK

9 <sup>b</sup>Department of Geography, Royal Holloway University of London, Egham, TW20 0EX, UK

10 <sup>c</sup>Faculty of Archaeology, Leiden University, Postbus 9514, 2300 RA, The Netherlands

11 <sup>d</sup>Department of History and Archaeology, University of Chester, Chester, CH1 4BJ, UK

12 <sup>e</sup>Archaeology (SAHC), University of Manchester, Mansfield Cooper Building, Oxford Road,  
13 Manchester M13 9PL, UK

14  
15 \*Corresponding author

16 hkrobson@hotmail.co.uk (H. K. Robson)

### 17 18 **Abstract**

19 This contribution directly relates to the paper published by Wheeler in 1978 entitled ‘Why  
20 were there no fish remains at Star Carr?’. Star Carr is arguably the richest, most studied and  
21 re-interpreted Mesolithic site in Europe but the lack of fish remains has continued to vex  
22 scholars. Judging from other materials, the preservation conditions at the site in the late  
23 1940s/early 1950s should have been good enough to permit the survival of fish remains, and  
24 particularly dentaries of the northern pike (*Esox lucius* L., 1758) as found on other European  
25 sites of this age. The lack of evidence has therefore been attributed to a paucity of fish in the  
26 lake. However, new research has provided multiple lines of evidence, which not only  
27 demonstrate the presence of fish, but also provide evidence for the species present, data on  
28 how and where fish were being processed on site, and interpretations for the fishing methods  
29 that might have been used. This study demonstrates that an integrated approach using a range  
30 of methods at landscape, site and microscopic scales of analysis can elucidate such questions.  
31 In addition, it demonstrates that in future studies, even in cases where physical remains are  
32 lacking, forensic techniques hold significant potential.

33  
34 **Keywords:** Mesolithic; Star Carr; Flixton Island, Fish remains; Seasonality; Use wear

35 **1.0. Introduction**

36

37 *Figure 1: location map of Star Carr.*

38

39 Grahame Clark excavated Star Carr from 1949-1951 (Clark 1954) (Figure 1). His discoveries  
40 have led to what has become known as one of Europe's most famous Mesolithic sites. This  
41 was due to the outstanding preservation of organic remains, including the discovery of a  
42 brushwood platform associated with an extensive faunal assemblage and extremely rare  
43 artefacts such as 'headdresses' made from the crania of red deer (*Cervus elaphus* L., 1758).  
44 Clark noted that:

45

46 'No remains of fish survived. Negative evidence is notoriously dangerous in prehistory, and  
47 never more so than when a substance so perishable as fish-bone is in question. Yet to judge  
48 from what was found on similar sites in different parts of northern Europe, traces might at  
49 least have been expected for the lower jaws of pike, had these been caught. It should be  
50 remembered though that the evidence for pike-fisheries among the later Maglemosian comes  
51 from sites occupied during the summer, in the early months of which the fishing was carried  
52 on with leisters in temperate Europe down to modern times. The absence of pike remains  
53 from Star Carr may therefore be a true reflection of the circumstance that the site was  
54 *abandoned during the summer months*' (Clark 1954, 16, our emphasis).

55

56 In the 1970s and 1980s a number of articles reinterpreted the evidence from Star Carr, in  
57 particular reconsidering the seasonality of the site and importantly, suggesting that the site  
58 had been occupied during the spring and summer (Carter 1998, Caulfield 1978; Jacobi 1978;  
59 Legge and Rowley-Conwy 1988; Mellars and Dark 1998). This overturned Clark's  
60 hypothesis, as set out above, and with new interpretations of summer occupation, it became  
61 even harder to account for a lack of pike at the site.

62

63 In 1978, Wheeler, wrote a seminal paper entitled 'Why were there no fish remains at Star  
64 Carr?'. Importantly, he drew attention to the fact that pike can be fished all year round, which  
65 negated the seasonality argument. Therefore, he suggested that there were probably no fish  
66 present in the lake throughout the course of site occupation. His hypothesis was that fish,  
67 attempting to colonise up the riverine systems, would not have permeated the Lake Flixton

68 basin because the water was too fast flowing at Kirkham Gorge, located roughly 40  
69 kilometres downstream.

70

71 However, Wheeler (1978) did not mention the presence of waterfowl, which can transport  
72 fish spawn via their feet. The Star Carr faunal assemblage contained at least seven species:  
73 white stork (*Ciconia ciconia* L., 1758), common crane (*Grus grus* L., 1758), red-breasted  
74 merganser (*Mergus serrator* L., 1758), red-throated diver (*Gavia stellata* P., 1763), great  
75 crested grebe (*Podiceps cristatus* L., 1758), little grebe (*Podiceps ruficollis* P., 1764) and a  
76 duck of similar size to the pintail (*Anas acuta* L., 1758) (Clark 1954). Thus fish could have  
77 colonised the lake via this method of passive dispersal.

78

79 The only potential (indirect) evidence for fishing is in the form of the barbed points. Clark  
80 found 190 barbed points and 1 harpoon at Star Carr (Figure 2). These were made out of red  
81 deer antler and manufactured so that they could be hafted onto wooden shafts for spearing or  
82 throwing. In some cases they may have been hafted in pairs or with the addition of a central  
83 bone point to provide a leister as has been observed at other sites in Europe: one such pairing  
84 of barbed points was observed by Clark *in situ* (Clark 1954, plate 12).

85

86 *Figure 2: A range of the different types of barbed points/harpoons found at Star Carr including the harpoon in*  
87 *the middle (scale: 5cm).*

88

89 Further evidence for the use of barbed points and harpoons related to fishing practices derives  
90 from a number of other Early Mesolithic ('Maglemosian') sites in north-west Europe:  
91 Holmegård, Lundby, Mullerup, Ulkestrup Lyng, Sværdborg, Vinde-Helsing and Ögaarde (in  
92 Denmark) and Duvensee, Friesack 4, Friesack 27a, Hohen Viecheln and Wustermark (in  
93 northern Germany) (Aaris-Sørensen 1976; Broholm 1924; Clark 1948; Gramsch and Beran  
94 2010; Groß 2014; Jessen *et al.* 2015; Noe-Nygaard 1995; Robson 2015; Rosenlund 1980;  
95 Schuldt 1961). In addition, fish remains were also encountered at the majority of these sites  
96 and are solely pike, or pike dominant. However, wels catfish (*Silurus glanis* L., 1758),  
97 European perch (*Perca fluviatilis* L., 1758), tench (*Tinca tinca* L., 1758), carp (Cyprinidae  
98 sp.), common bream (*Abramis brama* L., 1758), common rudd (*Scardinius erythrophthalmus*  
99 L., 1758) and European eel (*Anguilla anguilla* L., 1758) have also been identified (Aaris-  
100 Sørensen 1976; Broholm 1924; Gramsch and Beran 2010; Groß 2014; Jessen *et al.* 2015;  
101 Noe-Nygaard 1995; Robson 2015; Rosenlund 1980; Wundschedt 1961).

102

103 In addition, there is a close correlation between pike remains and barbed points, similar to  
104 those found at Star Carr. For instance, at Sværdborg, Denmark, 80 upper and 64 lower pike  
105 jaw bones were found along with 274 leister prongs and 11 hooks (Clark 1952, 47). There are  
106 also sites where barbed points have been found in association with pike bones within the lake  
107 bed. Clark (1948, appendix 1) lists Calbe (Germany), Esperöds Mosse (Scania), and Kunda  
108 (Estonia). In two cases at the latter site barbed points were found impaling pike skeletons:  
109 one in the back of a large pike and the other in the skull (Clark 1952, 47).

110

111 In comparison, there is very little evidence in Britain for freshwater fishing, particularly in  
112 the Early Mesolithic. The only comparable example to the European evidence appears to  
113 derive from nearby Holderness: in 1903 an antler harpoon was found at Atwick, East  
114 Yorkshire and in 1932 further investigations were carried at the nearby site at Skipsea by  
115 Godwin and Godwin (1933, 39) who found ‘fragments of *Pinus* bark, fins of pike (*Esox*  
116 *lucius*) and flint artifacts’.

117

118 In the later part of the British Mesolithic evidence for fishing freshwater species exists but  
119 these specimens are not found alongside barbed points: for example a single precaudal  
120 vertebrae of a pike was found at Bouldnor Cliff on the Isle of Wight (Momber *et al.* 2011, 52)  
121 and from the Severn Estuary Mesolithic sites a total of 513 identifiable fragments of fish  
122 were found including Salmonidae (salmon family), eel and a possible cyprinid (Cyprinidae  
123 sp.) as well as coastal species (Bell 2007, 166-168).

124

125 The reason for the lack of fish remains at Star Carr has therefore remained a mystery that has  
126 intrigued scholars and members of the public alike. A possible explanation for the absence of  
127 fish remains at Star Carr could be that Clark did not sieve the sediments, meaning that small  
128 specimens may have been missed. Sieving was not a common practice at the time; peat is  
129 extremely difficult to sieve because it is highly organic and does not easily pass through a  
130 sieve and therefore it is perhaps unsurprising that this was not attempted.

131

132 Renewed research since 2004 (Conneller *et al.* 2012; Milner *et al.* 2013) has provided further  
133 opportunities to test for the presence of fish remains at the site. Initially, it was considered  
134 highly unlikely that any fish remains would be found, even with sieving, due to the extremely  
135 acidic sediments that have formed over the last couple of decades (Boreham *et al.* 2011; High

136 *et al.* 2015). Some bone and antler has become ‘jellybone’: the mineral has dissolved in the  
137 acidic peat and the collagen has turned to gelatin (Milner *et al.* 2011a). Furthermore,  
138 quantities of bone and antler are extremely low when compared to Clark’s faunal collection,  
139 suggesting that much of this material has completely disappeared (Milner *et al.* 2011a).

140

141 During the last four years, three different lines of evidence have at last provided definitive  
142 evidence that not only a range of fish species were present in the lake, but that they were  
143 caught and processed by humans. Significantly, these lines of evidence came from  
144 completely different scales of analyses:

- 145 1. landscape scale: coring the lake sediments for environmental and climate records
- 146 2. site scale: excavations at Star Carr and at Flixton Island Site 2
- 147 3. microscopic scale: microwear traces on flint tools from Star Carr

148

### 149 **1.1. Background to the sites**

150 Star Carr and Flixton Island Site 2 are two of a number of Early Mesolithic sites that have  
151 been recorded in the area around the palaeo-Lake Flixton, in the eastern Vale of Pickering,  
152 North Yorkshire, UK (Figure 3). The palaeo-lake formed at the start of the Windermere  
153 Interstadial (c. 14,700-12,800 cal BP; 12,700-10,800 cal BC), a warm phase at the end of the  
154 last Ice Age before the final cold period of the Younger Dryas (12,700-11,600 cal BP;  
155 10,700-9600 cal BC), and it persisted as a water body until the end of the Mesolithic (c. 6000  
156 cal BP; 4000 cal BC).

157

158 John Moore, a local amateur archaeologist first carried out investigations in the area in the  
159 late 1940s and identified 10 sites around the lake. Moore excavated a trench at Star Carr in  
160 1948, and from 1949-1951 Grahame Clark from the University of Cambridge conducted  
161 three further seasons of fieldwork (Clark 1954). Moore also conducted fieldwork at Site 2 on  
162 Flixton Island which was published as a three-page summary at the end of the Star Carr  
163 monograph (Clark 1954).

164

165 Further work in the area has been carried out by the Vale of Pickering Research Trust since  
166 the 1980s, with the aim of mapping the extent of the lake and identifying further sites (for a  
167 full account see Milner *et al.* (2011b)). Since 2004, NM, CC and BT have been co-directing  
168 excavations at Star Carr and in 2012 the POSTGLACIAL project commenced: this is a five  
169 year, European Research Council funded project with the aim ‘To implement an

170 interdisciplinary, high-resolution approach to understanding hunter-gatherer lifeways within  
171 the context of climate and environment change during the early part of the post-glacial period  
172 (c. 10,000-8000 BC)'. In order to address this aim, further excavations have been carried out  
173 at Flixton Island Site 2 (2012-14) and Star Carr (2013-2015), in conjunction with a  
174 programme of coring lake sediments in order to reconstruct local climate and environmental  
175 change.

176

177 *Figure 3: The locations of Star Carr and Flixton Island Site 2 as well as all other known Mesolithic sites within*  
178 *the Lake Flixton basin (lake area a reconstruction of the water at its maximum during the Holocene). Key: 1,*  
179 *Star Carr; 2, Ling Lane; 3, Seamer Carr Site F; 4, Seamer Carr Sites L and N; 5, Seamer Carr Site K; 6,*  
180 *Seamer Carr Site D; 7, Seamer Carr Site B (Rabbit Hill); 8, Seamer Carr Site C; 9, Manham Hill; 10–12,*  
181 *Cayton Carr; 13, Lingholme Site B; 14, Killerby Carr; 15, Lingholme Site A; 16, Barry's Island; 17, Flixton*  
182 *School Field; 18, Flixton School House Farm; 19, Woodhouse Farm; 20, VP Site E; 21, VP Site D; 22, Flixton*  
183 *Site 9; 23, Flixton Island Site 1; 24, Flixton Island Site 2; 25, No Name Hill.*

184

## 185 **2.0. Methodology**

### 186 **2.1. Sediment coring**

187 Between 2010-2013, alongside the lake-edge excavations, a sediment core-based  
188 palaeoenvironmental study was undertaken as part of the POSTGLACIAL project. The  
189 preliminary results of this research were reported by Palmer *et al.* (2015). During coring, a  
190 single fish scale was identified toward the base of sediment core C (Palmer *et al.* 2015).

191

### 192 **2.2. Excavations at Flixton Island Site 2**

193 In 2012, excavations were carried out at Flixton Island Site 2, approximately 500 metres east  
194 of Star Carr. A programme of sieving the archaeological sediments was carried out and a  
195 50% sample was sieved using a 4 mm mesh in order to retrieve small pieces of bone or flint  
196 debitage, which otherwise would have been missed through trowelling. As the focus was on  
197 the lakeshore deposits, where fewer archaeological remains are generally found, a very small  
198 proportion of finds were retrieved in the sieve. However, a fish vertebra was discovered in  
199 the sieve from context 1003 (Mesolithic reed peat).

200

### 201 **2.3. Excavations at Star Carr and the flotation of the dryland deposits**

202 During the excavations in 2008 at Star Carr, the remains of a structure were discovered on the  
203 dryland, which has become known as the 'Earliest house in Britain' (Conneller *et al.* 2012).  
204 The structure was composed of a hollow in the ground, infilled with organic rich sediment,

205 and surrounded by postholes. It should be noted, that although significant quantities of  
206 mammal bone have been found on the dry land deposits, and in the structure itself, the  
207 preservation of the bone is generally extremely poor and most remains are unidentifiable  
208 either in terms of genus/species or in many cases element.

209

210 Sediment was sampled from the structure and processed in the laboratory using the bucket  
211 flotation method. Volumes ranged from 1.5 to 2.5 litres. A 300  $\mu$  sieve was used to collect the  
212 flot and a 1 mm sieve was used to catch the residue. Samples were dried in a drying cabinet  
213 for approximately 12 hours at 40°C. The flot was examined primarily for plant remains and  
214 the residue for other biological remains. The content of each sieve was examined a small  
215 fraction at a time under a stereomicroscope. Plant remains and any other biological remains  
216 retrieved were extracted using tweezers and stored in a sealed glass tube, clearly labelled with  
217 the site code and sample number in preparation for identification. A total of 11 fish remains  
218 were recovered from grid square I3, context 149, by ET, and analysed by HR using the  
219 modern reference collection housed at the University of York.

220

221 More recently, in 2015, excavations were carried out in the vicinity of Clark's trenches  
222 including a small area of unexcavated baulk and here two fish remains were recovered by  
223 hand during the careful excavation of context 312 (Mesolithic reed peat).

224

#### 225 **2.4. Microwear analysis**

226 Microwear analysis is the study of traces that are visible on a tool and which develop through  
227 the course of the tool's use. Traces can vary depending on the contact material that is worked  
228 and the direction in which the tool is used. (Vaughan 1985). This is determined by comparing  
229 archaeological traces with those on experimentally used tools. It is this comparison that  
230 allows an analyst to make an informed inference about the function of artefacts (van Gijn and  
231 Little in press).

232

233 Identifying fish within prehistoric assemblages is recognised as difficult due to their  
234 vulnerability of the traces and a frequent lack of distinction from other animal processing  
235 activities (van Gijn 1986, 23). However, experimental work replicating different aspects of  
236 fish processing, such as filleting, gutting and scaling (see García Díaz and Clemente Conte  
237 2011; van Gijn 1986) has enabled similar traces to be identified on archaeological lithic  
238 assemblages (Clemente *et al.* 2010; Högberg 2009). Recent research has even identified fish



239 microwear traces on Neanderthal stone tools (Hardy and Moncel 2011), showing that fish  
 240 processing tools have great antiquity.

241

242 **3.0. Results**

243 **3.1. Sediment coring**

244 The scale recovered from the sediment core had a characteristic ctenoid form and was  
 245 identified by DS as a scale from a perch. The scale was identified at c. 17.1 m.a.s.l around  
 246 705 cm below the current ground surface. The deposits in which it lay are carbonate-rich lake  
 247 sediments (Palmer *et al.*'s lithofacies 2a), associated with the Windermere interstadial  
 248 (equivalent to the European Bølling/Allerød chronozone and dating between c. 14,700-  
 249 12,800 cal BP, or 12,700-10,800 cal BC). The scale sat within the earliest sediments of this  
 250 interval and reflects the presence of fish within the lake soon after the commencement of  
 251 climatic warming. Fish scales of Lateglacial age are relatively uncommon finds within  
 252 sediment cores in the UK, being first reported for Esthwaite Water in the English Lake  
 253 District by Pennington and Frost (1961), and this chance find was not repeated in any other  
 254 samples.

255

256 **3.2. Flixton Island Site 2 and Star Carr ichthyoarchaeological results**

257 A total of 14 fish remains have been recovered from Flixton Island Site 2 and Star Carr  
 258 (Table 1). Two of the remains were vertebral fragments that could not be determined to  
 259 species. The following provides a brief summary of these data.

260

<b>Taxon/skeletal element</b>	<i>Esox lucius</i>	<i>Esox lucius</i> / Salmonidae	Cyprinidae	<i>Perca fluviatilis</i>	Unidentifiable	<b>Totals</b>
<b><i>Lake Flixton</i></b>						
Ctenoid scale				1		<b>1</b>
<b><i>Flixton Island Site 2</i></b>						
Caudal vertebra	1					<b>1</b>
<b><i>Star Carr</i></b>						
Caudal vertebra			2			<b>2</b>
Posterior abdominal vertebra	1					<b>1</b>
Pharyngeal tooth			2			<b>2</b>
Premaxilla	1					<b>1</b>
Rib			1			<b>1</b>
Tooth		3				<b>3</b>
Vertebral fragment				1	2	<b>3</b>
<b>Totals</b>	<b>3</b>	<b>3</b>	<b>5</b>	<b>2</b>	<b>2</b>	<b>15</b>

261 *Table 1: Represented skeletal elements of the various taxa found during the excavations and in the post*  
262 *excavation processing of bulk samples from Flixton Island Site 2 and Star Carr and also the ctenoid scale from*  
263 *the sediment core.*

264

265 Although neural and haemal spines were absent, the bone recovered from Flixton Island Site  
266 2 was a caudal vertebra which was identified by HR and AKGJ as pike (Figure 4). The total  
267 length (hereafter TL) of the pike was estimated as a function of bone size according to the  
268 methods as set out by Morales and Rosenlund (1979) utilizing the regression equations as  
269 stated by Zabilska (2013). The size of the vertebra, 12.1 mm across the greatest medio-lateral  
270 breadth of the centrum, corresponded to that of an individual approximately 815 mm in TL.  
271 Since adult pike normally range from 400 to 1000 mm in TL (Davies *et al.* 2004), this falls  
272 well within that range.

273

274 The vertebra was sent to Oxford Radiocarbon laboratory for direct AMS radiocarbon dating  
275 but unfortunately failed to produce a date due to a lack of collagen. However, a sample of  
276 willow (*Salix* sp.) from the same level was successfully dated to 9170-8570 cal BC (95.4%  
277 probability: OxA-X-2495-12, 9480 ± 90 BP). This date is contemporary with the dates  
278 yielded for Star Carr.

279

280 *Figure 4: SEM image of the pike caudal vertebra from Flixton Island Site 2.*

281

282 All of the fish remains recovered from the bucket flotation method were less than 1 cm in  
283 size, and all but one was calcined. It is likely that the one that was not visibly calcined had  
284 been subjected to some burning in order for it to have survived in this dryland context.

285

286 Although largely incomplete, a premaxilla that could not be sided was identified as that of  
287 pike. This specimen was calcined and was dark grey, almost black, in colour. Comparison  
288 with modern specimens suggested that it derived from an individual that was less than 200  
289 mm in TL.

290

291 Three isolated teeth were identified as likely to derive from Salmonidae: however, it must be  
292 noted that salmon and pike have very similar teeth and there is the possibility that these also  
293 belonged to pike. If they are Salmonidae there is a possibility that these remains could have  
294 belonged to either the anadromous brown trout (*Salmo trutta* L., 1758) or Atlantic salmon

295 (*Salmo salar* L., 1758). All specimens were calcined, ranging from light grey almost white to  
296 dark grey in colour.

297

298 A fragile, isolated pharyngeal tooth was identified as deriving from Cyprinidae (carp and  
299 minnow family). It was not possible to identify the specimen to species level. The specimen  
300 was calcined and was dark grey in colour. Although neural and haemal spines were absent,  
301 two caudal vertebrae were identified as Cyprinidae. The specimens were calcined and were  
302 light grey, almost white and white in colour respectively. It was not possible to estimate the  
303 TLs, although they derived from an individual that was less than 200 mm in TL by  
304 comparison with modern specimens. In addition, one rib was identified as Cyprinidae. The  
305 specimen was incomplete, as only the proximal end was present.

306

307 In addition, one vertebra was identified as that of perch. The specimen was incomplete, as  
308 only half of the vertebral centrum was present. It was calcined, and was light to dark grey,  
309 almost black in colour. It was not possible to estimate the TL, although it derived from an  
310 individual that was less than 100 mm in TL based on comparison with modern taxa.

311

312 In 2015, excavations at Star Carr yielded a further two fish remains. Although fragile, the  
313 first specimen, an isolated tooth with a portion of the pharyngeal bone was identified by HR  
314 as deriving from Cyprinidae. It was not possible to identify the specimen to the lower species  
315 level or estimate the TL. The second bone, a posterior abdominal vertebra (Figure 5) was  
316 identified by HR as pike. The TL of this specimen was estimated according to the criteria  
317 outlined above (Morales and Rosenlund 1979; Zabilska 2013). The size of the vertebra, 11.4  
318 mm across the greatest medio-lateral breadth of the centrum, corresponded to that of an  
319 individual approximately 873 mm in TL.

320

### 321 **3.3. Evidence that fish had been processed at Star Carr from microwear analysis**

322 The current programme of microwear analysis of flint tools from Star Carr is the first since  
323 Dumont's studies (Dumont 1983, 1988). Microwear analysis of the flint in and around the  
324 structure has only just begun. Two flints with fish processing polish have already been  
325 identified a short distance from the structure (Figure 6). Neither of the tools (92184 and  
326 91949) are retouched. One (92184) is classified typologically as a fragment; the other  
327 (91949) is a proximal blade fragment. The fish processing polish are presented in Figure 7.

328

329 The two tools from Star Carr were taken to the Laboratory for Artefact Studies at Leiden  
330 University to be blind tested by three experienced microwear analysts. None of the analysts  
331 knew the original identification was fish processing. After independently analysing tool  
332 91949, two analysts identified the wear traces as resulting from fish processing; the other said  
333 'possibly fish processing'. The test was repeated for tool 92184, with two analysts identifying  
334 fish processing and the third saying the polish was 'indeterminate'. When presented with the  
335 conclusions from the other two analysts, this third analyst accepted that fish processing was a  
336 strong possibility.

337

338 *Figure 5: Photograph of the pike vertebra in situ.*

339

340 Fish processing traces consist of randomly distributed lines of matt, dull polish often  
341 described as having a corrugated appearance (Figure 7a) sometimes located away from the  
342 edge (van Gijn 1986). Edge damage in the form of small flake scalar scars, which form a  
343 repeated but irregular appear along the edge of the flint, are visible on some tools (van Gijn  
344 1986; Högberg 2009). Lately, analysts have observed another feature to fish polish: areas  
345 where fish scales adhering to the surface of the tool have prevented polish forming, resulting  
346 in rounded, scalar areas of unpolished surface (García Díaz 2014, 98; see also Figure 7c).

347

348 Although not located within the structure, the fish processing tools were located a short  
349 distance from it and are from the same context as the calcined fish remains from within the  
350 structure. Due to taphonomic processes such as bioturbation and the palimpsests of activity  
351 on the dryland, we cannot be sure that the tools and fish remains are contemporary. Future  
352 microwear analysis of the flint assemblage from within the structure will determine whether a  
353 contextual relationship exists between the remains and tools. As this analysis is ongoing it  
354 also remains to be seen if future microscopic studies of the flint assemblages from other areas  
355 of the site will reveal similar evidence for fish processing tools.

356

357 *Figure 6: Structure at Star Carr showing the distribution of finds. Key: triangle, location of fish remains within  
358 the structure; stars, lithics exhibiting fish processing traces.*

359

360 *Figure 7: Flint 92184 (above) and 91949 (below), both of which display fish processing polish. All micrographs  
361 20x.*

362

363 **4.0. Discussion**

364 **4.1. The earliest evidence of fish in Lake Flixton**

365 A particularly significant discovery of this research is that the evidence of perch in the lake  
366 originates in the Windermere interstadial, roughly between 2000-4000 years before  
367 settlement commenced at Star Carr. Native to Britain, perch are presently distributed across  
368 the northern Palearctic, with the exception of the Iberian Peninsula, central Italy and the  
369 Adriatic basin (Freyhof and Kottelat 2008). They occur across a diverse range of habitats,  
370 including estuarine lagoons, lakes and medium-sized streams, spawning in NW Europe in  
371 early spring when water temperatures reach a minimum of c. 10°C (Gillet and Dubois 2007;  
372 Hokanson 1977) and the photoperiod conditions increase. As opportunistic feeders, they prey  
373 on a wide range of aquatic invertebrates, with larger individuals becoming piscivorous once  
374 they reach 120 mm in length (Freyhof and Kottelat 2008).

375

376 The spread of *Perca fluviatilis* across Europe after the last Ice Age has been documented  
377 using molecular techniques (Nesbø *et al.* 1999). This suggests that perch found in modern  
378 day western Europe dispersed along the major river systems centred around the Vistula, Elbe,  
379 Rhine Rhône, Saône and Thames, with British perch originating from a southern glacial  
380 refugium, probably located in France although the exact position remains unclear (Nesbø *et al.*  
381 *al.* 1999). A similar southern French refugium has been suggested for other freshwater taxa  
382 today found in Britain, such as barbel (*Barbus* sp.) (Persat and Berrebi 1990), chub  
383 (*Leuciscus cephalus* L., 1758) (Durand *et al.* 1999) and brown trout (García-Marín *et al.*  
384 1999), with the molecular data implying rapid northward expansion through northern  
385 European riverine systems since the last Glacial Maximum in the UK. Here, it is suggested  
386 that the presence of perch in Lake Flixton during the early Windermere interstadial could  
387 indicate colonisation through the Derwent river system.

388

389 **4.2. The fish trophic system in Lake Flixton**

390 This research also demonstrates the presence of pike, known from a number of Early  
391 Mesolithic sites in Denmark. Pike can be found throughout Asia, eastern North America and  
392 the majority of Europe. In Britain, it is the largest predatory freshwater fish, and consumes  
393 invertebrates, lesser fish, aquatic birds, amphibians and small mammals. Although it occurs  
394 in lakes to larger ornamental ponds and from canals and slow flowing rivers to streams, it can  
395 also reside in bogs as well as brackish lagoons and shallow, protected bays (Davies *et al.*  
396 2004). It has a high tolerance to changes in pH and is capable of surviving in polluted waters,  
397 including those with low oxygen content (Noe-Nygaard 1995). Whilst it is a solitary

398 carnivore, it is not territorial, often congregating in shoals to rest. Aided by its camouflaged  
399 appearance, it hides in submerged vegetation, where it lurks near its prey. Mating takes place  
400 between March and April when pike congregate and seek shallow water; it is during this time  
401 when they can almost be caught by hand (Noe-Nygaard 1995). Both the presence of perch  
402 and pike in Lake Flixton indicates a mature water body with an established trophic system.

403

404 Five of the fish remains have been identified to Cyprinidae. Although these specimens could  
405 not be identified to the species level, other species routinely identified in contemporaneous  
406 European faunal assemblages include the following: bream, white bream (*Abramis bjoerkna*  
407 L., 1758), Crucian carp (*Carassius carassius* L., 1758), common carp (*Cyprinus carpio* L.,  
408 1758), roach (*Rutilus rutilus* L., 1758), rudd and tench. Cyprinidae are, in general, lower  
409 down in the trophic level hierarchy, compared to perch and pike, and mainly feed on benthic  
410 invertebrates, including worms, molluscs and insect larvae (Maitland and Linsell 2009).

411

412 Although we cannot be 100% certain at the present that we have trout or salmon, it is  
413 important to discuss them in case further discoveries prove their presence. Both are  
414 anadromous species (migrate from the sea into freshwater to spawn) (Wheeler and Jones  
415 2009). During their spawning runs, they can often become concentrated, albeit sometimes for  
416 a short period, making them abundant and prime targets for fishing. Atlantic salmon enter  
417 freshwater from the sea and migrate upstream at different times throughout the year. Whilst  
418 this is largely dependent upon the flow of the river and the water temperature, they arrive at  
419 the spawning grounds from November through February (Mills 1971). Here they are  
420 extremely vulnerable to predation since they are in shallow waters and occupied by  
421 spawning.

422

### 423 **4.3. Fishing and consumption**

424 Apart from the barbed points, there is no other evidence of fishing gear, such as nets, hooks,  
425 or traps, found at any of the sites along the former lake edge. However, some suggestions can  
426 be made as to the possible techniques that might have been used. Since pike are known to  
427 congregate within the littoral zone of a lake during their spawning period in the spring, they  
428 could have been more easily exploited at this time (Noe-Nygaard 1995). The pike may also  
429 have been attracted to some of the food waste, such as bones, which were deposited at the  
430 lake margins and from these areas it may have been possible to spear the pike using the  
431 barbed points found at the site. However, it should be noted that the barbed points found in

432 the lakeshore deposits are not hafted and are therefore unlikely to have been lost during  
433 fishing activities. In addition, it is possible that bows and arrows were used or blows via  
434 clubs, and then collected (Aaris-Sørensen 1976).

435

436 Clark (1952, 48) recounts how Lapps spear pike from boats by targeting them when they sun  
437 themselves in the upper part of the water body. The use of boats is well established at this  
438 time, since people used the islands on the lake and presumably accessed them by boat. Clark  
439 (1954) also found what he thought was a wooden paddle at Star Carr: this is very thin but  
440 broken at both ends so difficult to estimate its full length. It has sometimes been dismissed as  
441 a paddle due to its thinness, however, paddles of this shape are known to be favoured in some  
442 place for navigating through reeds. Boats and paddles are also well documented for the  
443 Mesolithic across Europe.

444

445 As well as spearing fish in the day, Clark describes how the Lapps also enticed fish to the  
446 surface of the water at night by burning dry wood at the prow of the boat. An argument for  
447 night fishing has been suggested for pine tapers found on Irish Midland waterways linked to  
448 night-time fishing/eeling activities (Little 2009). At Star Carr, numerous burnt birch bark  
449 rolls have been found and analysed (Figure 8). Experiments have shown that plain rolls of  
450 bark burn for only a matter of minutes because the lack of oxygen in the middle of the roll  
451 suffocates the fire. However, if strips of bark are positioned within the rolls the torches burn  
452 for longer though tend not to survive as a burnt roll (Figure 9). Research is ongoing to  
453 determine whether the birch bark rolls from Star Carr could have been used for torches.

454

455 *Figure 8: Birch bark roll with evidence of burning from Lake Flixton.*

456

457 Clark (1952, 48) further mentions that the Lapps also catch pike using dragnets between two  
458 boats. Although nets have not been found at Star Carr, there are a large number of birch bark  
459 rolls some of which have not been burnt, and from ethnographical analogues, it can be  
460 demonstrated that they can be used as net floats (Figure 10).

461

462 *Figure 9: experimental torch made from a birch bark roll.*

463

464 *Figure 10: Picture of birch bark rolls used as net floats (from The National Museum of Finland).*

465

466 The site has yielded a tantalising glimpse of fish processing, and presumably consumption,  
467 from the use-wear traces on two flint tools. What is interesting is that this has taken place  
468 near to the structure where the calcined fish remains have been found. Future analysis will  
469 help to determine whether there is further patterning to these processes; however, the nature  
470 of the remains means that we will not be able to say how extensive fish consumption was.

471

472 Excluding the scale recovered from the sediment core, a total of 14 fish remains have been  
473 recovered from two archaeological sites located within the palaeo-Lake Flixton basin: Flixton  
474 Island Site 2 and Star Carr. In comparison with contemporaneous Maglemose sites in  
475 southern Scandinavia the assemblage is small. However, we believe that there are several  
476 reasons as to why so few have been recovered, and in particular from Star Carr: (1) Clark did  
477 not sieve the sediments; (2) the increase in the acidity of the sediments since the 1950s is  
478 likely to have destroyed any fish remains due to their delicate nature; (3) fish bones may have  
479 been deposited on the dry land and it is highly likely that they would not have survived unless  
480 burnt, as found in the structure; (4) we have previously demonstrated that ca. 5% of the site  
481 has been excavated, and so the possibility exists that fish remains (and possibly associated  
482 technology) may be present elsewhere on the site, possibly around the lake edge or  
483 discarded in the deeper deposits further into the lake as has been noted elsewhere in southern  
484 Scandinavia, such as Ringkloster (Andersen 1998).

485

## 486 **5.0. Conclusion**

487 This paper has yielded a number of important conclusions:

- 488 1. The evidence demonstrates that different species of fish were available in the lake  
489 during the Early Mesolithic and that these were exploited by the inhabitants of Star  
490 Carr. This data significantly adds to our understanding of Early Mesolithic economy  
491 in Britain.
- 492 2. The research demonstrates the importance of conducting flotation on sediments from  
493 dry land deposits and particularly from contexts such as structures. Although bone  
494 from this part of the site is generally very poorly preserved, the fish bone has been  
495 subjected to heating (in all likelihood it was thrown on a fire) which has meant that it  
496 has survived normal destructive taphonomic processes.
- 497 3. The application of use-wear analysis is of great importance for Mesolithic sites where  
498 faunal evidence is lacking. It holds the potential to uncover archaeological



499 information that is invisible to the naked eye, and answer economic questions that  
500 would otherwise remain unanswered.

501 4. By 3D plotting all artefacts using a Total Station and mapping using GIS important  
502 spatial relationships come to light that otherwise might be missed: the current  
503 evidence suggests people were processing the fish outside the structure, and then  
504 possibly cooking or consuming it within the structure before throwing the bones on a  
505 fire.

506

507 In sum, fish and fish-processing activities would have remained a mystery for Star Carr if it  
508 were not for the multi-scalar approach applied to the study of artefacts and the  
509 palaeoenvironment. Thus we believe the importance of combining micro- and macro-  
510 methods is critical when investigating hunter-gatherer settlement sites.

511

#### 512 **Acknowledgements**

513 The excavations at Star Carr have been funded by: European Research Council  
514 (POSTGLACIAL project, No 283938), The British Academy (SG-44333, SG-47081, SG-  
515 50217), English Heritage (grant numbers 5536, 6064, 6793 and 6796), NERC  
516 (NE/I015191/1), the Vale of Pickering Research Trust, University of Manchester, University  
517 of York, University of Chester, The McDonald Institute, Cambridge, and UCL. We would  
518 like to thank the landowners of Star Carr and Flixton Island and to English Heritage and  
519 Natural England for granting permission to excavate. We thank the team at the Laboratory for  
520 Artefact Studies at Leiden University for participating in blind tests of the fish processing  
521 tools. Finally, we are indebted to everyone who participated in the excavations since 2004.

522

#### 523 **References cited**

524 Andersen, S. H. 1998. Ringkloster. Ertebølle Trapper and Wild Boar Hunters in Eastern  
525 Jutland. *Journal of Danish Archaeology*, 12, 1994-1995, 13-59.

526

527 Aaris-Sørensen, K. 1976. A zoological investigation of the bone material from Sværdborg I-  
528 1943. *Arkæologiske Stud*, 3, 137-148.

529

530 Bell, M. 2007. *Prehistoric Coastal Communities: The Mesolithic in western Britain*. York:  
531 Council for British Archaeology, CBA Report 149.

532

533 Boreham, S., Conneller, C., Milner, N., Taylor, B., Needham, A., Boreham, J., and Rolfe, C.  
534 J. 2011. Geochemical Indicators of Preservation Status and Site Deterioration at Star Carr.  
535 *Journal of Archaeological Science*, 38(10), 2833–57.

536

537 Broholm, H. C. 1924. Nye Fund fra den ældste stenalder. Holmegård-og Sværdborg fundene.  
538 *Årbøger fra nordisk oldkyndighed og historie*, 1-144.

539

540 Carter, R. 1998. Reassessment of Seasonality at the Early Mesolithic Site of Star Carr,  
541 Yorkshire Based on Radiographs of Mandibular Tooth Development in Red Deer (*Cervus*  
542 *elaphus*). *Journal of Archaeological Science*, 25(9), 851–856.

543

544 Caulfield, S. 1978. Star Carr – an alternative view. *Irish Archaeological Research Forum*, 5,  
545 15-22.

546

547 Clark, J. G. D. 1948. The Development of Fishing in Prehistoric Europe. *The Antiquaries*  
548 *Journal*, 28(1-2), 45–85.

549

550 Clark, J. G. D. 1952. *Prehistoric Europe: the economic basis*. London: Methuen.

551

552 Clark, J. D. G. 1954. *Excavations at Star Carr: An early Mesolithic site at Seamer near*  
553 *Scarborough, Yorkshire*. Cambridge: Cambridge University Press.

554

555 Clemente, I., García, V., Ramos, J., Domínguez-Bella, S., Pérez, M., Vijande, E., Cantillo, J.  
556 J., Soriguer, M., Zabala, C., and Hernando, J. The Lithic Tools of the La Esparragosa Site  
557 (Chiclana de la Frontera, Cádiz, Spain, fourth Millennium BC): A Methodological  
558 contribution to the Study of Lithic Tools for the Consumption of Fisch. *Ancient Nets and*  
559 *Fishing Gear*, 275-285.

560

561 Conneller, C., Milner, N., Taylor, B., and Taylor, M. 2012. Substantial settlement in the  
562 European Early Mesolithic: new research at Star Carr. *Antiquity*, 86(334), 1004-1020.

563

564 Davies, C., Shelley, J., Harding, P., McLean, I., Gardiner, R., and Peirson, G. 2004.  
565 *Freshwater fishes in Britain – the species and their distribution*. Colchester: Harley Books.

566

567 Dumont, J. V. 1983. An interim report of the Star Carr microwear study. *Oxford Journal of*  
568 *Archaeology*, 2(2), 127–45.

569

570 Dumont, J. V. 1988. *A Microwear Analysis of Selected Artefact Types from the Mesolithic*  
571 *Sites of Star Carr and Mount Sandel*. Oxford: Archaeopress, Volume 187.

572

573 Durand, J. D., Persat, H., and Bouvet, Y. 1999. Phylogeography and postglacial dispersion of  
574 the chub (*Leuciscus cephalus*) in Europe. *Molecular Ecology*, 8, 989–997.

575

576 Freyhof, J., and Kottelat, M. 2008. *Perca fluviatilis*. The IUCN Red List of Threatened  
577 Species. Version 2015.2. [www.iucnredlist.org](http://www.iucnredlist.org). Downloaded on 18 August 2015.

578

579 García-Marín, J.-L., Utter, F. M., and Pla, C. 1999. Postglacial colonisation of brown trout in  
580 Europe based on distribution of allozyme variants. *Heredity*, 82, 46–56.

581

582 García Díaz, V. 2014. Flint, stones and bones: raw material selection, typology, technology  
583 and use-wear analysis. In E. M. Theunissen, S. M. Beckerman, D. C. Brinkhuizen, V. García  
584 Díaz, L. Kubiak-Martens, and D. C. M. Raemaekers (Eds.) *Synthesis-A mosaic of habitation*  
585 *at Zeewijk*, 85-117. Amersfoort: Nederlandse Archeologische Rapporten 10.

586

587 García Díaz, V., and Clemente Conte, I. 2011. Procesando pescado: reproducción de las  
588 huellas de uso en cuchillos de sílex experimentales. In A. Morgado, J. Baena Preysler, and G.  
589 González (Eds.) *La Investigación Experimental aplicada a la Arqueología*, 163-169. Málaga:  
590 Imprenta Galindo.

591

592 Gillet, C., and Dubois, J. P. 2007. Effect of water temperature and size of females on the  
593 timing of spawning of perch *Perca fluviatilis* L. in Lake Geneva from 1984 to 2003. *Journal*  
594 *of Fish Biology*, 70, 1001–1014.

595

596 Godwin, H., and Godwin, M. E. 1933. British Maglemose harpoon sites. *Antiquity*, 7(25), 36-  
597 48.

598

599 Gramsch, B., and Beran, J. 2010. Spätaltsteinzeitliche Funde von Wustermark, Fundplatz 22,  
600 Lkr. Havelland. *Veröffentlichungen zur brandenburgischen Landesarchäologie*, Band 41/42  
601 – 2007/2008, 95-142.

602

603 Groß, D. 2014. *Welt und Umwelt frühmesolithischer Jäger und Sammler. Mensch-Umwelt-*  
604 *Interaktion im frühen Holozän in der mitteleuropäischen Tiefebene*. Unpublished PhD  
605 Thesis: Christian-Albrechts-Universität zu Kiel.

606

607 Hardy, B. L., and Moncel, M-H. 2011. Neanderthal Use of Fish, Mammals, Birds, Starchy  
608 Plants and Wood 125-250,000 Years Ago. *PloS One* 6, e23768.

609

610 High, K., Milner, N., Panter, I., and Penkman, K. E. H. 2015. Apatite for destruction:  
611 Investigating bone degradation due to high acidity at Star Carr. *Journal of Archaeological*  
612 *Science*, 59, 159-168, 10.1016/j.jas.2015.04.001.

613

614 Högberg, A., Puseman, K., and Yost, C. 2009. Integration of Use-Wear with Protein Residue  
615 Analysis—a Study of Tool Use and Function in the South Scandinavian Early Neolithic.  
616 *Journal of Archaeological Science*, 36, 1725-1737.

617

618 Hokanson, K. E. F. 1977. Temperature requirements of some percids and adaptations to the  
619 seasonal temperature cycle. *Journal of the Fisheries Research Board of Canada*, 34, 1524–  
620 1550.

621

622 Jacobi, R. 1978. Northern England in the eighth millennium bc: an essay. In P. Mellars (Ed.)  
623 *The early postglacial settlement of Northern Europe*, 295-332. London: Duckworth.

624

625 Jessen, C. A., Pedersen, K. B., Christensen, C., Olsen, J., Mortensen, M. F., and Hansen, K.  
626 M. 2015. Early Maglemosian culture in the Preboreal landscape: Archaeology and vegetation  
627 from the earliest Mesolithic site in Denmark at Lundby Mose, Sjælland. *Quaternary*  
628 *International*, 378, 73-87, doi:10.1016/j.quaint.2014.03.056.

629

630 Legge, A., and Rowley-Conwy, P. 1988. *Star Carr revisited: a re-analysis of the large*  
631 *mammals*. London: Centre for Extra-Mural Studies.

632

633 Little, A. 2009. Fishy settlement patterns and their social significance: a case study from the  
634 northern Midlands of Ireland. In S. McCartan, R. Schulting, G. Warren, and P. Woodman  
635 (Eds.) *Mesolithic Horizons: Papers Presented at the Seventh International Conference on the*  
636 *Mesolithic in Europe, Belfast 2005*, 698–705. Oxford: Oxbow Books.

637

638 Maitland, P. S., and Linsell, K. 2009. *Guide to freshwater fish of Britain and Europe*.  
639 London: Philip's.

640

641 Mellars, P., and Dark, P. 1998. *Star Carr in context: new archaeological and palaeological*  
642 *investigations at the Early Mesolithic site of Star Carr, North Yorkshire*. Cambridge:  
643 MacDonal Institute.

644

645 Mills, D. M. 1971. *Salmon and trout: a resource, its ecology, conservation and management*.  
646 Edinburgh: Oliver and Boyd.

647

648 Milner, N., Conneller, C., Taylor, B., Koon, H., Penkman, K., Elliott, B., Panter, I., and  
649 Taylor, M. 2011a. From Riches to Rags: Organic Deterioration at Star Carr. *Journal of*  
650 *Archaeological Science*, 38(10), 2818-2832.

651

652 Milner, N., Lane P., Taylor, B., Conneller, C., and Schadla-Hall, T. 2011b. Star Carr in a  
653 Postglacial Lakescape: 60 Years of Research. *Journal of Wetland Archaeology*, 11(1), 1–19.

654

655 Milner, N., Taylor, B., Conneller, C., and Schadla-Hall, R. T. 2013. *Star Carr: Life in Britain*  
656 *after the Ice Age*. York: Council for British Archaeology.

657

658 Momber, G.,Tomalin, D., Scaife, R., Satchell, J., and Gillespie, J. 2011. *Mesolithic*  
659 *occupation at Bouldnor Cliff and the submerged prehistoric landscapes of the Solent*. York:  
660 Council for British Archaeology Research Report 164.

661

662 Morales, A., and Rosenlund, K. 1979. *Fish bone Measurements: an attempt to standardize*  
663 *the measuring of fish bones from archaeological sites*. København: Zoologisk Museum.

664

665 Nesbø, C. L., Fossheim, T., Vollestad, L. A., and Jakobsen, K. S. 1999. Genetic divergence  
666 and phylogeographic relationships among European perch (*Perca fluviatilis*) populations  
667 reflect glacial refugia and postglacial colonization. *Molecular Ecology*, 8, 1387-1404.  
668

669 Noe-Nygaard, N. 1995. *Ecological, Sedimentary, and Geochemical Evolution of the Late-*  
670 *glacial to Postglacial Åmose Lacustrine Basin, Denmark*. Oslo: Scandinavian University  
671 Press.  
672

673 Palmer, A. P., Matthews, I. P., Candy, I., Blockley, S. P. E., Macleod, A., Darvill, C. M.,  
674 Milner, N., Conneller, C., and Taylor, B. 2015. The evolution of Palaeolake Flixton and the  
675 environmental context of Star Carr, NE. Yorkshire: stratigraphy and sedimentology of the  
676 Last Glacial-Interglacial Transition (LGIT) lacustrine sequences. *Proceedings of the*  
677 *Geologists Association*, 126(1), 50-59, 10.1016/j.pgeola.2014.10.002.  
678

679 Pennington, W., and Frost, W. E. 1961. Fish vertebrae and scales in a sediment core from  
680 Esthwaite Water (English Lake District). *Hydrobiologia*, 17, 183–190.  
681

682 Persat, H., and Berrebi, P. 1990. Relative ages of present populations of *Barbus barbus* and  
683 *Barbus meridionalis* (Cyprinidae) in southern France: preliminary considerations. *Aquatic*  
684 *Living Resources*, 3, 253–263.  
685

686 Robson, H. K. 2015. *Identification, preliminary analyses and interpretation of the fish*  
687 *remains from the early Holocene site, Friesack IV*. Unpublished report: University of York.  
688

689 Rosenlund, K. 1980. Knoglematerialet fra bopladsen Lundby II. In B. B. Henriksen (Ed.)  
690 *Lundby-holmen: Pladser af Maglemose-type i Sydsjælland*, 128-142. København: Nordiske  
691 Fortidsminder Serie B in quarto 6.  
692

693 Schuldt, E. 1961. *Hohen Viecheln: ein mittelsteinzeitlicher Wohnplatz in Mecklenburg*.  
694 Berlin: Akademie-Verlag.  
695

696 Van Gijn, A. L. 1986. Fish Polish, Fact and Fiction. *Early Man News*, 9-10-11, 197–218.  
697

698 Van Gijn, A., and Little, A. in press. Tools, Use-Wear and Experimentation: Extracting  
699 Plants from Stone and Bone. In K. Hardy, and L. Kubiak-Martens (Eds.) *Wild Harvest:  
700 Plants and People in the Pre-Agricultural and Non-Agricultural World*. Oxford: Oxbow.  
701


702 Vaughan, P. 1985. *Use-Wear Analysis of Flaked Stone Tools*. Tucson: University of Arizona  
703 Press.  
704

705 Wheeler, A. 1978. Why Were There No Fish Remains at Star Carr? *Journal of  
706 Archaeological Science*, 5, 85-89.  
707

708 Wheeler, A., and Jones, A. K. G. 2009. *Fishes*. Cambridge: Cambridge University Press.  
709

710 Wumdsch, H. H. 1961. Die Fischreste. In E. Schuldt (Ed.) *Hohen Viecheln: ein  
711 mittelsteinzeitlicher Wohnplatz in Mecklenburg*, 70-74. Berlin: Akademie-Verlag.  
712

713 Zabilska, M. 2013. Srodowiskowe i kulturowe uwarunkowania rybołówstwa ludów wybrzeży  
714 Bałtyku w V-IV tysiącleciu przed Chrystusem na przykładzie osady w Dąbkach, stan. 9 (gm.  
715 Darłowo), cz. 1. Unpublished PhD thesis: Nicolaus Copernicus University.  
716  
717  
718  
719




720  
721 Figure 1  
722



723  
724 Figure 2  
725





726

727 Figure 3

728



729

730 Figure 4


731



732

733 Figure 5

734






735



736 Figure 6

737



738

739 Figure 7

740



741

742 Figure 8

743



744

745 Figure 9

746



747

748 Figure 10

749

750