



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/100764/>

Version: Accepted Version

Proceedings Paper:

Martel, A., Arvaneh, M., Dockree, P. et al. (2016) Neurophysiological correlates of mind-wandering, towards a predictive BCI-based neurofeedback. In: Müller-Putz, G.R., Huggins, J.E. and Steyrl, D., (eds.) Proceedings of the Sixth International Brain-Computer Interface Meeting: BCI Past, Present, and Future. 6th International BCI Meeting : BCI Past, Present and Future, 30 May - 03 Jun 2016, Pacific Grove, CA, USA. Graz University of Technology, p. 131. ISBN: 9783851254679.

<https://doi.org/10.3217/978-3-85125-467-9-131>

© 2016 The Authors. Article available under the terms of the CC-BY-NC-ND licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Neurophysiological correlates of mind-wandering, towards a predictive BCI-based Neurofeedback

A. Martel^{1*}, M. Arvaneh², P. Dockree¹, I. Roberston¹

¹Trinity Institute of Neuroscience, Trinity College Dublin, Dublin, Ireland; ²Department of Automatic Control and Systems Engineering, The University of Sheffield, Sheffield, U.K.

*A. Martel, Trinity College Dublin, Dublin, Ireland. E-mail: amartel@tcd.ie

Introduction: A brain computer interface (BCI) able to predict the disengagement of attention, e.g. mind-wandering episodes, regardless of the task being performed bears useful applications. Converging neuroscientific evidence has determined that our ability to remain attentive to a task, over a prolonged period of time, is subject to strong fluctuations. Recent electroencephalography (EEG) studies have identified neurophysiological signals reflecting inadequate task engagement preceding attentional lapses, in particular modulation of P300 amplitude and α -activity [1, 2, 3]. The present study investigated the evolution of neurophysiological signals preceding the report of 5 different levels of attention, from on-task to mind-wandering without awareness, during a breath-counting task and a sustained attention to response task (SART, [4]).

Material and Methods: Twenty-six healthy subjects (12 female; 23.4 ± 3.2 years) performed two tasks, a breath-counting task and a fixed version of the SART. The former asked participant to fixate a cross, count each breath from 1 to 9 and press the left mouse button for the first eight counts and the right mouse button for the final count before starting anew. The SART presented subjects with numbers from 1 to 9 subsequently and participants were required to respond with a button press to each number except the target (here number 6). Subjects were instructed to monitor their attention and interrupt the task if they noticed their thoughts stray from the task. Thought probes also interrupted the task at pseudo-random times. The probes presented subjects with a 5-step scale to categorize their level of attention/mind-wandering just prior to the interruption, (1) on-task (no mind-wandering), (2) on thoughts pertaining to the task, (3) distracted by internal sensations or external distractions, (4) on reminiscing or planning thoughts, (5) daydreaming (mind-wandering without awareness).




Figure 1. Smoothed average α band power measured over channel POz for all participants with standard errors of the mean. Left; the average α band power for the 3000ms period prior to the on-task (blue) and planning/reminiscing (magenta) reports during the breath counting task. Right; the average α band power for the 3500ms period prior to the on-task (blue) and daydreaming (red) reports during the SART (the sudden desynchronisation at around -1900ms is the result of the appearance of the SART stimulus).

Preliminary results: Analysis of the 64-channel EEG recorded data revealed that average α activity prior to thought probes differentiates between the on-task and planning/reminiscing condition during the breath counting task and between the on-task and daydreaming with awareness reports during the SART (Fig. 1). These preliminary results are encouraging for further analysis to determine whether mind-wandering could be detected online for the development of a Neurofeedback training (NFT).

References

- [1] Martel, A., Dähne, S., Blankertz, B. (2014). EEG Predictors of Covert Vigilant Attention. *Journal of Neural Engineering*, 11(3):035009. doi: 10.1088/1741-2560/11/3/035009
- [2] Smallwood, J., Beach, E., Schooler, J. W., & Handy, T. C. (2008). Going AWOL in the brain: mind wandering reduces cortical analysis of external events. *Journal of cognitive neuroscience*, 20(3), 458–469. doi:10.1162/jocn.2008.20037
- [3] O’Connell, R. G.; Dockree, P. M.; Robertson, I. H.; Bellgrove, M. A.; Foxe, J. J.; Kelly, S. P. (2009): Uncovering the Neural Signature of Lapsing Attention: Electrophysiological Signals Predict Errors up to 20 s before They Occur. In: *Journal of Neuroscience* 29 (26), S. 8604–8611
- [4] Robertson, I. H., Manly, T., Andrade, J., Baddeley, B. T., & Yiend, J. (1997). Oops: Performance correlates of everyday attentional failures in traumatic brain injured and normal subjects. *Neuropsychologia*, 35, 747–758.