



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/101592/>

Version: Accepted Version

Article:

Fordy, AP (2017) A note on some superintegrable Hamiltonian systems. *Journal of Geometry and Physics*, 115. pp. 98-103. ISSN: 0393-0440

<https://doi.org/10.1016/j.geomphys.2016.06.005>

© 2016 Elsevier B.V. This is an author produced version of a paper published in the *Journal of Geometry and Physics*. Uploaded in accordance with the publisher's self-archiving policy. This version licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

A note on some superintegrable Hamiltonian systems

Allan Fordy, School of Mathematics,
University of Leeds
e-mail a.p.fordy@leeds.ac.uk

January 14, 2016

Abstract

We consider some examples of superintegrable system which were recently isolated through a differential Galois group analysis. The identity of these systems is clarified and the corresponding Poisson algebras derived.

Keywords: Hamiltonian system, super-integrability, Poisson algebra, Euclidean algebra.
MSC: 17B63,37J15,37J35,70H06,70H20

1 Introduction

A Hamiltonian system of n degrees of freedom, Hamiltonian H , is said to be *completely integrable in the Liouville sense* if we have n independent functions I_n , which are *in involution* (mutually Poisson commuting), with H being a function of these and typically just one of them. Whilst n is the maximal number of independent functions which can be *in involution*, it is possible to have further integrals of the Hamiltonian H , which necessarily generate a non-Abelian algebra of integrals of H . The maximal number of additional *independent* integrals is $n - 1$, since the “level surface” of $2n - 1$ integrals (meaning the intersection of individual level surfaces) is just the (unparameterised) integral curve. Well known elementary examples are the isotropic harmonic oscillator, the Kepler system and the Calogero-Moser system. The quadratures of complete integrability are often achieved through the separation of variables of the Hamilton-Jacobi equation. The solution of a maximally super-integrable system can also be calculated purely algebraically (albeit implicitly), requiring just the solution of the equations $I_k = c_k$, $k = 1, \dots, 2n - 1$. Maximally superintegrable systems have a number of interesting properties: they can be separable in more than one coordinate system; all bounded orbits are closed; they give rise to interesting Poisson algebras with polynomial Poisson relations. The idea can be extended to *quantum integrable systems*, with first integrals replaced by commuting differential operators. For some examples of superintegrable quantum systems it is possible to use the additional commuting operators to build sequences of eigenfunctions [2, 3]. There is a large literature on the classification and analysis of superintegrable systems (see the review [5]) and they naturally occur in many applications in physics (additional integrals being referred to as “hidden symmetries”) [1].

A superintegrable system may occur in some other context and not be written in any recognisable form, so requires some judicious change of coordinates. In [7] the authors considered Hamiltonian functions of the form

$$H = \frac{1}{2}r^{m-k} \left(p_r^2 + \frac{p_\varphi^2}{r^2} \right) + r^m U(\varphi), \quad (1)$$

using differential Galois theory to determine necessary conditions for complete integrability. Restricting attention to the case $U(\varphi) = -\cos \varphi$, they found 8 cases, four of which were not only completely integrable, but also superintegrable (see their list (3.8)). In fact, for each case, they also presented 2 first integrals, F_1

and F_2 , which are quadratic in momenta, which means that each of these systems separates in 2 different coordinate systems.

In this paper we consider these 4 examples in more detail. The motivation of [7] was to extend the ideas of “homogeneous systems” to curved spaces, although the metric implied by the kinetic energy of (1) is, in fact, flat. A transformation to flat coordinates is derived in Section 2, giving us 3 “geometric” first integrals of the kinetic energy (related to the Killing vectors of the Euclidean metric), which are used to build quadratic integrals (specifically chosen to reproduce the examples of [7]). In fact, they share a common first integral, which allows us to use parabolic coordinates as common separation variables. The existence of the additional first integral in the superintegrable cases restricts the general separable potential to a finite parameter family. The 4 examples belong to two such superintegrable reductions, discussed in Section 3. It is shown that 2 of the examples actually have a first order integral. The results of Section 3 are translated back into the original coordinates in Section 4 and compared with [7].

2 Flat Coordinates for the Metric of (1)

It is easy to check that the inverse of the kinetic energy tensor of (1) defines a flat metric. Rather than compute the curvature, we calculate flat (Euclidean) coordinates (Q_1, Q_2) . We first summarise the algebraic properties of the Euclidean algebra (the algebra of Killing vectors of the Euclidean metric), working within the Hamiltonian framework.

2.1 The Euclidean Algebra

In a Euclidean space, with coordinates (Q_1, Q_2) and conjugate momenta (P_1, P_2) , the kinetic energy takes the form

$$T_e = \frac{1}{2} (P_1^2 + P_2^2). \quad (2)$$

The coordinates Q_i satisfy the simple relations

$$\{\{Q_i, T_e\}, T_e\} = 0, \quad P_i = \{Q_i, T_e\}, \quad \{Q_i, \{Q_i, T_e\}\} = 1. \quad (3)$$

The Hamiltonian T_e has 3 first degree integrals

$$K_1 = P_1, \quad K_2 = P_2, \quad K_3 = Q_2 P_1 - Q_1 P_2, \quad (4a)$$

which satisfy the usual Euclidean algebra Poisson relations:

$$\{K_1, K_2\} = 0, \quad \{K_1, K_3\} = K_2, \quad \{K_2, K_3\} = -K_1, \quad (4b)$$

with Casimir $T_e = \frac{1}{2}(K_1^2 + K_2^2)$, and correspond to the 3 Killing vectors of the 2D Euclidean metric:

$$v_i = \{Q_1, K_i\} \frac{\partial}{\partial Q_1} + \{Q_2, K_i\} \frac{\partial}{\partial Q_2} \quad \Rightarrow \quad [v_i, v_j] = -v_{\{K_i, K_j\}}.$$

2.2 Transformation between (r, φ) and (Q_1, Q_2)

Written in terms of any other coordinate system, K_i must satisfy the same relations. Starting with

$$T_r = \frac{1}{2} r^n \left(p_r^2 + \frac{p_\varphi^2}{r^2} \right), \quad (5)$$

we may consider $\{\{Q, T_r\}, T_r\} = 0$ as a system of *partial differential equations* for function $Q(r, \varphi)$ (the coefficients of $p_r^2, p_r p_\varphi, p_\varphi^2$):

$$\left(r^{\frac{n}{2}} Q\right)_r = 0, \quad \left(r^{\frac{n-2}{2}} Q\right)_{r\varphi} = 0, \quad 2Q_{\varphi\varphi} - (n-2)rQ_r = 0.$$

This is an overdetermined system whose integrability conditions are guaranteed by flatness. The solution contains two essential parameters, so we define

$$Q_1 = b_1 r^{1-n/2} \cos \frac{(n-2)\varphi}{2}, \quad Q_2 = b_2 r^{1-n/2} \sin \frac{(n-2)\varphi}{2}, \quad n \neq 2. \quad (6a)$$

The relation $\{Q_i, \{Q_i, T_r\}\} = 1$ implies $b_1 = b_2 = 2/(2-n)$, and

$$\begin{aligned} P_1 &= \{Q_1, T_r\} = r^{n/2} \left(p_r \cos \frac{(n-2)\varphi}{2} + \frac{p_\varphi}{r} \sin \frac{(n-2)\varphi}{2} \right), \\ P_2 &= \{Q_2, T_r\} = r^{n/2} \left(p_r \sin \frac{(n-2)\varphi}{2} - \frac{p_\varphi}{r} \cos \frac{(n-2)\varphi}{2} \right). \end{aligned} \quad (6b)$$

In these coordinates, $K_1 = P_1$ and $K_2 = P_2$ (now defined by (6b)) and $K_3 = \frac{2p_\varphi}{(2-n)}$, which, of course, satisfy the same Poisson relations (4b), with Casimir $T_r = \frac{1}{2}(K_1^2 + K_2^2)$.

Remark 2.1 *There is an alternative solution for the case $n = 2$, but this is not of interest to us here.*

3 Separability and Superintegrability

We start with the standard total energy in Euclidean space:

$$H = \frac{1}{2}(P_1^2 + P_2^2) + h(Q_1, Q_2) \quad (7a)$$

and require a quadratic (in momenta) first integral

$$F = T_f + f(Q_1, Q_2) = (F_{11}(Q_1, Q_2)P_1^2 + 2F_{12}(Q_1, Q_2)P_1P_2 + F_{22}(Q_1, Q_2)P_2^2) + f(Q_1, Q_2). \quad (7b)$$

The formula $\{H, F\} = 0$ has cubic and linear parts, which must separately vanish:

$$\{T_e, T_f\} = 0, \quad \{T_e, f\} + \{h, T_f\} = 0.$$

The first of these is equivalent to asking for the matrix F_{ij} to define a second order Killing tensor of the metric corresponding to the kinetic energy T_e . For a general kinetic energy, finding such a Killing tensor could be a difficult (and generally impossible) task. However, it is well known that for a flat metric, all second (and higher) order Killing tensors are built from tensor products of Killing vectors (see [4]). In our context, this means that the quadratic part of F is just some *quadratic form* of the functions K_i , given in (4), so there is a 6-parameter family of them. Two such quadratic forms are equivalent if they differ by a multiple of T_e .

The existence of a single quadratic integral corresponds to *separability* (in some coordinate system which diagonalises *both* H and F). In such a case, $h(Q_1, Q_2)$ (and the corresponding $f(Q_1, Q_2)$) depends upon two arbitrary functions (each of a single variable). The existence of two quadratic integrals (the *superintegrable case*) restricts these arbitrary functions, so that the potential function is *fixed* up to a finite number of arbitrary *parameters*.

3.1 Parabolic Coordinates

The choice of separation variables is purely a property of the quadratic part of F . The corresponding pair of quadratic forms are simultaneously diagonalised in these separation coordinates. The examples in this paper are chosen in order to include those of [7] as special cases. They each have two quadratic integrals, one of which has the form

$$F_1 = T_1 + f_1(Q_1, Q_2), \quad \text{where} \quad T_1 = K_2 K_3 = Q_2 P_1 P_2 - Q_1 P_2^2, \quad (8)$$

corresponding to (see [6]) parabolic coordinates (u, v) , defined by

$$u = \frac{1}{2}(Q_1 + \sqrt{Q_1^2 + Q_2^2}), \quad v = \frac{1}{2}(-Q_1 + \sqrt{Q_1^2 + Q_2^2}), \quad \text{with inverse} \quad Q_1 = u - v, \quad Q_2 = 2\sqrt{uv}. \quad (9a)$$

The corresponding canonical transformation is generated by

$$S = (u - v)P_1 + 2\sqrt{uv} P_2 \quad \Rightarrow \quad Q_1 = u - v, \quad Q_2 = 2\sqrt{uv}, \quad P_1 = \frac{up_u - vp_v}{u + v}, \quad P_2 = \frac{\sqrt{uv}(p_u + p_v)}{u + v}, \quad (9b)$$

which leads to

$$K_1 = \frac{up_u - vp_v}{u + v}, \quad K_2 = \frac{\sqrt{uv}(p_u + p_v)}{u + v}, \quad K_3 = \sqrt{uv}(p_u - p_v), \quad (10a)$$

and

$$T_h = \frac{1}{2}(K_1^2 + K_2^2) = \frac{1}{2} \left(\frac{up_u^2 + vp_v^2}{u + v} \right), \quad T_1 = K_2 K_3 = \frac{uv(p_u^2 - p_v^2)}{u + v}, \quad (10b)$$

which are evidently simultaneously diagonalised. It is a simple matter to determine the separable potentials, satisfying $\{T_h + h(u, v), T_1 + f_1(u, v)\} = 0$. The results are well known (see [6]):

$$H = \frac{1}{2} \left(\frac{up_u^2 + vp_v^2}{u + v} + \frac{A(u) + B(v)}{u + v} \right), \quad F_1 = \frac{uv(p_u^2 - p_v^2)}{u + v} + \frac{vA(u) - uB(v)}{u + v}, \quad (11)$$

where $A(u), B(v)$ are arbitrary functions of their respective single variables.

3.2 Superintegrable Restriction with $F_2 = K_1 K_3 + f_2(u, v)$

If we require that a second (independent) function Poisson commutes with H , then the arbitrary functions $A(u)$ and $B(v)$ are specified up to a finite number of parameters. In the case

$$F_2 = K_1 K_3 + f_2(u, v) = \frac{\sqrt{uv}(p_u - p_v)(up_u - vp_v)}{u + v} + f_2(u, v), \quad (12a)$$

the coefficients of p_u and p_v , in $\{H, F_2\} = 0$, give formulae for the first derivatives of $f_2(u, v)$, with integrability condition:

$$2uA''(u) + A'(u) = 2vB''(v) + B'(v).$$

This separable equation is easily solved, after which the (now consistent) overdetermined system for $f_2(u, v)$ can also be solved. The result is

$$h(u, v) = \frac{1}{2} \left(\frac{2k_0 + k_1\sqrt{u} + k_2\sqrt{v}}{u + v} \right), \quad f_1(u, v) = \frac{k_0(v - u) + k_1v\sqrt{u} - k_2u\sqrt{v}}{u + v}, \quad (12b)$$

$$f_2(u, v) = \frac{2k_0\sqrt{uv} + \frac{1}{2}(u - v)(k_1\sqrt{v} - k_2\sqrt{u})}{u + v}.$$

With these potentials we have

$$\{H, F_1\} = 0, \quad \{H, F_2\} = 0, \quad \text{and define} \quad F_3 = \{F_1, F_2\}, \quad (13a)$$

which is a cubic integral, which cannot be written as a linear (or polynomial) combination of H, F_1 and F_2 (but cannot, of course, be independent of them). However,

$$\{F_1, F_3\} = 2F_2H + \frac{1}{4}k_1k_2, \quad \{F_2, F_3\} = -2F_1H + \frac{1}{8}(k_1^2 - k_2^2), \quad (13b)$$

forming a polynomial Poisson algebra. As a consequence of these Poisson relations (ie not using the specific representation), we find the Casimir function:

$$\mathcal{C} = 2H(F_1^2 + F_2^2) - F_3^2 + \frac{1}{4}(k_2^2 - k_1^2)F_1 + \frac{1}{2}k_1k_2F_2. \quad (13c)$$

Since the entire algebra is built as a symmetry algebra of H , this Casimir should be a function of H . Indeed, using the specific representation, we find

$$\mathcal{C} = 2k_0^2H + \frac{1}{4}k_0(k_1^2 + k_2^2),$$

which furnishes us with the expected algebraic relation between the four functions H, F_i .

Remark 3.1 (A second parabolic coordinate system) *Clearly, if we consider the pair T_h and $T_2 = K_1K_3$, we obtain another parabolic coordinate system (\bar{u}, \bar{v}) , corresponding to the interchange $Q_1 \leftrightarrow Q_2$, and now*

$$T_h = \frac{1}{2}(K_1^2 + K_2^2) = \frac{1}{2} \left(\frac{\bar{u}p_{\bar{u}}^2 + \bar{v}p_{\bar{v}}^2}{\bar{u} + \bar{v}} \right), \quad T_2 = K_1K_3 = \frac{\bar{u}\bar{v}(p_{\bar{v}}^2 - p_{\bar{u}}^2)}{\bar{u} + \bar{v}}.$$

If we compose these transformations, we obtain

$$\bar{u} = \frac{1}{2}(\sqrt{u} + \sqrt{v})^2, \quad \bar{v} = \frac{1}{2}(\sqrt{u} - \sqrt{v})^2,$$

and the corresponding canonical transformation is a concrete realisation of the Lie algebraic automorphism

$$\bar{K}_1 = K_2, \quad \bar{K}_2 = K_1, \quad \bar{K}_3 = -K_3.$$

As well as correctly transforming the quadratic parts of H, F_i , this transformation also correctly transforms the potential functions, so

$$(H, F_1, F_2) \mapsto (H, -F_2, -F_1) \quad \text{with} \quad (k_0, k_1, k_2) \mapsto \left(k_0, \frac{1}{\sqrt{2}}(k_1 + k_2), \frac{1}{\sqrt{2}}(k_1 - k_2) \right),$$

which implies $F_3 \mapsto -F_3$ and represents an automorphism of the above quadratic algebra.

3.3 Superintegrable Restriction with $F_2 = K_1K_2 + f_2(u, v)$

In the case

$$F_2 = K_1K_2 + f_2(u, v) = \frac{\sqrt{uv}(p_u + p_v)(up_u - vp_v)}{(u + v)^2} + f_2(u, v), \quad (14a)$$

a similar calculation of $\{H, F_2\} = 0$, with H given by (11), leads to

$$h(u, v) = \frac{1}{2}k(u - v), \quad f_1(u, v) = kuv, \quad f_2(u, v) = k\sqrt{uv}, \quad (14b)$$

from which we immediately see that $\{K_2, H\} = 0$. We then find

$$\{F_1, F_2\} = 2K_2(H - K_2^2), \quad \{K_2, F_1\} = -F_2, \quad \{K_2, F_2\} = -\frac{k}{2}. \quad (15a)$$

The functions H, K_2, F_1, F_2 cannot, of course, be independent. They satisfy the algebraic relation

$$F_2^2 = 2HK_2^2 - K_2^4 + kF_1 \quad (15b)$$

4 Canonical Transformation to (r, φ) Coordinates

We now write the results of Section 3 in the original (r, φ) coordinates and compare them with the systems derived in [7].

Combining the coordinate transformations (6) and (9a), we obtain

$$u = \frac{2r^{1-n/2}}{2-n} \cos^2 \frac{(n-2)\varphi}{4}, \quad v = \frac{2r^{1-n/2}}{2-n} \sin^2 \frac{(n-2)\varphi}{4}, \quad (16a)$$

with the corresponding canonical transformation generated by

$$S = \frac{2r^{1-n/2}}{2-n} \left(p_u \cos^2 \frac{(n-2)\varphi}{4} + p_v \sin^2 \frac{(n-2)\varphi}{4} \right). \quad (16b)$$

In these coordinates, $K_1 = P_1$ and $K_2 = P_2$ (defined by (6b)) and $K_3 = \frac{2p_\varphi}{(2-n)}$. We can now write the superintegrable systems (12) and (14) in these coordinates.

The system (12) takes the form

$$\begin{aligned} H &= \frac{1}{2} r^n \left(p_r^2 + \frac{p_\varphi^2}{r^2} \right) + h(r, \varphi), \\ F_1 &= \frac{2}{2-n} \left(r^{n/2-1} p_\varphi \left(r p_r \sin \left(\frac{(n-2)\varphi}{2} \right) - p_\varphi \cos \left(\frac{(n-2)\varphi}{2} \right) \right) + f_1(r, \varphi) \right), \\ F_2 &= \frac{2}{2-n} \left(r^{n/2-1} p_\varphi \left(r p_r \cos \left(\frac{(n-2)\varphi}{2} \right) + p_\varphi \sin \left(\frac{(n-2)\varphi}{2} \right) \right) + f_2(r, \varphi) \right), \end{aligned}$$

where

$$\begin{aligned} h(r, \varphi) &= c_1 r^{(n-2)/2} + r^{(n-2)/4} \left(c_2 \cos \frac{(n-2)\varphi}{4} + c_3 \sin \frac{(n-2)\varphi}{4} \right), \\ f_1(r, \varphi) &= -c_1 \cos \frac{(n-2)\varphi}{2} - r^{(2-n)/4} \sin \frac{(n-2)\varphi}{2} \left(c_3 \cos \frac{(n-2)\varphi}{4} - c_2 \sin \frac{(n-2)\varphi}{4} \right), \\ f_2(r, \varphi) &= c_1 \sin \frac{(n-2)\varphi}{2} + \frac{1}{2} r^{(2-n)/4} \cos \frac{(n-2)\varphi}{2} \left(2c_2 \sin \frac{(n-2)\varphi}{4} - 2c_3 \cos \frac{(n-2)\varphi}{4} \right). \end{aligned}$$

Remark 4.1 Cases 1 and 2 of [7] (see their list (3.8)) correspond to $c_1 = c_3 = 0$, with $n = 6$ and $n = -2$, respectively.

The system (14) takes the form

$$\begin{aligned} H &= \frac{1}{2} r^n \left(p_r^2 + \frac{p_\varphi^2}{r^2} \right) + h(r, \varphi), \\ F_1 &= \frac{2}{2-n} \left(r^{n/2-1} p_\varphi \left(r p_r \sin \frac{(n-2)\varphi}{2} - p_\varphi \cos \frac{(n-2)\varphi}{2} \right) + f_1(r, \varphi) \right), \\ F_2 &= \frac{1}{2} r^n \left(\left(p_r^2 - \frac{p_\varphi^2}{r^2} \right) \sin(n-2)\varphi - \frac{p_r p_\varphi}{r} \cos(n-2)\varphi \right) + f_2(r, \varphi), \end{aligned}$$

where

$$h = c_1 r^{1-n/2} \cos \frac{(n-2)\varphi}{2}, \quad f_1 = c_1 r^{1-n/2} \sin \frac{(n-2)\varphi}{2}, \quad f_2 = -\frac{1}{2} c_1 r^{2-n} \sin^2 \frac{(n-2)\varphi}{2}.$$

Remark 4.2 Cases 3 and 4 of [7] (see their list (3.8)) correspond to $n = 0$ and $n = 4$, respectively.

Remark 4.3 The selection of the 4 values of n is just a consequence of the restriction to the case $U(\varphi) = -\cos \varphi$, in [7]. Comparing with the general formulae above, this just means choosing n so that $(n-2)/4 = \pm 1$ or $(n-2)/2 = \pm 1$.

References

- [1] M. Cariglia. Hidden symmetries of dynamics in classical and quantum physics. *Rev. Mod. Phys.*, 86:1283–1333, 2014.
- [2] A.P. Fordy. Quantum super-integrable systems as exactly solvable models. *SIGMA*, 3:025, 10 pages, 2007. <http://dx.doi.org/10.3842/SIGMA.2007.025>.
- [3] A.P. Fordy and M.J. Scott. Recursive procedures for Krall-Sheffer operators. *J Math Phys*, 54:043516, 2013. <http://arxiv.org/abs/1211.3075>.
- [4] R. Gilmore. *Lie Groups, Lie Algebras and Some of Their Applications*. Wiley, New York, 1974.
- [5] W. Miller Jr, S. Post, and P. Winternitz. Classical and quantum superintegrability with applications. *J. Phys. A*, 46:423001 (97pp), 2013.
- [6] A M Perelomov. *Integrable system of classical mechanics and Lie algebras*. Birkhäuser, Basel, 1990.
- [7] W. Szuminski, A.J. Maciejewski, and M. Przybylska. Note on integrability of certain homogeneous Hamiltonian systems. *Phys.Lett.A*, 379:2970–6, 2015.