

This is a repository copy of *Pathogen invasion indirectly changes the composition of soil microbiome via shifts in root exudation profile*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/103275/>

Version: Accepted Version

Article:

Gu, Yian, Wei, Zhong, Wang, Xueqi et al. (7 more authors) (2016) Pathogen invasion indirectly changes the composition of soil microbiome via shifts in root exudation profile. *Biology and fertility of soils*. 997–1005. ISSN: 1432-0789

<https://doi.org/10.1007/s00374-016-1136-2>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Pathogen invasion indirectly changes the composition of soil microbiome via shifts in root**
2 **exudation profile**

3 Yian Gu ¹, Zhong Wei ^{1*}, Xueqi Wang ¹, Ville-Petri Friman ², Jianfeng Huang ³, Xiaofang Wang ¹,
4 Xinlan Mei ¹, Yangchun Xu ^{1**}, Qirong Shen ¹, and Alexandre Jousset ^{1,4}

5

6 1. Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, National Engineering Research
7 Center for Organic-based Fertilizers, Nanjing Agricultural University, Nanjing, 210095, PR China

8 2. University of York, Department of Biology, Wentworth Way, York, YO10 5DD, UK

9 3 Institute of Agricultural Resources and Environment, Guangdong Key Laboratory of Nutrient
10 Cycling and Farmland Conservation, Guangdong Academy of Agricultural Sciences, Guangzhou
11 510640, PR China

12 4. Institute for Environmental Biology, Ecology & Biodiversity, Utrecht University, Padualaan 8,
13 3584CH Utrecht, The Netherlands

14

15 *Corresponding author

16 **Corresponding author

17 E-mail address: ycxu@njau.edu.cn (Yangchun Xu), Tel.: +86 025 84396824; fax: +86 025 84396260;

18 weizhong@njau.edu.cn (Zhong Wei), Tel.: +86 025 84396864; fax: +86 025 84396260

19

20 **Acknowledgements**

21 We thank Wu Xiong from Nanjing Agricultural University for help with bioinformatic analysis. Joana
22 Falcao Salles from University of Groningen is acknowledged for providing helpful advices. This
23 research was financially supported by the National Key Basic Research Program of China
24 (2015CB150503), the National Natural Science Foundation of China (31501837, 41301262, 41471213),
25 the Natural Science Foundation of Jiangsu Province (BK20130677), the 111 project (B12009), the
26 Priority Academic Program Development (PAPD) of Jiangsu Higher Education Institutions (Qirong
27 Shen), and the Qing Lan Project (Yangchun Xu).

28

Abstract

Plant-derived root exudates modulate plant-microbe interactions and may play an important role in pathogen suppression. Root exudates may for instance directly inhibit pathogens or alter microbiome composition. Here we tested if plants modulate their root exudation in the presence of a pathogen, and if these shifts alter the rhizosphere microbiome composition. We added exudates from healthy and *Ralstonia solanacearum*-infected tomato plants to an unplanted soil and followed changes in bacterial community composition. The presence of pathogen changed the exudation of phenolic compounds and increased the release of caffeic acid. The amendment of soils with exudates from the infected plants led to a development of distinct and less diverse soil microbiome communities. Crucially, we could reproduce similar shift in microbiome composition by adding pure caffeic acid into the soil. Caffeic acid further suppressed *R. solanacearum* growth *in vitro*. We conclude that pathogen-induced changes in root exudation profile may serve to control pathogen both by direct inhibition and by indirectly shifting the composition of rhizosphere microbiome.

Keywords

Amplicon sequencing; Phenolics; *Ralstonia solanacearum*; Root exudation; Root-pathogen interaction; Soil microbiome

Introduction

Plants invest a considerable fraction of their photosynthesized carbon into root exudates, a collection of low-molecular-weight compounds released into the rhizosphere (Bais et al. 2006). These exudates mediate complex interactions between plants and soil microbes and are essential in structuring the composition of soil microbiome (Carvalhais et al. 2015; Chaparro et al. 2013; Lagos et al. 2014). One key function of root exudates is to suppress pathogenic microorganisms (Bais et al. 2005) which is largely mediated by phenolic compounds (Badri et al. 2013; Lanoue et al. 2009). This function can be direct, for instance by inhibiting the growth of pathogen (Ling et al. 2013). Alternatively, phenolic compounds could affect the pathogen indirectly, for instance by modulating the expression of antibiotics-related genes of non-pathogenic soil microbes (de Werra et al. 2011). Such indirect effects could be very important, as the rhizosphere is enriched with mutualistic microbes that can protect plants against diseases (Li et al. 2015; Qiu et al. 2013; Trivedi et al. 2011) by producing antimicrobial compounds and lytic enzymes, stimulating plant immunity and intensifying competition for resources with the pathogen (Berendsen et al. 2012; Yu et al. 2014).

Disruption of the pathogen response-related jasmonic acid pathway alters root exudation patterns and the composition of rhizosphere microbial communities (Carvalhais et al. 2015) in *Arabidopsis thaliana*, confirming that shifts in exudation may be an integral part of plant response to pathogens. In this study, we addressed whether challenging plants with a pathogen alters the composition of soil microbiome via shifts in root exudation profile.

Exudation is very dynamic and depends on the plant growth stage (Chaparro et al. 2013) and the presence of pathogen. For example, the presence of the pathogenic fungus *Fusarium graminearum* in the rhizosphere of barley triggers the exudation of many phenolic compounds that prevent spore

germination (Lanoue et al. 2009). Similarly, alterations of phenolic compound exudation in barley infected with the oomycete *Pythium ultimum* induce expression of antibiotics-related genes in *Pseudomonas protegens* (Jousset et al. 2011).

In this study we assessed whether challenging plants with a pathogen leads to shifts in exudation patterns. We further assessed whether pathogen-induced exudates could inhibit growth of the pathogen and alter microbiome composition. We challenged tomato plants with *Ralstonia solanacearum*, a cosmopolite pathogen which causes bacterial wilt in more than 200 host species (Salanoubat et al. 2002). In order to disentangle the plant-mediated effects from pathogen-induced disturbance, we collected tomato exudates in the absence and presence of *R. solanacearum*. We sterilized the exudates and added them to an unplanted soil to mimic rhizosphere condition without direct pathogen influence. We then compared the effects of the different exudates on soil microbiome composition and linked them to changes in exudate composition.

Materials and methods

Bacterial strain and plant species

The bacterial pathogen *R. solanacearum* strain QL-Rs1115 (Wei et al. 2011) was routinely cultivated in NB medium (10.0 g of glucose, 5.0 g of peptone, 0.5 g of yeast extract, and 3.0 g of beef extract in 1 L of H₂O at pH 7.0). Overnight-grown bacteria were harvested by centrifugation (10,000×g for 6 min), washed twice with sterile saline solution (0.9% NaCl) and diluted to appropriate concentrations based on their optical density (OD₆₀₀).

Solanum lycopersicum cv. ‘Micro-Tom’ tomato was used as a model plant species. Seeds were surface sterilized with NaClO (3% v:v) for 10 min and rinsed four times with sterile distilled water. Surface-sterilized seeds were then plated on Murashige and Skoog agar medium (Murashige and Skoog

1962) supplemented with 1% sucrose and incubated in the dark at 28°C for 2 days, until the emergence of roots.

Collection of root exudates during the pathogen challenge

Root exudates were collected based on a previously published methodology (Badri et al. 2009) with minor modifications. Briefly, 2-day-old tomato seedlings were transferred to 6-well culture plates (Corning, CA, USA); each well contained two seedlings in 2 ml of liquid Murashige and Skoog medium amended with 1% sucrose. Plates were incubated on an orbital shaker at 90 rpm and exposed to white fluorescent light ($50 \mu\text{mol m}^{-2} \text{s}^{-1}$) with a 16:8 h light:dark photoperiod at $25 \pm 2^\circ\text{C}$. After ten days, plants were gently washed with sterile double-distilled water to remove the remaining exudates and transferred to new 6-well culture plates containing 2 ml of sterile double-distilled water per well. Sterilized double-distilled water was used to prevent the medium from interfering with the subsequent high-performance liquid chromatography (HPLC) analyses (Badri et al. 2013). We set up three treatments: a) *R. solanacearum* grown alone, b) Tomato plants grown alone and c) Tomato plants grown with *R. solanacearum*. Plants were inoculated with 20 μl of a bacterial suspension ($\text{OD}_{600} = 0.5$; $2 \times 10^8 \text{ CFU ml}^{-1}$ in 0.9% NaCl), or 20 μl of 0.9% NaCl. To obtain secretion of *R. solanacearum*, sterilized double-distilled water was inoculated with 20 μl of a bacterial suspension ($\text{OD}_{600} = 0.5$; $2 \times 10^8 \text{ CFU ml}^{-1}$ in 0.9% NaCl). Each treatment had three replicates and each replicate contained pooled exudate from 12 wells (i.e., 24 plants). Liquid medium was collected 72 h after pathogen inoculation and the pathogen survival was measured by serial dilution plating on NA medium (10.0 g of glucose, 5.0 g of peptone, 0.5 g of yeast extract, 3.0 g of beef extract, and 15 g of agar in 1 L of H_2O at pH 7.0). Pooled samples were centrifuged ($10,000\times g$ for 6 min) and sterile-filtered ($0.22 \mu\text{m}$) to remove the pathogen and root cells. Samples were then lyophilized and redissolved in 300 μl of solvent

(methanol:water = 30:70; v:v). Similar to experiments conducted with *A. thaliana* (Badri et al. 2013; Rudrappa et al. 2008), the tomato plants incubated in sterile double-distilled water did not show any visible nutrient deficiency symptoms or toxicity symptoms during the 3-day sampling period. We collected exudates over a short time period corresponding to the latence phase of the infection, during which infection remained asymptomatic (Jacobs et al. 2012; Milling et al. 2011). This allowed us to measure the plant response while avoiding biases due to disease onset.

HPLC analyses

In this study, we focused on antimicrobial phenolics present in the tomato root exudates. An 20 μ l aliquot of each sample was injected for HPLC analysis following a previously described method (Ling et al. 2010). Briefly, separation was performed by gradient elution using an Agilent 1200 system (Agilent Technologies, CA, USA) with an XDB-C18 column (4.6 mm \times 250 mm). The solvent system consisted of acetonitrile and 2% (v:v) acetic acid. The UV-visible photodiode detector was set to 280 nm wavelength. Peaks were identified by comparing their retention times with standards that were run under the same conditions (Rudrappa et al. 2008). The standard phenolic compounds used in this study were caffeic acid, cinnamic acid, coumaric acid, syringic acid, ferulic acid, β -hydroxybenzoic acid, gallic acid, benzoic acid, phthalic acid, and vanillic acid.

Effects of root exudates on microbiome composition in plant-free soil microcosms

The soil used in this assay was collected from a tomato field in Qilin (118°57'E, 32°03'N), Nanjing, China, which shows a high bacterial wilt incidence. The soil is a yellow-brown earth (Udic Argosol) with following properties: pH 5.4, organic matter (OM) content of 24.6 g kg⁻¹, total N of 6.3 g kg⁻¹, available P of 172.9 mg kg⁻¹, and available K of 178 mg kg⁻¹. The soil was first cleared of plant debris, sieved (< 2 mm), homogenized thoroughly, and transferred to 24-well culture plates, with each well

receiving 1.8 g of soil (dry weight). The exudates of 60 plants for each replicate were collected by the method described above and redissolved in 7.5 ml of solvent (methanol:water = 30:70; v:v) before being used to supplement the soil. We set up four different exudate treatments (250 μ l d⁻¹): a) solvent only (methanol:water = 30:70; v:v; control), b) exudates produced in the absence of a pathogen, c) exudates produced in the presence of a pathogen, and d) caffeic acid (3.6 mM; i.e., 0.5 μ mol g⁻¹ d⁻¹). Each treatment had three replicates and each replicate well received 250 μ l of exudate solution per day for a total of 30 days. The caffeic acid treatment was set up to assess its specific role in modulating the composition of soil bacterial community when tomato roots were challenged with *R. solanacearum*. The amount of caffeic acid entering the soil (i.e., 0.5 μ mol g⁻¹ d⁻¹) was based on previous studies (Eilers et al. 2010; Paterson et al. 2007; Qu and Wang 2008; Zhou and Wu 2012). The 24-well plates were weighed each day and were replenished with sterile distilled water to maintain the soil moisture at 60% of its maximum water holding capacity. Plates were incubated in a growth chamber with a 16:8 h light:dark photoperiod at 25 \pm 2°C to mimic natural conditions. At the end of the soil microcosm experiment, all the soils were collected, thoroughly homogenized, and stored at -80°C.

Total DNA extraction and 16S rRNA amplicons sequencing

To characterize changes in the soil microbiome composition, soil DNA was extracted from 300 mg soil using the PowerSoil DNA Isolation Kit (Mo Bio, Carlsbad, CA, USA) according to the manufacturer's instructions. Three DNA extracts of each replicate were pooled and quantified using a NanoDrop (ThermoScientific, Wilmington, DE, USA). The V4 hypervariable regions of the bacterial 16S rRNA gene were PCR-amplified using the primers pairs 563F (5'-AYTGGGYDTAAAGVG-3') and 802R (5'-TACNVGGGTATCTAATCC-3') (Cardenas et al. 2010) with the following PCR conditions: the reaction mix (20 μ l) contained 4 μ l of 5X FastPfu buffer, 2 μ l of 2.5 mM dNTPs, 0.4 μ l of each primer

(5 μ M), 0.5 μ l of DNA sample, and 0.4 μ l of FastPfu polymerase (TransGen Biotech, Beijing, China). PCR amplification included 30 cycles of 95°C for 30 s, 55°C for 30 s, and 72°C for 30 s in an Applied Biosystems thermal cycler (GeneAmp PCR system 9700, Applied Biosystems, Foster City, CA, USA). For each DNA sample, three independent PCRs were performed and the triplicate products were pooled to minimize the bias of PCR amplification. The amplicon products were purified using an AxyPrep PCR Clean-up Kit (Axygen Biosciences, Union City, CA, USA) before performing agarose gel electrophoresis. The concentrations of the purified PCR products were determined with QuantiFluorTM-ST (Promega, WI, USA) before subjecting them to 250-nucleotide paired-end sequencing using an Illumina MiSeq platform at Shanghai Majorbio Bio-pharm Bio-technology Co., Ltd.

16S rRNA sequencing analysis

The sequence data were processed following the UPARSE pipeline (Edgar 2013). Briefly, read pairs from each sample were assembled, low-quality nucleotides (maximal expected error of 0.25) were removed, and reads shorter than 200 bp were discarded. After elimination of singletons, sequence reads were clustered into operational taxonomic units (OTUs) at a threshold of 97% similarity, followed by removal of chimeras using the UCHIME method (Edgar et al. 2011). The representative sequences and OTU tables obtained using the UPARSE pipeline were then analyzed using Mothur (Schloss et al. 2009). Sampling depth was equalized to the depth of the smallest sample (31,200 reads). The taxonomies of each OTUs were annotated using the RDP 16S rRNA classifier (Wang et al. 2007) with a confidence threshold of 80%. The composition of the bacterial community was clustered based on unweighted UniFrac distance metrics (Lozupone et al. 2007).

Influence of caffeic acid on the growth of *R. solanacearum*

Bacteria were grown in 96-well culture plates with each well containing 188 µl of diluted (1:5) NB medium, 2 µl of bacterial suspension ($OD_{600} = 0.5$) and 10 µl of caffeic acid (prepared in pure ethanol; Sigma-Aldrich, St. Louis, MO, USA) at a final concentration of 0, 5, 10, 20, 40, 80, 120 or 160 µM. Plates were incubated at 30°C with shaking (170 rpm). Bacterial growth was determined by measuring the optical density at 600 nm using a SpectraMax M5 (Molecular Devices, Sunnyvale, CA, USA). The percentage of growth inhibition was calculated according to the following formula: Percentage of growth inhibition = (Bacterial growth in the absence of caffeic acid - Bacterial growth in the presence of caffeic acid) \times Bacterial growth in the absence of caffeic acid⁻¹ \times 100%.

Statistical analysis and sequence accession number

Analysis of variance (ANOVA, Duncan's multiple range test) and Student's t-test were used to compare mean differences between the treatments by using SPSS (v. 19). Redundancy analysis (RDA) was performed using CANOCO (ETTEN 2005). Effect of caffeic acid on the growth of *R. solanacearum* QL-Rs1115 was assessed with a linear model. All of the raw sequences have been deposited in the DDBJ SRA under the accession number SRP068343.

Results

The effect of *Ralstonia solanacearum* presence on tomato root exudate profile

Redundancy analysis (RDA) showed clearly different root exudate profiles in the three different treatments (Fig. 1a). The first two principal coordinates explained 88.9% of the total variation of secretions among the individual samples. Redundancy analysis and Monte Carlo permutation test (499 unrestricted permutations) were used to identify the HPLC peaks that significantly influenced the overall chromatographic profile. Our results revealed that compounds 2 ($F = 24.0$, $p = 0.004$), 4 ($F = 23.9$, $p = 0.002$), 10 ($F = 15.9$, $p = 0.006$) and 13 ($F = 21.2$, $p = 0.002$) were found at significantly

higher concentrations in the tomato exudates compared to pathogen-only samples. In contrast, compounds 1 ($F = 22.3$, $p = 0.002$), 3 ($F = 22.3$, $p = 0.02$), 7 ($F = 22.3$, $p = 0.008$) and 12 ($F = 5.4$, $p = 0.016$) were more abundant in pathogen-only samples compared to plant-only or plant-and-pathogen samples. Crucially, pathogen presence changed the tomato root exudate profile by favoring the secretion of compounds 5 ($F = 4.6$, $p = 0.008$) and 15 ($F = 7.5$, $p = 0.008$). The compound 15 was further determined to be caffeic acid by comparing its retention time with known standards. Chromatographic profiles revealed a significant increase ($p = 0.002$, Student's t test) of caffeic acid under pathogen infection (Fig. 1b and S1). While caffeic acid was also detected from plant-only samples, it was never detected in pathogen-only samples, suggesting that it was produced by the plant and not the pathogen. Compound 5 was detected only in the plant-and-pathogen samples. Unfortunately, we were not able to identify all other compounds except caffeic acid. This could be due to the lack of suitable reference standards or potentially low compound solubility in the solvent (Carvalhais et al. 2015).

The effect of tomato exudates and caffeic acid on the soil microbiome composition

At the end of the experiment, soils inoculated with the solvent (control) had the highest bacterial OTU richness (Fig. 2a). Bacterial OTU richness of soil treated with *R. solanacearum*-infected plant root exudates was significantly ($p = 0.007$, Student's t test) lower than samples treated with non-infected plant exudates. Cluster analyses based on the unweighted Unifrac metrics showed that bacterial community replicates from the same treatment clustered together (Fig. 2b). Non-infected plant exudate treatment clustered with the control (soil treated with 30% methanol) and *R. solanacearum*-infected plant root exudate treatment clustered with the caffeic acid treatment suggesting that addition of caffeic acid could mimic the effects of pathogen-induced shifts in root exudates. We further classified the soil

bacterial communities into phylotypes consisting of eight major bacterial phyla (Fig. S2). In contrast to non-infected plant exudate treatment, *R. solanacearum*-infected plant exudate treatment was associated with an increase abundance of *Proteobacteria* and *Actinobacteria* and reduced abundance of *Firmicutes*, *Acidobacteria*, *Verrucomicrobia*, *Bacteroidetes*, *Gemmatimonadetes*, and *Candidatus* *Saccharibacteria* (Fig. 2c). Caffeic acid and infected plant exudates had fairly similar effects on microbial community composition. For example, caffeic acid treatment was also associated with an increased abundance of *Proteobacteria* and *Actinobacteria* and reduced abundance of *Firmicutes*, *Acidobacteria*, and *Verrucomicrobia* compared to control (Fig. 2d).

Correlation between bacterial community composition and root exudates

Changes in the abundance of 20 major bacterial genera (47.9% by average relative abundance) were associated with changes in the concentration of specific exudates (Fig. 3). RDA ordination by vector fitting revealed that compound 5 ($F = 13.4$, $p = 0.002$, Mont Carlo) and caffeic acid ($F = 12.5$, $p = 0.002$, Mont Carlo) as significantly correlated exudate components. For example, positive correlations were observed between caffeic acid, the unidentified compound 5 and the genera *Brachybacterium*, *Janibacter*, *Dyella*, *Rhodanobacter*, and *Intrasporangium*, and these bacterial genera showed higher abundances in the plant-and-pathogen exudate treatment. In contrast, negative correlations were observed between caffeic acid or unidentified compound 5 and the genera *Saccharibacteria*, *Arthrobacter*, *Phycococcus*, *Gaiella*, and *Subdivision 3*, and these bacterial genera showed higher abundance in the plant-only exudate treatment.

The effect of caffeic acid on *R. solanacearum* growth

To validate the role of caffeic acid in plant-pathogen interactions, the effects of pure caffeic acid on the growth of *R. solanacearum* QL-Rs1115 were measured *in vitro*. Caffeic acid moderately reduced the

growth of *R. solanacearum* QL-Rs1115 in a dose-dependent way that could be well modeled with Michaelis-Menten kinetics ($R^2 = 0.66$, $p < 0.001$; Fig. 4). At concentrations above 80 μM , the effects of caffeic acid on the growth of *R. solanacearum* QL-Rs1115 reached plateau.

Discussion

Root exudates are crucial for modulating the interactions between plants and soil microbes (Bais et al. 2006). One of the main functions of these exudates is to directly suppress soil-borne pathogenic microorganisms (Bais et al. 2005). However, root exudates may also have indirect negative effects on the pathogens via changes in commensal rhizosphere microbiome composition. Here we demonstrated shifts in root exudate profile and an elevated secretion of caffeic acid triggered by *R. solanacearum* invasion (Fig. 1 and S1). The shifts in root exudate profile further affect the composition of soil bacterial community (Fig. 2 and 3). We found that increased caffeic acid directly inhibited the growth of *R. solanacearum* QL-Rs1115 (Fig. 4). Together these results suggest that pathogen invasion can activate plant defences that inhibit pathogen growth directly and change the composition of soil microbiome indirectly via shifts in root exudation profile

Correlation analyses have revealed close relationship between root exudate profiles and the activity, biomass and composition of the rhizosphere microbiome (Badri et al. 2009; Haichar et al. 2008; Paterson et al. 2007). For example, an ATP-binding cassette transporter mutant of *Arabidopsis*, which increased the secretion of phenolics while reducing sugars secretion, showed changes in the composition of rhizosphere bacterial and fungal communities compared to the wild type (Badri et al. 2009). Other studies have directly demonstrated that root exudates can considerably mediate the activity, biomass and composition of soil microbiome through the application of artificial or natural blends of root secretions to soil samples in the absence of plants (Badri et al. 2013; Eilers et al. 2010;

Paterson et al. 2007). In this study, we added exudates from healthy and *Ralstonia solanacearum*-infected tomato plants to a reference soil. Root exudates from healthy and *Ralstonia solanacearum*-infected tomato plants can help to assemble distinct soil microbiomes (Fig. 2b).

In contrast to non-infected plant exudate treatment, *R. solanacearum*-infected plant exudate treatment was associated with an increase abundance of *Proteobacteria* and *Actinobacteria* and reduced abundance of *Firmicutes*, *Acidobacteria*, *Verrucomicrobia*, *Bacteroidetes*, *Gemmatimonadetes*, and *Candidatus Saccharibacteria* (Fig. 2c). Crucially, the effects of infected plant exudates on the composition of soil bacterial community could be mimicked by adding pure caffeic acid to the soil (Fig. 2b); even though the effect of caffeic acid was not identical, it was also associated with an increased abundance of *Proteobacteria* and *Actinobacteria* and reduced abundance of *Firmicutes*, *Acidobacteria*, and *Verrucomicrobia* compared to the control (Fig. 2d). In addition, changes in the concentration of caffeic acid and unidentified compound 5 were also significantly (positively or negatively) correlated with changes in the abundances of several major bacterial genera (Fig. 3). These observations are in agreement with the results of Badri et al. (2013) where phenolic-related compounds were found to significantly (positively or negatively) correlate with a higher number of OTUs when compared with other classes of compounds such as sugars, amino acids, and sugar alcohols. These results thus suggest that caffeic acid may have played an important role in modulating the composition of soil bacterial community when tomato roots are challenged with *R. solanacearum* and phenolics in exudates may be the predominant modulators of soil bacterial community composition.

In this study, we show that the presence of different exudates can act as a filter and decrease the bacterial OTU richness of the soil bacterial community (Fig. 2a). Previous studies showed similar results that plant roots drive a reduction in the bacterial richness of the rhizosphere (Peiffer et al. 2013;

Shi et al. 2015). In contrast to bulk soil, rhizospheric soil is generally considered to be enriched in fast-growing microbes which respond positively to carbon sources (Fierer et al. 2007; Peiffer et al. 2013). Reductions in bacterial community diversity in our work may due to the selection or enrichment of specific fast-growing taxa. For example, *Proteobacteria* and *Actinobacteria*, which have generally been characterized as fast-growing phyla (Goldfarb et al. 2011), respond positively to caffeic acid addition compared to control (Fig. 2d). However, we still do not know whether reductions in OTU abundance correspond to reductions in bacterial functional diversity.

Accumulating evidence suggests that plant roots can secrete diverse protective metabolites upon pathogen infection and phenolics in root exudates may function as general antimicrobial agents (Bais et al. 2004; 2005; Ling et al. 2010). For instance, barley, when challenged with the pathogen *F. graminearum*, rapidly induces the *de novo* synthesis of phenolic compounds that inhibit the germination of *F. graminearum* (Lanoue et al. 2009). Caffeic acid secreted by grafted watermelon is associated with resistance to *F. oxysporum* (Ling et al. 2013). On the other hand, many phenolics in the root exudates can act as metal chelators and may change the availability of metallic soil micronutrients (Bais et al. 2006). For example, caffeic acid can chelate Cu (II) and alleviate its phytotoxicity (Garau et al. 2015), while the possibility remains that pathogenic bacteria use chelators as a strategy to acquire micronutrients essential for virulence and pathogenicity (Hood and Skaar 2012; Oide et al. 2006). Therefore, the complex impacts of root exudates and caffeic acid on the pathogen might have, in turn, multiple effects on plant health.

In this study, we only concentrated on bacterial interactions in this study. This excludes many other important soil microbial interactions with protist predators, phages, nematodes and fungi (Berendsen et al. 2012). For example, mycorrhizal fungi have been shown to also affect the

composition of rhizosphere bacterial community (Lioussanne et al. 2010), having effects on the functioning of the soil ecosystem (Van der Heijden et al. 1998; Vogelsang et al. 2006). As a result, we still need a better understanding of plant-pathogen interactions in more complex soil microbial communities. This information would be especially helpful from the applied perspective to guide how to manipulate the soil microbiome composition in order to improve the plant health and the crop yield (Chaparro et al. 2012; Xue et al. 2015). For example, in addition to adding artificial plant exudates, one could apply symbiotic mycorrhizal fungi into the soil to increase the suppression of pathogens (Borowicz 2001; Rodriguez and Sanders 2015).

In conclusion, here we show that pathogen invasion causes clear changes in tomato root exudation profile by specifically increasing the secretion of phenolic compounds. This change had important effects on the composition of soil microbial community. The increased release of caffeic acid in root exudate had negative effect on the pathogen growth. Together these results suggest that pathogen-plant interactions can have community-wide effects on the composition of soil microbial communities.

References

- Badri DV, Chaparro JM, Zhang R, Shen Q, Vivanco JM (2013) Application of Natural Blends of Phytochemicals Derived from the Root Exudates of Arabidopsis to the Soil Reveal That Phenolic-related Compounds Predominantly Modulate the Soil Microbiome. *J Biol Chem* 288:4502-4512. doi:10.1074/jbc.M112.433300
- Badri DV, Quintana N, El Kassis EG, Kim HK, Choi YH, Sugiyama A, Verpoorte R, Martinoia E, Manter DK, Vivanco JM (2009) An ABC transporter mutation alters root exudation of phytochemicals that provoke an overhaul of natural soil microbiota. *Plant Physiol*

151:2006-2017. doi:10.1104/pp.109.147462

Bais HP, Park S-W, Weir TL, Callaway RM, Vivanco JM (2004) How plants communicate using the underground information superhighway. *Trends Plant Sci* 9:26-32. doi:10.1016/j.tplants.2003.11.008

Bais HP, Prithiviraj B, Jha AK, Ausubel FM, Vivanco JM (2005) Mediation of pathogen resistance by exudation of antimicrobials from roots. *Nature* 434:217-221. doi:10.1038/nature09809

Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM (2006) The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu Rev Plant Biol* 57:233-266. doi:10.1146/annurev.arplant.57.032905.105159

Berendsen RL, Pieterse CM, Bakker PA (2012) The rhizosphere microbiome and plant health. *Trends Plant Sci* 17:478-486. doi:10.1016/j.tplants.2012.04.001

Borowicz VA (2001) Do arbuscular mycorrhizal fungi alter plant-pathogen relations? *Ecology* 82:3057-3068. doi:10.1890/0012-9658(2001)082[3057:DAMFAP]2.0.CO;2

Cardenas E, Wu WM, Leigh MB, Carley J, Carroll S, Gentry T, Luo J, Watson D, Gu B, Ginder-Vogel M, Kitanidis PK, Jardine PM, Zhou J, Criddle CS, Marsh TL, Tiedje JM (2010) Significant association between sulfate-reducing bacteria and uranium-reducing microbial communities as revealed by a combined massively parallel sequencing-indicator species approach. *Appl Environ Microbiol* 76:6778-6786. doi:10.1128/AEM.01097-10

Carvalhais LC, Dennis PG, Badri DV, Kidd BN, Vivanco JM, Schenk PM (2015) Linking Jasmonic Acid Signaling, Root Exudates, and Rhizosphere Microbiomes. *Mol Plant-Microbe Interact* 28:1049-1058. doi:10.1094/MPMI-01-15-0016-R

Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM (2013) Root exudation of

354 phytochemicals in *Arabidopsis* follows specific patterns that are developmentally programmed
 355 and correlate with soil microbial functions. *PLoS One* 8:e55731.
 356 doi:10.1371/journal.pone.0055731
 357 Chaparro JM, Sheflin AM, Manter DK, Vivanco JM (2012) Manipulating the soil microbiome to
 358 increase soil health and plant fertility. *Biol Fertil Soils* 48:489-499.
 359 doi:10.1007/s00374-012-0691-4
 360 de Werra P, Huser A, Tabacchi R, Keel C, Maurhofer M (2011) Plant- and microbe-derived compounds
 361 affect the expression of genes encoding antifungal compounds in a *Pseudomonas* with
 362 biocontrol activity. *Appl Environ Microbiol* 77:2807-2812. doi:10.1128/AEM.01760-10
 363 Edgar RC (2013) UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat*
 364 *Methods* 10:996-998. doi:10.1038/nmeth.2604
 365 Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and
 366 speed of chimera detection. *Bioinformatics* 27:2194-2200. doi:10.1093/bioinformatics/btr381
 367 Eilers KG, Lauber CL, Knight R, Fierer N (2010) Shifts in bacterial community structure associated
 368 with inputs of low molecular weight carbon compounds to soil. *Soil Biol Biochem* 42:896-903.
 369 doi:10.1016/j.soilbio.2010.02.003
 370 ETTEEN EV (2005) Multivariate analysis of ecological data using CANOCO. *Austral Ecol* 30:486-487.
 371 doi:10.1111/j.1442-9993.2005.01433.x
 372 Fierer N, Bradford MA, Jackson RB (2007) Toward an ecological classification of soil bacteria.
 373 *Ecology* 88:1354-1364. doi:10.1890/05-1839
 374 Garau G, Mele E, Castaldi P, Lauro GP, Deiana S (2015) Role of polygalacturonic acid and the
 375 cooperative effect of caffeic and malic acids on the toxicity of Cu(II) towards triticale plants

376 (× Triticosecale Wittm). Biol Fertil Soils 51:535-544. doi:10.1007/s00374-015-0999-y

377 Goldfarb KC, Karaoz U, Hanson CA, Santee CA, Bradford MA, Treseder KK, Wallenstein MD, Brodie

378 EL (2011) Differential growth responses of soil bacterial taxa to carbon substrates of varying

379 chemical recalcitrance. Front Microbiol 2:94. doi:10.3389/fmicb.2011.00094

380 Haichar FZ, Marol C, Berge O, Rangel-Castro JI, Prosser JI, Balesdent J, Heulin T, Achouak W (2008)

381 Plant host habitat and root exudates shape soil bacterial community structure. ISME J

382 2:1221-1230. doi:10.1038/ismej.2008.80

383 Hood MI, Skaar EP (2012) Nutritional immunity: transition metals at the pathogen-host interface. Nat

384 Rev Microbiol 10:525-537. doi:10.1038/nrmicro2836

385 Jacobs JM, Babujee L, Meng F, Milling A, Allen C (2012) The in planta transcriptome of Ralstonia

386 solanacearum: conserved physiological and virulence strategies during bacterial wilt of tomato.

387 mBio 3:e00114-00112. doi:10.1128/mBio.00114-12

388 Jousset A, Rochat L, Lanoue A, Bonkowski M, Keel C, Scheu S (2011) Plants respond to pathogen

389 infection by enhancing the antifungal gene expression of root-associated bacteria. Mol

390 Plant-Microbe Interact 24:352-358. doi:10.1094/MPMI-09-10-0208

391 Lagos LM, Navarrete OU, Maruyama F, Crowley DE, Cid FP, Mora ML, Jorquera MA (2014)

392 Bacterial community structures in rhizosphere microsites of ryegrass (Lolium perenne var.

393 Nui) as revealed by pyrosequencing. Biol Fertil Soils 50:1253-1266.

394 doi:10.1007/s00374-014-0939-2

395 Lanoue A, Burlat V, Henkes GJ, Koch I, Schurr U, Röse US (2009) De novo biosynthesis of defense

396 root exudates in response to Fusarium attack in barley. New Phytol 185:577-588.

397 doi:10.1111/j.1469-8137.2009.03066.x

398 Li X, Zhang Yn, Ding C, Jia Z, He Z, Zhang T, Wang X (2015) Declined soil suppressiveness to
 399 *Fusarium oxysporum* by rhizosphere microflora of cotton in soil sickness. *Biol Fertil Soils*
 400 51:935-946. doi:10.1007/s00374-015-1038-8
 401 Ling N, Huang Q, Guo S, Shen Q (2010) *Paenibacillus polymyxa* SQR-21 systemically affects root
 402 exudates of watermelon to decrease the conidial germination of *Fusarium oxysporum* f.sp.
 403 *niveum*. *Plant Soil* 341:485-493. doi:10.1007/s11104-010-0660-3
 404 Ling N, Zhang W, Wang D, Mao J, Huang Q, Guo S, Shen Q (2013) Root exudates from grafted-root
 405 watermelon showed a certain contribution in inhibiting *Fusarium oxysporum* f. sp. *niveum*.
 406 *PLoS One* 8:e63383. doi:10.1371/journal.pone.0063383
 407 Lioussanne L, Perreault F, Jolicoeur M, St-Arnaud M (2010) The bacterial community of tomato
 408 rhizosphere is modified by inoculation with arbuscular mycorrhizal fungi but unaffected by
 409 soil enrichment with mycorrhizal root exudates or inoculation with *Phytophthora nicotianae*.
 410 *Soil Biol Biochem* 42:473-483. doi:10.1016/j.soilbio.2009.11.034
 411 Lozupone CA, Hamady M, Kelley ST, Knight R (2007) Quantitative and Qualitative Diversity
 412 Measures Lead to Different Insights into Factors That Structure Microbial Communities. *Appl*
 413 *Environ Microbiol* 73:1576-1585. doi:10.1128/aem.01996-06
 414 Milling A, Babujee L, Allen C (2011) *Ralstonia solanacearum* extracellular polysaccharide is a specific
 415 elicitor of defense responses in wilt-resistant tomato plants. *PLoS One* 6:e15853.
 416 doi:10.1371/journal.pone.0015853
 417 Murashige T, Skoog F (1962) A Revised Medium for Rapid Growth and Bio Assays with Tobacco
 418 Tissue Cultures. *Physiol Plant* 15:473-497. doi:10.1111/j.1399-3054.1962.tb08052.x
 419 Oide S, Moeder W, Krasnoff S, Gibson D, Haas H, Yoshioka K, Turgeon BG (2006) NPS6, encoding a

420 nonribosomal peptide synthetase involved in siderophore-mediated iron metabolism, is a
 421 conserved virulence determinant of plant pathogenic ascomycetes. *Plant Cell* 18:2836-2853.
 422 doi:10.1105/tpc.106.045633

423 Paterson E, Gebbing T, Abel C, Sim A, Telfer G (2007) Rhizodeposition shapes rhizosphere microbial
 424 community structure in organic soil. *New Phytol* 173:600-610.
 425 doi:10.1111/j.1469-8137.2006.01931.x

426 Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL, Buckler ES, Ley RE (2013) Diversity and
 427 heritability of the maize rhizosphere microbiome under field conditions. *Proc Natl Acad Sci U*
 428 *S A* 110:6548-6553. doi:10.1073/pnas.1302837110

429 Qiu M, Li S, Zhou X, Cui X, Vivanco JM, Zhang N, Shen Q, Zhang R (2013) De-coupling of root–
 430 microbiome associations followed by antagonist inoculation improves rhizosphere soil
 431 suppressiveness. *Biol Fertil Soils* 50:217-224. doi:10.1007/s00374-013-0835-1

432 Qu XH, Wang JG (2008) Effect of amendments with different phenolic acids on soil microbial biomass,
 433 activity, and community diversity. *Appl Soil Ecol* 39:172-179.
 434 doi:10.1016/j.apsoil.2007.12.007

435 Rodriguez A, Sanders IR (2015) The role of community and population ecology in applying
 436 mycorrhizal fungi for improved food security. *ISME J* 9:1053-1061.
 437 doi:10.1038/ismej.2014.207

438 Rudrappa T, Czymmek KJ, Pare PW, Bais HP (2008) Root-secreted malic acid recruits beneficial soil
 439 bacteria. *Plant Physiol* 148:1547-1556. doi:10.1104/pp.108.127613

440 Salanoubat M, Genin S, Artiguenave F, Gouzy J, Mangenot S, Arlat M, Billault A, Brottier P, Camus J,
 441 Cattolico L (2002) Genome sequence of the plant pathogen *Ralstonia solanacearum*. *Nature*

442 415:497-502. doi:10.1038/415497a

443 Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB,
 444 Parks DH, Robinson CJ (2009) Introducing mothur: open-source, platform-independent,
 445 community-supported software for describing and comparing microbial communities. *Appl*
 446 *Environ Microbiol* 75:7537-7541. doi:10.1128/AEM.01541-09

447 Shi S, Nuccio E, Herman DJ, Rijkers R, Estera K, Li J, da Rocha UN, He Z, Pett-Ridge J, Brodie EL,
 448 Zhou J, Firestone M (2015) Successional Trajectories of Rhizosphere Bacterial Communities
 449 over Consecutive Seasons. *mBio* 6:e00746. doi:10.1128/mBio.00746-15

450 Trivedi P, He Z, Van Nostrand JD, Albrigo G, Zhou J, Wang N (2011) Huanglongbing alters the
 451 structure and functional diversity of microbial communities associated with citrus rhizosphere.
 452 *ISME J* 6:363-383. doi:10.1038/ismej.2011.100

453 Van der Heijden MG, Klironomos JN, Ursic M, Moutoglis P, Streitwolf-Engel R, Boller T, Wiemken A,
 454 Sanders IR (1998) Mycorrhizal fungal diversity determines plant biodiversity, ecosystem
 455 variability and productivity. *Nature* 396:69-72. doi:10.1038/23932

456 Vogelsang KM, Reynolds HL, Bever JD (2006) Mycorrhizal fungal identity and richness determine the
 457 diversity and productivity of a tallgrass prairie system. *New Phytol* 172:554-562.
 458 doi:10.1111/j.1469-8137.2006.01854.x

459 Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naive Bayesian classifier for rapid assignment of
 460 rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 73:5261-5267.
 461 doi:10.1128/AEM.00062-07

462 Wei Z, Yang X, Yin S, Shen Q, Ran W, Xu Y (2011) Efficacy of *Bacillus*-fortified organic fertiliser in
 463 controlling bacterial wilt of tomato in the field. *Appl Soil Ecol* 48:152-159.

464 doi:10.1016/j.apsoil.2011.03.013

465 Xue C, Penton CR, Shen Z, Zhang R, Huang Q, Li R, Ruan Y, Shen Q (2015) Manipulating the banana

466 rhizosphere microbiome for biological control of Panama disease. *Sci Rep* 5:11124.

467 doi:10.1038/srep11124

468 Yu Z, Zhang Y, Luo W, Wang Y (2014) Root colonization and effect of biocontrol fungus *Paecilomyces*

469 lilacinus on composition of ammonia-oxidizing bacteria, ammonia-oxidizing archaea and

470 fungal populations of tomato rhizosphere. *Biol Fertil Soils* 51:343-351.

471 doi:10.1007/s00374-014-0983-y

472 Zhou X, Wu F (2012) P-coumaric acid influenced cucumber rhizosphere soil microbial communities

473 and the growth of *Fusarium oxysporum* f. Sp. *Cucumerinum* owen. *PLoS One* 7:e48288.

474 doi:10.1371/journal.pone.0048288

475




Fig. 1 Variation in the composition of exudates originating from pathogen-only (RS), plant-only (Tomato) and plant-and-pathogen (Tomato + RS) treatments as detected by HPLC. (a) The relationship between individual exudate compounds and root exudate composition as determined by RDA. The percentage of the explained variation is indicated on X and Y axes. Individual exudate compounds that were significantly ($p < 0.05$) correlated with the exudate composition are presented as arrows. Numbers indicate peaks of on the HPLC chromatogram. (b) The difference in the caffeic acid exudation in the absence and presence of *R. solanacearum*. Bars show mean values \pm SE ($n = 3$). ** indicate $p < 0.01$




Fig. 2 The effect of plant exudates and caffeic acid on the soil microbiome composition. (a) Bacterial OTU richness in control, caffeic acid, plant-only (Tomato) and plant-and-pathogen (Tomato + RS) treatments. Different letters indicate significant differences. (b) Community similarity based on the cluster analysis of unweighted Unifrac metrics. (c) The fold change of bacterial phyla in the plant-and-pathogen exudate treatment relative to the plant-only exudate treatment (d) The fold change of bacterial phyla in caffeic acid treatment relative to the control. Bars show mean values \pm SE (n=3)




Fig. 3 RDA ordination summarizing the correlations between the top 20 genus of soil bacterial communities and selected exudate compounds in the plant-only (Tomato) and plant-and-pathogen (Tomato + RS) treatments (superimposed as fitted vectors). The red arrows show the magnitude (length) and correlation (angle) of individual exudate components that were significantly ($p < 0.05$) correlated with the ordination




Fig. 4 The effect of caffeic acid on the growth of *R. solanacearum*. Growth of *R. solanacearum* QL-Rs1115 in 20% NB media (after 24 h at 30°C) was determined by measuring the absorbance at 600 nm (OD₆₀₀). The regression curve is based on Michaelis-Menten fitting. Bars show mean values \pm SE (n = 3)




Fig. S1 High-performance liquid chromatography (HPLC) profile of exudates originating from pathogen-only (RS; blue line), plant-only (Tomato; red line) and plant-and-pathogen together (Tomato + RS; green line) treatments




Fig. S2 The relative abundance of the major bacterial phyla in the control, caffeic acid, plant-only (Tomato) and plant-and-pathogen (Tomato + RS) treatments