

This is a repository copy of *Long-term experiments to investigate irreversibility in sorption of pesticides to soil*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/103689/>

Version: Accepted Version

Article:

Suddaby, Laura, Beulke, Sabine, van Beinum, Wendy et al. (3 more authors) (2016) Long-term experiments to investigate irreversibility in sorption of pesticides to soil.

CHEMOSPHERE. pp. 40-47. ISSN: 0045-6535

<https://doi.org/10.1016/j.chemosphere.2016.07.062>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Title

Long-term experiments to investigate irreversibility in sorption of pesticides to soil

Authors

Suddaby, L.A.^{1,2,3*}, Beulke, S.^{2,4}, van Beinum, W.^{2,5}, Oliver, R.G.^{3,6}, Kuet, S.^{3,7} and Brown, C.D.^{1,8}

¹*Environment Department, University of York;* ²*Food and Environment Research Agency (FERA);* ³*Syngenta Ltd*

*Corresponding author details: e-mail laura.suddaby@tsgeconsulting.com; phone: + 44 (0) 1423 799647; present address: TSGE Consulting Ltd, Concordia House, Grimbold Crag Court, St. James Business Park, Knaresborough, North Yorkshire, HG5 8QB, UK.

Abstract

Experiments investigated irreversibility in pesticide sorption to soil. Sorption behaviour under abiotic conditions was quantified for chlorotoluron, prometryn and hexaconazole in three soils over periods of up to 274 days. An isotope-exchange procedure was used whereby sorption of ¹²C- and ¹⁴C-pesticide isotopes in shaken suspensions of three soils (56-168 days shaking) was followed by substitution of the isotopes in the liquid phase and a 14-day exchange phase. This was followed by forced isotope exchange where the sorbed ¹⁴C material was exchanged by adding an excess of non-radiolabelled compound. Experiments were concluded with solvent extraction and soil combustion to determine remaining radioactivity. Under conditions of continuous shaking, the pesticide-soil systems took around four months to approach sorption equilibrium, resulting in strong asymmetry between the profiles of exchange for isotopes of all three compounds. Physically entrapped residues were released back into solution under the steep concentration gradient of forced isotope exchange and

¹ *Environment Department, University of York, Heslington, York, YO10 5DD, UK.*

² *Food and Environment Research Agency, Sand Hutton, York, YO41 1LZ, UK.*

³ *Syngenta, Jealott's Hill International Research Station, Bracknell, Berkshire, RG42 6EY, UK.*

⁴ sabine.beulke@enviresearch.com

⁵ wendy.vanbeinum@enviresearch.com

⁶ robin.oliver@syngenta.com

⁷ sui.kuet@syngenta.com

⁸ colin.brown@york.ac.uk

small amounts of radioactivity were still being released at the termination of the experiment. The profiles of exchange did not deviate markedly from ideal behaviour based on the assumption that sorption is fully reversible. Whilst the timescales for release of sorbed residues back into solution were very long, soil combustion at study termination only yielded <1-2% of applied radioactivity; this confirms that sorption processes under abiotic soil conditions were overwhelmingly reversible for this set of compounds and soils.

Key words

Pesticide, sorption, desorption, irreversible sorption, bound residue, non-extractable residue

Introduction

Sorption is a key process controlling the fate of a pesticide in soil as it determines the distribution between soil solid phase and soil water phase. Sorption (both adsorption and absorption) and desorption can be kinetically-controlled, typically involving multiple domains of sorption kinetics ranging from instantaneous to very slow with true equilibrium approached over months or years [1]. Pesticide sorption to soils is often reported to be only partially reversible, with a fraction of the sorbed pesticide apparently resistant to desorption at the tails of kinetically-controlled release [2]. The formation of non-extractable (bound) residues is important for the fate and transport of organic contaminants in environmental systems as it limits the bioavailability of pesticides in surface soil systems and has potential to reduce pesticide mobility in the environment [3].

Physical entrapment within the soil solid phase and covalent bonding to soil organic matter are two processes known to give rise to bound residues [4, 5]. Some parent compounds will be directly subject to covalent bonding [6, 7], but in many cases, the formation of bound

residues has been found to arise as a result of biodegradation of the parent compound to one or more metabolites that are prone to covalent bonding to organic matter. Several studies have demonstrated a large reduction in formation of bound residues in sterile soil where biodegradation to metabolites is inhibited [8]. Recent work has demonstrated that biodegradation of parent compound can also be followed by assimilation into biogenic material by soil microorganisms (*e.g.* as fatty acids and amino acids) [9]. Such biogenic material is excluded from conventional definitions of bound residues, but is indistinguishable in studies based solely on recovery of radiolabelled chemical from treated soil. Whilst the formation of bound residues effectively reduces bioavailability and transport of pesticide in soil, questions have been raised about long-term reversibility and possible implications for human and ecosystem health. In this context, the parent compound is generally most toxicologically active and thus of greatest concern [10]. For chemicals that are not prone to covalent bonding to soil organic matter, bound residues of the parent compound will be most frequently associated with physical entrapment. The current paper investigates whether or not there is an irreversibly bound fraction for pesticide residues sorbed to soil over extended periods of time.

Experimental approaches and instrumental methods used to characterise bound residues in soil were reviewed by Northcott and Jones [11]. Classical extraction procedures aimed to recover as much of the pesticide as possible using exhaustive extraction techniques [12]. A modification to the IUPAC definition of bound residues by Führ *et al.* [13] stipulated that “the extraction method must not substantially change the compounds themselves or the structure of the matrix”, meaning that bound residues now typically refer to those residues that cannot be extracted from soil by an organic solvent and without alteration to the chemical structure of the compound.

Celis and Koskinen [14, 15] presented an isotope exchange technique using ^{14}C to characterise pesticide exchange kinetics. The principle of the approach is pre-equilibration of duplicate tubes with soil-water slurry and dosed with either ^{12}C - or ^{14}C -pesticide; the tubes are centrifuged and the supernatants switched to allow characterisation of the kinetics of pesticide exchange and estimation of amounts of sorbed pesticide that do not participate in the exchange. The authors proposed that their method eliminated inherent experimental artefacts of other approaches such as the specific effectiveness of the extracting method and changes to the soil solid phase. Despite the promise of the approach, it is unlikely that true irreversible sorption could be measured during Celis and Koskinen's [14, 15] three-day tests. Sander and Pignatello [16, 17] used equilibration times of up to 140 days with the objective of ensuring sorption equilibrium in their forward isotope exchange experiment with the persistent hydrocarbons naphthalene and 1,4-dichlorobenzene. Therefore, sorption and desorption periods should be extended to widen the scope and environmental applicability of isotope exchange studies.

An isotope exchange study was carried out using three test compounds in three soils to characterise irreversibility in pesticide sorption-desorption to soils over time and to determine whether any irreversibly bound fraction of pesticide changes over time and/or differs for differing pesticides and soil types. The three pesticides were selected to have a range in sorption properties; they comprised chlorotoluron [*N'*-(3-Chloro-4-methylphenyl)-*N,N*-dimethylurea], prometryn [*N,N'*-bis(1-methylethyl)-6-(methylthio)-1,3,5-triazine-2,4-diamine], and hexaconazole [α -butyl- α -(2,4-dichlorophenyl)-1*H*-1,2,4-triazole-1-ethanol].

Materials and methods

Soils. Three soils from the UK with differing physico-chemical properties were selected for study. Fresh soils from 0-15 cm depth of a Blackwood loamy sand, Andover clay loam and

Salop sandy clay loam were collected, air-dried and passed through a 2-mm sieve (sampling sites and dates are given in Table 1). Soil characterisation was carried out by NRM Laboratories Ltd (Bracknell, UK; Table 1). Soils were gamma irradiated at 35.40 kGy by Isotron Ltd (Bradford, UK) to sterilise and thus inhibit microbial degradation. Irradiated soils were subsequently stored under sterile conditions (at 4°C in darkness) until use. It is assumed throughout that any impact of irradiation on the soil constituents did not materially influence pesticide sorption behaviour.

Chemicals. Analytical-grade ^{12}C -pesticides were purchased from Sigma-Aldrich Ltd (Dorset, UK). All ^{14}C -pesticides were supplied by Syngenta Ltd (Jealott's Hill International Research Centre, Bracknell, UK). Details of purity and specific activity are given in Table S1 in the Supporting Information. Preliminary batch-slurry experiments with ^{12}C -pesticides quantified sorption isotherms in the three soils after 24-h shaking and for seven initial concentrations of each pesticide. All chemicals were either received sterile or sterilised prior to use by autoclave.

Isotope exchange. Experiments comprised a sequence of adsorption (periods from 7 to 168 d), isotope exchange and re-equilibration over 14 days, forced isotope exchange (periods from 145 to 204 days), then finally solvent extraction, analysis for metabolites and soil combustion. These different procedures are described in turn below and any differences in methodology for experiments with the three pesticides are given in Table S1 in the Supporting Information. Long-term experiments commenced with the isotope exchange technique based on the method described by Celis and Koskinen [14, 15], but longer timescales were used here. The experiment involved independent application of two isotopes of pesticide (^{12}C and ^{14}C) to the three study soils. The study was performed in triplicate and over four sorption periods (7, 14, 28 and 56 d for chlorotoluron; 28, 56, 112 and 168 days for prometryn and hexaconazole). Soil was irradiated and all materials were sterilised prior to use

in order to inhibit microbial degradation, but no additional sterility controls were applied during subsequent experimental manipulations.

Preliminary experiments determined that the optimal soil:solution ratio (*i.e.* giving 30-70% sorption over 96 hours) varied between 1:4 (chlorotoluron) and 1:50 (hexaconazole). Test vessels were 50 mL Teflon® centrifuge tubes, irradiated to sterilise for 20 min prior to use in a CL-1000 ultraviolet cross-linker (UVP Ltd, CA, USA). Tubes had screw-topped lids and we assumed negligible losses of liquid via evaporation between experimental steps. Soils were weighed into vessels to give the appropriate soil:solution ratio on an oven-dry basis (Supporting Information Table S1) and then 19.0 mL of sterile 0.01M CaCl₂ was added (24.0 mL for hexaconazole). The experiment was performed in the dark in an incubator at 4°C (Sanyo Fitotron RS232 incubator) to minimise degradation, with continuous shaking at 150 rpm (HS 501 Digital IKA®-Werke reciprocal shaker). Soil suspensions were pre-equilibrated by shaking overnight. Sterile pesticide treatment solutions (¹²C and ¹⁴C) were prepared in 0.01M CaCl₂, ensuring equivalence in concentration. This was applied (1.0 mL) to the pre-equilibrated soils to give 20 mL total pesticide solution (25 mL for hexaconazole) with an initial concentration of 0.70, 0.78 and 0.77 µg mL⁻¹ for chlorotoluron, prometryn and hexaconazole, respectively. For each soil-pesticide combination, half of the samples contained only ¹²C-pesticide and half contained only ¹⁴C-pesticide.

After the respective sorption period, samples were centrifuged at 3500 rpm for 10 min (Hermle Z513K, LaborTechnik, Bench Top Centrifuge). The supernatant from each sample was removed by weight (Supporting Information Table S1) using a Pasteur pipette, taking care not to disturb the soil. A 200-µL aliquot was taken from each ¹²C supernatant to determine percentage sorption by HPLC analysis. The same sample volume was also removed from each ¹⁴C supernatant but instead mixed with 10 mL of EcoScint A for quantification by LSC. Supernatants were then exchanged between corresponding tubes, with

initially ^{12}C samples receiving the parallel ^{14}C supernatant and vice versa. This substitution did not disturb sorption equilibrium when expressed for the combination of the two isotopes as the same sorption equilibrium was reached in both tubes prior to the exchange, only with different carbon isotopes. Samples were then shaken for a further 14 days for isotope exchange to occur and sampled during this time on days 1, 3, 7 and 14 (centrifuged and a 200- μL aliquot of supernatant removed for LSC quantification) to measure exchange between ^{12}C - and ^{14}C -pesticide isotopes over time.

Forced isotope exchange. A forced isotope exchange procedure involving the addition of a high-concentration ^{12}C -pesticide solution in 0.01M CaCl_2 was carried out following the 14-day isotope exchange phase. Repeated influx of high-concentration ^{12}C -pesticide over time ensures that competition for sorption to soil between ^{12}C - and ^{14}C -pesticide is increasingly biased towards the former; supply of ^{12}C -pesticide to the soil solid phase surfaces to take part in sorption is essentially instantaneous with exchange with ^{14}C -pesticide the rate-limiting step. Thus, as ^{12}C -pesticide occupies all available sorption sites by out-competing any available ^{14}C -pesticide, it is possible to identify, through measurement of ^{14}C -pesticide in solution, the proportion of sorbed ^{14}C -pesticide available for desorption and hence the proportion of ^{14}C -pesticide not taking part in the sorption-desorption process.

Only initially ^{14}C samples (three soils, three replicates) adsorbed for 56 days (all pesticides) or for 112 and 168 days (prometryn and hexaconazole) were used in this part of the study as these had the greatest mass of ^{14}C -pesticide sorbed to soil after the 14-day exchange phase. There were 12-17 sampling points over periods of 145-204 days (Supporting Information Table S1). Samples were shaken between sampling points (150 rpm, 4°C). At each sampling point, samples were centrifuged and the supernatant was removed by weight (Table S1). A 250- μL aliquot was taken for quantification by LSC to measure release of ^{14}C from soil over

time. The supernatant that was removed was then replaced with fresh, high-concentration (Table S1) ^{12}C -pesticide solution to maintain the competition for sorption sites.

At the end of forced isotope exchange, soils were extracted with acidified methanol (0.1% H_3PO_4) to determine whether the ^{14}C -pesticide residue remaining sorbed to the soil was available by harsher extraction. Extractions involved addition of 20 mL of methanol acidified with 0.1% H_3PO_4 before shaking at 250 rpm for 24 hours at room temperature. The process was repeated until further removal of ^{14}C -pesticide was insignificant (<1% of initial-applied). Supernatants and soil extracts were analysed by radio-HPLC for parent compound and metabolites, and soils were finally combusted to complete the mass balance.

Theoretical development for isotope exchange and forced exchange. Theoretical development is provided for treatments with initially ^{14}C -pesticide as these were taken through the full procedure of isotope exchange followed by forced exchange. Equivalent relationships (equations 1 to 3) for isotope exchange of treatments with initially ^{12}C -pesticide can be obtained by reversing the isotope nomenclature. At the end of the equilibration phase prior to isotope exchange and for tubes with initially ^{14}C -pesticide, the partition coefficient (K_d , mL g^{-1}) can be defined as:

$$K_d = \frac{{}^{14}C_s}{{}^{14}C_e} = \frac{{}^{14}M_s}{{}^{14}M_e} \cdot \frac{V}{S} \quad (1)$$

where ${}^{14}C_s$ and ${}^{14}C_e$ are the concentration of ^{14}C -pesticide in soil and solution ($\mu\text{g g}^{-1}$ and $\mu\text{g mL}^{-1}$), respectively; ${}^{14}M_s$ and ${}^{14}M_e$ are the mass of ^{14}C -pesticide in soil and solution (μg), respectively; V is the volume of solution (mL); and S is the mass of soil (g). At this point, the supernatants are exchanged between initially ^{14}C and initially ^{12}C tubes. The total mass of ^{14}C -pesticide present in the initially ^{14}C tube after exchange is now ${}^{14}M_s$ and all of this will be sorbed at the point of exchange. We can define the quantity of ^{12}C -pesticide as ${}^{12}M_e$ which

will be numerically equivalent to $^{14}M_e$ and will be wholly in the solution phase at the point of exchange.

It is assumed that sorbed and equilibrium pesticide concentrations are identical for tubes with the same treatment and differing only in having initially ^{14}C - or initially ^{12}C -pesticide and that the sorption equilibrium is maintained after isotope exchange when expressed on the basis of the total pesticide (*i.e.* ^{14}C and ^{12}C) in the system. This assumption is supported by the research of Sander & Pignatello [17] who showed complete reversibility of sorption at low concentrations in their long-term isotope exchange experiments. In this circumstance and assuming that all sorbed ^{14}C -pesticide is available for exchange, the K_d will be maintained at the pre-exchange value and there will be a net transfer of ^{12}C -pesticide out of solution into the sorbed phase and a numerically equivalent net transfer of ^{14}C -pesticide in the opposite direction. Theoretically, these net fluxes will continue until the point where the two isotopes each attain the characterising equilibrium (nomenclature as previously, with equilibrium for each individual isotope designated with prime symbols (')):

$$K_d = \frac{{}^{12}M'_s}{{}^{12}M'_e} \cdot \frac{V}{S} = \frac{{}^{14}M'_s}{{}^{14}M'_e} \cdot \frac{V}{S} \quad (2)$$

The amount of ^{14}C -pesticide released into solution following isotope exchange is given by $^{14}M'_e$. As it is assumed that K_d is constant throughout the exchange process and no degradation occurs, and as $^{14}M'_s$ is given by $(^{14}M_s - ^{14}M'_e)$:

$$^{14}M'_e = (^{14}M_s - ^{14}M'_e) \cdot \left(\frac{{}^{14}M_e}{{}^{14}M_s}\right)$$

which can be rearranged to:

$$^{14}M'_e = \frac{{}^{14}M_s \cdot {}^{14}M_e}{(^{14}M_s + ^{14}M_e)} \quad (3)$$

For the condition where there is no irreversible sorption, we can thus calculate theoretical values for $^{14}M'_e$ as a function of the partitioning measured immediately prior to isotope exchange. This relationship is plotted as the dashed line in Figure 3.

The sorption equilibrium is completely changed during forced exchange due to the addition of concentrated ^{12}C -pesticide solution. At this point in the experiment, we are interested in how much of the ^{14}C -pesticide that is still sorbed after isotope exchange is participating in reversible sorption equilibrium. Assuming zero irreversible sorption, all of the ^{14}C -pesticide should be released back into solution over successive exchange steps due to the huge imbalance in presence of the two isotopes. The ^{14}C released during forced exchange is thus defined as:

$$^{14}M''_s = (^{14}M_s - ^{14}M'_e) \quad (4)$$

For the condition where there is no irreversible sorption, we can thus calculate theoretical values for $^{14}M''_e$ as a function of the partitioning measured immediately prior to isotope exchange. This relationship is plotted as the solid line in Figure 3.

Testing for degradation. Degradation was measured in parallel to the isotope exchange studies. The three soils were treated with ^{12}C -pesticide (chlorotoluron) or ^{14}C -pesticide (prometryn and hexaconazole); there were again three replicates, but this time with five sorption periods (7, 14, 28, 56 and 70 days for chlorotoluron; 28, 56, 112, 168 and 182 days for prometryn and hexaconazole) to include the isotope exchange period. Soil suspensions were prepared and shaken exactly as in the main study using the same soil:solution ratio and initial mass of pesticide. After each sorption period, samples were centrifuged, the supernatant was removed by weight (Supporting Information Table S1) and a 1-mL aliquot was taken for analysis by HPLC or radio-HPLC. The soils were then extracted with solvent by adding 20 mL methanol (acidified with 0.1% H_3PO_4) and shaking at 250 rpm for one hour

at room temperature. After centrifuging to separate the soil extract, a 1-mL aliquot was taken for analysis by HPLC or radio-HPLC to derive the mass balance and assess presence/absence of any degradation products. Supernatants and soil extracts taken from chlorotoluron experiments at 28, 56 and 70 days were analysed by LC-TOF-MS to confirm that mass balances were solely attributable to parent compound and not to any degradation products. This step was not necessary for the other two compounds due to use of radio-HPLC.

Chemical analysis. All ^{12}C -pesticide samples in solution were analysed by HPLC (Agilent 1100 Series, Agilent Technologies UK Ltd); full details are given in Table S2 in the Supporting Information. ^{14}C -pesticide samples in solution were analysed by LSC (LS 6500 Beckman Coulter Inc., Fullerton, USA). The LSC method counted each sample three times for a total of 15 min (5 min each). Duplicate blanks were used to account for the background radioactivity and results were corrected for quench and luminescence. Limits of quantification were 0.00011, 0.00012 and 0.00022 Bq mL⁻¹ for chlorotoluron, prometryn and hexaconazole, respectively. Radio-HPLC analysis was carried out using a Hewlett Packard 1100 Series HPLC with Perkin-Elmer Radiomatic 625TR Flow Scintillation Analyser (see Table S3 in the Supporting Information for details).

To prepare soil samples for combustion, soils were air-dried (7 days) and then ground using a pestle and mortar. Approximately 200 mg of dry, ground soil was weighed into a combustion cone sandwiched between two combustion caps. Soil samples were then oxidised in a Perkin-Elmer Oximate 80, Model 370 and finally analysed by LSC (Perkin-Elmer Tri-Carb 2810TR Liquid Scintillation Analyser). Limit of quantification was 2.5 Bq g⁻¹ sample.

LC-TOF-MS analysis of supernatants and soil extracts containing chlorotoluron was performed using an Agilent 1200 Series LC with G120 Time of Flight Mass Spectrometer (Santa Clara, CA, USA). LC was performed using a Waters Acquity BEH C₁₈ (2.1 x 50 mm, 1.7 µm) column (at 35.0°C) with 0.2 µm in-line filter. Mobile phases were 5 mM ammonium

acetate in water (Channel A) and methanol (Channel B). A gradient method and flow rate of 0.6 mL min⁻¹ was used. The initial ratio of ammonium acetate:methanol was 98:2, changing to 2:98 over 5 min, held for 3.1 min before then returning to original conditions after 8.1 min (total run time 9 min). Retention time of parent chlorotoluron was 3.01 min. Injection volume was 3 µL in acetonitrile. TOF-MS analysis was carried out in positive or negative mode electrospray with a nebulizer pressure of 45 psi, capillary of 4000 V, gas temperature of 450°C, drying gas flow at 15 L min⁻¹, skimmer of 60 V, fragmentor of 150 V and octopole RF voltage of 250 V. The mass range measured was 100-1100 m/z with resolving power of 5000. Total ion chromatographs were generated using Agilent Masshunter.

Results and discussion

Degradation and 24-h sorption isotherms. LC-TOF-MS analysis did not identify chlorotoluron metabolites in any supernatants or soil extracts analysed following 28, 56 or 70 days of sorption; ¹²C-chlorotoluron samples did not differ (except for chlorotoluron peak) from the blank supernatants and soil extracts. Hence, it was assumed that negligible degradation of chlorotoluron occurred during the 70-day experiment. Radio-HPLC analysis confirmed that there was no quantifiable degradation of hexaconazole over the 182 days investigated. Small amounts of degradation of prometryn occurred with the proportion of radioactivity attributable to parent decreasing from 97 ± 0.3%, 97 ± 0.1% and 98 ± 1.9% after 56 days to 92, 94 and 97% after 182 days in the Blackwood, Andover and Salop soils, respectively; there were no replicates at the latter date due to combination and concentration of supernatants. Table S4 in the Supporting Information characterises sorption isotherms for 24-hour batch-slurry experiments with all pesticide-soil combinations to act as a reference point for data from long-term sorption experiments. These batch values are in line with ranges reported in regulatory databases (e.g. www.sitem.herts.ac.uk/aeru/ppdb).

Isotope exchange. Figure 1 characterises isotope exchange in the study soils during the 14-day exchange phase for the three compounds subjected to 56 days of sorption. This sorption period is discussed in detail as it was common for all pesticides; corresponding results for other sorption periods are shown in Figures S1-S3 in the Supporting Information. For initially ^{12}C samples, Figure 1 shows a significant reduction of ^{14}C -pesticide in solution over the first day after exchange. Meanwhile, for the initially ^{14}C samples, a significant increase of ^{14}C -pesticide in solution is evident during the same period. The influx of ^{12}C - (initially ^{14}C sample) or ^{14}C - (initially ^{12}C sample) pesticide at 0 days (supernatant exchange) gave large differences between the ratios of the two isotopes in the soil and solution in both tubes. Thus, rapid sorption (initially ^{12}C sample) and release (initially ^{14}C sample) of ^{14}C -pesticide in one direction dominates as the system responds to accommodate for the new conditions.

The remainder of the isotope exchange phase (between 1 and 14 days) sees any further net exchange between ^{12}C - and ^{14}C -pesticide become much slower. For both initially ^{12}C and ^{14}C samples, the disparity in isotope concentration has lessened as the majority of isotope exchange has already occurred. For initially ^{12}C samples, there is an overall decrease of ^{14}C -pesticide in solution over time as this sorbs to the soil, displacing ^{12}C -pesticide. For initially ^{14}C samples, the anticipated increase in ^{14}C -pesticide in solution over time is short-lived. After the initial (0 to 1 day) increase due to release of sorbed material, there is a net decrease (chlorotoluron) or relatively constant amounts (prometryn, hexaconazole) of ^{14}C -pesticide in solution between 1 and 14 days. Any decrease in concentration of ^{14}C -pesticide in solution is seen most strongly in those instances when the shaking period prior to isotope exchange was shorter (*e.g.* for shaking periods of 7-56 days; Supporting Information Figures S1-S3). This phenomenon is likely the result of continued sorption, demonstrating that the system had not reached equilibrium at the point of supernatant exchange, even after 56 days (Figure 1). None of the pairs of isotope exchange curves shown are symmetrical around a horizontal line,

indicating that none of the systems reached true equilibrium during the experimental period, even where the sorption phase lasted 168 days for prometryn and hexaconazole (Supporting Information Figures S1-S3). There is no trend of different behaviour for the different soils. If sorption equilibrium had been reached and all pesticide was in exchange between the soil and solution, meaning sorption was a fully reversible process, then Figure 1 would show the measured radioactivity in solution (solid lines) reach the expected radioactivity in solution (dashed lines) during the 14-day isotope exchange phase. Expected radioactivity lines were calculated based on the proportion of initial ^{14}C -pesticide in solution after the respective sorption period, so the same proportion of ^{14}C -pesticide was expected to be in solution after the 14-day exchange phase if all sorbed pesticide was participating in exchange. This is clearly not the case for the initially ^{14}C samples, as continued sorption of ^{14}C -pesticide that was released into solution immediately after supernatant exchange means the measured radioactivity in solution actually deviates further from that expected over time (Figure 1). Central to the explanation of Figure 1 is that the sorption of pesticide to soil is time-dependent. Extent of sorption increased between 28 and 56 days for all pesticide and soil combinations except prometryn in the Blackwood and Andover soils (Supporting Information Table S5). There were no measurements of sorption of chlorotoluron beyond 56 days, but data for prometryn in the Salop soil and hexaconazole in all three soils show further increases in sorption between 56 and 112 days. All systems with prometryn and hexaconazole were apparently very close to true sorption equilibrium after 112 days as there was little or no increase in sorption through to 168 days (Supporting Information Table S5). Gao *et al.* [18] also found chlorotoluron sorption to soil to be time-dependent and suggested that the time-range to reach equilibrium was likely to be months or years. In contrast, Celis and Koskinen [14, 15] considered that triadimefon and imidacloprid-guanidine had reached sorption equilibrium during their three-day isotope exchange tests. However, the authors observed

asymmetry in the behaviour of initially ^{12}C and ^{14}C samples which was similar to that observed here and may suggest that true equilibrium was not attained [19]. The fact that sorption was not at equilibrium at the point of isotope exchange meant that results from this phase of experimentation could not be used to calculate the irreversibly sorbed fraction of pesticide as proposed by Celis and Koskinen [14]. This is because exchange patterns for the two isotopes should be symmetrical in Figure 1.

Overall, the patterns of asymmetry exhibited in Figures 1 and S1-S3 are relatively consistent between the different pesticides and across the three different soils. One exception is for chlorotoluron in the Salop soil for shorter equilibration periods (Figure S1, 7- and 14-d equilibration) where the effect of continued increase in strength of sorption after isotope exchange is particularly marked. This indicates that sorption of chlorotoluron was further from the equilibrium in Salop soil after 7-14 days than in the other two soils at the equivalent time; however, the reasons for this deviation are unclear.

Forced isotope exchange. Figure 2 shows the result of the forced isotope exchange over time for samples sorbed for 56 days with ^{14}C -chlorotoluron then subjected to isotope exchange over 14 days (data for prometryn and hexaconazole showed a similar form and are given in the SI as Figures S4 and S5). The bulk of total recoverable ^{14}C -chlorotoluron was extracted with the first addition of ^{12}C -chlorotoluron (average 25% Blackwood, 42% Andover and 21% Salop after 1 day; Figure 2). This was followed by a gradual decline in ^{14}C -chlorotoluron recovery over time, which finally reached <1% recovery of initial for the Blackwood and Andover soils and 2% for the Salop soil between 161 and 204 days.

Although exposure to high concentrations of pesticide in solution is far from normal field conditions, the approach is anticipated to leave the soil solid phase essentially unchanged and thus may be a useful indication of long-term availability of residues to be biodegraded or leached in soils. Undoubtedly, there is some effect on the soil solid phase from shaking with

aqueous solution over a protracted period (*e.g.* a possible increase in activity of the sorption sites) and this has not been quantified here.

The amount of ^{14}C -pesticide released into solution from the solid soil phase following both isotope exchange and forced exchange phases was 79-87%, 93-96% and 88-96% of that sorbed immediately prior to exchange for chlorotoluron, prometryn and hexaconazole, respectively. It should be noted that in each case the forced exchange was still releasing small amounts of ^{14}C -pesticide from the soil solid phase at the end of the process, albeit at very slow rates (Figure 2; Figures S4 and S5 in the Supporting Information).

Mass balance. The complete mass balance of the three ^{14}C -pesticides for an initial sorption period of 56 days is given in Table 2, whilst those for prometryn and hexaconazole sorbed for 112 and 168 days are given in the Supporting Information (Table S6). Repeated solvent extraction after the forced isotope exchange released small amounts of the initially applied ^{14}C -pesticide from the three soils (2-11%, 0.6-1.9% and 0.5-2.5% of initial radioactivity for chlorotoluron, prometryn and hexaconazole, respectively). Whilst this material had not been released by forced isotope exchange, it would not be included within a standard definition of bound residues [13]. Radio-HPLC analysis of forced exchange solutions identified that a small proportion of the initial ^{14}C -chlorotoluron applied to the soils had degraded to transformation products during this phase of the study. The extent of degradation was $4.0 \pm 1.6\%$ (Blackwood), $3.9 \pm 2.3\%$ (Andover) and $1.6 \pm 1.5\%$ (Salop) of initial ^{14}C -chlorotoluron mass per sample. The half-life of chlorotoluron under laboratory conditions has been reported by Gao *et al.* [18] to be 30 days, but soil sterilisation and experimental conditions of 4°C in the dark effectively inhibited degradation despite the long duration of the study (274 days). Combustion of the extracted soils released radioactivity equivalent to 1.1-2.3%, 0.3-1.4% and 0.1-0.5% of initial radioactivity for chlorotoluron, prometryn and hexaconazole, respectively. These small fractions could not be identified, but fit with a classical definition of bound

residues. It should be noted that these percentages would not reflect environmentally-relevant irreversible fractions, which are likely to be greater than those reported here; this is because these figures were obtained through alteration of the soil solid phase via solvent-extraction and because the microbial community was eliminated. Finally, 0.6-4.6%, 0.2-1.4% and 0.9-3.6% of initial radioactivity was unaccounted for chlorotoluron, prometryn and hexaconazole, respectively. This may result from mineralisation to $^{14}\text{CO}_2$, volatilisation or cumulative errors over the course of the experiment due to the multiple solute exchanges undertaken.

Comparison with ideal behaviour. Under conditions of ideal, fully reversible sorption of pesticides to soil, the fraction of ^{14}C -pesticide that is in the solid phase after the initial period of sorption will exchange completely with ^{12}C -pesticide and be released into solution by the combination of isotope exchange and forced exchange. Perfect isotope exchange will be maximal at 50% initial sorption where 25% of the initial ^{14}C -pesticide (*i.e.* 50% of that sorbed) should exchange with ^{12}C -pesticide and be released back into solution (Equation 3); release of ^{14}C -pesticide will be a smaller proportion of initial radioactivity for both weaker and stronger sorption (dashed line in Figure 3). In contrast, ^{14}C released by forced exchange will increase non-linearly with proportion of initial radioactivity sorbed (Equation 4; solid line in Figure 3). Figure 3 compares this ideal behaviour with measurements for the three pesticides in the three soils following 56-days sorption. The measurements closely mirror the expected form of relationships indicating that there is no major deviation from ideal behaviour. ^{14}C -pesticide released by isotope exchange is smaller than predicted in all cases and this can be attributed to continued increase in sorption during the exchange phase and/or some small component of sorbed pesticide that does not participate in exchange. Presence of dissolved organic carbon and/or colloidal material in solution is a potential confounding factor as in most sorption studies. Any such material would be directly exchanged during

isotope exchange, whereas presence of dissolved organic carbon and colloids would be reduced during forced exchange due to repeated replacement of the soil solution.

In conclusion, study of the long-term fate of pesticide residues in soil provides complex experimental challenges. This study demonstrates that earlier proposals for use of isotope exchange are constrained by the extremely long timescales needed to approach true sorption equilibrium. Chlorotoluron and prometryn have typical degradation half-lives in soil in the range of one to a few months which is of the same order as the times needed to approach sorption equilibrium [20]. Time-dependence in pesticide sorption changes the availability of residues for leaching over time and it is important that this effect is included within models used to characterise risks to the environment from pesticide use [21]. The steep concentration gradient established within the forced isotope exchange element of this study acted to accelerate release of sorbed pesticide from the soil solid phase. Nevertheless, pesticide was still being released from the soil solid phase after more than six months exposure to this process. The forced exchange procedure was intermediate in efficiency of releasing sorbed pesticide between field conditions, where desorption would occur more slowly [22], and a conventional solvent extraction. This is confirmed by the release of further sorbed chlorotoluron on extraction with methanol. The accepted definition of pesticide bound residues excludes any fraction that can be extracted by a solvent [13]. Thus, the radioactivity released by soil combustion at study termination provides the purest measure of truly irreversible sorption (1-2% of applied radioactivity). This implies that abiotic processes of bound residue formation are overwhelmingly reversible for the three compounds and the set of soils studied here.

433 Acknowledgements

434 The Engineering and Physical Sciences Research Council and Syngenta Ltd. are gratefully
435 acknowledged for funding this research.

436 Supporting Information Available

437 Supplementary data are provided relating to experimental methods (Table S1), analytical
438 methods (Tables S2 and S3), classical sorption isotherms measured after 24 h (Table S4),
439 pesticide sorption over time (Table S5), mass balances for initial sorption periods of 112 and
440 168 d (Table S6 supplementary to Table 2), isotope exchange for different sorption periods
441 (Figures S1-S3 supplementary to Figure 1), and forced exchange over time (Figures S4 and
442 S5 supplementary to Figure 2).

References

1. Altfelder, S.; Streck, T. Capability and limitations of first-order and diffusion approaches to describe long term sorption of chlorotoluron in soil. *Journal of Contaminant Hydrology*. **2006**, *86*, 279-298.
2. Chen, Y.-X.; Chen, H.-L.; Xu, Y.-T.; Shen, M.-W. Irreversible sorption of pentachlorophenol to sediments: experimental observations. *Environment International*. **2004**, *30* (1), 31-37.
3. Barriuso, E.; Benoit, P.; Dubus, I.G. Formation of non-extractable (bound) residues in soil: magnitude, controlling factors and reversibility. *Environmental Science and Technology*. **2008**, *42* (6), 1845-1854.
4. Dec, J.; Haider, K.; Schäffer, A.; Fernandes, E.; Bollag, J.M. Use of a silylation procedure and ¹³C-NMR spectroscopy to characterise bound and sequestered residues of cyprodinil in soil. *Environmental Science and Technology*. **1997**, *31*, 2991-2997.
5. Kan, A.T.; Chen, W.; Tomson, M.B. Desorption kinetics of neutral hydrophobic organic compounds from field-contaminated sediment. *Environmental Pollution*. **2000**, *108* (1), 81-89.
6. Bollag, J.-M.; Myers, C.J.; Minard, R.D. Biological and chemical interactions of pesticides with soil organic matter. *Science of the Total Environment*. **1992**, *123-124* (0), 205-217.
7. Senesi, N. Binding mechanisms of pesticides to soil humic substances. *Science of the Total Environment*. **1992**, *123-124* (0), 63-76.

8. Rice, P.J.; Anderson, T.A.; Coats, J.R. Degradation and persistence of metolachlor in soil: effects of concentration, soil moisture, soil depth, and sterilization. *Environmental Toxicology and Chemistry*. **2002**, *21* (12), 2640-2648.
9. Nowak, K.M.; Miltner, A.; Gehre, M.; Schäffer, A.; Kästner, M. Formation and fate of bound residues from microbial biomass during 2,4-D degradation in soil. *Environmental Science and Technology*. **2010**, *45* (3), 999-1006.
10. Sinclair, C.J.; Boxall, A.B.A. Assessing the ecotoxicity of pesticide transformation products. *Environmental Science & Technology*. **2003**, *37* (20), 4617-4625.
11. Northcott, G.L.; Jones, K.C. Experimental approaches and analytical techniques for determining organic compound bound residues in soil and sediment. *Environmental Pollution*. **2000**, *108* (1), 19-43.
12. Gevaio, B.; Jones, K.C.; Semple, K.T. Formation and release of non-extractable ¹⁴C-Dicamba residues in soil under sterile and non-sterile regimes. *Environmental Pollution*. **2005**, *133* (1), 17-24.
13. Führ, F.; Ophoff, H.; Burauel, P.; Wanner, U.; Haider, K., *Pesticide bound residues in soils*, in *Modification of definition of bound residues*, F. Fuhr, et al., Editors. 1998, Wiley-VCH: Weinheim. p. pp 175-176.
14. Celis, R.; Koskinen, W.C. An isotope exchange method for the characterisation of the irreversibility of pesticide sorption-desorption in soil. *Journal of Agriculture and Food Chemistry*. **1999a**, *47*, 782-790.
15. Celis, R.; Koskinen, W.C. Characterisation of pesticide desorption from soil by the isotopic exchange technique. *Soil Science Society of America Journal*. **1999b**, *63*, 1659-1666.
16. Sander, M.; Pignatello, J.J. An isotope exchange technique to assess mechanisms of sorption hysteresis applied to naphthalene in kerogenous organic matter. *Environmental Science and Technology*. **2005**, *39* (19), 7476-7484.
17. Sander, M.; Pignatello, J.J. Sorption irreversibility of 1,4-dichlorobenzene in two natural organic matter-rich geosorbents. *Environmental Toxicology and Chemistry*. **2009**, *28* (3), 447-457.
18. Gao, M.; Dai, S.; Ma, Y.; Zhang, P. Impact of atrazine and nitrogen fertilizers on the sorption of chlorotoluron in soil and model sorbents. *Journal of Environmental Sciences*. **2007**, *19* (3), 327-331.
19. Suddaby, L.A.; Beulke, S.; van Beinum, W.; Celis, R.; Koskinen, W.C.; Brown, C.D. Reanalysis of experiments to quantify irreversibility of pesticide sorption-desorption in soil. *Journal of Agricultural and Food Chemistry*. **2013**, *61* (9), 2033-2038.
20. Pignatello, J.J.; Xing, B. Mechanisms of slow sorption of organic chemicals to natural particles. *Environmental Science and Technology*. **1996**, *30* (1), 1-11.
21. Beulke, S.; van beinum, W.; Suddaby, L.S. Interpretation of aged sorption studies for pesticides and their use in European Union regulatory leaching assessments. *Integrated Environmental Assessment and Management*. **2015**, *11*(2), 276-286.
22. Renaud, F.G.; Leeds-Harrison, P.B.; Brown, C.D.; van Beinum, W. Determination of time-dependent partition coefficients for several pesticides using diffusion theory. *Chemosphere*. **2004**, *57*, 1525-1535.

Tables and Figures

443 **TABLE 1.** Characterisation of the three study soils.

Soil type	Texture	Sample location (OS grid ref.)	Date collected	pH (in CaCl ₂)	Content (% w/w)			
					Sand	Silt	Clay	Organic matter
Blackwood	Loamy sand	SE50917051	29/05/2009	5.6	83	8	9	5.6
Andover	Clay loam	SE99105892	28/05/2009	6.9	49	32	19	5.1
Salop	Clay loamy sand	SP26616611	28/05/2009	5.8	54	26	20	5.2

444 **TABLE 2.** Mass balances for tubes containing initially ^{14}C -pesticide and with a 56-day sorption period (all values are % of initially applied
445 radioactivity with one standard deviation in parentheses)

Pesticide	Soil	In solution after sorption	Released by isotope exchange	Released by forced exchange	Released by solvent extraction	Residue remaining	Degradation	Unaccounted for
Chlorotoluron	Blackwood	23.58 (1.05)	11.21 (1.04)	50.79 (1.68)	5.48 (0.66)	2.27 (0.36)	3.96 (1.60)	2.71 (0.93)
	Andover	39.23 (1.93)	14.19 (1.22)	38.98 (0.66)	2.03 (0.23)	1.07 (0.12)	3.90 (2.31)	0.60 (0.86)
	Salop	10.84 (0.71)	5.09 (0.37)	64.99 (0.72)	10.77 (0.73)	2.19 (0.22)	1.57 (1.49)	4.55 (0.09)
Prometryn	Blackwood	46.96 (0.95)	18.72 (0.35)	32.37 (0.99)	1.09 (0.06)	0.29 (0.10)	2.94 (0.32)	0.57 (0.39)
	Andover	67.92 (1.16)	16.19 (0.56)	14.60 (0.29)	0.56 (0.05)	0.33 (0.06)	2.76 (0.11)	0.40 (0.57)
	Salop	63.10 (1.11)	14.80 (0.28)	19.49 (0.34)	1.21 (0.11)	0.34 (0.01)	1.95 (1.93)	1.06 (0.44)
Hexaconazole	Blackwood	42.94 (1.70)	20.74 (0.39)	34.18 (1.70)	0.89 (0.04)	0.18 (0.03)	0 (-)	1.07 (0.72)
	Andover	68.64 (0.76)	18.44 (1.67)	11.47 (0.49)	0.45 (0.09)	0.14 (0.01)	0 (-)	0.87 (2.58)
	Salop	64.36 (0.60)	18.05 (0.31)	13.36 (0.48)	1.23 (0.11)	0.19 (0.02)	0 (-)	2.81 (0.58)




FIGURE 1. Pesticide isotope exchange over time following 56 days of sorption. The solid lines show the measured change in total ^{14}C -pesticide in solution over 14 days for the Blackwood, Andover and Salop soils for (A) chlorotoluron; (B) prometryn; and (C) hexaconazole. Diamonds represent the initially ^{14}C -pesticide samples, squares represent the initially ^{12}C -pesticide samples. Data are shown as the mean value with standard deviation of three replicates included. The dotted lines represent the radioactivity in solution that the measured radioactivities are expected to reach if there is perfect exchange, *i.e.* if sorption is a fully reversible process.




FIGURE 2. Cumulative percentage recovery of ^{14}C -chlorotoluron during forced isotope exchange. Recovery is expressed as a percentage of the ^{14}C -chlorotoluron that is sorbed at the start of the forced exchange procedure. Samples are initially ^{14}C with 56-days sorption and isotope exchange for 14-days (3 replicates per soil).




FIGURE 3. Relationship between the amount of ^{14}C -pesticide sorbed immediately prior to isotope exchange (i.e. after 56, 112 or 168 d) and the amount of ^{14}C -pesticide released during subsequent phases of isotope exchange (open symbols) and forced exchange (solid symbols). Data are averages of triplicate measurements in three soils and for the three equilibration times for chlorotoluron (diamonds), prometryn (squares) and hexaconazole (triangles); standard deviations are given in Tables 2 and S6. The lines show theoretical behaviour assuming ideal and fully reversible exchange for the isotope exchange (dashed line) and forced exchange (solid line), respectively.

464 **Title**

465 Long-term experiments to investigate irreversibility in sorption of pesticides to soil

466 **Authors**

467 Suddaby, L.A.^{1,2,3*}, Beulke, S.^{2,4}, van Beinum, W.^{2,5}, Oliver, R.G.^{3,6}, Kuet, S.^{3,7} and
468 Brown, C.D.^{1,8}

469 ¹*Environment Department, University of York*; ²*Food and Environment Research Agency*
470 *(FERA)*; ³*Syngenta Ltd*

471 *Corresponding author details: e-mail laura.suddaby@tsgeconsulting.com; phone: + 44 (0)
472 1423 799647; present address: TSGE Consulting Ltd, Concordia House, Grimbald Crag
473 Court, St. James Business Park, Knaresborough, North Yorkshire, HG5 8QB, UK.

474 **Supporting Information Table of Contents**

475	TABLE S1. Points of difference in experimental methodology for the three compounds.....	S3
476	TABLE S2. HPLC methods for analysis of ¹² C samples.....	S4
477	TABLE S3. Radio-HPLC methods for analysis of ¹⁴ C samples.....	S5
478	TABLE S4. Freundlich sorption coefficient (k _f) and Freundlich exponent (n _f) from	
479	preliminary experiments to determine 24-h sorption isotherms for all pesticide-soil	
480	combinations.....	S6
481	TABLE S5. Sorption of ¹⁴ C-pesticide to the three soils over time.....	S7
482	TABLE S6. Mass balances for tubes containing initially ¹⁴ C pesticide and with 112- and	
483	168-d sorption period.....	S8
484	FIGURE S1. Isotope exchange over time for chlorotoluron.....	S9
485	FIGURE S2. Isotope exchange over time for prometryn.....	S10
486	FIGURE S3. Isotope exchange over time for hexaconazole.....	S11
487	FIGURE S4. Cumulative recovery of ¹⁴ C-prometryn (% of initially applied) using the forced	
488	isotope exchange method.....	S12
489	FIGURE S5. Cumulative recovery of ¹⁴ C-hexaconazole (% of initially applied) using the	
490	forced isotope exchange method.....	S13

491 **Tables and Figures**492 **TABLE S1. Points of difference in experimental methodology for the three compounds**

Parameter	Chlorotoluron	Prometryn	Hexaconazole
^{12}C pesticide grade	Fluka® analytical; 99.7% pure	Riedel-de-Haën® analytical; 99.2% pure	Fluka® analytical; 99.7% pure
^{14}C label position	Phenyl-U- ^{14}C	Triazinyl-U- ^{14}C	Triazolyl-U- ^{14}C
^{14}C pesticide purity and activity	99.2% pure; 4.57 MBq mg $^{-1}$	98.1% pure; 4.16 MBq mg $^{-1}$	99.6% pure; 2.26 MBq mg $^{-1}$
Initial pesticide concentration (µg mL $^{-1}$)	0.70	0.78	0.77
Soil:solution ratio	1:4 (5 g soil, 20 mL solution)	1:10 (2 g soil, 20 mL solution)	1:50 (0.5 g soil, 25 mL solution)
Initial mass of pesticide per tube (µg)	13.90	15.55	19.53
Initial radioactivity per tube (kBq; ^{14}C - pesticide only)	62.87	64.60	42.34
Initial sorption periods (d)	7, 14, 28, 56	28, 56, 112, 168	28, 56, 112, 168
Volume of supernatant exchanged (mL)	17.00	18.00	24.00
Isotope exchange period (d)	1, 3, 7, 14	1, 3, 7, 14	1, 3, 7, 14
Forced exchange period	17 sampling points over 204 days	14 sampling points over 175 days	12 sampling points over 145 days
Concentration of ^{12}C pesticide solution for forced exchange (µg mL $^{-1}$)	40.01	29.85	14.91

493 TABLE S2. HPLC methods for analysis of ¹²C samples

HPLC method parameter	Chlorotoluron	Prometryn	Hexaconazole
Method type	Gradient (Initial proportion of methanol at 65%, increasing to 100% after 10 minutes)	Gradient (Initial proportion of methanol at 20%, increasing to 100% after 15 minutes)	Gradient (Initial proportion of methanol at 20%, increasing to 100% after 15 minutes)
Mobile phases	A: HPLC-grade water (0.1% H ₃ PO ₄)* B: HPLC-grade methanol	A: HPLC-grade water (0.1% HCOOH) ^a B: HPLC-grade methanol	A: HPLC-grade water (0.1% HCOOH) ^a B: HPLC-grade methanol
Flow rate (mL min ⁻¹)	0.8	1.0	1.0
Total run time (min)	22	23	23
Column	C ₁₈ Supelco Discovery (15 cm x 4.6 mm x 5 μm)	C ₁₈ Supelco Discovery (15 cm x 4.6 mm x 5 μm)	Agilent Technologies CN Zorbax (25 cm x 4.6 mm x 5 μm)
Column temperature (°C)	30.0	30.0	30.0
Injection volume (μL)	25.0	25.0	25.0
Retention time (min)	8.91	11.79	12.59
Wavelength (nm)	210	254	205
Limit of detection (μg mL ⁻¹)	0.01 (in 0.01M CaCl ₂) 0.03 (in MeOH)	0.003 (in 0.01M CaCl ₂) 0.006 (in MeOH)	0.18 (in 0.01M CaCl ₂) 0.21 (in MeOH)

494 * Acidified with ortho-phosphoric acid (H₃PO₄)

495 ^aAcidified with formic acid (HCOOH)

496 TABLE S3. Radio-HPLC methods for analysis of ¹⁴C samples

HPLC method parameter	Chlorotoluron	Prometryn	Hexaconazole
Method type	Gradient (initial proportion of acetonitrile at 20%, increasing to 100% after 15 minutes)	Gradient (initial proportion of acetonitrile at 20%, increasing to 100% after 15 minutes)	Gradient (initial proportion of acetonitrile at 20%, increasing to 100% after 15 minutes)
Mobile phases	A: HPLC-grade water (0.1% HCOOH)* B: HPLC-grade acetonitrile + Flowlogic 1:1 scintillation cocktail	A: HPLC-grade water (0.1% HCOOH)* B: HPLC-grade acetonitrile + Flowlogic 1:1 scintillation cocktail	A: HPLC-grade water (0.1% HCOOH)* B: HPLC-grade acetonitrile + Flowlogic 1:1 scintillation cocktail
Flow rate (mL min ⁻¹)	1.0	1.0	1.0
Total run time (min)	20	20	20
Column	Agilent Eclipse XDB-C ₁₈ (5 µm x 4.6 mm x 150 mm)	Agilent Eclipse XDB-C ₁₈ (5 µm x 4.6 mm x 150 mm)	Agilent Eclipse XDB-C ₁₈ (5 µm x 4.6 mm x 150 mm)
Column temperature (°C)	30.0	30.0	30.0
Injection volume (µL)	100.0	100.0	100.0
Retention time (min)	9.42	9.54	12.42
Wavelength (nm)	210	254	205
Limit of detection (µg mL ⁻¹)	0.22	0.24	0.44

497 * Acidified with formic acid (HCOOH)

498 TABLE S4. Freundlich sorption coefficient (k_f) and Freundlich exponent (n_f) from preliminary experiments to determine 24-h sorption isotherms
 499 for all pesticide-soil combinations
 500

Pesticide	Blackwood soil		Andover soil		Salop soil	
	K_f (ml g ⁻¹)	n_f	K_f (ml g ⁻¹)	n_f	K_f (ml g ⁻¹)	n_f
Chlorotoluron	5.07	0.81	2.81	0.82	4.50	0.76
Prometryn	9.13	0.88	3.81	0.84	3.47	0.85
Hexaconazole	43.5	0.79	21.8	0.82	21.7	0.72

501
 502 All experiments used ¹²C-pesticide and the soil:solution ratios given in Table S1. Initial concentrations in solution were 0.5, 2.5, 5, 10, 20, 40 and
 503 50 µg mL⁻¹ for chlorotoluron, 0.3, 0.6, 1.25, 2.5, 5, 10 and 15 µg mL⁻¹ for prometryn, and 0.2, 0.5, 1, 2, 4, 8 and 12 µg mL⁻¹ for hexaconazole. All
 504 data were fitted to the Freundlich isotherm: $C_s = k_f \cdot C_e^{n_f}$, where C_s is concentration in the solid phase (µg g⁻¹) and C_e is concentration in the
 505 liquid phase (µg mL⁻¹).
 506

507 TABLE S5. Sorption of ^{14}C -pesticide to the three soils over time (values expressed as % of initially applied pesticide \pm 1 s.d.; n = 3)

Pesticide	Sorption period (d)	Blackwood soil	Andover soil	Salop soil
Chlorotoluron	7	71.10 \pm 0.48	54.61 \pm 1.57	77.14 \pm 0.34
	14	70.28 \pm 0.85	57.38 \pm 1.38	81.39 \pm 0.86
	28	71.19 \pm 0.54	57.90 \pm 0.84	85.72 \pm 0.11
	56	76.42 \pm 1.05	60.77 \pm 1.93	89.16 \pm 0.71
Prometryn	28	52.33 \pm 1.63	31.90 \pm 1.33	35.30 \pm 1.70
	56	53.04 \pm 0.95	32.08 \pm 1.16	36.90 \pm 1.11
	112	51.71 \pm 1.13	32.61 \pm 0.68	40.00 \pm 1.34
	168	51.46 \pm 0.18	32.76 \pm 1.20	38.09 \pm 0.33
Hexaconazole	28	53.58 \pm 1.13	32.40 \pm 1.88	33.33 \pm 1.55
	56	57.06 \pm 1.70	31.36 \pm 0.76	35.64 \pm 0.60
	112	57.34 \pm 0.30	35.00 \pm 2.40	41.22 \pm 2.86
	168	58.26 \pm 1.11	35.24 \pm 0.50	41.28 \pm 6.03

508 TABLE S6. Mass balances for tubes containing initially ^{14}C pesticide and with 112- and 168-d sorption period (all values are % of initially applied
509 radioactivity as mean of three replications and with one standard deviation in parentheses)

Pesticide	Sorption period (d)	Soil	In solution after sorption	Released by isotope exchange	Released by forced exchange	Released by solvent extraction	Residue remaining	Degradation	Unaccounted for
Prometryn	112	Blackwood	48.29 (1.13)	18.35 (0.20)	30.93 (0.91)	1.14 (0.07)	0.54 (0.11)	0 (-)	0.75 (1.10)
		Andover	67.39 (0.68)	15.47 (0.30)	15.36 (0.48)	0.72 (0.07)	0.58 (0.07)	0 (-)	0.48 (0.61)
		Salop	60.00 (1.34)	15.12 (1.32)	21.12 (1.40)	1.89 (0.19)	0.48 (0.06)	0 (-)	1.39 (0.86)
	168	Blackwood	48.54 (0.18)	18.19 (0.27)	30.06 (0.56)	1.27 (0.03)	0.83 (0.27)	0 (-)	1.11 (0.22)
		Andover	67.24 (1.20)	15.13 (1.12)	15.19 (1.06)	0.89 (0.13)	1.35 (0.60)	0 (-)	0.20 (0.44)
		Salop	61.91 (0.33)	14.32 (0.42)	20.23 (0.26)	1.78 (0.14)	0.77 (0.04)	0 (-)	0.99 (0.44)
	112	Blackwood	42.66 (0.30)	20.43 (0.34)	32.16 (0.40)	1.28 (0.04)	0.29 (0.00)	0 (-)	3.18 (0.25)
		Andover	65.00 (2.40)	18.76 (1.38)	11.77 (0.70)	0.68 (0.23)	0.25 (0.03)	0 (-)	3.55 (0.49)
		Salop	58.78 (2.86)	18.59 (0.73)	16.33 (1.93)	2.47 (0.62)	0.36 (0.03)	0 (-)	3.45 (0.46)
		Blackwood	41.74 (1.11)	19.48 (0.63)	33.08 (0.82)	1.89 (0.26)	0.39 (0.11)	0 (-)	3.43 (0.35)
		Andover	64.76 (0.50)	17.54 (0.63)	13.21 (1.30)	0.70 (0.06)	0.28 (0.01)	0 (-)	3.52 (0.63)
		Salop	58.72 (6.03)	17.22 (1.36)	18.07 (4.34)	2.18 (0.67)	0.50 (0.06)	0 (-)	3.31 (0.35)




FIGURE S1. Isotope exchange over time for chlorotoluron. The solid lines show the change in total ^{14}C -chlorotoluron in solution over 14 days for the Blackwood, Andover and Salop soils after adsorption times of (A) 7 d; (B) 14 d; (C) 28 d; (D) 56 d. Diamonds represent the initially ^{14}C -chlorotoluron samples, squares represent the initially ^{12}C -chlorotoluron samples. The dotted lines represent the radioactivity in solution that the measured radioactivities are expected to reach if there is perfect exchange, i.e. if sorption is a fully reversible process.




FIGURE S2. Isotope exchange over time for prometryn. The solid lines show the change in total ^{14}C -prometryn in solution over 14 days for the Blackwood, Andover and Salop soils after adsorption times of (A) 28 d; (B) 56 d; (C) 112 d; (D) 168 d. Diamonds represent the initially ^{14}C -prometryn samples, squares represent the initially ^{12}C -prometryn samples. The dotted lines represent the radioactivity in solution that the measured radioactivities are expected to reach if there is perfect exchange, i.e. if sorption is a fully reversible process.




FIGURE S3. Isotope exchange over time for hexaconazole. The solid lines show the change in total ^{14}C -hexaconazole in solution over 14 days for the Blackwood, Andover and Salop soils after adsorption times of (A) 28 d; (B) 56 d; (C) 112 d; (D) 168 d. Diamonds represent the initially ^{14}C -hexaconazole samples, squares represent the initially ^{12}C -hexaconazole samples. The dotted lines represent the radioactivity in solution that the measured radioactivities are expected to reach if there is perfect exchange, i.e. if sorption is a fully reversible process.




FIGURE S4. Cumulative recovery of ^{14}C -prometryn (% of initial-sorbed) using the forced isotope exchange method. Samples are initially ^{14}C with sorption for (A) 56 d, (B) 112 d, and (C) 168 d followed by isotope exchange for 14-days (three replicates per soil).




FIGURE S5. Cumulative recovery of ^{14}C -hexaconazole (% of initial-sorbed) using the forced isotope exchange method. Samples are initially ^{14}C with sorption for (A) 56 d, (B) 112 d, and (C) 168 d followed by isotope exchange for 14-days (three replicates per soil).

Highlights

- Isotope exchange method is extended to study sorption over periods up to 9 months
- Three pesticides in three soils took ca. 4 months to approach sorption equilibrium
- Release of sorbed pesticide after exchange approximated ideal behaviour
- Sorption processes under abiotic conditions were overwhelmingly reversible