

This is a repository copy of *Probing ultrafast dynamics in a solid-density plasma created by an intense femtosecond laser*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/103708/>

Version: Published Version

---

**Article:**

Adak, Amitava, Blackman, Dave, Chatterjee, Gourab et al. (6 more authors) (2016)  
Probing ultrafast dynamics in a solid-density plasma created by an intense femtosecond laser. Journal of Physics: Conference Series. 012001. ISSN: 1742-6596

<https://doi.org/10.1088/1742-6596/688/1/012001>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

Probing ultrafast dynamics in a solid-density plasma created by an intense femtosecond laser

This content has been downloaded from IOPscience. Please scroll down to see the full text.

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 144.32.224.203

This content was downloaded on 12/08/2016 at 12:44

Please note that [terms and conditions apply](#).

# Probing ultrafast dynamics in a solid-density plasma created by an intense femtosecond laser

Amitava Adak<sup>1</sup>, Dave Blackman<sup>2</sup>, Gourab Chatterjee<sup>1</sup>, Prashant Kumar Singh<sup>1</sup>, Amit D. Lad<sup>1</sup>, P. Brijesh<sup>1</sup>, A. P. L. Robinson<sup>3</sup>, John Pasley<sup>2,3</sup> and G. Ravindra Kumar<sup>\*1</sup>

<sup>1</sup>Tata Institute of Fundamental Research, Dr. Homi Bhabha Road, Colaba, Mumbai-400005, India

<sup>2</sup>York Plasma Institute, University of York, Heslington, York, YO10 5DQ, United Kingdom

<sup>3</sup>Central Laser Facility, RutherfordAppleton Laboratory, Chilton, Didcot, OX10 0QX, United Kingdom

\*Email: grk@tifr.res.in


**Abstract.** We report a study on the dynamics of a near-solid density plasma using an ultraviolet (266 nm) femtosecond probe laser pulse, which can penetrate to densities of  $\sim 10^{22} \text{ cm}^{-3}$ , nearly an order of magnitude higher than the critical density of the 800 nm, femtosecond pump laser. Time-resolved probe-reflectivity from the plasma shows a rapid decay (picosecond-timescale) while the time-resolved reflected probe spectra show red shifts at early temporal delays and blue shifts at longer delays. This spectral behaviour of the reflected probe can be explained by a laser-driven shock moving inward and a subsequent hydrodynamic free expansion in the outward direction.

## 1. Introduction

Ultra-intense lasers can create extremely hot, high-density plasma while interacting with a solid. Extreme states of matter, achieved on a laboratory table-top by such interactions are very interesting to study, the measurement of the equation of state similar to that of an astrophysical object being a striking example[1]. Relativistic energy electron beams produced at the plasma critical surface in such interactions are crucial to the success of fast ignition[2] of the fusion pellet in inertial confinement fusion(ICF). A lot of experimental and theoretical studies have been performed in recent decades for the basic understanding of intense short-pulse-laser matter interaction[3] and for applications like particle acceleration[4, 5]. The hot dense plasma created via various laser absorption mechanisms[6] evolves very rapidly (on femtosecond and picosecond time-scales) and these dynamics can be monitored using the pump-probe technique[7, 8, 9].

In this paper, we investigate the temporal dynamics of a highly dense electron layer( $n_e = 10^{22} \text{ cm}^{-3}$ ) inside a plasma created on an aluminium-coated BK7 glass target by a laser at relativistic light intensities. Doppler spectrometry of the reflected probe enables the observation of the ultrafast motion of its critical surface (high density layer) inside the hot dense plasma. A numerical simulation is performed which reproduces the results of the experiment.





**Figure 1.** Experimental setup: T-target, BS-Beamsplitter, M-mirror, L-lens, SM-spectrometer, PD- Photo diode.

## 2. Experiment


The experiment (Fig. 1) was carried out with a chirped-pulse-amplification-based 20 TW laser system (800 nm, 40 fs, 10 Hz) at Tata Institute of Fundamental Research. An extra Pockels' cell was used to obtain a  $10^{-6}$  nanosecond intensity contrast. The pump pulse was focussed on a solid slab at  $45^\circ$  angle of incidence to a focal spot of  $17 \mu\text{m}$  (FWHM) to obtain intensities of  $\sim 10^{18} \text{ W/cm}^2$ . A small portion of the laser pulse (5 %) was extracted using a beam-splitter, up-converted to 266 nm, and focussed to a spot of  $60 \mu\text{m}$  at the interaction point at near-normal incidence using a fused-silica lens. The reflected probe pulse was then split into two parts and fed to a photodiode and a high-resolution ultraviolet spectrometer. A delay line was introduced in the path of the probe to change the relative temporal delay between the pump and the probe. The focussed probe intensity was  $\sim 10^{11} \text{ W/cm}^2$ . Spatial and temporal overlap was achieved by looking at the reflected probe intensity from a plasma created on a dielectric slab at relatively lower pump intensity ( $10^{17} \text{ W/cm}^2$ ). We define the temporal zero where the reflectivity shows a sudden spike. In this experiment we observed the probe reflectivity and spectrum from a solid-density plasma on aluminium-coated BK-7 target.

## 3. Results & Discussions

Figure 2(a) shows reflection of the probe as a function of probe delay with respect to the pump from a super critical layer ( $n_e=10^{22} \text{ cm}^{-3}$ ) in the plasma. Target: Al-coated 5 mm BK-7 glass target. The probe reflection shows an exponential decay ( $\tau = 5 \text{ ps}$ ) as the plasma evolves after excitation by pump. The reflectivity of the normally incident probe can be written as [10]

$$R \propto \exp\left(-\frac{8\nu_{ei}^* L}{3c}\right) \quad (1)$$

where  $\nu_{ei}^*$  is the effective electron ion collision frequency. L is the spatial scale length of plasma over which the probe gets absorbed. The collision frequency is a function of electron density



**Figure 2.** (a) Reflected probe intensity as a function of probe delay. The blue line is the exponential fit to the experimental data (blue asterisks). The decay time is measured to be 5 ps from the fit. (b) A similar plot for the velocity of the probe-critical-layer is shown. These velocities (black open circles) are calculated from the Doppler shifts of the ultraviolet probe measured from pump-probe Doppler spectrometry.

and temperature. In this context of fast time scale (few picosecond) probe reflectivity is mainly dependent on  $L$ , if we assume quasi-static values of density and temperature.


The velocity of the supercritical layer was measured by pump-probe Doppler spectrometry[8]. The velocity can be expressed as,

$$v_{exp} = -0.5c \frac{\Delta\lambda}{\lambda} \quad (2)$$

Where  $\Delta\lambda$  is the Doppler shift from the experiment and  $\lambda = 266$  nm in our case. Figure 2(b) shows the velocity of the probe-critical-layer, calculated from the Doppler shift at various probe delays. At initial few picoseconds, the probe critical-surface moves deeper into the plasma (negative velocity) riding on a non-relativistic shock. At subsequent times, the critical surface moves towards vacuum with the freely expanding plasma (positive velocity).

#### 4. Simulation

Figure 3 shows the velocity of the probe critical-layer results from 1-D hybrid simulations. First, the HYADES code was run on a  $500 \mu\text{m}$  silicate target. The output from this (ion, mass and electron density) was then interpolated onto a regular grid for use in an 1-D PIC code (ELPS) to find the hotspot formed by the pump laser. The code was run with 800 nm, 30 fs,  $2 \times 10^{18} \text{W}/\text{cm}^2$  laser pulse. A density spike was observed and its motion was simulated by a Lagrangian hydro code. These 1-D simulations calculate a slightly higher velocity of the critical-layer of the 266 nm probe beam than the experimental observation. The sign reversal of the velocity is around 6 ps, which is close to the experimental result.



**Figure 3.** The simulated velocity of the probe-critical-layer as a function of the probe delay. The sign convention: outward motion corresponds to positive velocity.

## 5. Conclusions

In this study, we observe ultrafast motion of a supercritical ( $10^{22} \text{ cm}^{-3}$ ) electron layer in a plasma created by a high-intensity, femtosecond laser pulse on a solid target. We see laser-driven density pile-up and propagation of a non-relativistic shock inside the solid. 1-D hybrid HYADES-PIC-HYDRO simulations support the experiment results.

## 6. Acknowledgments

AA and ADL thank the ‘Strong Field Science’ (11P-1401) programme of the DAE, Government of India for support for participation in IFSA2013. GRK acknowledges support from a J.C. Bose Fellowship grant from DST, Government of India.

## References

- [1] Paul McKenna *et al.*, *Laser-Plasma Interactions and Applications* (Springer Switzerland, Heidelberg, New York, Dordrecht, London, 2013).
- [2] M. Tabak *et al.*, *Phys. Plasmas* **1**, 1626 (1994).
- [3] P. Gibbon *et al.*, *Plasma Phys. Control. Fusion* **38**, 769-793 (1996).
- [4] T. Tajima *et al.*, *Phys. Rev. Lett.* **43**, 267 (1979).
- [5] V. Malka *et al.*, *Science* **298**, 1596 (2002).
- [6] S. C. Wilks *et al.*, *IEEE JOURNAL OF QUANTUM ELECTRONICS*, **33**, 1954 (1997).
- [7] A. S. Sandhu *et al.*, *Phys. Rev. Lett.* **89**, 225002 (2002).
- [8] S. Mondal *et al.*, *Phys. Rev. Lett.* **105**, 105002 (2010).
- [9] Y. Ping *et al.*, *Phys. Rev. Lett.* **109**, 145006 (2012).
- [10] W. L. Kruer, *The physics of laser plasma interactions* (Boulder, Colorado, 2003).