

This is a repository copy of *Hyperglycaemia and risk of adverse perinatal outcomes:A systematic review and meta-analysis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/104564/>

Version: Accepted Version

Article:

Farrar, Diane, Simmonds, Mark Crawford orcid.org/0000-0002-1999-8515, Bryant, Maria orcid.org/0000-0001-7690-4098 et al. (5 more authors) (2016) Hyperglycaemia and risk of adverse perinatal outcomes:A systematic review and meta-analysis. BMJ. i4694. pp. 1-11. ISSN: 1756-1833

<https://doi.org/10.1136/bmj.i4694>


Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.


Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.


Combining 75g and 100g OGTT tests – Fasting



50g OGCT



Two-hour postload 75g OGTT



Two-hour postload 100g OGTT

