

This is a repository copy of *Lifetime measurement of the 41+ state of Ni 58 with the recoil distance method*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/106479/>

Version: Accepted Version

---

**Article:**

Loelius, C., Iwasaki, H., Brown, B. A. et al. (24 more authors) (2016) Lifetime measurement of the 41+ state of Ni 58 with the recoil distance method. Physical Review C. 024340. ISSN 2469-9993

<https://doi.org/10.1103/PhysRevC.94.024340>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Lifetime measurement of the $4_1^+$ state of $^{58}\text{Ni}$ with the recoil distance method

C. Loelius,<sup>1,2</sup> H. Iwasaki,<sup>1,2</sup> B. A. Brown,<sup>1,2</sup> M. Honma,<sup>3</sup> T. Braunroth,<sup>4</sup> V. M. Bader,<sup>1,2</sup> T. Baugher,<sup>1,2</sup> D. Bazin,<sup>1</sup> J. S. Berryman,<sup>1</sup> C. M. Campbell,<sup>5</sup> A. Dewald,<sup>4</sup> A. Gade,<sup>1,2</sup> N. Kobayashi,<sup>1</sup> C. Langer,<sup>1,6</sup> I. Y. Lee,<sup>5</sup> A. Lemasson,<sup>1</sup> E. Lunderberg,<sup>1,2</sup> C. Morse,<sup>1,2</sup> F. Recchia,<sup>1</sup> D. Smalley,<sup>1</sup> S. R. Stroberg,<sup>1,2</sup> R. Wadsworth,<sup>7</sup> C. Walz,<sup>1,8</sup> D. Weisshaar,<sup>1</sup> A. Westerberg,<sup>9</sup> K. Whitmore,<sup>1,2</sup> and K. Wimmer<sup>1,9</sup>

<sup>1</sup>*National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, Michigan 48824, USA*

<sup>2</sup>*Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA*

<sup>3</sup>*Center for Mathematical Science, University of Aizu, Aizu-Wakamatsu, Fukushima 965-8580, Japan*

<sup>4</sup>*Institut für Kernphysik der Universität zu Köln, D-50937 Köln, Germany*

<sup>5</sup>*Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California, 94720, USA*

<sup>6</sup>*Joint Institute for Nuclear Astrophysics, Michigan State University, East Lansing, Michigan 48824, USA*

<sup>7</sup>*Department of Physics, University of York, Heslington, York YO10 5DD, United Kingdom*

<sup>8</sup>*Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany*

<sup>9</sup>*Department of Physics, Central Michigan University, Mount Pleasant, Michigan 48859, USA*

(Dated: August 5, 2016)

The quadrupole transition rate for the  $4_1^+ \rightarrow 2_1^+$  transition of  $^{58}\text{Ni}$  was determined from an application of the recoil distance method with the GRETINA array. The present result of the  $B(E2; 4_1^+ \rightarrow 2_1^+)$  was found to be  $50_{-6}^{+11} e^2\text{fm}^4$ , which is about three times smaller than the literature value, indicating substantially less collectivity than previously believed. Shell model calculations performed with the GXPF1A effective interaction agree with the present data and the validity of the standard effective charges in understanding collectivity in the Nickel isotopes is discussed.

## I. INTRODUCTION

The quadrupole transition rates of excited nuclear states serve as a useful indicator of collectivity or deformation in atomic nuclei and provide experimental inputs to validate theoretical predictions. Recent progress in Coulomb excitation and lifetime measurements with rare isotope beams has provided  $B(E2)$  data for well deformed nuclei with  $N=40-50$  [1][2][3][4] and in the vicinity of doubly-magic nuclei  $^{56}\text{Ni}$  and  $^{100,132}\text{Sn}$ , away from the valley of stability [5][6][7][8]. Concurrently, the development of new effective interactions in model spaces with larger dimensions has expanded the shell model's predictive power into heavier mass regions, including a new frontier around  $A \approx 60-80$  at the  $pf$  shell extending into the  $sdg$  shells [9][10][11].

The Ni isotopic chain spans over three doubly-magic nuclei, from the most neutron-deficient doubly-magic nucleus  $^{48}\text{Ni}$  [12] via the self-conjugate  $^{56}\text{Ni}$  [5] to the very neutron-rich  $^{78}\text{Ni}$  [13][14]. As such, the Ni isotopes have attracted much interest as a benchmark for shell model calculations, for example as in [15][16]. Nuclides  $^{56}\text{Ni}$  at  $N = 28$  and  $^{68}\text{Ni}$  at  $N = 40$ , a closed shell for the harmonic oscillator potential, demonstrate a significant increase of the  $2_1^+$  excitation energy, which is associated with magic nuclei. Likewise, the  $B(E2; 2_1^+ \rightarrow 0_1^+)$  values of nuclei between  $^{56}\text{Ni}$  and  $^{68}\text{Ni}$  exhibit the parabolic shape expected between two magic nuclei, with minima at  $^{56}\text{Ni}$  and  $^{68}\text{Ni}$  and a maximum at midshell. However,  $^{56}\text{Ni}$  and  $^{68}\text{Ni}$  are of particular interest, as shell model calculations result in ground state wavefunctions with substantial contributions from excitations across the  $Z = 28$  and  $N = 28, 40$  gaps, which could alter the behavior of collectivity of nuclei between these two iso-

topes [11][17][18].

In this paper, we report on a new lifetime measurement of the  $4_1^+$  state of  $^{58}\text{Ni}$ . Lying two neutrons above  $^{56}\text{Ni}$ , the structure of  $^{58}\text{Ni}$  is expected to be sensitive to core excitations over the  $N = Z = 28$  shell gap. The lifetime of the  $4_1^+$  state of  $^{58}\text{Ni}$  has been measured previously using the Doppler Shift Attenuation Method (DSAM), which resulted in a  $B(E2; 4_1^+ \rightarrow 2_1^+)$  of  $148_{-15}^{+18} e^2\text{fm}^4$ . [19]. In fact, this is the only lifetime data for this transition incorporated into the NNDC evaluation [20] and the result turns out to be approximately three times larger than the prediction of  $55 e^2\text{fm}^4$  from the GXPF1 shell model interaction, which has demonstrated good predictive power in this mass region [9].

The potential enhancement of collectivity in the  $4_1^+ \rightarrow 2_1^+$  transition could indicate that the importance of core excitations across the  $N = Z = 28$  gap is underestimated or could even signify mixing in an extended model space including the lower-lying  $sd$  or higher-lying  $sdg$  shells, impacting our understanding of the evolution of structure in the Ni isotopic chain. On the other hand, neither the aforementioned DSAM measurement nor recent Coulomb excitation measurements of  $^{58}\text{Ni}$  exhibit an enhanced  $B(E2; 2_1^+ \rightarrow 0_1^+)$  compared to the GXPF1 shell model [19][21]. In fact, the previous DSAM measurement of the  $4_1^+$  state was sensitive primarily to lifetimes on the order of 1 ps or less, which does not cover the theoretical  $4_1^+$  lifetime of 14 ps predicted by the GXPF1 shell model. In order to confirm or reject the enhanced collectivity for the  $4_1^+$  state of  $^{58}\text{Ni}$ , an independent measurement is needed which is capable of distinguishing between the literature value and the theoretical prediction.

In this work, the lifetime of the  $4_1^+$  state was determined using a modified version of the recoil distance method [22][23]. This method is based on the use of




FIG. 1. (color online). (a) Doppler-shift corrected and (b) laboratory frame  $\gamma$ -ray spectra for  $^{58}\text{Ni}$  at forward angles using a 1-mm separation between the target and degraders are shown. In (a) the arrows indicate the observed  $^{58}\text{Ni}$  transitions and show the fast, reduced, and slow components for the lifetime measurement. In (a) and (b) the  $\gamma$ -ray peaks from neutron-induced reactions are labeled.

three foils, a target and two degraders. This configuration is suited for covering a large range of lifetimes in a single setup. In this experiment, the excellent position resolution needed to resolve the three different Doppler-shifted peaks associated with the three foil configuration was provided by the Gamma-Ray Energy Tracking In-beam Nuclear Array (GRETINA) [24].

## II. EXPERIMENT

The experiment was performed at the National Superconducting Cyclotron Laboratory, using the same experimental setup of [25]. A primary beam of  $^{78}\text{Kr}$  was accelerated to 150 MeV per nucleon by the K500 and K1200 coupled cyclotrons and incident on a  $^9\text{Be}$  target to produce a secondary cocktail beam including  $^{74}\text{Kr}$ .




FIG. 2. A partial level scheme for  $^{58}\text{Ni}$  showing states observed in the  $\gamma$ -ray spectra measured in the experiment is presented. The arrow thicknesses for the  $2_1^+$ ,  $4_1^+$  and  $4_3^+$  transitions correspond to the observed intensity of the transition relative to the  $2_1^+ \rightarrow 0_1^+$  transition. The mean lifetime shown in parentheses for the  $4_1^+$  state is from the current measurement, while those for the  $2_1^+$  and  $4_3^+$  states are from the NNDC evaluation [20].

The A1900 fragment separator [26] was then used to separate the fragments by their rigidities. The secondary  $^{74}\text{Kr}$  beam at 93 MeV per nucleon had an intensity averaging at  $1 \times 10^5$  particles per second with a purity of approximately 40%. The secondary beam was then sent to the experimental area, where lifetime measurements were performed with the Triple PLunger for EXotic Beams (TRIPLEX) [23] coupled with GRETINA [24] and the S800 spectrograph [27].

The TRIPLEX [23] is a device which enables sensitive lifetime measurements based on the recoil distance method by the placement of up to three foils into the beam line. The first foil was a  $750 \mu\text{m}$   $^9\text{Be}$  target foil. The second foil was a  $125 \mu\text{m}$  Ta degrader foil. The third foil was a  $90 \mu\text{m}$  Ta degrader foil. The excited states of  $^{58}\text{Ni}$  were populated by multi nucleon removal reactions of  $^{74}\text{Kr}$  on the  $^9\text{Be}$  target. The second two foils were used as degraders to enable the recoil distance method. In this setup, there are three different Doppler-shifted  $\gamma$ -ray peaks, a “fast” peak for decays occurring before the first degrader, a “reduced” peak for decays occurring between the first and second degraders, and a “slow” peak corresponding to decays after both degraders. The TRIPLEX plunger was configured to provide a measurement of lifetimes in the range of 10 ps, and therefore a separation of 1 mm was used between both the target and first degrader and between the first degrader and second degrader.

To measure background contributions from reactions in the degraders, an additional measurement was performed by setting the separation between the target and the first degrader to 10 mm while keeping the distance between the degraders at 1 mm. Due to the long distance between the target and degraders, nearly all of the excited states produced in the target will decay prior to reaching the degraders. Therefore, the ratio of counts in the three Doppler-shifted regions provides a measurement of relative yields of reactions in the target to those in the degraders.

The reaction products and beam energy were identified using the S800 spectrometer. The S800 spectrom-




FIG. 3. The results of the Recoil Distance Method analysis for the  $^{58}\text{Ni}$   $4_1^+ \rightarrow 2_1^+$  transition using (a) 1-mm and (b) 10-mm target to degrader separations are shown. The fit includes the  $2_1^+ \rightarrow 0_1^+$  transition, which is sensitive to the  $4_1^+$  lifetime. The data are compared to the fit spectra, which are decomposed into the GEANT4 simulation, the  $\gamma$ -rays from neutron-induced reactions in  $^{27}\text{Al}$ , and the exponential background. In (a), fits for 5 ps (dotted), consistent with the literature value [19], and the present result of 16 ps (solid curve) are shown.

eter uses time of flight and energy loss measurements to separate out and identify the reaction residues of interest [27]. The energies of the reaction products after passing through the TRIPLEX were determined from the S800 magnetic rigidity to be 40 MeV per nucleon. Based on the final  $^{58}\text{Ni}$  energy, the  $^{58}\text{Ni}$  recoil velocities relative to the speed of light ( $v/c$ ) after each foil are estimated to be 0.36 (target), 0.32 (first degrader), and 0.28 (second degrader), respectively.

The de-excitation  $\gamma$ -rays were detected using GRETINA in coincidence with the identified  $^{58}\text{Ni}$  nuclei. The GRETINA configuration consisted of 7 detector modules made up of 4 HPGe crystals, each of which was divided into 36 segments. Signal decomposition provides subsegment position resolution [24]. The array was set up in the GRETINA frame with the TRIPLEX target positioned 13 cm upstream of the pivot point to increase the effective area of the array at small angles from the first degrader. The forward-most four detectors were placed at laboratory angles of  $20^\circ$  to  $50^\circ$  with respect to the first degrader, and were used to measure lifetimes. Additionally, three detectors were placed around  $70^\circ$ , which provided information to identify higher-lying states and coincidence relationships.

The Doppler-shift correction for each event was performed by using the angle from the degrader position to the largest energy deposit in GRETINA and the average velocity of the  $^{58}\text{Ni}$  particles between the degraders ( $v/c=0.32$ ). Furthermore, an addback of all  $\gamma$ -ray events which occurred within a distance of 7.5 cm from the first interaction point was employed. The 7.5 cm was chosen to balance the gain in peak-to-background from addback against a potential loss from accidental coincidences, which were not negligible due to the large background of this measurement.

### III. DATA ANALYSIS AND RESULTS

The Doppler-shift corrected spectrum for  $^{58}\text{Ni}$  with the 1-mm separation is shown in Figure 1 (a). The three peak structure characteristic of decays with lifetimes on the order of 10 ps is evident for the  $4_1^+$  decay at 1005 keV populating the  $2_1^+$  state. A similar feature is seen for the  $2_1^+$  decay to the  $0_1^+$  state at 1454 keV. In addition, a hint of a three peak structure is seen at around 1160 keV, which is consistent with the  $4_3^+$  decay to the  $4_1^+$  state, as shown in the level scheme of  $^{58}\text{Ni}$  in Figure 2. A peak at around 870 keV is consistent with the fast component of the  $7_2^+ \rightarrow 6_2^+$  transition of  $^{58}\text{Ni}$ , but it was not identified in the  $\gamma$ - $\gamma$  coincidence as part of a decay scheme. As the Doppler-reconstruction is optimized for decays occurring just after the first degrader, the reduced (middle) peak corresponds to the transition energy.

In Figure 1 (a), several peaks in the low energy region between 300 keV and 800 keV are also observed. These are associated with background  $\gamma$ -rays from neutron-induced reactions in the surrounding materials, such as the Al beam pipe and the Ge detectors, which are clearly identified in the laboratory frame spectrum as shown in Figure 1 (b). Contrary to the usual expectation, these laboratory frame  $\gamma$ -ray transitions manifest a peak structure after Doppler-correction, as only forward-angle data from GRETINA are used for the present analysis. Most of the background peaks appear at lower energies, but the peaks at 1368 keV and 1808 keV from the reactions of neutrons onto  $^{27}\text{Al}$  overlap with the  $^{58}\text{Ni}$  transitions after the Doppler-shift correction and require a careful analysis as discussed below.

In order to analyze the lifetime of the  $4_1^+$  state in  $^{58}\text{Ni}$ , a software package based on GEANT4 [28] incorporating the present experimental setup was employed, which created simulated spectra to be compared with the experimental data [29]. The simulated spectra were fit to the data using a  $\chi^2$  minimization, with the lifetime of the  $4_1^+$  state, the amplitude of the simulated spectra, and the




FIG. 4. (color online). The  $\gamma$ -ray spectrum for the  $4_3^+ \rightarrow 4_1^+$  transition of  $^{58}\text{Ni}$  is shown. The data are compared to simulated spectra with  $\tau(4_3^+)$  of 0 ps (dotted), 3 ps (solid), and 8 ps (dashed curve).

exponential background as variable parameters. Other parameters were estimated and fixed in the fit, including the feeding contributions from higher-lying states and the degrader reaction contribution. The systematic errors associated with these parameters were separately evaluated and are included in the final result. The simulation performs the addback of all points within 7.5 cm of the highest energy deposit, as is done in the data analysis. The lab-frame background contribution is incorporated by simulating the neutron-induced background peaks at 1368 keV and 1808 keV with the yields observed in the experiment and converting them into the projectile frame as shown by the filled spectra in Figure 3 (a).

To constrain the relative population strength and the reaction ratio, the data from the 10-mm separation was analyzed and the best fit is shown in Figure 3 (b). The proportion of  $^{58}\text{Ni}$  states directly populated in the reaction was found to be  $25\% \pm 5\%$  for the  $2_1^+$  state,  $50\% \pm 5\%$  for the  $4_1^+$  state, and  $25\% \pm 5\%$  for the  $4_3^+$  state, where the error is statistical only. The reaction ratio of the target to the degraders was determined to be  $13_{-2}^{+4}$ , where the error is statistical only. Note that the apparent fast to slow yield ratio in the spectrum is smaller than that determined above, as the S800 rigidity is tuned for the lower-momentum  $^{58}\text{Ni}$  products, which favors reactions in the degraders.

The lifetime of the  $4_1^+$  state was determined from a fit to the 1-mm separation data as shown in Figure 3 (a). The adopted value of the lifetime of the  $2_1^+$  state is 0.94 ps [20]. Due to the short  $2_1^+$  state lifetime and the significant feeding from the  $4_1^+$  state, the spectral shape for the  $2_1^+$  decay should reflect the lifetime of the  $4_1^+$  state, which is expected to be in the 10-ps range. Therefore, the region for the fit was chosen to include both the  $2_1^+$  and  $4_1^+$  peaks. The best fit result is shown in Figure 3 (a), where the  $4_1^+$  lifetime of  $16 \pm 1$  ps was obtained.

The main systematic errors in the present measurement are from ambiguities in the target to degrader reaction ratio and in feeding contributions from higher-lying states. The feeding effects were studied by assuming that

all of the higher-lying feeding states decay first to the  $4_3^+$  state. The lifetime of the  $4_3^+$  state has previously been deduced to be 0.16 ps [20]. However, the three peak structure observed for the 1161-keV transition (Figure 1 (a)) indicates a longer effective lifetime for this decay. The effective lifetime was determined by analyzing the spectrum between 1080 keV and 1300 keV with the amplitude as a free parameter and with a fixed exponential background. The best fit  $4_3^+$  effective lifetime was found to be  $3_{-3}^{+5}$  ps. The 1-mm spectrum focused around 1161 keV is shown in Figure 4 together with simulations incorporating the various lifetimes.

The systematic error for the  $4_1^+$  lifetime due to the uncertainties in the degrader reactions and feeding effects was found by analyzing the dependence of the lifetime to the modification of the corresponding parameters in the fit. The uncertainty in the  $4_1^+$  lifetime from the degrader reaction ratio and the feeding lifetime of the  $4_3^+$  state were found to be  $\pm 1$  ps and  $_{-2.5}^{+1}$  ps respectively. The additional systematic error from uncertainties in the direct population of states was found to be negligible. By adding the statistical and systematic errors in quadrature, the present result of the  $4_1^+$  mean lifetime is determined to be  $16_{-3}^{+2}$  ps, which corresponds to a  $B(E2; 4_1^+ \rightarrow 2_1^+)$  of  $50_{-6}^{+11} e^2\text{fm}^4$ .

#### IV. DISCUSSION

The experimental and theoretical results for the energy levels,  $B(E2; 2_1^+ \rightarrow 0_1^+)$  and  $B(E2; 4_1^+ \rightarrow 2_1^+)$  values of Nickel isotopes in the  $pf$  shell are presented in Figure 5. The experimental data for  $^{58}\text{Ni}$  are from the current measurement and the previous measurement by Kenn *et al* [19], while the remaining data are the adopted values from the NNDC [20]. The theoretical values come from the GXPF1A [30] and KB3G [31] Hamiltonians for protons and neutrons in the full  $pf$  model space, with a portion of the shell-model calculations carried out using the code MSHELL64 [32]. The GXPF1A interaction is a modification of the GXPF1 interaction, where discrepancies between experimental data and shell model calculations for unstable neutron-rich isotopes of Ca, Ti, and Cr were addressed by changing the values of five two-body matrix elements [30]. Two sets of proton ( $e_p$ ) and neutron ( $e_n$ ) effective charges are used in calculating the  $B(E2)$  values with the GXPF1A interaction: the “standard” values of  $e_p = 1.5$  and  $e_n = 0.5$  as used in [30] and the values  $e_p = 1.12$  and  $e_n = 0.67$  determined from the  $B(E2)$  values of the mirror  $27/2^-$  to  $23/2^-$  transitions in  $^{51}\text{Fe}$  and  $^{51}\text{Mn}$  in [21].

The  $A_p$  and  $A_n$  values presented in Table I are the bare proton and neutron  $E2$  matrix elements in the  $pf$  model space. The radial integrals were evaluated with harmonic oscillator wave functions with  $\hbar\omega = 45A^{-1/3} - 25A^{-2/3}$ . The  $B(E2)$  values are obtained from the bare  $E2$  matrix elements by incorporating the effective charges as:  $B(E2; I_i \rightarrow I_f) = (A_p e_p + A_n e_n)^2 / (2J_i + 1)$ . The



$B(E2)$  values are calculated using both sets of effective charges previously discussed with the bare  $E2$  matrix elements determined from shell model calculations. Although  $N = 28$  and  $N = 40$  are not identified as good shell closures in the Ni isotopes, shell effects remain in the systematics of the Ni isotopes, with the energy of the  $2_1^+$  state attaining a local maximum at  $^{56}\text{Ni}$  and  $^{68}\text{Ni}$ . Likewise, a similar decrease in the  $B(E2; 2_1^+ \rightarrow 0_1^+)$  can be observed at the same nuclei, with the  $B(E2)$  attaining a maximum near midshell. While the previous  $B(E2; 4_1^+ \rightarrow 2_1^+)$  data suggested a potential increase in collectivity in  $^{58}\text{Ni}$  relative to the  $B(E2; 4_1^+ \rightarrow 2_1^+)$  of  $^{60}\text{Ni}$ , the current data restores a decrease in collectivity away from midshell consistent with the other systematics and shell model calculations.

As can be seen in Figure 5 and Table I, while the GXPF1A and KB3G interactions both agree with the  $B(E2; 4_1^+ \rightarrow 2_1^+)$  of  $^{58}\text{Ni}$ , there is a significant difference between the results for the GXPF1A and the KB3G interactions for  $B(E2; 2_1^+ \rightarrow 0_1^+)$  values throughout the rest of the isotopic chain, with the GXPF1A being in much better agreement with experimental data [30][31]. The main reason for this is that the effective  $N = Z = 28$  shell gap is larger in the KB3G interaction, resulting in a smaller amount of proton and neutron excitations across the  $N$  or  $Z = 28$  gap and thus reduced collectivity. On the other hand, the  $B(E2)$  values predicted by the GXPF1A interaction using the two different sets of effective charges are similar to each other, and in both cases reproduce the data well.

The present result for the  $B(E2; 4_1^+ \rightarrow 2_1^+)$  of  $50_{-6}^{+11} e^2\text{fm}^4$  can be seen in Figure 5 (bottom) to agree more closely with the GXPF1A shell model calculation than previous DSAM results. In addition, the present result restores the clear behavior of the  $B(E2; 4_1^+ \rightarrow 2_1^+)$  systematics towards a single large peak at  $^{62}\text{Ni}$ , as predicted by the GXPF1A shell model. This strong peak in the  $B(E2; 4_1^+ \rightarrow 2_1^+)$  values at  $^{62}\text{Ni}$  arises from the increased occupation of the neutron  $f_{5/2}$  orbital relative to  $^{58}\text{Ni}$ . The neutron  $f_{5/2}$  contribution to the  $E2$  matrix element adds coherently with the contributions from the neutron  $p_{3/2}$  and neutron  $p_{1/2}$  orbitals, and so the overall  $E2$  matrix element becomes much larger at  $^{62}\text{Ni}$ .

In conclusion, a new measurement of the lifetime of the  $4_1^+$  state in  $^{58}\text{Ni}$  has been performed using the Recoil Distance Method, which resulted in a value of  $16_{-3}^{+2}$  ps. The resulting  $B(E2; 4_1^+ \rightarrow 2_1^+)$  for  $^{58}\text{Ni}$  of  $50_{-6}^{+11} e^2\text{fm}^4$  obtained in this experiment resolves a discrepancy with the GXPF1A theory that arose from the previous DSAM results. The new value is more consistent with the GXPF1A shell model interaction prediction of  $66 e^2\text{fm}^4$  [30] than with the literature value of  $148_{-15}^{+18} e^2\text{fm}^4$  and so suggests much less collectivity in  $^{58}\text{Ni}$  than previously indicated.




FIG. 5. (color online). The systematics of the Nickel isotopes are presented, including the  $4_1^+$  and  $2_1^+$  energies (top),  $B(E2; 2_1^+ \rightarrow 0_1^+)$  (middle), and  $B(E2; 4_1^+ \rightarrow 2_1^+)$  (bottom). The experimental values are compared with shell model calculations using the GXPF1A and KB3G shell model interactions. For the GXPF1A shell model,  $B(E2)$  values are presented with the two sets of effective charges discussed in the text.

## ACKNOWLEDGEMENTS

The authors would like to thank C. Bancroft, D. Barofsky, and J. Lloyd from Central Michigan University for help during the experiment. This work is supported by the National Science Foundation(NSF) under Phy-1102511 and Phy-1404442, by the Department of Energy(DOE) National Nuclear Security Administration (NNSA) under award No. DE-NA0000979, by the UK STFC under ST/F000124 and ST/L005727, by the Bundesministerium für Bildung und Forschung (BMBF, Germany) under Contract No.

Nucleus	$J_i \rightarrow J_f$	GXPF1A				KB3G			
		$A_p [e\text{fm}^2]$	$A_n [e\text{fm}^2]$	$B(E2)^a [e^2\text{fm}^4]$	$B(E2)^b [e^2\text{fm}^4]$	$A_p$	$A_n$	$B(E2)^a [e^2\text{fm}^4]$	$B(E2)^b [e^2\text{fm}^4]$
$^{56}\text{Ni}$	$2_1^+ \rightarrow 0_1^+$	14.19	14.19	161	129	13.33	13.33	142	114
$^{58}\text{Ni}$	$2_1^+ \rightarrow 0_1^+$	10.11	17.97	117	109	6.13	17.49	64	69
$^{60}\text{Ni}$	$2_1^+ \rightarrow 0_1^+$	12.88	22.17	185	171	7.45	22.72	102	111
$^{62}\text{Ni}$	$2_1^+ \rightarrow 0_1^+$	15.00	23.38	234	211	7.71	22.58	104	113
$^{64}\text{Ni}$	$2_1^+ \rightarrow 0_1^+$	10.50	21.08	138	134	5.16	20.29	64	75
$^{56}\text{Ni}$	$4_1^+ \rightarrow 2_1^+$	18.58	18.58	153	123	11.59	11.59	60	48
$^{58}\text{Ni}$	$4_1^+ \rightarrow 2_1^+$	9.88	19.01	66	63	6.11	17.81	36	39
$^{60}\text{Ni}$	$4_1^+ \rightarrow 2_1^+$	15.40	24.74	140	127	5.63	14.13	27	28
$^{62}\text{Ni}$	$4_1^+ \rightarrow 2_1^+$	22.54	33.38	283	252	7.86	19.98	53	55
$^{64}\text{Ni}$	$4_1^+ \rightarrow 2_1^+$	7.44	9.87	29	25	1.28	6.04	2.7	3.3

<sup>a</sup>  $e_p = 1.5$ ,  $e_n = 0.5$

<sup>b</sup>  $e_p = 1.12$ ,  $e_n = 0.67$

TABLE I. The bare  $E2$  matrix element  $A_p$  and  $A_n$  values calculated using the GXPF1A and KB3G shell model Hamiltonians are presented for the  $2_1^+ \rightarrow 0_1^+$  and  $4_1^+ \rightarrow 2_1^+$  transitions in the Ni isotopes. In addition, the  $B(E2)$  values for those transitions are calculated using the standard effective charges  $e_p = 1.5$ ,  $e_n = 0.5$  and modified effective charges  $e_p = 1.12$ ,  $e_n = 0.67$ .

05P15PKFNA. GREITINA was funded by the US DOE Office of Science. Operation of the array at the NSCL

is supported by the NSF under Cooperative Agreement PHY-1102511(NSCL) and DOE under Grant No. DE-AC02-05CH11231.

- [1] W. Rother, A. Dewald, H. Iwasaki, S. M. Lenzi, K. Starosta, D. Bazin, T. Baugher, B. A. Brown, H. L. Crawford, C. Fransen, A. Gade, T. N. Ginter, T. Glasmacher, G. F. Grinyer, M. Hackstein, G. Ilie, J. Jolie, S. McDaniel, D. Miller, P. Petkov, T. Pissulla, A. Ratkiewicz, C. A. Ur, P. Voss, K. A. Walsh, D. Weisshaar, and K.-O. Zell, *Phys. Rev. Lett.* **106**, 022502 (2011).
- [2] H. L. Crawford, R. M. Clark, P. Fallon, A. O. Macchiavelli, T. Baugher, D. Bazin, C. W. Beausang, J. S. Berryman, D. L. Bleuel, C. M. Campbell, M. Cromaz, G. de Angelis, A. Gade, R. O. Hughes, I. Y. Lee, S. M. Lenzi, F. Nowacki, S. Paschalis, M. Petri, A. Poves, A. Ratkiewicz, T. J. Ross, E. Sahin, D. Weisshaar, K. Wimmer, and R. Winkler, *Phys. Rev. Lett.* **110**, 242701 (2013).
- [3] M. Niikura, B. Mouginot, S. Franchoo, I. Matea, I. Stefan, D. Verney, F. Azaiez, M. Assie, P. Bednarczyk, C. Borcea, A. Burger, G. Burgunder, A. Buta, L. Cáceres, E. Clément, L. Coquard, G. de Angelis, G. de France, F. de Oliveira Santos, A. Dewald, A. Dijon, Z. Dombradi, E. Fiori, C. Fransen, G. Friessner, L. Gaudefroy, G. Georgiev, S. Grévy, M. Hackstein, M. N. Harakeh, F. Ibrahim, O. Kamalou, M. Kmiecik, R. Lozeva, A. Maj, C. Mihai, O. Möller, S. Myalski, F. Negoita, D. Pantelica, L. Perrot, T. Pissulla, F. Rotaru, W. Rother, J. A. Scarpaci, C. Stodel, J. C. Thomas, and P. Ujic, *Phys. Rev. C* **85**, 054321 (2012).
- [4] T. Braunroth, A. Dewald, H. Iwasaki, S. M. Lenzi, M. Albers, V. M. Bader, T. Baugher, T. Baumann, D. Bazin, J. S. Berryman, C. Fransen, A. Gade, T. Ginter, A. Gottardo, M. Hackstein, J. Jolie, A. Lemasson, J. Litzinger, S. Lunardi, T. Marchi, V. Modamio, C. Morse, D. R. Napoli, A. Nichols, F. Recchia, S. R. Stroberg, R. Wadsworth, D. Weisshaar, K. Whitmore, and K. Wimmer, *Phys. Rev. C* **92**, 034306 (2015).
- [5] K. L. Yurkewicz, D. Bazin, B. A. Brown, C. M. Campbell, J. A. Church, D. C. Dinca, A. Gade, T. Glasmacher, M. Honma, T. Mizusaki, W. F. Mueller, H. Oliver, T. Otsuka, L. A. Riley, and J. R. Terry, *Phys. Rev. C* **70**, 054319 (2004).
- [6] J. M. Allmond, A. E. Stuchbery, J. R. Beene, A. Galindo-Uribarri, J. F. Liang, E. Padilla-Rodal, D. C. Radford, R. L. Varner, A. Ayres, J. C. Batchelder, A. Bey, C. R. Bingham, M. E. Howard, K. L. Jones, B. Manning, P. E. Mueller, C. D. Nesaraja, S. D. Pain, W. A. Peters, A. Ratkiewicz, K. T. Schmitt, D. Shapira, M. S. Smith, N. J. Stone, D. W. Stracener, and C.-H. Yu, *Phys. Rev. Lett.* **112**, 172701 (2014).
- [7] P. Doornenbal, S. Takeuchi, N. Aoi, M. Matsushita, A. Obertelli, D. Steppenbeck, H. Wang, L. Audirac, H. Baba, P. Bednarczyk, S. Boissinot, M. Ciemala, A. Corsi, T. Furumoto, T. Isobe, A. Jungclaus, V. Lapoux, J. Lee, K. Matsui, T. Motobayashi, D. Nishimura, S. Ota, E. C. Pollacco, H. Sakurai, C. Santamaria, Y. Shiga, D. Sohler, and R. Taniuchi, *Phys. Rev. C* **90**, 061302 (2014).
- [8] V. M. Bader, A. Gade, D. Weisshaar, B. A. Brown, T. Baugher, D. Bazin, J. S. Berryman, A. Ekström, M. Hjorth-Jensen, S. R. Stroberg, W. B. Walters, K. Wimmer, and R. Winkler, *Phys. Rev. C* **88**, 051301 (2013).
- [9] M. Honma, T. Otsuka, B. A. Brown, and T. Mizusaki, *Phys. Rev. C* **69**, 034335 (2004).

- [10] S. M. Lenzi, F. Nowacki, A. Poves, and K. Sieja, *Phys. Rev. C* **82**, 054301 (2010).
- [11] Y. Tsunoda, T. Otsuka, N. Shimizu, M. Honma, and Y. Utsuno, *Phys. Rev. C* **89**, 031301 (2014).
- [12] B. Blank, M. Chartier, S. Czajkowski, J. Giovinazzo, M. S. Pravikoff, J.-C. Thomas, G. de France, F. de Oliveira Santos, M. Lewitowicz, C. Borcea, R. Grzywacz, Z. Janas, and M. Pfützner, *Phys. Rev. Lett.* **84**, 1116 (2000).
- [13] Z. Y. Xu, S. Nishimura, G. Lorusso, F. Browne, P. Doornenbal, G. Gey, H.-S. Jung, Z. Li, M. Niikura, P.-A. Söderström, T. Sumikama, J. Taprogge, Z. Vajta, H. Watanabe, J. Wu, A. Yagi, K. Yoshinaga, H. Baba, S. Franchoo, T. Isobe, P. R. John, I. Kojouharov, S. Kubono, N. Kurz, I. Matea, K. Matsui, D. Mengoni, P. Morfouace, D. R. Napoli, F. Naqvi, H. Nishibata, A. Odahara, E. Sahin, H. Sakurai, H. Schaffner, I. G. Stefan, D. Suzuki, R. Taniuchi, and V. Werner, *Phys. Rev. Lett.* **113**, 032505 (2014).
- [14] P. T. Hosmer, H. Schatz, A. Aprahamian, O. Arndt, R. R. C. Clement, A. Estrade, K.-L. Kratz, S. N. Liddick, P. F. Mantica, W. F. Mueller, F. Montes, A. C. Morton, M. Ouellette, E. Pellegrini, B. Pfeiffer, P. Reeder, P. Santi, M. Steiner, A. Stolz, B. E. Tomlin, W. B. Walters, and A. Wöhr, *Phys. Rev. Lett.* **94**, 112501 (2005).
- [15] T. Marchi, G. de Angelis, J. J. Valiente-Dobón, V. M. Bader, T. Baugher, D. Bazin, J. Berryman, A. Bonaccorso, R. Clark, L. Coraggio, H. L. Crawford, M. Doncel, E. Farnea, A. Gade, A. Gadea, A. Gargano, T. Glasmacher, A. Gottardo, F. Gramegna, N. Itaco, P. R. John, R. Kumar, S. M. Lenzi, S. Lunardi, S. McDaniel, C. Michelagnoli, D. Mengoni, V. Modamio, D. R. Napoli, B. Quintana, A. Ratkiewicz, F. Recchia, E. Sahin, R. Stroberg, D. Weisshaar, K. Wimmer, and R. Winkler, *Phys. Rev. Lett.* **113**, 182501 (2014).
- [16] K. Kolos, D. Miller, R. Grzywacz, H. Iwasaki, M. Al-Shudifat, D. Bazin, C. R. Bingham, T. Braunroth, G. Cerizza, A. Gade, A. Lemasson, S. N. Liddick, M. Madurga, C. Morse, M. Portillo, M. M. Rajabali, F. Recchia, L. L. Riedinger, P. Voss, W. B. Walters, D. Weisshaar, K. Whitmore, K. Wimmer, and J. A. Tostevin, *Phys. Rev. Lett.* **116**, 122502 (2016).
- [17] S. M. Lenzi, F. Nowacki, A. Poves, and K. Sieja, *Phys. Rev. C* **82**, 054301 (2010).
- [18] O. Sorlin, S. Leenhardt, C. Donzau, J. Duprat, F. Azaiez, F. Nowacki, H. Grawe, Z. Dombrádi, F. Amorini, A. Astier, D. Baiborodin, M. Belleguic, C. Borcea, C. Bourgeois, D. M. Cullen, Z. Dlouhy, E. Dragulescu, M. Górska, S. Grévy, D. Guillemaud-Mueller, G. Hagemann, B. Herskind, J. Kiener, R. Lemmon, M. Lewitowicz, S. M. Lukyanov, P. Mayet, F. de Oliveira Santos, D. Pantalica, Y.-E. Penionzhkevich, F. Pougheon, A. Poves, N. Redon, M. G. Saint-Laurent, J. A. Scarpaci, G. Sletten, M. Stanoiu, O. Tarasov, and C. Theisen, *Phys. Rev. Lett.* **88**, 092501 (2002).
- [19] O. Kenn, K.-H. Speidel, R. Ernst, J. Gerber, P. Maier-Komor, and F. Nowacki, *Phys. Rev. C* **63**, 064306 (2001).
- [20] S. G. C. Nesaraja and B. Singh, **111** (2010).
- [21] J. M. Allmond, B. A. Brown, A. E. Stuchbery, A. Galindo-Uribarri, E. Padilla-Rodal, D. C. Radford, J. C. Batchelder, M. E. Howard, J. F. Liang, B. Manning, R. L. Varner, and C.-H. Yu, *Phys. Rev. C* **90**, 034309 (2014).
- [22] A. Dewald, O. Mller, and P. Petkov, *Progress in Particle and Nuclear Physics* **67**, 786 (2012).
- [23] H. Iwasaki, A. Dewald, T. Braunroth, C. Fransen, D. Smalley, A. Lemasson, C. Morse, K. Whitmore, and C. Loelius, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **806**, 123 (2016).
- [24] S. Paschalis, I. Lee, A. Macchiavelli, C. Campbell, M. Cromaz, S. Gros, J. Pavan, J. Qian, R. Clark, H. Crawford, D. Doering, P. Fallon, C. Lionberger, T. Loew, M. Petri, T. Stezelberger, S. Zimmermann, D. Radford, K. Lagergren, D. Weisshaar, R. Winkler, T. Glasmacher, J. Anderson, and C. Beausang, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **709**, 44 (2013).
- [25] H. Iwasaki, A. Lemasson, C. Morse, A. Dewald, T. Braunroth, V. M. Bader, T. Baugher, D. Bazin, J. S. Berryman, C. M. Campbell, A. Gade, C. Langer, I. Y. Lee, C. Loelius, E. Lunderberg, F. Recchia, D. Smalley, S. R. Stroberg, R. Wadsworth, C. Walz, D. Weisshaar, A. Westerberg, K. Whitmore, and K. Wimmer, *Phys. Rev. Lett.* **112**, 142502 (2014).
- [26] D. Morrissey, B. Sherrill, M. Steiner, A. Stolz, and I. Wiedenhoever, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* **204**, 90 (2003), 14th International Conference on Electromagnetic Isotope Separators and Techniques Related to their Applications.
- [27] D. Bazin, J. Caggiano, B. Sherrill, J. Yurkon, and A. Zeller, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* **204**, 629 (2003), 14th International Conference on Electromagnetic Isotope Separators and Techniques Related to their Applications.
- [28] S. Agostinelli, J. Allison, K. Amako, J. Apostolakis, H. Araujo, P. Arce, M. Asai, D. Axen, S. Banerjee, G. Barrand, F. Behner, L. Bellagamba, J. Boudreau, L. Broglia, A. Brunengo, H. Burkhardt, S. Chauvie, J. Chuma, R. Chytrcek, G. Cooperman, G. Cosmo, P. Degtyarenko, A. Dell'Acqua, G. Depaola, D. Dietrich, R. Enami, A. Feliciello, C. Ferguson, H. Fesefeldt, G. Folger, F. Foppiano, A. Forti, S. Garelli, S. Giani, R. Giannitrapani, D. Gibin, J. G. Cadenas, I. Gonzlez, G. G. Abril, G. Greeniaus, W. Greiner, V. Grichine, A. Grossheim, S. Guatelli, P. Gumplinger, R. Hamatsu, K. Hashimoto, H. Hasui, A. Heikkinen, A. Howard, V. Ivanchenko, A. Johnson, F. Jones, J. Kallenbach, N. Kanaya, M. Kawabata, Y. Kawabata, M. Kawaguti, S. Kelner, P. Kent, A. Kimura, T. Kodama, R. Kokoulin, M. Kossov, H. Kurashige, E. Lamanna, T. Lampn, V. Lara, V. Lefebure, F. Lei, M. Liendl, W. Lockman, F. Longo, S. Magni, M. Maire, E. Medernach, K. Minamimoto, P. M. de Freitas, Y. Morita, K. Murakami, M. Nagamatsu, R. Nartallo, P. Nieminen, T. Nishimura, K. Ohtsubo, M. Okamura, S. O'Neale, Y. Oohata, K. Paech, J. Perl, A. Pfeiffer, M. Pia, F. Ranjard, A. Rybin, S. Sadilov, E. D. Salvo, G. Santin, T. Sasaki, N. Savvas, Y. Sawada, S. Scherer, S. Sei, V. Sirotenko, D. Smith, N. Starkov, H. Stoecker, J. Sulkimo, M. Takahata, S. Tanaka, E. Tcherniaev, E. S. Tehrani, M. Tropeano, P. Truscott, H. Uno, L. Urban, P. Urban, M. Verderi, A. Walkden, W. Wander, H. Weber, J. Wellisch, T. Wenaus, D. Williams, D. Wright,



- T. Yamada, H. Yoshida, and D. Zschesche, [Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment](#) **506**, 250 (2003).
- [29] P. Adrich, D. Enderich, D. Miller, V. Moeller, R. Norris, K. Starosta, C. Vaman, P. Voss, and A. Dewald, [Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment](#) **598**, 454 (2009).
- [30] M. Honma, T. Otsuka, B. Brown, and T. Mizusaki, [The European Physical Journal A - Hadrons and Nuclei](#) **25**, 499 (2005).
- [31] A. Poves, J. Snchez-Solano, E. Caurier, and F. Nowacki, [Nuclear Physics A](#) **694**, 157 (2001).
- [32] T. Mizusaki, N. Shimizu, Y. Utsuno, and M. Honma, “Code mshell64,” Unpublished.