

This is a repository copy of *Embedding in factorisable restriction monoids*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/109893/>

Version: Accepted Version

Article:

Gould, Victoria Ann Rosalind orcid.org/0000-0001-8753-693X, Hartmann, Miklos and Szendrei, Maria (2017) Embedding in factorisable restriction monoids. *Journal of Algebra*. 476. pp. 216-237. ISSN 1090-266X

<https://doi.org/10.1016/j.jalgebra.2016.11.033>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

EMBEDDING IN FACTORISABLE RESTRICTION MONOIDS

VICTORIA GOULD, MIKLÓS HARTMANN, AND MÁRIA B. SZENDREI

ABSTRACT. Each restriction semigroup is proved to be embeddable in a factorisable restriction monoid, or, equivalently, in an almost factorisable restriction semigroup. It is also established that each restriction semigroup has a proper cover which is embeddable in a semidirect product of a semilattice by a group.

1. INTRODUCTION

Restriction semigroups are non-regular generalisations of inverse semigroups. They are semigroups equipped with two additional unary operations which satisfy certain identities. In particular, each inverse semigroup determines a restriction semigroup where the unary operations assign the idempotents aa^{-1} and $a^{-1}a$, respectively, to any element a . The class of restriction semigroups is just the variety of algebras generated by these restriction semigroups obtained from inverse semigroups, see [3]. Restriction semigroups (formerly called weakly E -ample semigroups) have arisen from a number of mathematical perspectives. For a historical overview of restriction semigroups and a detailed introduction to their fundamental properties the reader is referred to [5].

So far, a number of important results of the rich structure theory of inverse semigroups have been recast in the broader setting of restriction semigroups. The current paper is a contribution to this body of work. Our primary goal is to prove the following (see Theorem 3.10).

Main result *Any restriction semigroup is embeddable in a factorisable restriction monoid as a restriction semigroup, that is, in a way that respects both unary operations.*

Earlier works [7], [8], [11], [12] have achieved embeddings of some or all restriction semigroups in members of wider classes that are inherently one-sided. The difficulty of our task is perhaps understood when we remark that it is undecidable whether a finite restriction semigroup embeds as a restriction semigroup into an inverse semigroup [6]. To this end we need to develop a proof strategy that is different in some crucial aspects from earlier ones.

In the theory of inverse semigroups, semidirect products of semilattices by groups play an important role. One of the reasons for this is that every inverse semigroup divides

Date: November 5th, 2016.

Research partially supported by the Hungarian National Foundation for Scientific Research grants no. K083219, K104251, PD115705, and by EPSRC grant no. EP/IO32312/1.

Mathematical Subject Classification (2010): 20M10, 20M05.

Key words: Restriction semigroup, weakly E -ample semigroup, (almost) factorisability, semidirect product.

such a semidirect product (see [9] for the main results in this area and for references). In fact, something stronger is true, namely that every inverse semigroup can be embedded into an (idempotent separating) homomorphic image of such a semidirect product. One way to see this result is through extensions of partial isomorphisms of semilattices: an inverse semigroup determines partial isomorphisms of its semilattice of idempotents, and constructing the corresponding semidirect product in fact includes embedding this semilattice into a bigger one, and finding a group acting on the bigger semilattice such that the original partial isomorphisms are restrictions of the automorphisms determined by the group action.

The situation for restriction semigroups is similar, but also different in some crucial respects. In this case groups are replaced by monoids, which gives rise to several problems. The first result analogous to that formulated above for inverse semigroups was obtained by the third author in [12]: she has shown that every restriction semigroup can be embedded into a (projection separating) homomorphic image of a so-called W -product of a semilattice by a monoid, that is, into an almost left factorisable restriction semigroup (cf. [4]). An alternative proof, based on the idea of extending partial isomorphisms to injective endomorphisms, has been presented by Kudryavtseva in [8].

In a W -product in general, the monoid acts by injective endomorphisms on the semilattice, and this was the case in the results of [12] and [8] mentioned. In particular, if the monoid acts by automorphisms on the semilattice then the W -product becomes a semidirect product.

In this paper we obtain the strongest possible result in this direction by showing that the monoid and the semilattice can be chosen such that the monoid acts on the semilattice by automorphisms, thus demonstrating that any restriction semigroup can be embedded into a (projection separating) homomorphic image of a semidirect product of a semilattice by a monoid where the monoid acts on the semilattice by automorphisms. In other words, we prove that any restriction semigroup can be embedded into an almost factorisable restriction semigroup, or, equivalently, into a factorisable restriction monoid (cf. [4], [12]). Furthermore, the construction applied in the proof of this result allows us to show that the proper cover embeddable in a W -product, which is provided for a restriction semigroup in [11] and [8], is embeddable also in a semidirect product of a semilattice by a group.

Note that restriction semigroups in general, proper restriction semigroups, almost factorisable restriction semigroups, factorisable restriction monoids, and the semidirect products of semilattices by monoids where monoids act on semilattices by automorphisms are defined in a left-right symmetric way. However, this is not the case with almost left factorisable restriction semigroups or W -products. Therefore the embedding results of the present paper reflect the left-right symmetry of restriction and proper restriction semigroups, while this was not the case with the foregoing results in [11] and [8].

In the last section, we strengthen some results in [8] and [7] by proving that each ultra F -restriction, or, equivalently, each perfect restriction monoid S whose greatest reduced factor is a free monoid is embeddable in a semidirect product of a semilattice by a monoid in such a way that the monoid acts on the semilattice by automorphisms, and all congruences of S extend to the semidirect product.

2. PRELIMINARIES

In general, mappings and partial mappings are considered as right operands, and so their products are calculated from the left to the right. This applies, amongst others, to the group of automorphisms $\text{Aut } Y$, to the Munn semigroup T_Y of a semilattice Y , and to the symmetric inverse monoid $I(X)$ on a set X . By writing $\text{Aut}^{\text{op}} Y$, T_Y^{op} , etc., we indicate that the (partial) mappings in them are considered left operands, and their products are calculated from the right to the left.

2.1. Restriction semigroups. [5] A *left restriction semigroup* is defined to be an algebra of type $(2, 1)$, more precisely, an algebra $S = (S; \cdot, +)$ where $(S; \cdot)$ is a semigroup and $+$ is a unary operation such that the following identities are satisfied:

$$(1) \quad x^+x = x, \quad x^+y^+ = y^+x^+, \quad (x^+y)^+ = x^+y^+, \quad xy^+ = (xy)^+x.$$

A *right restriction semigroup* is defined dually, that is, it is an algebra $S = (S; \cdot, *)$ satisfying the duals of the identities (1). Finally, if $S = (S; \cdot, +, *)$ is an algebra of type $(2, 1, 1)$ where $S = (S; \cdot, +)$ is a left restriction semigroup, $S = (S; \cdot, *)$ is a right restriction semigroup and the identities

$$(2) \quad (x^+)^* = x^+, \quad (x^*)^+ = x^*$$

hold, then it is called a *restriction semigroup*. Notice that the defining properties of a restriction semigroup are left-right dual. Therefore in the sequel dual definitions and statements will not be explicitly formulated.

Among restriction semigroups, the notions of subalgebra, homomorphism, congruence and factor algebra are understood in type $(2, 1, 1)$, which is emphasised by using the expressions $(2, 1, 1)$ -subsemigroup, $(2, 1, 1)$ -morphism, $(2, 1, 1)$ -congruence and $(2, 1, 1)$ -factor semigroup, respectively. A restriction semigroup with identity element is also called a *restriction monoid*.

In particular, each monoid M becomes a restriction monoid by defining $a^+ = a^* = 1$ for any $a \in M$. It is easy to see that these restriction semigroups are just those with both unary operations being constant. Such a restriction monoid is called *reduced*. Notice that the submonoids, congruences, etc. of monoids and the $(2, 1, 1)$ -subsemigroups, $(2, 1, 1)$ -congruences, etc. of the reduced restriction monoids obtained from them coincide. Therefore we often consider reduced restriction monoids just as monoids, and vice versa. Similarly, each semilattice Y becomes a restriction semigroup by defining $a^+ = a^* = a$ for every $a \in Y$. These restriction semigroups are just those where both unary operations equal the identity map. Clearly, the $(2, 1, 1)$ -subsemigroups, $(2, 1, 1)$ -congruences, etc. of such a restriction semigroup Y are just the subsemilattices, congruences, etc. of the semilattice Y . Therefore such a restriction semigroup is simply considered and called a semilattice.

Let S be any restriction semigroup. By (2), we have $\{x^+ : x \in S\} = \{x^* : x \in S\}$. This set is called *the set of projections of S* , and is denoted by $P(S)$. By (1) and its dual, $P(S)$ can be seen to be a $(2, 1, 1)$ -subsemigroup in S which is a semilattice. Notice that

a restriction semigroup S is reduced if and only if $P(S)$ is a singleton, and, if this is the case, then the unique element of $P(S)$ is the identity element of S .

Given a restriction semigroup S , we define a relation \leq on S such that, for every $a, b \in S$,

$$a \leq b \quad \text{if and only if} \quad a = a^+b.$$

It is easy to see that $a \leq b$ if and only if $a = eb$ for some $e \in P(S)$. Observe that the dual of this relation is the same since $a = a^+b$ implies $a = b(a^+b)^* = ba^*$, and the dual implication is also valid. The relation \leq is a compatible partial order on S , and it extends the natural partial order of the semilattice $P(S)$. It is called the *natural partial order on S* .

We also consider a relation on S , denoted by σ_S , or simply σ : for any $a, b \in S$, let

$$a \sigma b \quad \text{if and only if} \quad ea = eb \quad \text{for some} \quad e \in P(S).$$

Again notice that if there exists $e \in P(S)$ with $ea = eb$ then there exists also $f \in P(S)$ with $af = bf$, and conversely. Therefore the relation defined dually to σ coincides with σ . The relation σ is the least congruence on $S = (S; \cdot)$ where $P(S)$ is in a congruence class, which we denote by $P(S)\sigma$. Consequently, it is the least $(2, 1, 1)$ -congruence ρ on $S = (S; \cdot, +, *)$ such that the $(2, 1, 1)$ -factor semigroup S/ρ is reduced. Therefore we call σ *the least reduced $(2, 1, 1)$ -congruence on S* . Obviously, $P(S)\sigma$ is the identity element of S/σ . The reduced restriction monoid S/σ is often considered just as a monoid $S/\sigma = (S/\sigma; \cdot, P(S)\sigma)$.

Let S and T be restriction semigroups. We say that S is *proper* if, for every $a, b \in S$, the relations $a^+ = b^+$ and $a \sigma b$ together imply $a = b$ and the dual property also holds. It is easy to see that each $(2, 1, 1)$ -subsemigroup of a proper restriction semigroup is proper. A $(2, 1, 1)$ -morphism $\varphi: T \rightarrow S$ is called *projection separating* if $e\varphi = f\varphi$ implies $e = f$ for every $e, f \in P(T)$. We say that T is a *proper cover* of S if T is proper and there exists a projection separating $(2, 1, 1)$ -morphism from T onto S .

2.2. Semidirect products and factorisability. [4], [12] Let T be a monoid and $\mathcal{X} = (\mathcal{X}; \wedge)$ a semilattice. We say that T *acts on \mathcal{X} (on the left) by automorphisms* if a monoid homomorphism $\alpha: T \rightarrow \text{Aut}^{\text{op}} \mathcal{X}$, $t \mapsto \alpha_t$ is given. The element $\alpha_t a$ ($t \in T$, $a \in \mathcal{X}$) is usually denoted by $t \cdot a$. The fact that T acts on \mathcal{X} by automorphisms is equivalent to requiring that the mapping $\alpha_t: \mathcal{X} \rightarrow \mathcal{X}$, $a \mapsto t \cdot a$ is bijective for every $t \in T$, and that the following equalities are valid for every $a, b \in \mathcal{X}$ and $t, u \in T$:

$$(3) \quad t \cdot (a \wedge b) = t \cdot a \wedge t \cdot b, \quad u \cdot (t \cdot a) = (ut) \cdot a, \quad 1 \cdot a = a.$$

The *semidirect product* $\mathcal{X} \rtimes T$ is the algebra of type $(2, 1, 1)$ defined on the set $\mathcal{X} \times T = \{(t \cdot a, t) : a \in \mathcal{X}, t \in T\}$ with the following operations:

$$(t \cdot a, t)(u \cdot b, u) = (t \cdot (a \wedge u \cdot b), tu), \quad (t \cdot a, t)^+ = (t \cdot a, 1) \quad \text{and} \quad (t \cdot a, t)^* = (a, 1).$$

Note that since T acts by automorphisms, the multiplication and the $+$ operation may be also expressed as

$$(a, t)(b, u) = (a \wedge t \cdot b, tu), \quad \text{and} \quad (a, t)^+ = (a, 1).$$

The reason for the unusual formulation is in the definition of the operation $*$: since T is not necessarily a group, one cannot use t^{-1} . It is routine to check that $\mathcal{X} \rtimes T$ is a

proper restriction semigroup where $P(\mathcal{X} \rtimes T) = \{(a, 1) : a \in \mathcal{X}\}$ is isomorphic to \mathcal{X} , the relation σ is just the kernel of the second projection $\mathcal{X} \rtimes T \rightarrow T$, $(t \cdot a, t) \mapsto t$, and so $(\mathcal{X} \rtimes T)/\sigma$ is isomorphic to T . Moreover, the natural partial order is the following relation: $(t \cdot a, t) \leq (u \cdot b, u)$ if and only if $a \leq b$ and $t = u$. The semidirect product $\mathcal{X} \rtimes T$ is a (proper restriction) monoid if and only if \mathcal{X} has an identity element. In this case the fact that T acts by automorphisms implies that $t \cdot e = e$ for any $t \in T$ where e is the identity element of \mathcal{X} . To avoid confusion, we call the attention to the fact that, from now on, when speaking about a semidirect product of a semilattice by a monoid, it is always meant to be a restriction semigroup just defined. In particular, the action of the monoid on the semilattice is taken to be an action by automorphisms.

Note that the semidirect product $\mathcal{X} \rtimes T$ is isomorphic to the reverse semidirect product $T \ltimes \mathcal{X}$ where the right action of T on \mathcal{X} is given by the monoid homomorphism $T \rightarrow \text{Aut } \mathcal{X}$, $t \mapsto \alpha_t^{-1}$. By definition (see [4]), the reverse semidirect product $T \ltimes \mathcal{X}$ is, in fact, a W -product $W(T, \mathcal{X})$. Therefore the same is the case with the semidirect product $\mathcal{X} \rtimes T$.

In this paper, an action of a monoid T on a semilattice \mathcal{X} by automorphisms will frequently come as a restriction of an action of a group G to its submonoid T . In this case, the $*$ operation may be expressed as

$$(a, t)^* = (t^{-1} \cdot a, 1),$$

and so we will prefer representing the elements of the semidirect product $\mathcal{X} \rtimes T$, as it is usual within the inverse semigroup $\mathcal{X} \rtimes G$, in the form (a, t) with $a \in \mathcal{X}$, $t \in T$. In this form, the natural partial order is $(a, t) \leq (b, u)$ if and only if $a \leq b$ and $t = u$.

Note that, instead of assuming that a monoid T is embeddable in a group and it acts on a semilattice \mathcal{X} by automorphisms, we can require without loss of generality that T is a submonoid in a group G which it generates, G acts on \mathcal{X} , and the action of T is the restriction of the action of G . For, recall first that if T is embeddable in a group then the inclusion map $\kappa: T \rightarrow G_T$ where $G_T = \langle T | \Xi \rangle$ is given by the defining relations $\Xi = \{tuv^{-1} : t, u, v \in T \text{ and } tu = v \text{ in } T\}$ is an embedding. Second, if the action of T on \mathcal{X} is defined by the homomorphism $\alpha: T \rightarrow \text{Aut}^{\text{op}} \mathcal{X}$ then $\alpha_t \alpha_u \alpha_v^{-1}$ is the identity automorphism for every $t, u, v \in T$ with $tu = v$, and so there exists a homomorphism $\beta: G_T \rightarrow \text{Aut}^{\text{op}} \mathcal{X}$ such that $\alpha = \kappa\beta$. Thus β defines an action of G_T on \mathcal{X} , and its restriction to T is just α .

Factorisable restriction monoids and almost factorisable restriction semigroups were introduced in [12] (see also [4]) analogously to factorisable inverse monoids and almost factorisable inverse semigroups, respectively. A restriction monoid F is called *factorisable* if $F = P(F)U(F)$ where $U(F) = \{u \in F : u^+ = u^* = 1\}$ is the reduced $(2, 1, 1)$ -subsemigroup of F analogous to the group of units in an inverse monoid. Note that $F = P(F)U(F)$ if and only if $F = U(F)P(F)$, therefore factorisability is a left-right symmetric property. The notion of an *almost factorisable* restriction semigroup is defined by means of permissible subsets but, instead of the definition, it suffices for the purposes of this paper to recall how they relate to factorisable restriction monoids ([12, Proposition 3.10 and Theorem 3.11]). If F is a factorisable restriction monoid then both F and the $(2, 1, 1)$ -subsemigroup $F \setminus U(F)$ of F are almost factorisable, and conversely, each almost factorisable restriction

semigroup is, up to $(2, 1, 1)$ -isomorphism, of the latter form. In particular, this implies that a restriction semigroup is $(2, 1, 1)$ -embeddable in a factorisable restriction monoid if and only if it is $(2, 1, 1)$ -embeddable in an almost factorisable restriction semigroup.

In the proof of the main result of the paper, we need the following description of these restriction semigroups by means of semidirect products.

Proposition 2.1. [12, Theorem 3.12] *A restriction monoid (semigroup) is factorisable (almost factorisable) if and only if it is a $(2, 1, 1)$ -morphic image of a semidirect product of a semilattice with identity (a semilattice) by a monoid.*

Given an almost factorisable restriction semigroup F^- in the above form, now we establish for later use, how to construct a factorisable restriction monoid F such that $F^- = F \setminus U(F)$.

Lemma 2.2. *Consider an almost factorisable restriction semigroup $F^- = (\mathcal{X} \rtimes T) / \rho$, where T is a monoid acting on the semilattice \mathcal{X} by automorphisms, and ρ is a $(2, 1, 1)$ -congruence on the restriction semigroup $\mathcal{X} \rtimes T$. Let \mathcal{X}^e be the semilattice obtained from \mathcal{X} by adjoining an identity element $e \notin \mathcal{X}$ even if \mathcal{X} has an identity element. Extend the action of T on \mathcal{X} to \mathcal{X}^e by putting $t \cdot e = e$ for every $t \in T$, and consider the relation $\rho_e = \rho \cup \iota$ on the semidirect product $\mathcal{X}^e \rtimes T$ where ι is the equality relation on $U = (\mathcal{X}^e \rtimes T) \setminus (\mathcal{X} \rtimes T)$. Then ρ_e is a $(2, 1, 1)$ -congruence on the restriction monoid $\mathcal{X}^e \rtimes T$ extending ρ , $F = (\mathcal{X}^e \rtimes T) / \rho_e$ is a factorisable restriction monoid with $U(F) = \{u\rho_e : u \in U\} = \{\{(e, t)\} : t \in T\}$, and the mapping $F^- \rightarrow F$, $(a, t)\rho \mapsto (a, t)\rho_e$ ($(a, t) \in \mathcal{X} \rtimes T$) is an injective $(2, 1, 1)$ -morphism with range $F \setminus U(F)$.*

Proof. Obviously, T acts on \mathcal{X}^e by automorphisms, the semidirect product $\mathcal{X}^e \rtimes T$ is a restriction monoid with $U(\mathcal{X}^e \rtimes T) = U$, and $\mathcal{X} \rtimes T$ is a $(2, 1, 1)$ -subsemigroup and ideal of $\mathcal{X}^e \rtimes T$. Moreover, it is also clear that the relation ρ_e is an equivalence whose restriction to $\mathcal{X} \rtimes T$ is ρ , and that it is compatible with both unary operations. It remains to check that ρ_e is compatible with the multiplication. We have to verify that if $(a, t)\rho(b, u)$ in $\mathcal{X} \rtimes T$ and $v \in T$ then both $(a, t)(e, v)\rho(b, u)(e, v)$ and $(e, v)(a, t)\rho(e, v)(b, u)$ hold, that is, both $(a, tv)\rho(b, uv)$ and $(v \cdot a, vt)\rho(v \cdot b, vu)$ are valid. We verify the first relation, the second being similar. Since ρ is a $(2, 1, 1)$ -congruence on $\mathcal{X} \rtimes T$, we have $(a \wedge t \cdot c, tv) = (a, t)(c, v)\rho(b, u)(c, v) = (b \wedge u \cdot c, uv)$ for every $c \in \mathcal{X}$. Choosing c to be the unique element of \mathcal{X} such that $t \cdot c = a$, we obtain that $(a, tv)\rho = (b \wedge u \cdot c, uv)\rho \leq (b, uv)\rho$ in F^- . Changing the roles of (a, t) and (b, u) the equality follows, proving the first relation. \square

2.3. Congruences on restriction semigroups generated by relations. [12] Given a set X of variables, by a (*restriction*) *term in X* we mean a formal expression built up from the elements of X by means of the operational symbols — the binary operational symbol \cdot and the unary operational symbols $^+$ and * — in finitely many steps. Since the binary operation \cdot is always interpreted in a semigroup we delete the unnecessary parentheses from the terms. If S is a fixed restriction semigroup then we introduce a nullary operational symbol for every element s in S , and, for simplicity, denote it also by s . By a *polynomial of S* we mean an expression obtained in a way similar to terms, but from variables and

these nullary operational symbols. A polynomial can also be interpreted in the way that such nullary operational symbols — briefly elements of S — are substituted for certain variables in a term. So a *unary polynomial of S* , that is, a polynomial in one variable x , is of the form $\mathbf{t}(x, s_1, \dots, s_{k-1})$ for some term \mathbf{t} in k variables and for some elements $s_1, \dots, s_{k-1} \in S$. To simplify our notation, we denote by S^* the set of finite (possibly empty) sequences of elements of S , and we use Greek letters to denote elements of S^* . The length of the sequences will be determined by the context where they appear. For example, if \mathbf{t} is a term in k variables and we are talking about $\mathbf{t}(x, \alpha)$ for some $\alpha \in S^*$ then the length of α is $k - 1$.

Let S be a restriction semigroup and let $\tau \subseteq S \times S$ be a symmetric relation. We denote by $\tau^\#$ the $(2, 1, 1)$ -congruence on S generated by τ . It is a basic fact of universal algebra that if $s, t \in S$ then $s \tau^\# t$ if and only if there exist a sequence $p_1(x), p_2(x), \dots, p_k(x)$ of unary polynomials of S and elements $c_1, d_1, \dots, c_k, d_k \in S$ such that $(c_i, d_i) \in \tau$ for all i ($1 \leq i \leq k$) and

$$s = p_1(c_1), p_1(d_1) = p_2(c_2), \dots, p_k(d_k) = t.$$

In general, unary polynomials over a restriction semigroup can be quite complicated. However, it is shown in [12] that the unary polynomials appearing in the sequence can be chosen to be simpler, and this will be key to our later arguments.

We define two sequences of terms in variables x, y, z, y_0, z_0, \dots in the following way: let

$$\mathbf{t}_+^{(0)}(x, y_0, z_0) = (y_0 x z_0)^+, \quad \mathbf{t}_*^{(0)}(x, y_0, z_0) = (y_0 x z_0)^*,$$

and, for every $i \in \mathbb{N}$, let

$$\begin{aligned} \mathbf{t}_+^{(i)}(x, y_0, z_0, \dots, y_{i-2}, z_{i-1}, y_i) &= (y_i \mathbf{t}_*^{(i-1)}(x, y_0, z_0, \dots, y_{i-2}, z_{i-1}))^+, \\ \mathbf{t}_*^{(i)}(x, y_0, z_0, \dots, z_{i-2}, y_{i-1}, z_i) &= (\mathbf{t}_+^{(i-1)}(x, y_0, z_0, \dots, z_{i-2}, y_{i-1}) z_i)^*. \end{aligned}$$

For convenience we note that

$$\begin{aligned} \mathbf{t}_+^{(1)}(x, y_0, z_0, y_1) &= (y_1 (y_0 x z_0)^+)^+, \\ \mathbf{t}_*^{(1)}(x, y_0, z_0, z_1) &= ((y_0 x z_0)^+ z_1)^*, \\ \mathbf{t}_+^{(2)}(x, y_0, z_0, z_1, y_2) &= (y_2 ((y_0 x z_0)^+ z_1)^+)^+, \\ \mathbf{t}_*^{(2)}(x, y_0, z_0, y_1, z_2) &= ((y_1 (y_0 x z_0)^+)^+ z_2)^*. \end{aligned}$$

We define the following sets of terms:

$$\mathbf{T}_{(+)} = \{\mathbf{t}_+^{(i)}, \mathbf{t}_*^{(i+1)} : i \in \mathbb{N}_0 \text{ is even}\}, \quad \mathbf{T}_{(*)} = \{\mathbf{t}_*^{(i)}, \mathbf{t}_+^{(i+1)} : i \in \mathbb{N}_0 \text{ is even}\},$$

$$\mathbf{T} = \{\mathbf{t} : \mathbf{t}(x, y, z) = yxz \text{ or } \mathbf{t}(x, y, z, y_0, z_0, \dots) = \mathbf{u}yz \text{ where } \mathbf{u} \in \mathbf{T}_{(+)} \cup \mathbf{T}_{(*)}\}.$$

Notice that $\mathbf{T}_{(+)}$ ($\mathbf{T}_{(*)}$) contains terms with $+$ ($*$) as innermost unary operation. The set $\mathbf{T}_{(+)}$ ($\mathbf{T}_{(*)}$) differs from the set \mathbf{T}_+ (\mathbf{T}_*) used in [12] since the latter consists of the terms with $+$ ($*$) as outermost unary operation. The reason for this modification (and slightly awkward notation) is that, in contrast to the proof of the main result of [12], we are going to distinguish cases depending on the innermost unary operation.

It is shown in [12] that the terms in \mathbf{T} are enough to determine a $(2, 1, 1)$ -congruence generated by a symmetric relation.

Proposition 2.3. [12, Proposition 4.5] *If τ is a symmetric relation on a restriction semigroup S then, for any $s, t \in S$, we have $s \tau^\# t$ if and only if $s = t$ or there exist $k \in \mathbb{N}$, $\mathbf{t}_1, \dots, \mathbf{t}_k \in \mathbf{T}$, $\alpha_1, \dots, \alpha_k \in S^*$ and $(c_1, d_1), \dots, (c_k, d_k) \in \tau$ such that*

$$s = \mathbf{t}_1(c_1, \alpha_1), \mathbf{t}_1(d_1, \alpha_1) = \mathbf{t}_2(c_2, \alpha_2), \dots, \mathbf{t}_k(d_k, \alpha_k) = t.$$

Let S and S' be restriction semigroups such that S is a $(2, 1, 1)$ -subsemigroup of S' , and let ρ be $(2, 1, 1)$ -congruence on S . A $(2, 1, 1)$ -congruence ρ' on S' is said to *extend* ρ if the restriction of ρ' to S equals ρ . We say that ρ *extends to* S' if there exists a $(2, 1, 1)$ -congruence ρ' on S' extending ρ . Notice that ρ extends to S' if and only if the $(2, 1, 1)$ -congruence on S' generated by ρ extends ρ .

2.4. Free restriction monoids. [3] A transparent model of the free restriction monoid on a set is given in [3] as a full subsemigroup in the free inverse monoid on the same set (cf. [9]).

Let Ω be a non-empty set, and denote by Ω^* and $\text{FG}(\Omega)$, respectively, the usual models of the free monoid and the free group on Ω , where Ω^* consists of all words (i.e., of finite (possibly empty) sequences) over Ω and $\text{FG}(\Omega)$ consists of all reduced words over $\Omega \cup \Omega^{-1}$. The empty word is denoted by 1. Obviously, Ω^* is a subsemigroup of $\text{FG}(\Omega)$, and 1 is the identity element in both.

Let \mathcal{C} be the Cayley graph of $\text{FG}(\Omega)$ which is well known to be a tree. Therefore each finite subset $A \subseteq \text{FG}(\Omega)$ determines a maximum subgraph of \mathcal{C} having A as its set of vertices. From now on, we identify A with this subgraph. Let \mathcal{X} be the set of all finite connected subgraphs of \mathcal{C} (i.e., of all finite subtrees), partially ordered by reverse inclusion. Then \mathcal{X} is a semilattice without an identity. Clearly, $\mathcal{Y} = \{A \in \mathcal{X} : 1 \in A\}$ is a principal order ideal of \mathcal{X} and a subsemilattice with identity element $\mathbf{1} = \{1\}$. Moreover, $(\text{FG}(\Omega), \mathcal{X}, \mathcal{Y})$ is a McAlister triple where $\text{FG}(\Omega)$ acts on the partially ordered set \mathcal{X} by multiplication: if $g \in \text{FG}(\Omega)$ and $A \in \mathcal{X}$ then $g \cdot A = \{ga : a \in A\}$. It is well known that the P -semigroup $P(\text{FG}(\Omega), \mathcal{X}, \mathcal{Y})$ is a free inverse monoid on Ω .

It is established in [3] that

$$\text{FR}(\Omega) = \{(A, w) \in P(\text{FG}(\Omega), \mathcal{X}, \mathcal{Y}) : w \in \Omega^*\}$$

is a $(2, 1, 1)$ -subsemigroup in $P(\text{FG}(\Omega), \mathcal{X}, \mathcal{Y})$, it is a free restriction monoid on Ω and $\text{FRS}(\Omega) = \text{FR}(\Omega) \setminus \{(\mathbf{1}, 1)\}$ is a free restriction semigroup on Ω . The main result of this paper depends on certain property of the action of the free monoid Ω^* involved in this model of $\text{FR}(\Omega)$. We are going to specify this property in the next section, but show here how $\text{FR}(\Omega)$ fits into the picture.

Since the free inverse monoid $P(\text{FG}(\Omega), \mathcal{X}, \mathcal{Y})$ is F -inverse, the partially ordered set \mathcal{X} , where the partial order will subsequently be denoted \leq , is a semilattice. Thus \mathcal{Y} is a principal ideal in it with identity element $\mathbf{1}$, and the action of $\text{FG}(\Omega)$ on the partially ordered set \mathcal{X} is, in fact, an action on the semilattice \mathcal{X} . Since \mathcal{C} is a tree, for every finite subgraph $A \subseteq \text{FG}(\Omega)$, there exists a minimum subtree A' containing A . The semilattice operation \wedge of \mathcal{X} can be given by the rule $A \wedge B = (A \cup B)'$ ($A, B \in \mathcal{X}$). Thus the semidirect products $\mathcal{X} \rtimes \text{FG}(\Omega)$ and $\mathcal{X} \rtimes \Omega^*$ are defined, and $\text{FR}(\Omega)$ can be given in this

context as the following $(2, 1, 1)$ -subsemigroup of $\mathcal{X} \rtimes \Omega^*$:

$$\text{FR}(\Omega) = \{(A, w) \in \mathcal{Y} \times \Omega^* : A \leq w \cdot \mathbf{1}\}.$$

Note that the free inverse monoid $P(\text{FG}(\Omega), \mathcal{X}, \mathcal{Y})$ is usually considered as an inverse subsemigroup of a semidirect product of another semilattice by $\text{FG}(\Omega)$, namely, of the semilattice of all finite subgraphs of \mathcal{C} with respect to the usual join. In the sequel, we need the approach in the previous paragraph, as it was the case with the W -product applied in [11] and [12].

3. MAIN RESULT

Our aim is to show that every restriction monoid, and consequently, every restriction semigroup is $(2, 1, 1)$ -embeddable in a factorisable restriction monoid, or equivalently in an almost factorisable restriction semigroup. We obtain a $(2, 1, 1)$ -embedding by taking a free restriction monoid $\text{FR}(\Omega)$, having the given restriction monoid S as its image via a $(2, 1, 1)$ -morphism, say, φ , and by considering the $(2, 1, 1)$ -embedding of $\text{FR}(\Omega)$ into the semidirect product $\mathcal{X} \rtimes \Omega^*$ described in the previous section where the monoid Ω^* acts on the semilattice \mathcal{X} by automorphisms. Since the kernel of $\varphi: \text{FR}(\Omega) \rightarrow S$ is a $(2, 1, 1)$ -congruence on $\text{FR}(\Omega)$, we consider the $(2, 1, 1)$ -congruence on the semidirect product $\mathcal{X} \rtimes \Omega^*$ it generates. If the restriction of the latter congruence to $\text{FR}(\Omega)$ equals the kernel, then we obtain a $(2, 1, 1)$ -embedding of S into a $(2, 1, 1)$ -morphic image of the semidirect product, that is, into an almost factorisable restriction semigroup. As it turns out, the fact that this approach works depends only on the fact that the actions of the free monoid Ω^* and of the free group $\text{FG}(\Omega)$ on the semilattice \mathcal{X} satisfy a simple condition. This allows us to slightly generalise our approach.

From now on, let \mathcal{X} be a semilattice and let \mathcal{Y} be a principal ideal of \mathcal{X} . Denote the identity element of \mathcal{Y} by $\mathbf{1}$. Let T be a monoid embeddable in a group, and suppose that T acts on \mathcal{X} (on the left) by automorphisms. We have seen formerly that, in this case, there exists a group G acting on \mathcal{X} such that T is a submonoid of G generating G , and the action of T is the restriction of the action of G . Let us choose and fix such a group G . In the sequel we will use boldface letters to represent the elements of the semidirect products $\mathcal{X} \rtimes G$ and $\mathcal{X} \rtimes T$, the corresponding capital letters and overlined letters to denote their first and second components, respectively — for example, $\mathbf{a} = (A, \bar{a})$.

Define the following subset of the semidirect product $\mathcal{X} \rtimes T$:

$$(4) \quad R = \{(A, \bar{a}) \in \mathcal{Y} \times T : A \leq \bar{a} \cdot \mathbf{1}\}.$$

Note that if $(A, \bar{a}) \in R$ and $A = \bar{a} \cdot B \leq \bar{a} \cdot \mathbf{1}$ where $B \in \mathcal{X}$, then as T acts by automorphisms we have $B \leq \mathbf{1}$. It is now easy to check that R is a $(2, 1, 1)$ -subsemigroup in $\mathcal{X} \rtimes T$ with identity element $(\mathbf{1}, \mathbf{1})$, and so it is a proper restriction monoid. Notice that, for any $(A, \bar{a}) \in \mathcal{X} \rtimes T$ there exists a maximum element of R less than or equal to (A, \bar{a}) which we denote by $(A, \bar{a})^\downarrow$. It is clear that $(A, \bar{a})^\downarrow = (\mathbf{1} \wedge A \wedge \bar{a} \cdot \mathbf{1}, \bar{a})$.

Observe that $G \cdot \mathcal{Y}$ is the smallest subsemilattice in \mathcal{X} containing \mathcal{Y} which is invariant under the action of G . Since, for any $g, h \in G$ and $A, B \in \mathcal{Y}$, we have $g \cdot A \wedge h \cdot B =$

$g \cdot (A \wedge g^{-1}h \cdot B)$, and \mathcal{Y} is an ideal in \mathcal{X} , we see that $G \cdot \mathcal{Y}$ is, indeed, a subsemilattice of \mathcal{X} . Since we have $R \subseteq \mathcal{Y} \times T$, we can assume without loss of generality that $\mathcal{X} = G \cdot \mathcal{Y}$.

A sequence w_1, w_2, \dots, w_n of elements of G is called *an alternating sequence in T* if, for every i ($1 \leq i \leq n$), we have $w_i = t_i^{\epsilon_i}$ for some $t_i \in T$, $\epsilon_i \in \{1, -1\}$, and, for every i ($1 \leq i < n$), we have $\epsilon_{i+1} = -\epsilon_i$. A factorisation $g = w_1 w_2 \cdots w_n$ of an element $g \in G$ is called *a nice factorisation in T with respect to $\mathbf{1}$* if w_1, w_2, \dots, w_n is an alternating sequence in T such that, for every i ($1 \leq i < n$), we have that

$$w_i \cdot \mathbf{1} \geq \mathbf{1} \wedge w_i \cdots w_n \cdot \mathbf{1}.$$

We say that the action of the monoid T on the semilattice \mathcal{X} (necessarily by automorphisms) is *nice over the group G with respect to the element $\mathbf{1} \in \mathcal{X}$* if every element $g \in G$ has a nice factorisation in T with respect to $\mathbf{1}$. Note that, in particular, the action of a group on a semilattice is always nice over the group itself with respect to any element of the semilattice. We say that $\mathcal{X} \rtimes T$ is a *nice semidirect product of a semilattice \mathcal{X} by a monoid T* if T is a submonoid of a group G acting on \mathcal{X} such that the following conditions are satisfied:

- (a) T generates G ,
- (b) the action of T on \mathcal{X} in $\mathcal{X} \rtimes T$ is the restriction of the action of G ,
- (c) the action of T on \mathcal{X} is nice over G with respect to an element $\mathbf{1} \in \mathcal{X}$ such that $\mathcal{X} = G \cdot \mathcal{Y}$ for the principal ideal \mathcal{Y} of \mathcal{X} with identity $\mathbf{1}$.

Example 3.1. The action of Ω^* on the semilattice \mathcal{X} defined in Subsection 2.4 is nice over $\text{FG}(\Omega)$ with respect to $\mathbf{1}$. It is easy to see that if $g \in \text{FG}(\Omega)$, then the unique reduced alternating factorisation $g = w_1 w_2 \cdots w_n$ where $w_1, w_2, \dots, w_n \in \Omega^* \cup (\Omega^*)^{-1}$ is a nice factorisation of g in Ω^* with respect to $\mathbf{1}$. Moreover, we have $\mathcal{X} = \text{FG}(\Omega) \cdot \mathcal{Y}$ where \mathcal{Y} is a principal ideal with identity $\mathbf{1}$, and so $\mathcal{X} \rtimes \Omega^*$ is a nice semidirect product.

Example 3.2. Let G and T be the free Abelian group and free commutative semigroup, respectively, on the set Ω . Then for every $g \in G$, there exist unique elements $u, t \in T$ such that $g = u^{-1}t$. We say that a subset $A \subseteq G$ is *min-closed* if, for every $g = \omega_1^{a_1} \cdots \omega_s^{a_s}$, $h = \omega_1^{b_1} \cdots \omega_s^{b_s} \in A$, where $\omega_1, \dots, \omega_s$ are pairwise distinct elements of Ω , we have that $\omega_1^{\min(a_1, b_1)} \cdots \omega_s^{\min(a_s, b_s)} \in A$. Let $\mathcal{X} = \{A \subseteq G : A \neq \emptyset, A \text{ is min-closed}\}$ and, for all $A, B \in \mathcal{X}$, let $A \wedge B$ be the smallest min-closed subset of G containing $A \cup B$. Then \mathcal{X} is a semilattice and G acts on \mathcal{X} by multiplication. Furthermore, $\mathcal{Y} = \{A \in \mathcal{X} : \mathbf{1} \in A\}$ is a principal ideal of \mathcal{X} with identity $\{\mathbf{1}\} = \mathbf{1}$. Note that, for any $g \in G$, the unique factorisation $g = u^{-1}t$ is nice in T with respect to $\mathbf{1}$, because if $g = \omega_1^{a_1} \cdots \omega_s^{a_s}$ then $u^{-1} = \omega_1^{\min(a_1, 0)} \cdots \omega_s^{\min(a_s, 0)} \in \{\mathbf{1}\} \wedge \{g\}$, showing that $u^{-1} \cdot \mathbf{1} \geq \mathbf{1} \wedge g \cdot \mathbf{1}$.

Let G be a group acting on a semilattice \mathcal{X} , and let ε be a congruence on \mathcal{X} . We say that ε is *G -invariant* if $x \varepsilon x'$ implies $g \cdot x \varepsilon g \cdot x'$ for every $x, x' \in \mathcal{X}$ and $g \in G$. In this case, G acts by automorphisms on \mathcal{X}/ε by the rule $g \cdot (x\varepsilon) = (g \cdot x)\varepsilon$. The following observation is straightforward.

Lemma 3.3. *If an action of a monoid T on a semilattice \mathcal{X} is nice over a group G with respect to an element $\mathbf{1} \in \mathcal{X}$ and ε is a G -invariant congruence on \mathcal{X} , then the restriction*

of the action of G on \mathcal{X}/ε to T is also nice over G with respect to $\mathbf{1}\varepsilon \in \mathcal{X}/\varepsilon$. Consequently, if $\mathcal{X} \rtimes T$ is a nice semidirect product then the same holds for $(\mathcal{X}/\varepsilon) \rtimes T$.

From now on, additionally to our former assumptions on \mathcal{X} , \mathcal{Y} , T and G , we suppose that the action of T on \mathcal{X} is nice over the group G with respect to $\mathbf{1}$.

Lemma 3.4. *Let $\mathbf{t}^{(i)} \in \mathbf{T}_{(+)} \cup \mathbf{T}_{(*)}$ and let $\alpha \in (\mathcal{X} \rtimes T)^*$. Then there exist $U, V \in \mathcal{X}$, $g \in G$ depending on α such that*

$$\mathbf{t}^{(i)}(\mathbf{c}, \alpha) = (U \wedge g \cdot \tilde{C} \wedge g\tilde{c} \cdot V, \mathbf{1})$$

for every $\mathbf{c} \in R$ where

$$\tilde{\mathbf{c}} = (\tilde{C}, \tilde{c}) = \begin{cases} (C, \bar{c}) & \text{if } \mathbf{t}^{(i)} \in \mathbf{T}_{(+)}, \\ (c^{-1} \cdot C, \bar{c}^{-1}) & \text{if } \mathbf{t}^{(i)} \in \mathbf{T}_{(*)}. \end{cases}$$

Proof. We proceed by induction on i . If $i = 0$ then $\mathbf{t}^{(i)} = (\mathbf{y}_0 \mathbf{xz}_0)^+$ or $\mathbf{t}^{(i)} = (\mathbf{y}_0 \mathbf{xz}_0)^*$, and $\alpha = ((A, \bar{a}), (B, \bar{b}))$ for some $(A, \bar{a}), (B, \bar{b}) \in \mathcal{X} \rtimes T$. If $\mathbf{t}^{(i)} \in \mathbf{T}_{(+)}$ then

$$\mathbf{t}^{(i)}(\mathbf{c}, \alpha) = ((A, \bar{a})(C, \bar{c})(B, \bar{b}))^+ = (A \wedge \bar{a} \cdot C \wedge \bar{a}\bar{c} \cdot B, \mathbf{1}),$$

so we can set $U = A$, $V = B$ and $g = \bar{a}$. Similarly, if $\mathbf{t}^{(i)} \in \mathbf{T}_{(*)}$ then

$$\mathbf{t}^{(i)}(\mathbf{c}, \alpha) = ((A, \bar{a})(C, \bar{c})(B, \bar{b}))^* = (\bar{b}^{-1}\bar{c}^{-1}\bar{a}^{-1} \cdot A \wedge \bar{b}^{-1}\bar{c}^{-1} \cdot C \wedge \bar{b}^{-1} \cdot B, \mathbf{1}),$$

so we can set $U = \bar{b}^{-1} \cdot B$, $V = \bar{a}^{-1} \cdot A$ and $g = \bar{b}^{-1}$.

Let $i \geq 1$ and $\mathbf{t}^{(i)} \in \mathbf{T}_{(+)} \cup \mathbf{T}_{(*)}$, and suppose that we have proven the statement for $i - 1$. Then $\mathbf{t}^{(i)} = (y_i \mathbf{t}^{(i-1)})^+$ or $\mathbf{t}^{(i)} = (\mathbf{t}^{(i-1)} z_i)^*$ for an appropriate $\mathbf{t}^{(i-1)} \in \mathbf{T}_{(+)} \cup \mathbf{T}_{(*)}$ and $\alpha = (\alpha', (A, \bar{a}))$ where $\alpha' \in (\mathcal{X} \rtimes T)^*$ and $(A, \bar{a}) \in \mathcal{X} \rtimes T$. Note that $\mathbf{t}^{(i)} \in \mathbf{T}_{(+)}$ if and only if $\mathbf{t}^{(i-1)} \in \mathbf{T}_{(+)}$, since $\mathbf{T}_{(+)}$ contains terms with $+$ as innermost unary operation. By the induction hypothesis there exist $U', V' \in \mathcal{X}$ and $g' \in G$ such that $\mathbf{t}^{(i-1)}(\mathbf{c}, \alpha') = (U' \wedge g' \cdot \tilde{C} \wedge g'\tilde{c} \cdot V', \mathbf{1})$ for all $\mathbf{c} \in R$. If $\mathbf{t}^{(i)} = (y_i \mathbf{t}^{(i-1)})^+$ then

$$\mathbf{t}^{(i)}(\mathbf{c}, \alpha) = ((A, \bar{a})(U' \wedge g' \cdot \tilde{C} \wedge g'\tilde{c} \cdot V', \mathbf{1}))^+ = (A \wedge \bar{a} \cdot U' \wedge \bar{a}g' \cdot \tilde{C} \wedge \bar{a}g'\tilde{c} \cdot V', \mathbf{1}),$$

so we can set $U = A \wedge \bar{a} \cdot U'$, $V = V'$ and $g = \bar{a}g'$. Similarly, if $\mathbf{t}^{(i)} = (\mathbf{t}^{(i-1)} z_i)^*$ then

$$\mathbf{t}^{(i)}(\mathbf{c}, \alpha) = ((U' \wedge g' \cdot \tilde{C} \wedge g'\tilde{c} \cdot V', \mathbf{1})(A, \bar{a}))^* = (\bar{a}^{-1} \cdot U' \wedge \bar{a}^{-1}g' \cdot \tilde{C} \wedge \bar{a}^{-1}g'\tilde{c} \cdot V' \wedge \bar{a}^{-1} \cdot A, \mathbf{1}),$$

so we can set $U = \bar{a}^{-1} \cdot U' \wedge \bar{a}^{-1} \cdot A$, $V = V'$ and $g = \bar{a}^{-1}g'$. \square

The previous lemma showed that for all terms $\mathbf{t} \in \mathbf{T}_{(+)} \cup \mathbf{T}_{(*)}$, the elements $\mathbf{t}(\mathbf{c}, \alpha)$ ($\alpha \in (\mathcal{X} \rtimes T)^*$) are of special form. Our aim is to replace in certain circumstances the sequence of constants α by $\beta \in R^*$ and the term \mathbf{t} by a term \mathbf{t}' such that $\mathbf{t}(\mathbf{c}, \alpha)^\downarrow = \mathbf{t}'(\mathbf{c}, \beta)$ for all $\mathbf{c} \in R$. The following lemma constitutes the first step in this direction.

Lemma 3.5. *Let $U, V \in \mathcal{X}$ and $g \in G$. Then there exist $\mathbf{t} \in \mathbf{T}_{(+)}$ and $\beta \in R^*$ such that*

$$(\mathbf{1} \wedge U \wedge g \cdot C \wedge g\bar{c} \cdot V, \mathbf{1}) = \mathbf{t}(\mathbf{c}, \beta)$$

for all $\mathbf{c} \in R$. Dually, there exists $\mathbf{t} \in \mathbf{T}_{(*)}$ and $\beta \in R^*$ such that

$$(\mathbf{1} \wedge U \wedge g\bar{c}^{-1} \cdot C \wedge g\bar{c}^{-1} \cdot V, \mathbf{1}) = \mathbf{t}(\mathbf{c}, \beta)$$

for all $\mathbf{c} \in R$.

Proof. Let $g = w_1 \cdots w_n$ be a nice factorisation. We prove the lemma by induction on n .

If $n = 1$ then there are four cases to consider (recall the definition of the element (\tilde{C}, \tilde{c}) from Lemma 3.4):

Conditions	$((\mathbf{1} \wedge U \wedge w_1 \cdot \tilde{C} \wedge w_1 \tilde{c} \cdot V, \mathbf{1}) =$
$w_1 \in T, \mathbf{t} \in \mathbf{T}_+$	$((\mathbf{1} \wedge U \wedge w_1 \cdot \mathbf{1}, w_1)(C, \bar{c})(\mathbf{1} \wedge V, \mathbf{1}))^+$
$w_1 \in T^{-1}, \mathbf{t} \in \mathbf{T}_+$	$((C, \bar{c})(\mathbf{1} \wedge V, \mathbf{1}))^+ (\mathbf{1} \wedge w_1^{-1} \cdot U \wedge w_1^{-1} \cdot \mathbf{1}, w_1^{-1})^*$
$w_1 \in T, \mathbf{t} \in \mathbf{T}_*$	$((\mathbf{1} \wedge U \wedge w_1 \cdot \mathbf{1}, w_1)((\mathbf{1} \wedge V, \mathbf{1})(C, \bar{c}))^*)^+$
$w_1 \in T^{-1}, \mathbf{t} \in \mathbf{T}_*$	$((\mathbf{1} \wedge V, \mathbf{1})(C, \bar{c})(\mathbf{1} \wedge w_1^{-1} \cdot U \wedge w_1^{-1} \cdot \mathbf{1}, w_1^{-1})^*)^*$

The equalities are easy to check and left to the reader. Note that throughout variations of the inequalities $w_1 \cdot \mathbf{1} \geq w_1 \cdot C$ and $c \cdot \mathbf{1} \geq C$ are used.

Let us suppose that for some $n \geq 2$ we have proven the lemma for every $g' \in G$ which admits a nice factorisation having less than n factors, and let us suppose that $g = w_1 \cdots w_n$ is a nice factorisation. Then $w_1^{-1}g = w_2 \cdots w_n$ is also a nice factorisation, so applying the induction hypothesis for $U' = w_1^{-1} \cdot U$, $V' = V$ and $g' = w_1^{-1}g$, there exists $\mathbf{t}' \in \mathbf{T}_{(+)}$ and $\beta' \in R^*$ such that

$$(\mathbf{1} \wedge w_1^{-1} \cdot U \wedge w_1^{-1}g \cdot C \wedge w_1^{-1}g\bar{c} \cdot V, \mathbf{1}) = \mathbf{t}'(\mathbf{c}, \beta')$$

for all $\mathbf{c} \in R$. Now there are two cases to consider. If $w_1 \in T$ then $w_1 \cdot \mathbf{1} \geq \mathbf{1} \wedge g \cdot \mathbf{1} \geq \mathbf{1} \wedge g \cdot C$, and so

$$\begin{aligned} (\mathbf{1} \wedge U \wedge g \cdot C \wedge g\bar{c} \cdot V, \mathbf{1}) &= ((\mathbf{1} \wedge w_1 \cdot \mathbf{1}, w_1)(\mathbf{1} \wedge w_1^{-1} \cdot U \wedge w_1^{-1}g \cdot C \wedge w_1^{-1}g\bar{c} \cdot V, \mathbf{1}))^+ \\ &= ((\mathbf{1} \wedge w_1 \cdot \mathbf{1}, w_1)\mathbf{t}'(\mathbf{c}, \beta'))^+ = \mathbf{t}(\mathbf{c}, \beta) \end{aligned}$$

for $\mathbf{t} = (y_i \mathbf{t}')^+$ and $\beta = (\beta', (\mathbf{1} \wedge w_1 \cdot \mathbf{1}, w_1))$. If $w_1 \in T^{-1}$ then by the same fact we have

$$\begin{aligned} (\mathbf{1} \wedge U \wedge g \cdot C \wedge g\bar{c} \cdot V, \mathbf{1}) &= ((\mathbf{1} \wedge w_1^{-1} \cdot U \wedge w_1^{-1}g \cdot C \wedge w_1^{-1}g\bar{c} \cdot V, \mathbf{1})(\mathbf{1} \wedge w_1^{-1} \cdot \mathbf{1}, w_1^{-1}))^* \\ &= (\mathbf{t}'(\mathbf{c}, \beta')(\mathbf{1} \wedge w_1^{-1} \cdot \mathbf{1}, w_1^{-1}))^* = \mathbf{t}(\mathbf{c}, \beta) \end{aligned}$$

for $\mathbf{t} = (\mathbf{t}'z_i)^*$ and $\beta = (\beta', (\mathbf{1} \wedge w_1^{-1} \cdot \mathbf{1}, w_1^{-1}))$. To finish this part of the proof, one has to note that the fact that the factorisation $g = w_1 \cdots w_n$ is alternating implies that the terms defined above are all contained in $\mathbf{T}_{(+)}$. This proves the first statement of the lemma. The induction step for the second statement is proven similarly, utilising the fact that $w_1 \cdot \mathbf{1} \geq \mathbf{1} \wedge g\bar{c}^{-1} \cdot C$. \square

Lemma 3.6. *Let $\mathbf{t} \in \mathbf{T}$ and let $\alpha \in (\mathcal{X} \rtimes T)^*$. Then there exist $\mathbf{t}' \in \mathbf{T}$ and $\beta \in R^*$ such that*

$$(5) \quad \mathbf{t}(\mathbf{c}, \alpha)^\downarrow = \mathbf{t}'(\mathbf{c}, \beta)$$

for all $\mathbf{c} \in R$.

Proof. There are three cases to consider: the first one is if $\mathbf{t} = yxz$. In this case let $\mathbf{t}' = \mathbf{t}$, and let $\beta = ((\mathbf{1} \wedge A \wedge \bar{a} \cdot \mathbf{1}, \bar{a}), (\mathbf{1} \wedge B \wedge \bar{b} \cdot \mathbf{1}, \bar{b}))$ if $\alpha = ((A, \bar{a}), (B, \bar{b}))$. Note that since $\mathbf{c} \in R$, the inequalities $\bar{a}c \cdot \mathbf{1}, \bar{a} \cdot \mathbf{1} \geq \bar{a} \cdot C$ imply that

$$\begin{aligned} \mathbf{t}(\mathbf{c}, \alpha)^\downarrow &= (\mathbf{1} \wedge A \wedge \bar{a} \cdot C \wedge \bar{a}c \cdot B \wedge \bar{a}c\bar{b} \cdot \mathbf{1}, \bar{a}c\bar{b}) \\ &= (\mathbf{1} \wedge A \wedge \bar{a} \cdot \mathbf{1}, \bar{a})(C, \bar{c})(\mathbf{1} \wedge B \wedge \bar{b} \cdot \mathbf{1}, \bar{b}) = \mathbf{t}'(\mathbf{c}, \beta). \end{aligned}$$

The second case is where $\mathbf{t} = \mathbf{y}\mathbf{r}\mathbf{z}$ for some $\mathbf{r} \in \mathbf{T}_{(+)}$. Then $\alpha = (\alpha', (A, \bar{a}), (B, \bar{b}))$ for some $(A, \bar{a}), (B, \bar{b}) \in \mathcal{X} \rtimes T$ and $\alpha' \in (\mathcal{X} \rtimes T)^*$. By Lemma 3.4 there exist $U, V \in \mathcal{X}$ and $g \in G$ such that $\mathbf{r}(\mathbf{c}, \alpha') = (U \wedge g \cdot C \wedge g\bar{c} \cdot V, \mathbf{1})$ for every $\mathbf{c} \in R$. In this case

$$\mathbf{t}(\mathbf{c}, \alpha) = (A, \bar{a})(U \wedge g \cdot C \wedge g\bar{c} \cdot V, \mathbf{1})(B, \bar{b}) = (A \wedge \bar{a} \cdot U \wedge \bar{a}g \cdot C \wedge \bar{a}g\bar{c} \cdot V \wedge \bar{a} \cdot B, \bar{a}\bar{b}).$$

By Lemma 3.5, there exist $\tilde{\mathbf{t}} \in \mathbf{T}_{(+)}$ and $\beta' \in R^*$ such that for all $\mathbf{c} \in R$ we have

$$(\mathbf{1} \wedge A \wedge \bar{a} \cdot U \wedge \bar{a}g \cdot C \wedge \bar{a}g\bar{c} \cdot V \wedge \bar{a} \cdot B, \mathbf{1}) = \tilde{\mathbf{t}}(\mathbf{c}, \beta').$$

So altogether

$$\begin{aligned} \mathbf{t}(\mathbf{c}, \alpha)^\downarrow &= (\mathbf{1} \wedge A \wedge \bar{a} \cdot U \wedge \bar{a}g \cdot C \wedge \bar{a}g\bar{c} \cdot V \wedge \bar{a} \cdot B \wedge \bar{a}\bar{b} \cdot \mathbf{1}, \bar{a}\bar{b}) \\ &= (\mathbf{1} \wedge A \wedge \bar{a} \cdot U \wedge \bar{a}g \cdot C \wedge \bar{a}g\bar{c} \cdot V \wedge \bar{a} \cdot B, \mathbf{1})(\mathbf{1} \wedge \bar{a}\bar{b} \cdot \mathbf{1}, \bar{a}\bar{b}) \\ &= \tilde{\mathbf{t}}(\mathbf{c}, \beta')(\mathbf{1} \wedge \bar{a}\bar{b} \cdot \mathbf{1}, \bar{a}\bar{b}), \end{aligned}$$

so $\mathbf{t}' = \mathbf{y}\tilde{\mathbf{t}}\mathbf{z}$ and $\beta = ((\mathbf{1}, \mathbf{1}), (\mathbf{1} \wedge \bar{a}\bar{b} \cdot \mathbf{1}, \bar{a}\bar{b}), \beta')$ satisfy the requirements of the lemma.

The third case, where $\mathbf{t} \in \mathbf{T}_{(*)}$, can be dealt with similarly to the previous case. \square

Lemma 3.7. *Let ρ be a $(2, 1, 1)$ -congruence on R and let $\rho^\#$ be the $(2, 1, 1)$ -congruence on $\mathcal{X} \rtimes T$ generated by ρ . Then the restriction of $\rho^\#$ to R equals ρ , i.e., $\rho^\#$ extends ρ . As a consequence, R/ρ is $(2, 1, 1)$ -embeddable in $(\mathcal{X} \rtimes T)/\rho^\#$.*

Proof. Let $(\mathbf{s}, \mathbf{t}) \in \rho^\# \cap (R \times R)$. Then there exists a ρ -sequence

$$\mathbf{s} = \mathbf{t}_1(\mathbf{c}_1, \alpha_1), \mathbf{t}_1(\mathbf{d}_1, \alpha_1) = \mathbf{t}_2(\mathbf{c}_2, \alpha_2), \dots, \mathbf{t}_n(\mathbf{d}_n, \alpha_n) = \mathbf{t}$$

connecting \mathbf{s} and \mathbf{t} in $\mathcal{X} \rtimes T$. By Lemma 3.6, for every k ($1 \leq k \leq n$), there exist $\mathbf{t}'_k \in \mathbf{T}$ and $\beta_k \in R^*$ satisfying (5). Thus for every k ($1 \leq k \leq n$), we have

$$\mathbf{t}_k(\mathbf{c}_k, \alpha_k)^\downarrow = \mathbf{t}'_k(\mathbf{c}_k, \beta_k), \mathbf{t}'_k(\mathbf{d}_k, \beta_k) = \mathbf{t}(\mathbf{d}_k, \alpha_k)^\downarrow,$$

showing that the sequence

$$\mathbf{s} = \mathbf{t}_1(\mathbf{c}_1, \alpha_1)^\downarrow = \mathbf{t}'_1(\mathbf{c}_1, \beta_1), \mathbf{t}'_1(\mathbf{d}_1, \beta_1) = \mathbf{t}'_2(\mathbf{c}_2, \beta_2), \dots, \mathbf{t}'_n(\mathbf{d}_n, \beta_n) = \mathbf{t}_n(\mathbf{d}_n, \alpha_n)^\downarrow = \mathbf{t}$$

connects \mathbf{s} and \mathbf{t} within R , proving that $(\mathbf{s}, \mathbf{t}) \in \rho$. \square

Although this lemma easily implies our embedding result referred in the title to, before formulating it, we show two lemmas in order to prepare our result on proper covers. Note that in the first lemma the action of T need not be nice.

Lemma 3.8. *Let $\rho \subseteq \sigma$ be a $(2, 1, 1)$ -congruence on the semidirect product $\mathcal{X} \rtimes T$. Then, for every $(A, \bar{a}), (B, \bar{b}) \in \mathcal{X} \rtimes T$, we have $(A, \bar{a}) \rho (B, \bar{b})$ if and only if $\bar{a} = \bar{b}$ and $(A, \mathbf{1}) \rho (B, \mathbf{1})$. As a consequence, ρ is generated by its restriction to the projections.*

Proof. Let $\tau = \{((A, \bar{a}), (B, \bar{b})) \in (\mathcal{X} \rtimes T) \times (\mathcal{X} \rtimes T) : \bar{a} = \bar{b} \text{ and } (A, 1) \rho (B, 1)\}$. It is clear that $\rho \subseteq \tau$. Conversely, let $((A, \bar{a}), (B, \bar{b})) \in \tau$. Then

$$(A \wedge B, \bar{a}) = (A, 1)(B, \bar{a}) \rho (B, 1)(B, \bar{a}) = (B, \bar{a}),$$

which together with its dual completes the proof. \square

Lemma 3.9. *Let ρ be a $(2, 1, 1)$ -congruence on R . Denote by ρ_P the restriction of ρ to $P \times P$, with $P = P(R)$, and by τ and $\tau^\#$ the $(2, 1, 1)$ -congruences on R and on $\mathcal{X} \rtimes T$, respectively, generated by ρ_P . Then R/τ $(2, 1, 1)$ -embeds into $(\mathcal{X}/\varepsilon) \rtimes T$ for an appropriate G -invariant congruence ε on \mathcal{X} , and $(\mathcal{X}/\varepsilon) \rtimes T$ is a nice semidirect product of \mathcal{X}/ε by T . Moreover, the $(2, 1, 1)$ -congruence ρ/τ on R/τ is projection separating, and the corresponding congruence on the image of R/τ in $(\mathcal{X}/\varepsilon) \rtimes T$ extends to a $(2, 1, 1)$ -congruence on $(\mathcal{X}/\varepsilon) \rtimes T$. Consequently, R/ρ is $(2, 1, 1)$ -embeddable in a $(2, 1, 1)$ -morphic image of $(\mathcal{X}/\varepsilon) \rtimes T$.*

Proof. Notice that $\tau \subseteq \sigma_R$ and $\tau^\# \subseteq \sigma_{\mathcal{X} \rtimes T}$. By definition, $\rho_P \subseteq \tau \subseteq \rho$, and as the restriction of ρ to P equals ρ_P the same is true for τ , thus the $(2, 1, 1)$ -congruence ρ/τ on R/τ is projection separating. Denote by $\rho^\#$ the $(2, 1, 1)$ -congruence on $\mathcal{X} \rtimes T$ generated by ρ . Since T acts nicely over G on \mathcal{X} , we see by Lemma 3.7 that $\rho^\#$ and $\tau^\#$ extend ρ and τ , respectively. Put $\varepsilon = \{(A, B) \in \mathcal{X} \times \mathcal{X} : (A, 1) \tau^\# (B, 1)\}$. Then ε is obviously a congruence on \mathcal{X} , and it is easily seen to be G -invariant. For, let $A, B \in \mathcal{X}$ with $A \varepsilon B$ and $\bar{a} \in T$. Then we obtain by Lemma 3.8 that $(A, \bar{a}) \tau^\# (B, \bar{a})$ which implies $(\bar{a}^{-1} \cdot A, 1) = (A, \bar{a})^* \tau^\# (B, \bar{a})^* = (\bar{a}^{-1} \cdot B, 1)$. Hence $\bar{a}^{-1} \cdot A \varepsilon \bar{a}^{-1} \cdot B$ follows. Moreover, we also see that $(\bar{a} \cdot A, 1) = (\bar{a} \cdot A, \bar{a})^+ = ((\bar{a} \cdot A, \bar{a})(A, 1))^+ \tau^\# ((\bar{a} \cdot A, \bar{a})(B, 1))^+ = (\bar{a}(A \wedge B), \bar{a})^+ = (\bar{a}(A \wedge B), 1)$, and so we deduce that $\bar{a} \cdot A \varepsilon \bar{a} \cdot (A \wedge B)$. Similarly, $\bar{a} \cdot B \varepsilon \bar{a} \cdot (A \wedge B)$ also holds, implying the relation $\bar{a} \cdot A \varepsilon \bar{a} \cdot B$. Since G is generated by T , one can see that $g \cdot A \varepsilon g \cdot B$ for every $g \in G$. Thus ε is, indeed, G -invariant, so the action of G on \mathcal{X} induces an action of G on \mathcal{X}/ε , and defines semidirect products $(\mathcal{X}/\varepsilon) \rtimes G$ and $(\mathcal{X}/\varepsilon) \rtimes T$. Lemma 3.3 implies that the latter semidirect product is nice.

By Lemma 3.8 we also have that $\iota: (\mathcal{X} \rtimes T)/\tau^\# \rightarrow (\mathcal{X}/\varepsilon) \rtimes T$, $(A, \bar{a})\tau^\# \mapsto (A\varepsilon, \bar{a})$ is an isomorphism. Let $\tilde{\rho}$ be the congruence on $(\mathcal{X}/\varepsilon) \rtimes T$ corresponding via ι to the congruence $\rho^\#/\tau^\#$ on $(\mathcal{X} \rtimes T)/\tau^\#$. The rest follows by basic universal algebra: since $\rho^\#$ and $\tau^\#$ extend ρ and τ , respectively, we obtain that $\rho^\#/\tau^\#$ extends ρ/τ , thus $(2, 1, 1)$ -embedding R/ρ , which is isomorphic to $(R/\tau)/(\rho/\tau)$, into $((\mathcal{X}/\varepsilon) \rtimes T)/\tilde{\rho}$. \square

Now we are ready to formulate the main result of the paper. It strengthens [12, Theorem 4.1] which establishes that every restriction semigroup is $(2, 1, 1)$ -embeddable in an almost left factorisable restriction semigroup, and [11, Theorem 4.11] which says that every restriction semigroup has a proper cover $(2, 1, 1)$ -embeddable in a W -product of a semilattice by a monoid.

Recall from Subsection 2.2 that a restriction semigroup (monoid) is almost factorisable (factorisable) if and only if it is a $(2, 1, 1)$ -morphic image of a semidirect product of a semilattice (semilattice with identity) by a monoid.

- Theorem 3.10.** (i) *Every restriction semigroup is $(2, 1, 1)$ -embeddable in an almost factorisable restriction semigroup, or, equivalently, in a factorisable restriction monoid.*
- (ii) *Every restriction semigroup has a proper cover which is $(2, 1, 1)$ -embeddable in a nice semidirect product of a semilattice by a monoid, and consequently, in a semidirect product of a semilattice by a group.*

Proof. (i) Let S be a restriction semigroup. By Lemma 2.2, it suffices to embed the restriction monoid S^1 into a $(2, 1, 1)$ -morphic image of a semidirect product of a semilattice by a monoid. Clearly, S^1 is isomorphic to $\text{FR}(\Omega)/\rho$ for some set Ω and $(2, 1, 1)$ -congruence ρ on $\text{FR}(\Omega)$. Here $\text{FR}(\Omega)$ is just the $(2, 1, 1)$ -subsemigroup R in (4) of the semidirect product $\mathcal{X} \rtimes \Omega^*$ given in Subsection 2.4. Example 3.1 shows that the action of Ω^* on \mathcal{X} is nice over $\text{FG}(\Omega)$ with respect to $\mathbf{1}$, and so Lemma 3.7 implies that $\text{FR}(\Omega)/\rho$ and its $(2, 1, 1)$ -isomorphic copy S^1 $(2, 1, 1)$ -embed into a $(2, 1, 1)$ -morphic image of $\mathcal{X} \rtimes \Omega^*$.

(ii) Applying Lemma 3.9 for $R = \text{FR}(\Omega)$ and the $(2, 1, 1)$ -congruence ρ in the previous paragraph, we obtain that $\text{FR}(\Omega)/\tau$ is $(2, 1, 1)$ -embeddable in a nice semidirect product $\mathcal{X}' \rtimes \Omega^*$ where \mathcal{X}' is a factor semilattice of \mathcal{X} over a $\text{FG}(\Omega)$ -invariant congruence. This implies that the semidirect product $\mathcal{X}' \rtimes \text{FG}(\Omega)$ is also defined, and $\mathcal{X}' \rtimes \Omega^*$ is a $(2, 1, 1)$ -subsemigroup in $\mathcal{X}' \rtimes \text{FG}(\Omega)$. The restriction monoid $\text{FR}(\Omega)/\tau$ is proper since the semidirect product $\mathcal{X}' \rtimes \Omega^*$ is. Moreover, Lemma 3.9 also establishes that ρ/τ is a projection separating $(2, 1, 1)$ -congruence on R/τ . This shows that $\text{FR}(\Omega)/\tau$ is a proper cover of $(\text{FR}(\Omega)/\tau)/(\rho/\tau)$, and so of $\text{FR}(\Omega)/\rho$ and S^1 . This completes the proof if $S = S^1$, that is, if S is a monoid. In the opposite case, it is routine to check that we can ensure the ρ -class of the identity element of $\text{FR}(\Omega)$ is a singleton, and the same property follows for τ . This implies that the restriction semigroup obtained from $\text{FR}(\Omega)/\tau$ by deleting its identity element is a proper cover of S with the required properties. \square

Note that, similarly to the embedding theorems in [12] and [8], the embedding given here is not a monoid embedding.

4. ADDITIONAL REMARKS

One can see from Lemmas 3.8 and 3.9 (see also the proof of Theorem 3.10(ii)) that if C is a restriction monoid which is a $(2, 1, 1)$ -factor monoid of a free restriction monoid over a $(2, 1, 1)$ -congruence contained in σ , then C is $(2, 1, 1)$ -embeddable in a nice semidirect product. Both Kudryavtseva [8] and Jones [7] have introduced a somewhat more general class of restriction monoids, called ultra F -restriction monoids in [8] and perfect restriction monoids in [7]. Moreover, if the greatest reduced factor of such a restriction monoid is a free monoid then it is proved to be $(2, 1, 1)$ -embeddable in a W -product in [8, Theorem 28]. In this section we strengthen this result by considering an even more general class of restriction monoids and showing that their members are $(2, 1, 1)$ -embeddable in a nice semidirect product in such a way that each of their $(2, 1, 1)$ -congruences extends to the semidirect product. Later on, we adopt the name ‘perfect’ from [7] for the class of restriction monoids mentioned, and we use the notation of [8].

Let T be a monoid and \mathcal{Y} a semilattice with identity element. Denote by $O_{\mathcal{Y}}$ the inverse submonoid of the symmetric inverse monoid $I(\mathcal{Y})$ consisting of all isomorphisms between the ideals of \mathcal{Y} . Notice that the Munn semigroup $T_{\mathcal{Y}}$ is an inverse submonoid in $O_{\mathcal{Y}}$. A *left partial action of T on \mathcal{Y}* is defined to be a dual monoid prehomomorphism $\alpha: T \rightarrow O_{\mathcal{Y}}^{\text{op}}, t \mapsto \alpha_t$, that is, α is required to have the properties that α_1 is the identity automorphism of \mathcal{Y} , and $\alpha_t \alpha_u$ is a restriction of α_{tu} for every $t, u \in T$. For any $t \in T$ and $A \in \mathcal{Y}$, we denote the element $\alpha_t A \in \mathcal{Y}$ by $t \diamond A$. Since $O_{\mathcal{Y}}$ is an inverse monoid, the mapping $\alpha': T \rightarrow O_{\mathcal{Y}}, t \mapsto \alpha_t^{-1}$ is a dual monoid prehomomorphism, and so α' defines a right partial action of T on \mathcal{Y} . For any $t \in T$ and $A \in \mathcal{Y}$, the element $A \alpha'_t \in \mathcal{Y}$ will be denoted by $A \circ t$. Clearly, the partial actions \diamond and \circ are reverse to each other, that is, for every $t \in T$ and $A \in \mathcal{Y}$, if $t \diamond A$ is defined then $(t \diamond A) \circ t$ is also defined and $(t \diamond A) \circ t = A$, and similarly, if $A \circ t$ is defined then $t \diamond (A \circ t)$ is also defined and $t \diamond (A \circ t) = A$.

Consider the set

$$M(T, \mathcal{Y}) = \{(A, t) \in \mathcal{Y} \times T : A \circ t \text{ is defined}\},$$

and define the following operations on it:

$$(A, t)(B, u) = (t \diamond ((A \circ t) \wedge B), tu), \quad (A, t)^+ = (A, 1) \quad \text{and} \quad (A, t)^* = (A \circ t, 1).$$

Then $M(T, \mathcal{Y})$ is a proper restriction monoid, and, conversely, each proper restriction monoid S is isomorphic to $M(S/\sigma, P(S))$ where the left partial action of S/σ on $P(S)$ is induced by S in a natural way, see [1].

A congruence ρ on a semigroup S is called *perfect* if for every $a, b \in S$ we have $(a\rho)(b\rho) = (ab)\rho$, where the left hand side of the equality is the set product of the classes $a\rho$ and $b\rho$. A restriction monoid S is called *perfect* if σ is a perfect congruence and each σ -class has a greatest element with respect to the natural partial order. It follows ([7, Theorem 1.1]) that a restriction monoid is perfect if and only if it is isomorphic to a restriction monoid $M(T, \mathcal{Y})$ where the left partial action of T on \mathcal{Y} is a monoid homomorphism from T into $T_{\mathcal{Y}}^{\text{op}}$.

Remark 4.1. (i) Note that [8] uses the term ‘left partial action’ in a more general sense. However, in a semilattice, the order ideals (resp. principal order ideals) and the ideals (resp. principal ideals) coincide. Moreover, the order isomorphisms and the isomorphisms between the ideals of a semilattice coincide. Therefore a left partial action of a monoid T on a semilattice \mathcal{Y} defined above is just what is called in [8] a ‘left partial action of T on \mathcal{Y} such that axioms (A),(B),(C) hold’.

(ii) The restriction monoid denoted by $W(T, \mathcal{Y})$ in [7] is the left-right dual of $M(T, \mathcal{Y})$ defined above, and so it is a much more general construction than what is usually meant by the notation $W(T, \mathcal{Y})$ (see [2], [4], [8], [11], [12]), in particular in this paper, called a W -product of a semilattice by a monoid and denoted $W(T, \mathcal{Y})$.

Suppose now that T is a monoid embeddable in a group, \mathcal{X} is a semilattice, and T acts on \mathcal{X} by automorphisms. As above, let us choose and fix a group G such that T is a submonoid in G generating G , G acts on \mathcal{X} , and the action of T on \mathcal{X} is just the restriction of the action of G on \mathcal{X} . Moreover, consider a principal ideal \mathcal{Y} of \mathcal{X} , and denote its greatest element by

1. It is routine to check that the restriction of the action of G on \mathcal{X} to \mathcal{Y} is a left partial action $\alpha: G \rightarrow T_{\mathcal{Y}}^{\text{op}}$ where the domain and range of α_g ($g \in G$) are the principal ideals of \mathcal{Y} with greatest elements $g^{-1} \cdot \mathbf{1} \wedge \mathbf{1}$ and $g \cdot \mathbf{1} \wedge \mathbf{1}$, respectively. Therefore the restriction monoids $M(G, \mathcal{Y})$ and $M(T, \mathcal{Y})$ are defined, and we have $M(T, \mathcal{Y}) \leq M(G, \mathcal{Y})$. Taking into account that all elements of \mathcal{X} appearing in $M(G, \mathcal{Y})$ belong to the subsemilattice $G \cdot \mathcal{Y}$, we can suppose without loss of generality that $\mathcal{X} = G \cdot \mathcal{Y}$. In this case, $(G, \mathcal{X}, \mathcal{Y})$ is a McAlister triple, and it is easy to see that $M(G, \mathcal{Y}) = P(G, \mathcal{X}, \mathcal{Y})$, the P -semigroup defined by it. This implies that $M(G, \mathcal{Y})$ is an F -inverse monoid, and it is an inverse subsemigroup in the semidirect product $\mathcal{X} \rtimes G$. Furthermore, it also follows that $M(T, \mathcal{Y})$ is equal to the $(2, 1, 1)$ -subsemigroup R of the semidirect product $\mathcal{X} \rtimes T$ defined in (4). Thus Lemmas 3.7 and 3.9 imply the following statement.

Proposition 4.2. *Suppose that a left partial action of a monoid T on a semilattice \mathcal{Y} with identity $\mathbf{1}$ can be extended to an action \cdot of T on a semilattice \mathcal{X} containing \mathcal{Y} as a principal ideal such that the action \cdot is nice over a group G with respect to $\mathbf{1}$, and we have $\mathcal{X} = G \cdot \mathcal{Y}$. Then the following hold:*

- (i) $M(T, \mathcal{Y}) \leq \mathcal{X} \rtimes T$, $M(G, \mathcal{Y}) \leq \mathcal{X} \rtimes G$ where $\mathcal{X} \rtimes T$ is a nice semidirect product and $M(G, \mathcal{Y})$ is an F -inverse monoid;
- (ii) every $(2, 1, 1)$ -congruence ρ on $M(T, \mathcal{Y})$ extends to $\mathcal{X} \rtimes T$;
- (iii) for every $(2, 1, 1)$ -congruence ρ of $M(T, \mathcal{Y})$ with $\rho \subseteq \sigma_{M(T, \mathcal{Y})}$, the restriction monoid $M(T, \mathcal{Y})/\rho$ embeds into a nice semidirect product of a factor semilattice of \mathcal{X} by T .

Since any proper restriction monoid S is $(2, 1, 1)$ -isomorphic to the monoid $M(S/\sigma, P(S))$ defined by means of the induced left partial action, we have the following consequence.

Corollary 4.3. *Let S be a proper restriction monoid such that the induced left partial action of S/σ on $P(S)$ can be extended to an action \cdot of S/σ on a semilattice \mathcal{X} containing $P(S)$ as a principal ideal in such a way that the action \cdot is nice over a group G with respect to the identity element of $P(S)$ and $\mathcal{X} = G \cdot P(S)$. Then each $(2, 1, 1)$ -factor monoid of S is embeddable in a $(2, 1, 1)$ -factor semigroup of the nice semidirect product $\mathcal{X} \rtimes (S/\sigma)$, which is a $(2, 1, 1)$ -subsemigroup in the inverse semigroup $\mathcal{X} \rtimes G$. Moreover, for every $(2, 1, 1)$ -congruence ρ of S with $\rho \subseteq \sigma$, the $(2, 1, 1)$ -factor monoid S/ρ embeds into a nice semidirect product of a factor semilattice of \mathcal{X} by S/σ .*

Now we establish that the perfect restriction monoids $M(\Omega^*, \mathcal{Y})$ studied in [8] and [7] satisfy the assumptions of Proposition 4.2 and Corollary 4.3.

Let Ω be a set, \mathcal{Y} a semilattice with identity element $\mathbf{1}$, and consider a left partial action of the free monoid Ω^* on \mathcal{Y} which is a monoid homomorphism $\alpha: \Omega^* \rightarrow T_{\mathcal{Y}}^{\text{op}}$. This defines a perfect restriction monoid $S = M(\Omega^*, \mathcal{Y})$. The free monoid Ω^* is a submonoid of the free group $\text{FG}(\Omega)$ such that $\text{FG}(\Omega)$ is generated by Ω^* , and each element of $\text{FG}(\Omega)$ can be uniquely written as a product of members of an alternating sequence, see Example 3.1. This allows us to extend α to a left partial action, also denoted by α , of $\text{FG}(\Omega)$ on \mathcal{Y} by putting $\alpha_{t^{-1}} = \alpha'_t$ for every $t \in \Omega^*$, and then by defining $\alpha_{w_1 w_2 \dots w_n} = \alpha_{w_1} \alpha_{w_2} \dots \alpha_{w_n}$ in $T_{\mathcal{Y}}^{\text{op}}$ for every alternating sequence w_1, w_2, \dots, w_n in Ω^* . Notice that the left partial action $\alpha: \text{FG}(\Omega) \rightarrow T_{\mathcal{Y}}^{\text{op}}$ is no longer a monoid homomorphism. However, if $g, h \in \text{FG}(\Omega)$ such

that g, h and gh (the concatenation of g and h) are reduced words then $\alpha_g \alpha_h = \alpha_{gh}$. For every $g \in \text{FG}(\Omega)$ denote the identity element of the range of α_g by M_g . By definition, we clearly have $M_1 = \mathbf{1}$ and, since $M_{g^{-1}}$ equals the identity element of the domain of α_g , we have

$$(6) \quad M_g = g \diamond M_{g^{-1}} \quad \text{for every } g \in \text{FG}(\Omega).$$

The restriction monoid $M(\text{FG}(\Omega), \mathcal{Y})$ is easily seen to be an F -inverse monoid, where the greatest element in the σ -class corresponding to $g \in \text{FG}(\Omega)$ is (M_g, g) . Hence $M(\text{FG}(\Omega), \mathcal{Y})$ can be represented as a P -semigroup $P(\text{FG}(\Omega), \mathcal{Y}, \mathcal{X})$ where \mathcal{X} is a semilattice and is, up to isomorphism, uniquely determined. Furthermore, by [10], \mathcal{X} can be constructed in the following manner. The relation χ on the set $\mathcal{Y} \times \text{FG}(\Omega)$, defined by $(A, g) \chi (B, h)$ if and only if $B = A \circ g^{-1}h$, is a preorder, and so it induces a partial order \leq on the set \mathcal{X} of all $\bar{\chi}$ -classes where $\bar{\chi}$ is the join of χ and its converse. Denoting the $\bar{\chi}$ -class of the element $(A, g) \in \mathcal{Y} \times \text{FG}(\Omega)$ by $[A, g]$, we can deduce from the proof of [9, Theorem 7.4.5] that

$$(7) \quad [A, g] \wedge [B, h] = [g^{-1}h \diamond ((A \wedge M_{g^{-1}h}) \circ g^{-1}h \wedge B), g]$$

for every element $[A, g], [B, h]$ in the semilattice \mathcal{X} . Hence $\mathcal{X}_0 = \{[A, 1] : A \in \mathcal{Y}\}$ is a principal ideal in \mathcal{X} , and the mapping $\mathcal{Y} \rightarrow \mathcal{X}_0$, $A \mapsto [A, 1]$ is an isomorphism. By identifying \mathcal{X}_0 with \mathcal{Y} along this isomorphism, \mathcal{Y} becomes a principal ideal of \mathcal{X} . The rule $h \cdot [A, g] = [A, hg]$ ($h \in \text{FG}(\Omega)$, $[A, g] \in \mathcal{X}$) defines an action of the group $\text{FG}(\Omega)$ on \mathcal{X} such that $\mathcal{X} = \text{FG}(\Omega) \cdot \mathcal{Y}$, and the restriction of this action to \mathcal{Y} is obviously the left partial action \diamond .

On the other hand, the action \cdot of $\text{FG}(\Omega)$ on \mathcal{X} naturally restricts to an action, also denoted by \cdot , of Ω^* on \mathcal{X} . Now we establish that this action of Ω^* on \mathcal{X} is nice over $\text{FG}(\Omega)$ with respect to the element $\mathbf{1}$. It suffices to show that if $g, h \in \text{FG}(\Omega)$ are reduced words such that there is no non-empty common prefix of g and h then the inequality $\mathbf{1} \geq g \cdot \mathbf{1} \wedge h \cdot \mathbf{1}$ holds in \mathcal{X} , or equivalently, $g \cdot \mathbf{1} \wedge h \cdot \mathbf{1} \in \mathcal{Y}$. Applying (7), (6) and that $M_g \in \mathcal{Y}$ for every $g \in \text{FG}(\Omega)$, we obtain that

$$\begin{aligned} [\mathbf{1}, g] \wedge [\mathbf{1}, h] &= [g^{-1}h \diamond ((\mathbf{1} \wedge M_{g^{-1}h}) \circ g^{-1}h \wedge \mathbf{1}), g] \\ &= [g^{-1}h \diamond (M_{g^{-1}h} \circ g^{-1}h), g] = [M_{g^{-1}h}, g] = [g^{-1}h \diamond M_{h^{-1}g}, g]. \end{aligned}$$

Since $g^{-1}h$ is a reduced word by assumption, we have $g^{-1}h \diamond M_{h^{-1}g} = \alpha_{g^{-1}h} M_{h^{-1}g} = \alpha_{g^{-1}}(\alpha_h M_{h^{-1}g})$, whence it follows that both $\alpha_h M_{h^{-1}g} = h \diamond M_{h^{-1}g}$ and $\alpha_{g^{-1}}(\alpha_h M_{h^{-1}g}) = (h \diamond M_{h^{-1}g}) \circ g$ are defined. This implies by the definition of the preorder χ that

$$[\mathbf{1}, g] \wedge [\mathbf{1}, h] = [(h \diamond M_{h^{-1}g}) \circ g, g] = [h \diamond M_{h^{-1}g}, 1] \in \mathcal{Y},$$

thus completing the proof that $g \cdot \mathbf{1} \wedge h \cdot \mathbf{1} \in \mathcal{Y}$.

Summarising, we have strengthened the results [8, Theorem 28] and [7, Proposition 8.4] in the following way:

Corollary 4.4. *Let S be a perfect restriction monoid such that S/σ is a free monoid on a set Ω . Then the induced left partial action of S/σ on $P(S)$ can be extended to an action \cdot of S/σ on a semilattice \mathcal{X} containing $P(S)$ as a principal ideal such that the action \cdot is nice over*

the free group $\text{FG}(\Omega)$ with respect to the identity element of $P(S)$ and $\mathcal{X} = \text{FG}(\Omega) \cdot P(S)$. Consequently, S is $(2, 1, 1)$ -embeddable in the nice semidirect product $\mathcal{X} \rtimes \Omega^*$, which is a $(2, 1, 1)$ -subsemigroup in the inverse semigroup $\mathcal{X} \rtimes \text{FG}(\Omega)$, and the following are valid:

- (i) each $(2, 1, 1)$ -congruence of S extends to $\mathcal{X} \rtimes \Omega^*$ (along the $(2, 1, 1)$ -embedding), and
- (ii) for every $(2, 1, 1)$ -congruence ρ of S with $\rho \subseteq \sigma$, the restriction monoid S/ρ embeds into a nice semidirect product of a factor semilattice of \mathcal{X} by Ω^* .

REFERENCES

- [1] C. Cornock, V. Gould, Proper restriction semigroups and partial actions, *J. Pure Appl. Algebra* **216** (2012) 936–949.
- [2] J. Fountain, G. M. S. Gomes, Proper left type- A monoids revisited, *Glasgow Math. J.* **35** (1993) 293–306.
- [3] J. Fountain, G. M. S. Gomes, V. Gould, The free ample monoid, *Internat. J. Algebra Comput.* **19** (2009) 527–554.
- [4] G. M. S. Gomes, M. B. Szendrei, Almost factorizable weakly ample semigroups, *Comm. Algebra* **35** (2007) 3503–3523.
- [5] V. Gould, V. Notes on restriction semigroups and related structures; formerly (Weakly) left E -ample semigroups <http://www-users.york.ac.uk/~varg1/gpubs.htm>
- [6] V. Gould, M. Kambites, Faithful functors from cancellative categories to cancellative monoids with an application to abundant semigroups, *Internat. J. Algebra Comput.* **15** (2005) 683–698.
- [7] P. Jones, Almost perfect restriction semigroups, *J. Algebra* **445** (2016) 193–220.
- [8] G. Kudryavtseva, Partial monoid actions and a class of restriction semigroups, *J. Algebra* **429** (2015) 342–370.
- [9] M. V. Lawson, *Inverse semigroups: The Theory of Partial Symmetries*. World Scientific, Singapore, 1998.
- [10] W. D. Munn, A note on E -unitary inverse semigroups, *Bull. London Math. Soc.* **8** (1976) 71–76.
- [11] M. B. Szendrei, Proper covers of restriction semigroups and W -products, *Internat. J. Algebra Comput.* **22** (2012) 1250024, 16 pp.
- [12] M. B. Szendrei, Embedding into almost left factorizable restriction semigroups, *Comm. Algebra* **41** (2013) 1458–1483.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK, YO10 5DD, UK
E-mail address: victoria.gould@york.ac.uk

BOLYAI INSTITUTE, UNIVERSITY OF SZEGED, ARADI VÉRTANÚK TERE 1, SZEGED, HUNGARY, H-6720
E-mail address: hartm@math.u-szeged.hu

BOLYAI INSTITUTE, UNIVERSITY OF SZEGED, ARADI VÉRTANÚK TERE 1, SZEGED, HUNGARY, H-6720
E-mail address: m.szendrei@math.u-szeged.hu