



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/110878/>

Version: Accepted Version

Article:

Hickson, N., Hampshire, D., Castaman, G. et al. (2011) Effect of the VWF promoter (GT)_n repeat and single-nucleotide polymorphism c.-2527G>A on circulating von Willebrand factor levels under normal conditions. *Journal of Thrombosis and Haemostasis*, 9 (3). pp. 603-605. ISSN: 1538-7933

<https://doi.org/10.1111/j.1538-7836.2010.04161.x>

This is the peer reviewed version of the following article: HICKSON, N., HAMPSHIRE, D., CASTAMAN, G., EIKENBOOM, J., RODEGHIERO, F., PEAKE, I., GOODEVE, A. and ON BEHALF OF THE MCMDM-1VWD AND ZPMCB-VWD STUDY GROUPS (2011), Effect of the VWF promoter (GT)_n repeat and single-nucleotide polymorphism c.-2527G>A on circulating von Willebrand factor levels under normal conditions. *Journal of Thrombosis and Haemostasis*, 9: 603–605., which has been published in final form at <https://doi.org/10.1111/j.1538-7836.2010.04161.x>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Published in final edited form as:

J Thromb Haemost. 2011 March ; 9(3): 603–605. doi:10.1111/j.1538-7836.2010.04161.x.

Effect of the *VWF* promoter (GT)_n repeat and SNP c.-2527G>A on circulating *VWF* levels under normal conditions

N HICKSON^{*}, D HAMPSHIRE^{*}, G CASTAMAN[†], J EIKENBOOM[‡], F RODEGHIERO[†], I PEAKE^{*}, and A GOODEVE^{*} ON BEHALF OF THE MCMDM-1VWD AND ZPMCB-VWD STUDY GROUPS

^{*} Haemostasis Research Group, Department of Cardiovascular Science, University of Sheffield, UK [†] Haematology Department, San Bortolo Hospital, Vicenza, Italy [‡] Einthoven Laboratory for Experimental Vascular Medicine, Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden, The Netherlands

Keywords

GT repeat; haplotype; promoter sequence variation; SNP; von Willebrand factor level

Circulating von Willebrand factor (*VWF*) protein levels in plasma (*VWF*:Ag) of healthy individuals can be affected by a number of factors, either physiological and environmental such as stress, or genetic such as ABO blood group [1]. It has previously been suggested that *VWF*:Ag levels could be partly determined by four single nucleotide polymorphisms (SNP) in the promoter region of *VWF*; c.-3268C>G, c.-2709T>C, c.-2661G>A and c.-2527G>A, segregating predominantly as two haplotypes, GCAG (haplotype 1) or CTGA (haplotype 2), with the latter being associated with lower *VWF*:Ag levels [2,3]. Shear stress has been shown to enhance *VWF* promoter activity, with a dinucleotide (GT)_n short tandem repeat (STR) located at c.-2144_-2105 in the *VWF* promoter exhibiting size-dependent mediation of this stress-induced transactivation [4]. Studies on Cushing's syndrome patients suggest that both the SNPs and STR influence a glucocorticoid-induced increase in plasma *VWF*:Ag level [5,6]. However, other studies have failed to reproduce this SNP and STR association [7,8]. The current study reports the largest investigation of the association between SNP c.-2527G>A, the (GT)_n STR, and *VWF*:Ag levels in healthy controls (HC) from the MCMDM-1VWD study [9] under normal conditions, while controlling for the effect of ABO blood group.

ABO blood group, age and *VWF*:Ag levels were available for 1115 European male and female healthy controls (HC) from nine countries [9]. None had been referred to a specialist centre due to bleeding problems. *VWF*:Ag levels were determined at a single centre [9]. Genotypes for the c.-2527G>A SNP were determined using TaqMan allelic discrimination (ABI 7900; Applied Biosystems, Europe BV, Warrington, UK). The *VWF* region containing the (GT)_n STR was amplified by PCR (primer sequences available on request), sequenced and analysed on an ABI PRISM™ 3730 DNA analyser (Applied Biosystems). Analysis of two homozygous samples showed that (GT)_n alleles 181bp and 185bp long corresponded to (GT)₁₉ and (GT)₂₁ respectively. Hardy-Weinberg equilibrium, allele frequencies and

linkage disequilibrium analyses were performed using SNPStats: <http://bioinfo.iconcologia.net/index.php?module=Snpstats> (accessed August 2010). Associations between genotypes and VWF:Ag levels were assessed using Kruskal Wallis tests (GraphPad Prism v5.00, GraphPad Software Inc, San Diego, CA, USA). *p* values <0.05 were considered to be statistically significant in all tests.

Genotype data for the c.-2527G>A SNP were used for investigation of a genotype-VWF:Ag relationship. There was no significant association between VWF:Ag level and SNP genotype before or after controlling for the effect of ABO blood group (Table 1A), thus contradicting previous results of a smaller study on blood group O Canadian HC (n=250) [3] which identified a significant association (*p*=0.006). Presence of an extended founder haplotype in the Canadian population with an effect on VWF level could be a possible explanation. 100 European HC from two countries were also genotyped for the four promoter SNP examined in the Canadian study and the SNP were shown to be in complete linkage disequilibrium (*D'*=0.999, *p*<0.0001), segregating as the two previously reported haplotypes GCAG (haplotype 1) and CTGA (haplotype 2). No association of genotype with VWF:Ag levels was found in this subgroup (data not shown).

Genotyping of the STR identified alleles ranging in size from (GT)₁₆ to (GT)₂₇ with a bimodal distribution, i.e. a peak at (GT)₁₉ (37.2%) and another at (GT)₂₁ (32.2%). The genotype (GT)₁₉/(GT)₂₁ was the most frequent (23.8%), followed by (GT)₁₉/(GT)₁₉ (13.3%) and (GT)₂₁/(GT)₂₁ (10.2%). GT repeat alleles were divided into two subclasses for ease of analysis [8]. Small (S) alleles included all those <(GT)₂₀ and large (L) included those ≥(GT)₂₀. Based on this approach, 42.8% of alleles were (GT)_S and 57.2% were (GT)_L; 18% HC had genotype S/S, 50% had S/L and 32% had L/L, while genotype distribution was consistent with Hardy-Weinberg equilibrium (*p*=0.42).

In contrast to the results of a smaller study (n=394 HC) reporting a lack of association between (GT)_n genotypes and VWF:Ag levels in HC [8], in this study VWF:Ag values were lower in HC homozygous for S alleles compared to HC heterozygous or homozygous for L alleles (S/S vs S/L vs L/L = 92.0 vs 97.0 vs 100.0 IU/dL, *p*=0.027) (Table 1B). This association was not maintained when HC of only blood group O were examined (S/S vs S/L vs L/L = 80.0 vs 80.0 vs 82.0 IU/dL, *p*=0.524), but was observed when examining HC of non-O blood groups (S/S vs S/L vs L/L = 101.0 vs 107.5 vs 114.0 IU/dL, *p*=0.020) (Table 1B). These data suggest an effect of GT repeat length on VWF levels of ~10% and are in agreement with recent findings suggesting that longer GT repeats induce a larger increase in VWF promoter activity than shorter ones under shear stress conditions [4]. A statistically significant association amongst O blood group HC may only be observed with analysis of a larger number of individuals in this subcategory.

A small degree of linkage was observed between the c.-2527G>A SNP and the (GT)_n repeat (*D'*=0.661, *R*²=0.597, *p*<0.0001). The two most frequent haplotypes (S/G and L/A) accounted for ~80% of HC (49.9% and 30.5% respectively). The most frequent genotypes resulting from homozygosity or heterozygosity for the above haplotypes were S/L-G/A (32.8%), L/L-A/A (23.6%) and S/S-G/G (10.1%). Other allele combinations comprised the remaining 33.5% of genotypes. The association between the three genotypes (S/L-G/A, L/L-A/A, S/S-G/G) and VWF:Ag levels was examined in order to identify a potential cooperative effect of the STR and SNP loci on VWF:Ag levels. The results shown in Table 1C suggest that these two polymorphic regions have no significant cooperative effect on VWF:Ag levels in the EU population, before or after controlling for the effect of ABO blood group on VWF:Ag levels (Table 1C). This suggests that an effect on VWF:Ag levels is mediated only by the (GT)_n repeat.

In conclusion, this study has provided evidence that the c.-2527G>A VWF promoter SNP does not have a significant effect on levels of VWF in the EU population. In contrast, the VWF (GT)_n STR exhibited a small effect on VWF levels, mediated by its length. A SNP/STR cooperative effect on the regulation of VWF:Ag levels in HC was excluded. This is the largest study examining the potential influence of these loci on VWF levels in HC while controlling for the effect of ABO blood group and is the first study to show a significant effect of the c.-2144_-2105 locus on VWF levels under normal conditions.

Acknowledgments

This work was supported by the European Community Fifth Framework Programme (QLG1-CT-2000-00387) and National Institutes of Health Zimmerman Program for Molecular and Clinical Biology of VWD (HL-081588). The authors thank all contributors to the MCMDM-1VWD study and John Anson, Sheffield University for excellent technical support.

References

1. Gill JC, Endres-Brooks J, Bauer PJ, Marks WJ Jr, Montgomery RR. The effect of ABO blood group on the diagnosis of von Willebrand disease. *Blood*. 1987; 69:1691–5. [PubMed: 3495304]
2. Keightley AM, Lam YM, Brady JN, Cameron CL, Lillicrap D. Variation at the von Willebrand factor (vWF) gene locus is associated with plasma vWF:Ag levels: identification of three novel single nucleotide polymorphisms in the vWF gene promoter. *Blood*. 1999; 93:4277–83. [PubMed: 10361125]
3. Harvey PJ, Keightley AM, Lam YM, Cameron C, Lillicrap D. A single nucleotide polymorphism at nucleotide -1793 in the von Willebrand factor (VWF) regulatory region is associated with plasma VWF:Ag levels. *Br J Haematol*. 2000; 109:349–53. [PubMed: 10848823]
4. Hough C, Cameron CL, Notley CR, Brown C, O'Brien L, Keightley AM, Berber E, Lillicrap D. Influence of a GT repeat element on shear stress responsiveness of the VWF gene promoter. *J Thromb Haemost*. 2008; 6:1183–90. [PubMed: 18485092]
5. Casonato A, Daidone V, Sartorello F, Albiger N, Romualdi C, Mantero F, Pagnan A, Scaroni C. Polymorphisms in von Willebrand factor gene promoter influence the glucocorticoid-induced increase in von Willebrand factor: the lesson learned from Cushing syndrome. *Br J Haematol*. 2008; 140:230–5. [PubMed: 18173757]
6. Daidone V, Pontara E, Romualdi C, Cattini MG, Scaroni C, Albiger N, Pagnan A, Casonato A. Microsatellite (GT)(n) is part of the von Willebrand factor (VWF) promoter region that influences the glucocorticoid-induced increase in VWF in Cushing's syndrome. *Thromb Res*. 2010; 125:e275–80. [PubMed: 20156642]
7. Lethagen S, Hillarp A, Ekholm C, Mattson E, Hallden C, Friberg B. Distribution of von Willebrand factor levels in young women with and without bleeding symptoms: influence of ABO blood group and promoter haplotypes. *Thromb Haemost*. 2008; 99:1013–8. [PubMed: 18521502]
8. Daidone V, Cattini MG, Pontara E, Sartorello F, Gallinaro L, Marotti A, Scaroni C, Pagnan A, Casonato A. Microsatellite (GT)(n) repeats and SNPs in the von Willebrand factor gene promoter do not influence circulating von Willebrand factor levels under normal conditions. *Thromb Haemost*. 2009; 101:298–304. [PubMed: 19190813]
9. Goodeve A, Eikenboom J, Castaman G, Rodeghiero F, Federici AB, Battle J, Meyer D, Mazurier C, Goudemand J, Schneppenheim R, Budde U, Ingerslev J, Habart D, Vorlova Z, Holmberg L, Lethagen S, Pasi J, Hill F, Hashemi Soteh M, Baronciani L, et al. Phenotype and genotype of a cohort of families historically diagnosed with type 1 von Willebrand disease in the European study, Molecular and Clinical Markers for the Diagnosis and Management of Type 1 von Willebrand Disease (MCMDM-1VWD). *Blood*. 2007; 109:112–21. [PubMed: 16985174]

Table 1

Association of genotype at *VWF* locations (A) c.-2527, B) (GT)_n and C) (GT)_n plus c.-2527 with median levels of VWF in healthy controls from the MCMDM-1VWD study controlling for blood group.

Analysis	Genotype and ABO blood group	Mean age (SD)	n*	Median VWF:Ag level (IU/dL) (Range [†])	p value
A) c.-2527G>A	All [‡]		1115		
	G/G	40.7 (12.5)	162	96.5 (74.8–119.5)	0.961
	G/A	40.2 (11.9)	521	98.0 (77.0–120.0)	
	A/A	40.5 (11.4)	432	96.0 (75.0–122.0)	
	O		410		0.747
	G/G	41.6 (11.3)	51	84.0 (66.0–106.0)	
	G/A	42.1 (12.5)	191	80.0 (64.0–98.0)	
	A/A	41.2 (10.6)	168	78.5 (66.0–99.0)	0.852
	Non-O		678		
	G/G	40.8 (13.1)	104	106.0 (81.0–133.0)	
G/A	39.3 (11.4)	319	107.0 (89.0–126.0)		
A/A	40.2 (12.0)	255	112.0 (86.0–129.0)		
B) (GT) _n	All		1105		
	S/S	41.0 (11.5)	196	92.0 (70.3–114.5)	0.027
	S/L	40.3 (11.8)	555	97.0 (76.0–120.0)	
	L/L	40.6 (12.3)	354	100.0 (78.0–125.0)	
	O		408		0.524
	S/S	41.8 (11.0)	71	80.0 (62.0–99.0)	
	S/L	42.0 (11.7)	199	80.0 (65.0–102.0)	
	L/L	40.9 (11.6)	138	82.0 (67.8–101.3)	0.020
	Non-O		569		
	S/S	41.1 (11.7)	118	101.0 (77.8–124.3)	
S/L	39.3 (11.8)	344	107.5 (86.0–127.0)		
L/L	40.7 (12.8)	207	114.0 (91.0–132.0)		
C) c.-2527G>A & (GT) _n	All		727		
	S/S-G/G	41.3 (12.0)	110	92.0 (71.8–116.3)	0.314

Analysis	Genotype and ABO blood group	Mean age (SD)	n*	Median VWF:Ag level (IU/dL) (Range [†])	p value
	S/L-G/A	39.7 (11.7)	358	96.5 (77.8–120.0)	0.714
	L/L-A/A	40.7 (11.9)	259	98.0 (76.0–125.0)	
	O		278		
	S/S-G/G	41.5 (10.6)	39	84.0 (66.0–101.0)	
	S/L-G/A	41.8 (12.0)	134	81.5 (65.8–104.0)	
	L/L-A/A	40.8 (10.7)	105	85.0 (67.0–103.0)	
	Non-O		432		0.387
	S/S-G/G	42.1 (12.5)	65	102.0 (76.5–134.5)	
	S/L-G/A	38.6 (11.4)	219	107.0 (87.0–126.0)	
	L/L-A/A	40.8 (12.9)	148	112.5 (88.0–130.0)	

* ABO blood group data were available for 1088 out of 1115 HC.

[†]Range = 25th–75th percentile.

[‡]All: all blood groups; O: blood group O only; Non-O: blood groups other than O.

SD: Standard deviation