



This is a repository copy of *Identification of new members of the Escherichia coli K-12 MG1655 SlyA regulon.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/110899/>

Version: Accepted Version

---

**Article:**

Curran, T., Abacha, F., Hibberd, S. et al. (3 more authors) (2017) Identification of new members of the Escherichia coli K-12 MG1655 SlyA regulon. *Microbiology*, 163 (3). pp. 400-409. ISSN 1350-0872

<https://doi.org/10.1099/mic.0.000423>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Microbiology

## Identification of new members of the Escherichia coli K-12 MG1655 SlyA regulon --Manuscript Draft--

<b>Manuscript Number:</b>	MIC-D-16-00406R1
<b>Full Title:</b>	Identification of new members of the Escherichia coli K-12 MG1655 SlyA regulon
<b>Article Type:</b>	Standard
<b>Section/Category:</b>	Regulation
<b>Corresponding Author:</b>	Jeffrey Green University of Sheffield Sheffield, UNITED KINGDOM
<b>First Author:</b>	Thomas Curran
<b>Order of Authors:</b>	Thomas Curran Fatima Abacha Stephen Hibberd Matthew Rolfe Melissa Lacey Jeffrey Green
<b>Abstract:</b>	<p>SlyA is a member of the MarR family of bacterial transcriptional regulators. Previously, SlyA has been shown to directly regulate only two operons in Escherichia coli K-12 MG1655, fimB and hlyE (clyA). In both cases SlyA activates gene expression by antagonizing repression by the nucleoid associated protein H-NS. Here the transcript profiles of aerobic glucose-limited steady-state chemostat cultures of E. coli K-12 MG1655, slyA mutant and slyA over-expression strains are reported. The transcript profile of the slyA mutant was not significantly different to that of the parent; however, that of the slyA expression strain was significantly different from that of the vector control. Transcripts representing 27 operons were increased in abundance, whereas 3 were decreased. Of the 30 differentially regulated operons, 24 have been previously associated with sites of H-NS binding, suggesting that antagonism of H-NS repression is a common feature of SlyA-mediated transcription regulation. Direct binding of SlyA to DNA located upstream of a selection of these targets permitted the identification of new operons likely to be directly regulated by SlyA. Transcripts of four operons coding for cryptic adhesins exhibited enhanced expression and this was consistent with enhanced biofilm formation associated with the SlyA over-producing strain.</p>

1 **Identification of new members of the *Escherichia coli* K-12**  
2 **MG1655 SlyA regulon**

3  
4 Thomas D. Curran<sup>1</sup>, Fatima Abacha<sup>2</sup>, Stephen P. Hibberd<sup>1</sup>, Matthew D. Rolfe<sup>1</sup>, Melissa M.  
5 Lacey<sup>2</sup> and Jeffrey Green<sup>1</sup>

6  
7 <sup>1</sup>Department of Molecular Biology and Biotechnology, University of Sheffield, Sheffield, S10  
8 2TN, UK

9 <sup>2</sup>Biomolecular Research Centre, Sheffield Hallam University, Sheffield, S1 1WB, UK

10

11 Key words: biofilm, gene expression, MarR family, transcription regulation

12

13 Abbreviations used: A<sub>600</sub>; absorbance at 600 nm; Amp<sup>R</sup>, ampicillin resistance; Btn, biotin;  
14 Cm<sup>R</sup>, chloramphenicol resistance; CRP, cyclic-AMP receptor protein; EMSA, electrophoretic  
15 mobility shift assay; Kan<sup>R</sup>, kanamycin resistance; OD<sub>600</sub>; optical density at 600 nm; Tet<sup>R</sup>,  
16 tetracycline resistance; T<sup>s</sup>, temperature sensitive

17

18 Running title: The SlyA regulon of *E. coli* K-12 MG1655

19

20 Subject category: Regulation

21

22 **Correspondence**

23 Melissa Lacey, Biomolecular Research Centre, Sheffield Hallam University, Sheffield, S1  
24 1WB, UK

25 Email: m.lacey@shu.ac.uk

26 Matthew Rolfe, Department of Molecular Biology and Biotechnology, University of Sheffield,  
27 Sheffield, S10 2TN, UK

28 Email: m.rolfe@sheffield.ac.uk

29

30 SlyA is a member of the MarR family of bacterial transcriptional regulators. Previously, SlyA  
31 has been shown to directly regulate only two operons in *Escherichia coli* K-12 MG1655, *fimB*  
32 and *hlyE* (*clyA*). In both cases SlyA activates gene expression by antagonizing repression  
33 by the nucleoid associated protein H-NS. Here the transcript profiles of aerobic glucose-  
34 limited steady-state chemostat cultures of *E. coli* K-12 MG1655, *slyA* mutant and *slyA* over-  
35 expression strains are reported. The transcript profile of the *slyA* mutant was not significantly  
36 different to that of the parent; however, that of the *slyA* expression strain was significantly  
37 different from that of the vector control. Transcripts representing 27 operons were increased  
38 in abundance, whereas 3 were decreased. Of the 30 differentially regulated operons, 24  
39 have been previously associated with sites of H-NS binding, suggesting that antagonism of  
40 H-NS repression is a common feature of SlyA-mediated transcription regulation. Direct  
41 binding of SlyA to DNA located upstream of a selection of these targets permitted the  
42 identification of new operons likely to be directly regulated by SlyA. Transcripts of four  
43 operons coding for cryptic adhesins exhibited enhanced expression and this was consistent  
44 with enhanced biofilm formation associated with the SlyA over-producing strain.

45

## 46 INTRODUCTION

47 The MarR family of transcription regulators are widespread throughout the Bacterial and Archeal  
48 kingdoms [1]. MarR family members are homodimeric and bind to palindromic DNA sequences  
49 within regulated promoters using a characteristic winged-helix-turn-helix DNA-binding domain [2].  
50 These regulators repress gene expression by promoter occlusion (e.g. MarR; [3]), or activate gene  
51 expression by stabilizing RNA polymerase-promoter DNA interactions (e.g. OhrR; [4]), or by  
52 antagonizing the action of repressors (e.g. RovA; [5]). These activities of MarR proteins are inhibited  
53 upon interaction with cognate signalling molecules, although for many members the natural ligand is  
54 unknown [6].

55 The *Salmonella enterica* serovar Typhimurium LT2 SlyA protein is one of the best  
56 characterized members of the MarR family. The *S. enterica* serovar Typhimurium *slyA* mutant is  
57 attenuated for virulence, is hypersensitive to oxidative stress and is impaired for survival in  
58 macrophages [7, 8]. A consensus DNA binding site has been proposed, TTAGCAAGCTAA [9, 10],  
59 and proteomic and transcriptomic comparisons of parent and *slyA* mutant strains suggest that SlyA  
60 can act as both a negative and positive regulator of gene expression, with significant overlap with  
61 genes of the PhoPQ regulon involved in cell envelope function, virulence, resistance to anti-microbial  
62 peptides and regulation of small RNAs [11-15]. *Salmonella enterica* serovar Typhimurium 14028s  
63 SlyA has also been linked to the stringent response by binding ppGpp resulting in enhanced DNA-  
64 binding [16, 17]. The expression of many SlyA-regulated genes is subject to H-NS-mediated silencing

65 and activation of these genes generally involves an element of antagonism of H-NS repression by  
66 SlyA; e.g. [11, 18-22].

67 The SlyA protein of *E. coli* MG1655 is 91% identical, 95% similar (over 142 amino acids) to  
68 the *S. enterica* serovar Typhimurium LT2 protein, but is much more poorly characterized. Only two  
69 genes, *hlyE* and *fimB* (as well as autoregulation of *slyA*) have been shown to be directly regulated by  
70 SlyA [19, 21, 23]. In some other *E. coli* strains, SlyA regulates capsule synthesis and lipid A  
71 palmitoylation in biofilms [18, 19, 22]. Here transcriptional profiling of parent, *slyA* mutant and *slyA*  
72 over-expression strains reveals the breadth of the *E. coli* MG1655 SlyA regulon, indicating roles in  
73 activating expression of cryptic fimbrial-like adhesins that contribute to enhanced biofilm formation.

74

## 75 **METHODS**

76 **Bacterial strains, plasmids, oligonucleotides and growth conditions.** The bacterial strains  
77 plasmids and oligonucleotides that were used are listed in Table 1. Bacterial strains were routinely  
78 cultured in Luria Bertani broth or on Luria Bertani agar plates [24]. Aerobic glucose-limited steady-  
79 state chemostat cultures of *E. coli* were established in Evans minimal medium [25] in Labfors 3  
80 fermentation vessels (Infors-HT, Switzerland) with a 1 L working volume, 0.2 h<sup>-1</sup> dilution rate, 37°C,  
81 pH 6.9, 400 rpm stirring rate and sparging with 1 L min<sup>-1</sup> air. Evans minimal medium consists of: 10  
82 mM NaH<sub>2</sub>PO<sub>4</sub>, 10 mM KCl, 1.25 mM MgCl<sub>2</sub>, 20 mM NH<sub>4</sub>Cl, 0.02 mM CaCl<sub>2</sub>, 0.1 mM Na<sub>2</sub>SeO<sub>3</sub>, 1.5  
83 mM monosodium nitrilotriacetate, 20 mM glucose and 100 ml trace element solution. The trace  
84 element solution consisted of (g L<sup>-1</sup>): ZnO (0.412), FeCl<sub>3</sub>·6H<sub>2</sub>O (5.4), MnCl<sub>2</sub>·4H<sub>2</sub>O (2.0), CuCl<sub>2</sub>·2H<sub>2</sub>O  
85 (0.172), CoCl<sub>2</sub>·6H<sub>2</sub>O (0.476), H<sub>3</sub>BO<sub>3</sub> (0.064), Na<sub>2</sub>MoO<sub>4</sub>·H<sub>2</sub>O (0.004) in 0.3% v/v HCl. For generation  
86 of cell paste for purification of His-tagged SlyA, *E. coli* BL21 (λDE3) transformed with pGS2469  
87 was grown in auto-induction medium supplemented with ampicillin (100 mg L<sup>-1</sup>) [26]. Resistance to  
88 chloramphenicol was tested by inoculating Luria Bertani broth (2 ml) containing kanamycin (30 µg  
89 ml<sup>-1</sup>) and either 0, 1, 2, 3 or 4 µg ml<sup>-1</sup> chloramphenicol with 10 µl of overnight starter cultures (*E. coli*  
90 K-12 MG1655 pET28a or *E. coli* K-12 MG1655 pGS2468). Triplicate cultures were grown under  
91 aerobic conditions for 6 h at 37°C before measuring OD<sub>600</sub> as a measure of growth. The experiment  
92 was carried out twice.

93

94 **Biofilm assay.** Biofilm assays were performed using 96-well plates essentially as described  
95 by Tagliabue *et al.* [27] using M9 minimal medium with 20% (w/v) glucose and 50 µg ml<sup>-1</sup>  
96 kanamycin. Wells containing 200 µl of medium were inoculated (1:10) from an overnight culture of *E.*  
97 *coli* K-12 MG1655 pET28a or *E. coli* K-12 MG1655 pGS2468 and then incubated for 16 h under  
98 aerobic conditions at 37°C. Growth of cultures was monitored by measuring OD<sub>600</sub>. The planktonic  
99 cells were removed and the remaining biofilm was stained for 5 min with 200 µl 1% (w/v) crystal  
100 violet solution. Excess stain was removed by three washes with deionized water before the plate was

101 air-dried. To quantify the extent of staining, 200  $\mu$ l ethanol:acetone (4:1) was added to each well, and  
102 after incubating for 20 min the amount of biofilm was estimated by measuring  $A_{600}$ . Adhesion units  
103 were calculated by dividing the  $A_{600}$  values for crystal violet-stained adhered cells by the  $OD_{600}$  values  
104 for the corresponding planktonic cells.

105

106 **Creation of *E. coli* K-12 MG1655 *slyA* mutant.** A PCR-amplified DNA fragment containing  
107 the kanamycin cassette from pKD4 flanked by 40 bp DNA homologous to regions surrounding the  
108 *slyA* gene was synthesized using oligonucleotide primers TC7 and TC8 (Table 1). The purified  
109 (QiaQuick PCR cleanup, Qiagen) PCR product (5  $\mu$ g) was used to transform *E. coli* JRG6072 by  
110 electroporation (Hybaid Cell Shock unit; 1800 V, 1 mm path length). The *E. coli* JRG6072 competent  
111 cells were prepared from aerobic Luria Bertani broth batch cultures supplemented with ampicillin  
112 (100 mg L<sup>-1</sup>) at 30°C that had been induced to express the  $\lambda$ red recombinase by addition of L-  
113 arabinose (1 mM). Kanamycin resistant mutants were selected on Luria Bertani agar plates  
114 containing kanamycin (30 mg L<sup>-1</sup>) at 37°C. Mutation of the *slyA* gene by insertion of the kanamycin  
115 resistance cassette was confirmed by colony PCR using oligonucleotides TC9 and TC10. The *slyA*  
116 mutation was then transduced using bacteriophage P1 to *E. coli* MG1655 [24].

117

118 **Transcriptional profiling.** Transcriptomic analyses were carried out as described by Rolfe *et al.*  
119 [28] using directly quenched samples from glucose-limited steady-state chemostat cultures (dilution  
120 rate 0.2 h<sup>-1</sup>) for the three *E. coli* K-12 MG1655 strains; parent, *slyA* mutant (JRG6457) and *slyA* over-  
121 producer (JRG6636). RNA samples were labelled with Cy5 and the reference *E. coli* K-12 MG1655  
122 genomic DNA was labelled with Cy3. In total, two independent biological replicates were performed  
123 that were hybridised in duplicate (technical replicates) giving four replicates. After hybridization and  
124 image capture, data were extracted from the raw image files using Agilent Feature Extraction v11.5  
125 software and analyzed using GeneSpring v7.3.1. Transcriptomic data have been deposited with  
126 ArrayExpress (accession E-MTAB-5220).

127

128 **Purification of SlyA and Western blotting.** Cultures (500 ml auto-induction medium  
129 supplemented with ampicillin in 2 L conical flasks) of *E. coli* BL21 ( $\lambda$ DE3) pGS2469 were grown at  
130 37°C for 24 h with shaking (250 rpm). Bacteria were collected by centrifugation, the pellet was re-  
131 suspended in 15 ml of breakage buffer (20 mM Tris-HCl, 500 mM NaCl, 5% v/v glycerol, pH 7.5),  
132 the bacteria were lysed by two passages through a French pressure cell (16,000 psi) and the extract  
133 clarified by centrifugation (27,000 g, 15 min, 4°C). The His-tagged SlyA protein was isolated from  
134 the cell-free extract by affinity chromatography on a HiTrap chelating column (1 ml) attached to an  
135 AKTA prime according to the standard manufacturer's protocol (GE Healthcare). The eluted SlyA  
136 was buffer exchanged into 20 mM Tris-HCl, pH 7.4 containing 200 mM NaCl by repeated dilution

137 and Vivaspin 6 concentration (Sartorius Stedim Biotech). The protein was judged to be >90% pure by  
138 Coomassie blue-stained SDS-PAGE and protein concentration was estimated by the BioRad protein  
139 reagent protocol [29]. SlyA protein was detected by Western blotting after separation of polypeptides  
140 by SDS-PAGE and electrophoretic transfer (100 V for 1 h; transfer buffer: 5.8 g L<sup>-1</sup> Tris, 2.9 g L<sup>-1</sup>  
141 glycine, 20% v/v methanol, 0.037% w/v SDS) to Hybond-C Extra nitrocellulose membranes (GE  
142 Healthcare). The membranes were soaked in a blocking solution, which contained 5% w/v dried  
143 skimmed milk in PBS (10 mM phosphate buffer, 137 mM NaCl, 2.7 mM KCl, pH 7.4) and 0.05% v/v  
144 Tween 20, for 16 h at 4°C. The blocking solution was then removed and the membranes washed in  
145 PBS containing 0.05% v/v Tween 20 before exposure to a 1:1000 dilution of the SlyA antibody  
146 (raised in rabbit and provided by Prof. Ian Blomfield, University of Kent) in blocking solution for 1 h  
147 at room temperature. After four washes with PBS containing 0.05% v/v Tween 20, the membranes  
148 were soaked in blocking solution containing anti-rabbit secondary antibody provided in the Pierce  
149 ECL Western Blotting kit and the presence of SlyA was visualized according to the manufacturer's  
150 standard protocol (Thermo Scientific).

151

152 **Electrophoretic mobility shift assays (EMSA).** The LightShift Chemiluminescent EMSA kit  
153 (Thermo Scientific) was used according the manufacturer's instructions. Biotin-labelled DNA of  
154 target promoter regions was amplified from genomic DNA using the appropriate oligonucleotide  
155 primer pairs (Table 1). The core binding assays (20 µl) contained: 2 µl 10x binding buffer (100 mM  
156 Tris-HCl, pH 7.5, containing 500 mM KCl and 10 mM dithiothreitol and 1 µg poly (dI•dC). The  
157 DNA concentration was ~1 nM and the concentration of SlyA ranged from 0 to 500 nM as indicated.  
158 Mixtures were incubated at 25°C for 30 min before separation of SlyA-DNA complexes by native gel  
159 electrophoresis, followed by transfer to Hybond-N+ nylon membranes, UV-crosslinking for 60 s at  
160 120 mJ cm<sup>-2</sup> and detection of labelled DNA using the Nucleic Acid Detection Module (Thermo  
161 Scientific).

162

## 163 **RESULTS AND DISCUSSION**

### 164 **Enhanced expression of *slyA* in *E. coli* K-12 MG1655 results in altered abundance of** 165 **transcripts from 30 operons**

166 Previous work has shown that SlyA directly activates the expression of two genes in *E. coli* K-12  
167 (*hlyE* and *fimB*) by antagonising H-NS repression [20, 21, 23]. However, in *S. enterica* serovar  
168 Typhimurium the influence of SlyA is much more extensive, with at least 31 regulated genes resulting  
169 in hypersensitivity to reactive oxygen species and attenuation in infection models [7, 8, 12]. The  
170 initial aim of this work was to apply transcript profiling to determine the extent of the *E. coli* K-12  
171 MG1655 SlyA regulon by comparison of steady-state glucose-limited aerobic chemostat cultures of  
172 wild-type and *slyA* mutant strains. Comparison of transcript profiles of wild-type and *slyA* mutant

173 cultures grown at a dilution rate of 0.5 h<sup>-1</sup> (equivalent to a doubling time of 1.4 h) revealed no  
174 significant ( $\geq 2$ -fold;  $p \leq 0.05$ ) changes in transcript abundance. Because SlyA translation might be  
175 enhanced at low growth rates, due to its unusual UUG start codon [21], steady-state cultures at  
176 dilution rates 0.2, 0.1 and 0.05 h<sup>-1</sup> were established (equivalent to doubling times of 3.5, 6.9 and 13.8  
177 h, respectively). However, once again, when the transcript profiles and growth characteristics of the  
178 wild-type and *slyA* mutant cultures were compared no significant differences were detected. These  
179 observations indicated that, under the conditions tested, deletion of the *slyA* gene had no significant  
180 effect on gene expression in *E. coli* K-12 MG1655, even at low growth rates.

181 Anti-SlyA serum was used to determine whether SlyA was detectable in *E. coli* K-12  
182 MG1655 cells grown in glucose-limited chemostats at a dilution rate of 0.2 h<sup>-1</sup>. In accordance with  
183 the transcript profiling, SlyA was not detected (Fig. 1). This suggests that the expression of SlyA is  
184 regulated and switched on under conditions other than those imposed here; for example, SlyA protein  
185 has been detected by Western blotting extracts from *E. coli* batch cultures grown in minimal medium  
186 with glycerol as the carbon and energy source [21]. To overcome any regulatory barrier to identifying  
187 genes potentially controlled by SlyA, a plasmid (pGS2468) to express *slyA* under the control of its  
188 own promoter was constructed. Western blotting showed that SlyA protein was now readily  
189 detectable in the transformed *E. coli* K-12 MG1655 cells grown in glucose-limited chemostats at a  
190 dilution rate of 0.2 h<sup>-1</sup> (Fig. 1). The growth characteristics of the vector control and the *slyA*  
191 expression strains were essentially the same, with similar yields ( $1.4 \pm 0.2$  g cell dry weight per litre)  
192 and no detectable glucose or over-metabolites in the culture supernatants. Therefore, the transcript  
193 profiling experiments were carried out with these strains grown in aerobic glucose-limited chemostats  
194 at a dilution rate of 0.2 h<sup>-1</sup>. The transcript profile of the SlyA over-production strain was significantly  
195 different from that of the vector control. Transcripts representing 27 operons were increased in  
196 abundance and 3 were decreased (Table 2; Fig. 1c). The transcripts exhibiting decreased abundance  
197 were: the *sgc* operon (*sgcXBCQAER*), which encodes a phosphotransferase system for the uptake of  
198 an unknown sugar; *fecIR* the membrane-bound sensor (FecR) that receives signals from the outer  
199 membrane ferric citrate uptake receptor (FecA) for transmission to FecI ( $\sigma^{19}$ ), which activates  
200 transcription of the *fecABCDE* operon encoding components of a cytoplasmic membrane bound ferric  
201 citrate uptake system; and *yecH*, which encodes a predicted protein of unknown function (Table 2)  
202 [30, 31].

203 Amongst the up-regulated transcripts were the previously identified SlyA-regulated gene *hlyE*  
204 and *slyA* itself (Table 2). The latter was not surprising as the *slyA* gene was present in multi-copy, but  
205 despite this the *slyA* transcript only increased ~3-fold in abundance, yet the SlyA protein level  
206 increased from being undetectable in the control to a level equivalent to ~1.5  $\mu\text{M}$  in the cytoplasm  
207 (based on the dry weight of *E. coli* being  $3 \times 10^{-13}$  g with an aqueous volume of  $7 \times 10^{-13}$  ml per cell;  
208 [32]). The relatively low level of induction of the *slyA* transcript when present in multi-copy but



209 much greater induction of SlyA protein suggests that the *slyA* promoter is subject to auto-regulation,  
210 consistent with the reported SlyA binding at the *slyA* promoter [19]. It was also notable that the *ydhI*-  
211 *K* operon, which is divergently transcribed from *slyA* and not present on the *slyA* expression plasmid,  
212 also exhibited enhanced transcript abundance, suggesting that SlyA is capable of activating expression  
213 from divergent promoters; an assertion supported by the enhanced abundances of the divergently  
214 transcribed *hlyE* and *C0299* (encodes a small RNA) genes in the presence of SlyA (Table 2).

215 Twenty-four of the 30 operons (~80%) that showed altered transcript abundance upon over-  
216 production of SlyA have also been shown to be associated with H-NS binding sites (Table 2). Thus, it  
217 appears that H-NS repressed genes are over-represented in the set of transcripts that increase in  
218 abundance when SlyA is expressed, suggesting that SlyA acts by antagonizing H-NS repression at the  
219 corresponding promoters; a mechanism that is established for *hlyE* [20]. H-NS binds DNA by  
220 recognizing the structure of A-T-rich minor grooves and silences the expression of horizontally  
221 acquired A-T-rich genes (reviewed by [33]). H-NS is thus considered crucial in permitting the  
222 acquisition of new genes whilst counteracting the potentially detrimental effects of inappropriate  
223 expression of these genes. Counter-silencing by H-NS antagonists, such as SlyA, provides a route to  
224 integrate expression of the genes into the regulatory circuits of *E. coli* under appropriate conditions.  
225 Horizontally acquired genes are located within genomic islands, which are regions of bacterial  
226 chromosomes containing that are often associated with drug resistance, metabolic adaptability, stress  
227 tolerance and pathogenesis. Genomic islands can be recognized by their sequence composition and  
228 increased transcript start point densities [32]. The analysis tools GIST (Genomic-island Identification  
229 by Signals of Transcription) and IslandViewer have been used to map the genomic islands of *E. coli*  
230 K-12 MG1655 [34]. Notably, 13 of the 30 differentially regulated operons overlapped predicted  
231 genomic islands, suggesting a general role for SlyA in the counter-silencing of H-NS repressed  
232 horizontally acquired genes under conditions when *slyA* is up-regulated (Table 2).

233 The H-NS-repressed *casABC* operon was up-regulated by SlyA (Table 2). This operon  
234 encodes proteins involved in maintaining and utilising the library of foreign genetic elements  
235 interspersed between CRISPR sequences which act as the immune system memory of Bacteria and  
236 Archaea [35]. CRISPR loci, in general, consist of closely spaced direct repeats separated by short  
237 spacer regions of variable sequence. Spacer regions mostly correspond to sections of foreign plasmid  
238 or viral sequences which have been integrated. The CRISPR loci are found adjacent to the *casABC*  
239 operon. The fact that the *casABC* operon was significantly up-regulated by SlyA suggests that this  
240 regulator may contribute to viral resistance and immunity in *E. coli* K-12 MG1655.

241 Other transcripts that exhibited increased abundance in the presence of SlyA were associated  
242 with uptake and metabolism of phenylacetic acid (*paaA-K*), utilization of alkanesulfonates as  
243 alternative sulfur sources (*ssuEADCB*; divergently transcribed from the *elf* operon; see below), a  
244 cryptic galactosamine transport and catabolism system (*agaS-I*) and a 2-O- $\alpha$ -mannosyl-D-glycerate  
245 phosphotransferase and  $\alpha$ -mannosidase (Table 2) [36-39]. Hence, it appears that SlyA plays a role in

246 regulating systems that expand the repertoire of substrates utilized by *E. coli*. Increased abundance of  
247 the *mdtM* transcript suggests a role for SlyA in enhancing expression of this multidrug transporter that  
248 confers resistance to ethidium bromide and chloramphenicol with mutants exhibiting attenuated  
249 growth at alkaline pH [40]. However, simple growth inhibition studies suggested that *slyA* expression  
250 led to increased sensitivity to chloramphenicol (growth yield after 6 h at 37°C in Luria Bertani broth  
251 was lowered to ~50% by 2 µg ml<sup>-1</sup> for the wild-type carrying the empty vector compared to 1 µg ml<sup>-1</sup>  
252 for the wild-type carrying the *slyA* expression plasmid), rather than increased resistance, perhaps  
253 reflecting the complexity of the phenotype of the *slyA* expression strain.

254 Several of the SlyA-regulated operons code for proteins involved in membrane function. In *S.*  
255 *enterica* serovar Typhimurium the majority of genes affected by SlyA encode proteins associated with  
256 the bacterial cell envelope and are important for virulence and survival within murine macrophages.  
257 Although it has been previously shown that the majority of genes regulated by SlyA in *S. enterica*  
258 serovar Typhimurium are not present in *E. coli* K-12 [12, 15], a similar propensity for cell envelope  
259 proteins being regulated by the *E. coli* SlyA was evident here. Thirteen (43%) of the 30 operons that  
260 exhibited altered expression in SlyA-expressing bacteria were associated with cell-surface/membrane  
261 functions (Table 2).

262 The *gspC-O* operon is cryptic membrane-associated, H-NS-repressed, transcription unit that  
263 was up-regulated by SlyA (Table 2). The *gspC-O* operon encodes a Type-II Secretion System (T2SS)  
264 for the export of endogenous proteins and formation of structural elements of the Gsp secreton, which  
265 is thought to facilitate the export of the endogenous endochitinase ChiA, a product of another H-NS  
266 silenced gene [41, 42].

267 Amongst the transcripts with increased abundance in the SlyA over-producing strain were  
268 four cryptic operons (*elfADCG-ycbUVF*, *sfmHF*, *yehDCBA* and *yadN*) encoding fimbrial-like  
269 adhesins (Table 2). These four operons were amongst seven putative chaperone-usher fimbrial  
270 systems shown to be poorly expressed under laboratory conditions by Korea *et al.* [43]. Nevertheless,  
271 when these operons were individually expressed by placing them under the control of a constitutive  
272 promoter six were shown to be functional and expression of the *elf* (*ycb*), *yad* and *yeh* operons  
273 resulted in enhanced biofilm formation on abiotic surfaces, whereas *sfm* promoted binding to  
274 eukaryotic cells [43]. Moreover, all four operons were repressed by H-NS. The increased  
275 abundances of the *elf*, *sfm*, *yad* and *yeh* transcripts upon expression of SlyA is consistent with the  
276 cryptic status of these genes under normal laboratory conditions and suggests that these chaperone-  
277 usher fimbriae are functional under environmental conditions that enhance *slyA* expression such that  
278 SlyA can operate as an H-NS antagonist (Table 2).

279

## 280 **SlyA over-production is associated with enhanced biofilm formation**

281 The observation that SlyA increased transcription of four cryptic fimbrial-like adhesins suggested that  
282 the SlyA over-producing strain should exhibit enhanced biofilm production. This was tested using

283 static cultures of *E. coli* K-12 MG1655 transformed with pET28a (control) or the *slyA* expression  
284 plasmid pGS2468 in conditions that mirrored the transcript profiling experiment. The data showed a  
285 4-fold increase in biofilm formation when *slyA* was over-expressed, consistent with the transcript  
286 profiling data (Fig. 2).

287

### 288 **Identification of new *E. coli* K-12 MG1655 operons that are directly regulated by SlyA**

289 The changes in transcript profiles that were observed upon over-production of SlyA could result from  
290 direct interaction of SlyA with the promoter regions of the corresponding genes or indirectly via  
291 SlyA-regulated factors. For example, one of the genes up-regulated upon SlyA over-production, *leuO*,  
292 encodes a transcriptional regulator that, like SlyA, operates by antagonising H-NS regulation [44, 45].  
293 Fourteen (52%) of the 27 transcripts that were increased in abundance when SlyA was expressed in *E.*  
294 *coli* K-12 MG1655 were associated with LeuO binding sites identified in the SELEX-chip study of  
295 Shimada *et al.* [44]. This strong correlation could arise from; (1) the positive effect SlyA has on the  
296 expression of *leuO* resulting in an increase in expression of the entire LeuO regulon, i.e. indirect  
297 regulation by SlyA; or (2) SlyA and LeuO have overlapping regulons as a consequence of the fact  
298 they both operate by antagonising H-NS-mediated repression. To further investigate the extent of  
299 direct SlyA-mediated regulation in *E. coli* K-12 MG1655 binding of SlyA to ten promoter regions  
300 was examined by electrophoretic mobility shift assays (EMSA).

301 Amongst the transcripts differentially regulated by over-production of SlyA there were three  
302 arranged as divergent operons (Fig. 3). Binding of SlyA at the *hlyE-C0299* intergenic region was  
303 shown previously (Fig. 3) [20]. Two other examples of SlyA-activated divergent operons (*slyA-*  
304 *ydhIJK* and *ssuE-B-elfADCG-ycbUVF*) were shown to bind SlyA in EMSA (Fig. 3). Furthermore,  
305 SlyA bound at the *casA*, *fecIR*, *gspCDEF*, *leuO*, *mdtM* and *paaA-K* promoters (Fig. 3). The  $K_{d(app)}$   
306 values for SlyA binding at these promoters were similar at ~50-100 nM. These experiments indicate  
307 that these operons are likely to be directly regulated by SlyA. The *sgcXBCQ-sgcAER* genes are  
308 separated by a sRNA *ryjB* on the opposite DNA strand (Fig. 3). It is suggested that the *sgcXBCQAER*  
309 is a single transcription unit, but there is no high quality evidence to support this suggestion [46].  
310 Therefore, both the region upstream of *sgcX* and the intergenic region between *sgcQ* and *sgcA* were  
311 used in EMSAs with the SlyA protein. No specific interaction was observed with the region upstream  
312 of *sgcA* but interaction, albeit weaker than that observed for the promoter regions analyzed above, was  
313 observed when the DNA upstream of *sgcX* was tested (Fig. 3). These observations suggest that  
314 *sgcXBCQAER* is a single SlyA-repressed transcription unit.

315 The EMSA experiments indicate that SlyA binds *Pssu*, *Pcas*, *Ppaa*, *Pelf*, *PleuO* and *Pgsp*, all  
316 of which are promoter regions of genes or operons proposed to be part of the LeuO regulon (Table 2).  
317 This suggests that, perhaps because of the similarity in their mode of action, i.e. antagonizing H-NS  
318 repression, the SlyA and LeuO regulons substantially overlap such that upon activation by their  
319 respective signals a similar transcriptional response is elicited.

320 A consensus binding site (TTAGCAAGCTAA) for the *Salmonella enterica* serovar  
321 Typhimurium LT2 SlyA protein was proposed based on footprinting and a limited SELEX analysis  
322 [10]. This consensus was further analyzed by site-directed mutagenesis, which suggested the  
323 consensus sequence TTAN<sub>6</sub>TAA [9]. All the DNA fragments that bound *E. coli* SlyA in EMSAs (Fig.  
324 3) possessed DNA sequences similar to the previously proposed consensus sequences (Table 3). Site-  
325 directed replacement amino acid residues of *Salmonella enterica* serovar Typhimurium LT2 SlyA  
326 identified 16 locations that impaired DNA-binding [9], all these amino acids are conserved in the *E.*  
327 *coli* SlyA protein, suggesting that these closely related proteins recognize similar DNA motifs.

328

### 329 **Concluding remarks**

330 SlyA proteins have been shown to play important roles in regulating gene expression in a wide range  
331 of bacterial species. The most common mechanism for SlyA-mediated activation of gene expression  
332 is through antagonism of H-NS repression. Here transcript profiling has revealed the breadth of the  
333 SlyA regulon (directly and indirectly regulated genes) in *E. coli* K-12 MG1655 cultures grown under  
334 precisely controlled conditions such that any potential effects associated with changes in growth  
335 rate/growth phase could not confound the interpretation of the data obtained. Enhanced transcript  
336 abundance for several cryptic fimbrial operons in a SlyA over-producing strain and an over-  
337 representation of H-NS repressed genes were consistent with the current model of SlyA-mediated  
338 gene activation. The SlyA protein was shown to bind at 9 intergenic regions controlling the  
339 expression of 11 operons, thus expanding the number of known directly SlyA-regulated genes in *E.*  
340 *coli* MG1655 from 2 to 13.

341

### 342 **FUNDING INFORMATION**

343 TDC was supported by a Biotechnology and Biological Sciences Research Council (BBSRC) UK  
344 PhD scholarship; SPH was supported by a Sheffield Undergraduate Research Experience (SURE)  
345 project.

346

### 347 **ACKNOWLEDGEMENTS**

348 The authors thank Ian Blomfield (University of Kent, UK) for providing the SlyA antiserum and Ruth  
349 Roberts (University of Sheffield) for technical support.

350

### 351 **CONFLICTS OF INTEREST**

352 The authors declare no conflicts of interest.

353

### 354 **REFERENCES**

355

- 356 **1. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY et al.** Pfam: the protein families  
357 database. *Nucl Acids Res* 2014;42:D222-D230.
- 358 **2. Perera IC, Grove A.** Molecular mechanisms of ligand-mediated attenuation of DNA-binding by  
359 MarR family transcription regulators. *J Mol Cell Biol* 2010;2:243-254.
- 360 **3. Martin RG, Rosen JL.** Binding of the purified multiple antibiotic-resistance repressor protein  
361 (MarR) to *mar* operator sequences. *Proc Natl Acad Scis U S A* 1995;92:5456-5460.
- 362 **4. Oh SY, Shin JH, Roe JH.** Dual role of OhrR as a repressor and an activator in response to organic  
363 hydroperoxides in *Streptomyces coelicolor*. *J Bacteriol* 2007;189:6284-6292.
- 364 **5. Cathelyn JS, Ellison DW, Hinchliffe SJ, Wren BW, Miller V.** The RovA regulons of *Yersinia*  
365 *enterocolitica* and *Yersinia pestis* are distinct: evidence that many RovA-regulated genes were  
366 acquired more recently than the core genome. *Mol Microbiol* 2007;66:189-205.
- 367 **6. Wilkinson, S. P. & Grove, A. (2006).** Ligand-responsive transcriptional regulation by members of  
368 the MarR family of winged helix proteins. *Curr Issues Mol Biol* **8**, 51-62.
- 369 **7. Buchmeier N, Bossie S, Chen, CY, Fang FC, Guiney DG et al.** SlyA, a transcriptional regulator  
370 of *Salmonella typhimurium*, is required for resistance to oxidative stress and is expressed in the  
371 intracellular environment of macrophages. *Infect Immun* 1997;65:3725-3730.
- 372 **8. Libby SJ, Goebel W, Ludwig A, Buchmeier N, Bowe F et al.** A cytolysin encoded by *Salmonella*  
373 is required for survival within macrophages. *Proc Natl Acad Sci U S A* 1994; 91:489-493.
- 374 **9. Haider F, Lithgow JK, Stapleton MR, Norte VA, Roberts RE et al.** DNA recognition by the  
375 *Salmonella enterica* serovar Typhimurium transcription factor SlyA. *Intl Microbiol* 2008;11:245-  
376 250.
- 377 **10. Stapleton MR, Norte VA, Read RC, Green J.** Interaction of the *Salmonella typhimurium*  
378 transcription and virulence factor SlyA with target DNA and identification of members of the SlyA  
379 regulon. *J Biol Chem* **2002;277**:17630-17637.
- 380 **11. Colgan AM, Kröger C, Diard M, Hardt WD, Puente J L et al.** The impact of 18 ancestral and  
381 horizontally-acquired regulatory proteins upon the transcriptome and sRNA landscape of  
382 *Salmonella enterica* serovar Typhimurium. *PLoS Genetics* 2016;12:e1006258.
- 383 **12. Navarre WW, Halsey TA, Walthers D, Frye J, McClelland M et al.** Co-regulation of  
384 *Salmonella enterica* genes required for virulence and resistance to antimicrobial peptides by SlyA  
385 and PhoP/PhoQ. *Mol Microbiol* 2005;56:492-508.
- 386 **13. Norte VA, Stapleton MR, Green J.** PhoP-responsive expression of the *Salmonella enterica*  
387 serovar Typhimurium *slyA* gene. *J Bacteriol* 2003;185:3508-3514.
- 388 **14. Perez JC, Latifi T, Groisman EA.** Overcoming H-NS-mediated transcriptional silencing of  
389 horizontally acquired genes by the PhoP and SlyA proteins in *Salmonella enterica*. *J Biol Chem*  
390 2008;283:0773-10783.

- 391 **15. Spory A, Bosserhoff A, von Rhein C, Goebel W, Ludwig A.** Differential regulation of multiple  
392 proteins of *Escherichia coli* and *Salmonella enterica* serovar Typhimurium by the transcriptional  
393 regulator SlyA. *J Bacteriol* 2002;184:3549-3559.
- 394 **16. Dalebroux ZD, Swanson MS.** ppGpp: magic beyond RNA polymerase. *Nat Rev Microbiol*  
395 2012;10:203-212.
- 396 **17. Zhao G, Weatherspoon N, Kong W, Curtiss R, Shi YX.** A dual-signal regulatory circuit  
397 activates transcription of a set of divergent operons in *Salmonella typhimurium*. *Proc Natl Acad*  
398 *Sci U S A* 2008;105:20924-20929.
- 399 **18. Chalabaev S, Chauhan A, Novikov A, Iyer P, Szczesny M et al.** Biofilms formed by gram-  
400 negative bacteria undergo increased lipid A palmitoylation, enhancing *in vivo* survival. *MBio*  
401 2014;5:pii:e01116-14.
- 402 **19. Corbett D, Bennett HJ, Askar H, Green J, Roberts IS.** SlyA and H-NS regulate transcription of  
403 the *Escherichia coli* K5 capsule gene cluster, and expression of *slyA* in *Escherichia coli* is  
404 temperature-dependent, positively autoregulated, and independent of H-NS. *J Biol Chem*  
405 2007;282:33326-33335.
- 406 **20. Lithgow JK, Haider F, Roberts IS, Green J.** Alternate SlyA and H-NS nucleoprotein complexes  
407 control *hlyE* expression in *Escherichia coli* K-12. *Mol Microbiol* 2007;66:685-698.
- 408 **21. McVicker G, Sun L, Sohanpal BK, Gashi K, Williamson RA et al.** SlyA protein activates *fimB*  
409 gene expression and type 1 fimbriation in *Escherichia coli* K-12. *J Biol Chem* 2011;286:32026-  
410 32035.
- 411 **22. Xue P, Corbett D, Goldrick M, Naylor C, Roberts IS.** Regulation of expression of the region 3  
412 promoter of the *Escherichia coli* K5 capsule gene cluster involves H-NS, SlyA and a large 5'  
413 untranslated region. *J Bacteriol* 2009;191:838-1846.
- 414 **23. Wyborn NR, Stapleton MR, Norte VA, Roberts RE., Grafton J et al.** Regulation of  
415 *Escherichia coli* hemolysin E expression by H-NS and *Salmonella* SlyA. *J Bacteriol*  
416 2004;186:1620-1628.
- 417 **24. Sambrook J, Russell DW.** Molecular cloning - a laboratory manual, 3rd ed., Cold Spring Harbor  
418 NY: Cold Spring Harbor Laboratory; 2001.
- 419 **25. Evans CGT, Herbert D, Tempest DW.** The continuous cultivation of micro-organisms. *Meth*  
420 *Microbiol* 1970;2:278-327.
- 421 **26. Studier FW.** Protein production by auto-induction in high-density shaking cultures. *Prot Express*  
422 *Purific* 2005;41:207-234.
- 423 **27. Tagliabue L, Antoniani D, Maciag A, Bocci P, Raffaelli N et al.** The diguanylate cyclase YddV  
424 controls production of the exopolysaccharide poly-N-acetylglucosamine (PNAG) through  
425 regulation of the PNAG biosynthetic *pgaABCD* operon. *Microbiology* 2010;156:2901-2911.

- 426 **28. Rolfe MD, Ter Beek A, Graham AI, Trotter EW, Shahzad Asif HM et al.** Transcript profiling  
427 and inference of *Escherichia coli* K-12 ArcA activity across the range of physiologically relevant  
428 oxygen concentrations. *J Biol Chem* 2011;286:10147-10154.
- 429 **29. Bradford MM.** Rapid and sensitive method for quantification of microgram quantities of protein  
430 utilizing principle of protein-dye binding. *Anal Biochem* 1976;72:248-254.
- 431 **30. Braun V, Mahren S, Sauter A.** Gene regulation by transmembrane signalling. *Biometals*  
432 2006;19:103-113.
- 433 **31. Reizer J, Reizer A, Saier MH.** Is the ribulose monophosphate pathway widely distributed in  
434 bacteria? *Microbiology* 1997;143:2519-2520.
- 435 **32. Sundararaj S, Guo A, Habibi-Nazhad B, Rouani M, Stothard P et al.** The CyberCell database  
436 (CCDB): a comprehensive, self-updating, relational database to coordinate and facilitate *in silico*  
437 modeling of *Escherichia coli*. *Nucl Acids Res* 2004;32:D293-D295.
- 438 **33. Navarre WW.** The impact of gene silencing on horizontal gene transfer and bacterial evolution.  
439 *Adv Microb Physiol* 2016;69:157-186.
- 440 **34. Huang Q, Cheng X, Cheung MK, Kiselev SS, Ozoline ON et al.** High-density transcriptional  
441 initiation signals underline genomic islands in bacteria. *PLoS One* 2012;7:e33759.
- 442 **35. Horvath P, Barrangou R.** CRISPR/Cas, the immune system of Bacteria and Archaea. *Science*  
443 2010;327:167-170.
- 444 **36. Brinkkötter A, Klöß, H, Alpert CA, Lengeler JW.** Pathways for the utilization of N-acetyl-  
445 galactosamine and galactosamine in *Escherichia coli*. *Mol Microbiol* 2000;37:125-135.
- 446 **37. Eichhorn E, van der Ploeg JR, Leisinger T.** Deletion analysis of the *Escherichia coli* taurine  
447 and alkanesulfonate transport systems. *J Bacteriol* 2000;182:2687-2695.
- 448 **38. Ismail W, Mohamed ME, Wanner BL, Datsenko KA, Eisenreich W et al.** Functional  
449 genomics by NMR spectroscopy – phenylacetate catabolism in *Escherichia coli*. *Eur J Biochem*  
450 2003;270:3047-3054.
- 451 **39. Sampaio MM, Chevance F, Dippel R, Eppler T, Schlegel A et al.** Phosphotransferase-mediated  
452 transport of the osmolyte 2-O-alpha-mannosyl-D-glycerate in *Escherichia coli* occurs by the  
453 product of the *mngA* (*hrsA*) gene and is regulated by the *mngR* (*farR*) gene product acting as  
454 repressor. *J Biol Chem* 2004;279:5537-5548.
- 455 **40. Holdsworth SR, Law CJ. (2013).** Multidrug resistance protein MdtM adds to the repertoire of  
456 antiporters involved in alkaline pH homeostasis in *Escherichia coli*. *BMC Microbiol* **13**, 113.
- 457 **41. Francetic O, Belin D, Badaut C, Pugsley AP.** Expression of the endogenous type II secretion  
458 pathway in *Escherichia coli* leads to chitinase secretion. *EMBO J* 2000;19:6697-6703.
- 459 **42. Francetic O, Pugsley AP.** The cryptic general secretory pathway (*gsp*) operon of *Escherichia coli*  
460 K-12 encodes functional proteins. *J Bacteriol* 1996;178:3544-3549.

- 461 **43. Korea CG, Badouraly R, Prevost MC, Ghigo JM, Beloin C.** *Escherichia coli* K-12 possesses  
462 multiple cryptic but functional chaperone-usher fimbriae with distinct surface specificities. *Env*  
463 *Microbiol* 2010;12:1957-1977.
- 464 **44. Shimada T, Bridier A, Briandet R, Ishihama A.** Novel roles of LeuO in transcription regulation  
465 of *E. coli* genome: antagonistic interplay with the universal silencer H-NS. *Mol Microbiol*  
466 2011;82:378-397.
- 467 **45. Stratmann T, Madhusudan S, Schnetz K.** Regulation of the *yjjQ-bglJ* operon, encoding LuxR-  
468 type transcription factors, and the divergent *yjjP* gene by H-NS and LeuO. *J Bacteriol*  
469 2008;190:926-935.
- 470 **46. Keseler IM, Mackie A, Peralta-Gil M, Santos-Zavaleta A, Gama-Castro S et al.** EcoCyc:  
471 fusing model organism databases with systems biology. *Nucl Acids Res* 2013;41:D605-D612.
- 472 **47. Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V et al.** The complete genome  
473 sequence of *Escherichia coli* K-12. *Science* 1997; 277:1453-1462.
- 474 **48. Datsenko KA, Wanner BL** One-step inactivation of chromosomal genes in *Escherichia coli* K-  
475 12 using PCR products. *Proc Natl Acad Sci U S A* 2000; 97:6640-6645.
- 476 **49. Grainger DC, Hurd D, Goldberg MD, Busby SJW.** Association of nucleoid proteins with  
477 coding and non-coding segments of the *Escherichia coli* genome. *Nucl Acids Res* 2006;34:4642-  
478 4652.
- 479 **50. Kahramanoglou C, Seshasayee ASN, Prieto AI, Ibberson D, Schmidt S et al.** Direct and  
480 indirect effects of H-NS and Fis on global gene expression control in *Escherichia coli*. *Nucl Acids*  
481 *Res* 2011; 39:2073-2091.
- 482 **51. Oshima T, Ishikawa S, Kurokawa K, Aiba H, Ogasawara N.** *Escherichia coli* histone-like  
483 protein H-NS preferentially binds to horizontally acquired DNA in association with RNA  
484 polymerase. *DNA Res* 2006;13:141-153.
- 485



Strain or plasmid	Relevant characteristics <sup>a</sup>	Reference or source
<b>Bacterial strain</b>		
<i>E. coli</i> BL21 (λDE3)	<i>E. coli</i> BL21 lysogen for inducible (IPTG) expression of the T7 RNA polymerase	Novagen
<i>E. coli</i> JRG6457	<i>E. coli</i> MG1655 <i>slyA</i>	This work
<i>E. coli</i> JRG6636	<i>E. coli</i> MG1655 pGS2468	This work
<i>E. coli</i> JRG6072	<i>E. coli</i> MG1655 pKD46	This work
<i>E. coli</i> MG1655	Genome sequenced parental strain	[47]
<b>Plasmid</b>		
pET28a	Multi-copy plasmid; Kan <sup>R</sup>	Novagen
pGS2468	pET28a derivative for expression of <i>slyA</i> from the <i>slyA</i> promoter; Kan <sup>R</sup>	This work
pGS2469	pLATE-51 derivative for over-production of SlyA; Amp <sup>R</sup>	This work
pKD4	Source of kanamycin resistance cassette; Amp <sup>R</sup> , Kan <sup>R</sup>	[48]
pKD46	Plasmid for inducible (L-arabinose) expression of the λred recombinase; Amp <sup>R</sup> , T <sup>s</sup>	[48]
pLATE-51	Expression vector for production of His-tagged proteins; Amp <sup>R</sup>	Thermo Scientific
<b>Oligonucleotide</b>		
TC7	TAAAGCCGCATAATATCTTAGCAAGCTAATTATAAGGAGATTA CACGTCTTGAGCGATT; creation of <i>slyA</i> mutant	This work
TC8	TTGCGTGTGGTCAGGTTACTGACCACACGCCCCCTTCATTCAT ATGAATATCCTCCTTAG; creation of <i>slyA</i> mutant	This work
TC9	CTGACGGTAACCAAATGCAG; PCR of <i>slyA</i> locus	This work
TC10	TTTGCCTGTGGTCAGGTTAC; PCR of <i>slyA</i> locus	This work
TC49	[Btn]ACTCTCTCCTTATAACCAATTG; forward primer for PCR of biotin (Btn)-labelled 355 bp intergenic region between <i>ssuE</i> and <i>elfA</i>	This work
TC50	CGTTATCATCCTGATCTCTT; reverse primer for use with TC49	This work
TC51	[Btn]TGGTGAATATTATTGATCAATTAAT; forward primer for PCR of biotin (Btn)-labelled 344 bp intergenic region between <i>leuO</i> and <i>leuL</i>	This work
TC52	ACTTAACTCCACTGTACACTTAA; reverse primer for use with TC51	This work
TC53	[Btn]TTGTTCTCCTTCATATGCTC; forward primer for PCR of biotin (Btn)-labelled 414 bp intergenic region between <i>casA</i> and <i>cas3</i>	This work
TC54	CTTCGGGAATGATTGTTATC; reverse primer for use with TC53	This work
TC55	[Btn]TGTTGCTAATAGTTAAATCGC; forward primer for PCR of biotin (Btn)-labelled 257 bp intergenic region between <i>paaA</i> and <i>paaZ</i>	This work
TC56	GTCATCACCTTTACGATTCC; reverse primer for use with TC55	This work
TC57	[Btn]AACAAACAACCTCTTGCCG; forward primer for PCR of biotin (Btn)-labelled 400 bp region upstream of <i>mdtM</i>	This work
TC58	CCCCGAGGCGCTTTCCAGGC; reverse primer for use with TC57	This work
TC59	[Btn]AGAACTTCCTGTTTTAATTATTG; forward primer for PCR of biotin (Btn)-labelled 179 bp intergenic region between <i>gspA</i> and <i>gspC</i>	This work
TC60	GATGTATGTTCTAATAAAATAGATTG; reverse primer for use with TC59	This work
TC61	[Btn]CCGTCGTTGACTCCATGC; forward primer for PCR of biotin (Btn)-labelled 130 bp intergenic region between <i>sgcA</i> and <i>sgcQ</i>	This work
TC62	GATGGGGATAAGCAGAGC; reverse primer for use with TC61	This work
TC63	[Btn]GCGGAGTGCATCAAAAGT; forward primer for PCR of biotin (Btn)-labelled 291 bp intergenic region between <i>fecI</i> and <i>insA-7</i>	This work
TC64	GCAAGCACCTTAAAATCAC; reverse primer for use with TC63	This work
TC65	[Btn]TTTCATCTCCTTATAATTAGCTT; forward primer for PCR of biotin (Btn)-labelled 200 bp intergenic region between <i>slyA</i> and <i>ydhI</i>	This work
TC66	AAAGTAGATTCCCTTACGACC; reverse primer for use with TC65	This work
TC70	[Btn]AGCTATCTCCGTAGACCGT; forward primer for PCR of biotin (Btn)-labelled 400 bp region upstream of <i>sgcX</i>	This work
TC71	GATTATCTATACTCCCTCTGAATC; reverse primer for use with TC70	This work

487 <sup>a</sup>Amp<sup>R</sup>, ampicillin resistant; Kan<sup>R</sup>, kanamycin resistant; T<sup>s</sup>, temperature sensitive replication

488

**Table 2** Transcripts exhibiting altered abundance upon over-expression of *slyA* in *E. coli* MG1655

Operon <sup>a</sup>	Fold-change <sup>b</sup>	Gene function <sup>c</sup>	H-NS regulon <sup>d</sup>	LeuO regulon <sup>e</sup>	Overlap with genomic island <sup>f</sup>
<i>ybeT</i>	4.1	conserved outer membrane protein	K	✓	
<i>trkG</i>	3.8	Rac prophage potassium transporter subunit	K,O		IV
<i>ssuEADCB</i>	3.6	aliphatic sulfonate transport and metabolism	G,K,O	✓	
<i>yehDCBA</i>	3.6	chaperone-usher fimbrial operon (cryptic)	K,O	✓	GIST
<i>mngAB</i>	3.4	2-O- $\alpha$ -mannosyl-D-glycerate PTS and $\alpha$ -mannosidase		✓	
<i>casABC</i>	3.3	CRISPR associated genes	K	✓	GIST, IV
<i>yghS</i>	3.1	predicted protein with nucleoside triphosphate hydrolase domain	K,O	✓	
<i>slyA</i>	3.0	DNA-binding transcriptional activator	O		
<i>yfbN</i>	2.8	predicted protein	K,O	✓	IV
<i>paaA-K</i>	2.8	phenylacetic acid degradation			
<i>ybeU-hscD</i>	2.8	predicted tRNA ligase and chaperone	K,O	✓	
<i>elfADCG-yebUVF</i>	2.7	predicted fimbrial-like adhesin protein (cryptic)	G,K,O	✓	
<i>ygeG</i>	2.7	predicted chaperone	G,K,O		GIST
<i>crfC-yjcZ</i>	2.6	clamp-binding sister replication fork co-localization protein and predicted protein	K,O		
<i>sfmHF</i>	2.6	predicted fimbrial-like adhesin protein (cryptic)	O		IV
<i>agaS-kbaY-agaBCDI</i>	2.5	predicted galactosamine-transport and metabolism (cryptic)		✓	
<i>ydhYV-T</i>	2.5	predicted oxidoreductase	G,K,O		GIST
<i>yiiE</i>	2.5	predicted transcriptional regulator	K,O		
<i>mdtM</i>	2.5	multidrug efflux system protein			
<i>leuO</i>	2.5	DNA-binding transcriptional activator	G,K,O	✓	GIST
<i>C0299</i>	2.4	sRNA C0299	O		
<i>ycjMN-V</i>	2.4	predicted sugar transporter and metabolism	K,O	✓	
<i>yadN</i>	2.4	predicted fimbrial-like adhesin protein (cryptic)	G,K,O	✓	GIST
<i>gspCDEF</i>	2.4	type II secretion system (cryptic)	K,O	✓	
<i>ydhIJK</i>	2.2	predicted proteins	O		
<i>yfdM</i>	2.1	CPS-53 (KpLE1) prophage predicted methyltransferase	O		GIST, IV

<i>hlyE</i>	2.0	hemolysin E (cryptic)	K,O	
<i>yecH</i>	0.5	predicted protein	O	GIST
<i>sgcXBCQAER</i>	0.5	predicted sugar transport and metabolism		IV
<b><i>fecIR</i></b>	0.4	transcription regulation of ferric citrate transport		IV

490 <sup>a</sup>The fold-change data shown are for the first gene in the operon except where indicated by bold typeface; note  
491 that all genes in the operons followed the same pattern of regulation.

492 <sup>b</sup>Fold-change ( $\geq 2$ -fold,  $p \leq 0.05$ ) is the product of dividing the transcript abundance for the *slyA* over-expression  
493 cultures by that for the control cultures.

494 <sup>c</sup>Gene functions as assigned in Ecocyc.org [46].

495 <sup>d</sup>Genes associated with H-NS binding were identified from Grainger *et al.* [49] (G), Kahramanoglou *et al.* [50] (K)  
496 and Oshima *et al.* [51] (O).

497 <sup>e</sup>Genes located up- or down-stream of a LeuO binding site identified by Shimada *et al.* [44].

498 <sup>f</sup>Genes the overlap with genomic islands in *E. coli* K-12 MG1655 identified by GIST and/or IslandViewer (IV) [34].

499 **Table 3** Candidate SlyA binding sites within the DNA fragments used for EMSA analyses

Promoter region	Possible SlyA binding sites	Location of site relative to start codon
<i>PcasA</i>	TTATTG <u>A</u> ATTAA	100 bp upstream of <i>casA</i>
<i>PssuE/elfA</i>	TCAGGAT <u>G</u> GATAA	8 bp upstream of <i>elfA</i>
<i>PgspC</i>	TTATATTAGTAA	79 bp upstream of <i>gspA</i>
<i>PpaaA</i>	TTAAATC <u>G</u> CGAA	239 bp upstream of <i>paaA</i>
	TTATA <u>A</u> AAATAG	136 bp upstream of <i>paaA</i>
	TTACTT <u>A</u> ACTAT	81 bp upstream of <i>paaA</i>
<i>PsgcX</i>	TTATGCT <u>G</u> GGAA	336 bp upstream of <i>sgcX</i>
	TTTCA <u>A</u> CCATAA	188 bp upstream of <i>sgcX</i>
<i>PfecI</i>	TTAG <u>A</u> AAAACAA	109 bp upstream of <i>fecI</i>
<i>PslyA</i>	TTAG <u>C</u> AAGCTAA	22 bp upstream of <i>slyA</i>
	TTAG <u>A</u> TTAATAA	161 bp upstream of <i>slyA</i>
<i>PleuO</i>	TTAATGCATTA <u>A</u>	305 bp upstream of <i>leuO</i>
	TTAAAT <u>A</u> TATAA	297 bp upstream of <i>leuO</i>
<i>PmdtM</i>	TATAC <u>A</u> CCTTAA	249 bp upstream of <i>mdtM</i>

500

501 Sequences shown are those with the greatest similarity to the previously proposed consensus for the  
 502 *Salmonella enterica* serovar Typhimurium LT2 SlyA protein (Haider *et al.* [9]; **TTAN<sub>6</sub>TAA**). Where  
 503 more than one possible site was present those with the greatest similarity to the consensus sequence  
 504 TTAGCAAGCTAA proposed by Stapleton *et al.* [10] are shown. Locations of sites are given as the  
 505 number of base pairs from the start codon of the specified gene to the centre of the proposed binding  
 506 site.

507

508 **Figure legends**

509

510 **FIG. 1.** Changes in the transcript profile of *E. coli* K-12 MG1655 over-producing SlyA. (a)  
511 Transformation of *E. coli* K-12 MG1655 with a multi-copy plasmid expressing *slyA* under the  
512 control of its own promoter results in detectable SlyA protein in lysed cell suspensions from  
513 aerobic steady-state glucose-limited chemostat cultures. The upper panel shows the  
514 Coomassie blue-stained SDS-polyacrylamide gel and the lower panel shows the relevant  
515 region of a Western blot prepared with the same samples and loadings developed with SlyA  
516 antiserum. The gels were loaded as follows: Lane M, SDS-PAGE markers (sizes, kDa, are  
517 indicated); lanes 1 and 2, extracts from independent cultures of *E. coli* K-12 MG1655  
518 transformed with the vector pET28a (SlyA<sup>WT</sup>); lanes 3 and 4, extracts from independent  
519 cultures of *E. coli* K-12 MG1655 transformed with the expression plasmid pGS2468 (SlyA<sup>+</sup>);  
520 lane 5, purified (His)<sub>6</sub>-SlyA (~10 ng protein loaded). (b) Western blot corresponding to the  
521 gel shown in (a). The locations of SlyA and purified (His)<sub>6</sub>-SlyA are indicated. (c) Graphical  
522 representation of the changes in transcript abundance occurring upon over-production of  
523 SlyA in *E. coli* K-12 MG1655. Comparison of the fold-changes in transcript abundance of  
524 aerobic steady-state glucose-limited chemostat cultures of *E. coli* K-12 MG1655 transformed  
525 with either the pET28a (SlyA<sup>WT</sup>) or pGS2468 (SlyA<sup>+</sup>). Each line represents a gene that  
526 exhibits a ≥2-fold change in transcript abundance ( $p \leq 0.05$ ) from two biological and two  
527 technical replicates i.e. four measurements.

528


529 **FIG. 2.** Biofilm formation by *E. coli* K-12 MG1655 is enhanced by elevated *slyA* expression.  
530 Wells containing M9 minimal medium with 20% w/v glucose as a carbon source were  
531 seeded with 1:10 inocula of overnight cultures and incubated at 37°C for 16 h. The OD<sub>600</sub> of  
532 the planktonic bacteria was measured before a biofilm assay was carried out. Values shown  
533 are the mean and standard deviation (n = 12) \*\*\* denotes  $p \leq 0.00001$  in a Student's *t*-test.


534

535 **FIG. 3.** Electrophoretic mobility shift assays show specific binding of SlyA to intergenic  
536 regions of selected operons. The dashed lines in the diagrams on the left indicate the DNA  
537 regions used in the EMSA shown on the right. The arrows indicate the polarity of the genes  
538 (names above the arrows). The numbers below the arrows representing genes are the fold-  
539 changes in transcript abundance observed upon over-production of SlyA (Table 2). SlyA-  
540 binding to the *hlyE-C0299* intergenic region has been reported previously [20]. For the  
541 EMSAs, biotin labelled intergenic DNA was prepared as described in the *Methods*. Labelled  
542 DNA was incubated with increasing concentrations of purified SlyA protein and protein-DNA  
543 complexes were separated by electrophoresis on native polyacrylamide gels. Lanes 1-8: 0, 1,  
544 5, 10, 50, 100, 200, 500 nM SlyA. The locations of the free DNA (D) and the SlyA-DNA

545 complexes (C) are indicated. Note that binding at the *sgcX* upstream region was only  
546 evident at the highest SlyA concentration tested and the complex (C) was located close to a  
547 contaminating DNA species.

548





*SlyA*<sup>WT</sup>

*SlyA*<sup>+</sup>



