



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/112160/>

Version: Accepted Version

Article:

Evison, S.E.F., Gallagher, J.D., Thompson, J.J.W. et al. (2016) Cuticular colour reflects underlying architecture and is affected by a limiting resource. *Journal of Insect Physiology*, 98. pp. 7-13. ISSN: 0022-1910

<https://doi.org/10.1016/j.jinsphys.2016.11.005>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Title: Cuticular colour reflects underlying architecture and is affected by a**
2 **limiting resource**

3

4

5 **Authors: Sophie E. F. Evison^{a*} Joe D. Gallagher^a, John J. W. Thompson^a, Michael T.**
6 **Siva-Jothy^a, Sophie A. O. Armitage^{a,1*}**

7

8 *a. University of Sheffield, Department of Animal and Plant Sciences, University of*
9 *Sheffield, Sheffield, South Yorkshire, S10 2TN, UK*

10 *1. Present address: Institute for Evolution and Biodiversity, University of Münster, 48149*
11 *Münster, Germany*

12 Email addresses:

13 j.d.gallagher@sheffield.ac.uk; jjwthompson@hotmail.com; m.siva-jothy@sheffield.ac.uk

14 ***Corresponding authors:** s.evison@sheffield.ac.uk; sophie.armitage@uni-muenster.de

15

16 **Abstract**

17 Central to the basis of ecological immunology are the ideas of costs and trade-offs between
18 immunity and life history traits. As a physical barrier, the insect cuticle provides a key
19 resistance trait, and *Tenebrio molitor* shows phenotypic variation in cuticular colour that
20 correlates with resistance to the entomopathogenic fungus *Metarhizium anisopliae*. Here we
21 first examined whether there is a relationship between cuticular colour variation and two
22 aspects of cuticular architecture that we hypothesised may influence resistance to fungal
23 invasion through the cuticle: its thickness and its porosity. Second, we tested the hypothesis
24 that tyrosine, a semi-essential amino acid required for immune defence and cuticular
25 melanisation and sclerotisation, can act as a limiting resource by supplementing the larval diet
26 and subsequently examining adult cuticular colouration and thickness. We found that stock
27 beetles and beetles artificially selected for extremes of cuticular colour had thicker less porous
28 cuticles when they were darker, and thinner more porous cuticles when they were lighter,
29 showing that colour co-varies with two architectural cuticular features. Experimental
30 supplementation of the larval diet with tyrosine led to the development of darker adult cuticle
31 and affected thickness in a sex-specific manner. However, it did not affect two immune traits.
32 The results of this study provide a mechanism for maintenance of cuticular colour variation in
33 this species of beetle; darker cuticles are thicker, but their production is potentially limited by
34 resource constraints and differential investments in resistance mechanisms between the sexes.

35

36

37 **Key words**

38 *Tenebrio molitor*; cuticular thickness; pore canal; fungal resistance; immunity; tyrosine

39

40

41 **Introduction**

42 Insect cuticle has a variety of functions, including skeletal support, camouflage,
43 thermoregulation, protection from ultraviolet light and sexual signalling (Wigglesworth,
44 1948). The cuticle is also a barrier to pathogens (Moret and Moreau, 2012) and is the direct
45 route of entry for some insect pathogenic fungi (e.g. some Entomophthoromycota and
46 Ascomycota; Boomsma et al., 2014). Resistance to important entomopathogens has been
47 related to the degree of cuticular darkness, where a darker cuticle is more resistant (e.g.
48 Barnes and Siva-Jothy, 2000; Dubovskiy et al., 2013; Mitsui and Kunimi, 1988; Reeson et al.,
49 1998; Wilson and Reeson, 1998; Wilson et al., 2001, 2002; Verhoog et al., 1996). Increased
50 resistance could be due to behavioural defences, but is more likely to be linked to physical or
51 chemical properties of the cuticle itself (Silva et al., 2016), and/or due to immune factors that
52 attack the pathogen once it is through the cuticle. The mechanical properties of the cuticle are
53 affected by many factors, including the degree of sclerotisation and melanisation, cuticular
54 thickness, protein composition, relative amounts of chitin and proteases, water content and
55 intracuticular pH (Andersen, 2010). All of these can influence the likelihood of invasion by
56 fungal pathogens (e.g. Pekrul and Grula, 1979; Lui et al., 2014; Schabel, 1978; St. Leger et al.,
57 1988). In addition, the cuticle is able to mount an active defence against pathogens (e.g.
58 Ashida and Brey, 1995; Golkar et al., 1993) through the action of proteases and chitinases that
59 prevent the attachment, penetration and degradation of the cuticle by fungal pathogens (e.g.
60 Kuo and Alexander, 1967; Söderhäll and Ajaxon, 1982; St. Leger et al., 1988). Several
61 studies have also found relationships between cuticular darkness and humoral (Reeson et al.,
62 1998; Wilson et al., 2002) and cellular (Cotter et al., 2004) immune function (but see Bailey,
63 2011; Karl et al., 2010).

64

65 In the mealworm beetle, *Tenebrio molitor*, adult cuticular colour is density dependent and can

66 range from ‘tan’ to ‘black’ (Thompson et al., 2002); larvae experiencing higher population
67 densities develop into adults with darker cuticles (Barnes and Siva-Jothy, 2000). This
68 response to ecological conditions also appears to have a heritable component (Prokkola et al.,
69 2013; Rolff et al., 2005), which is amenable to selection. The physiochemical properties of
70 the cuticle are known to vary with colour in *T. molitor* (Silva et al., 2016), and selection for a
71 darker cuticle has been shown to correlate with both higher activity of phenoloxidase (PO)
72 and increased haemocyte density (Armitage and Siva-Jothy, 2005). The process of cuticle
73 hardening and darkening involves (1) sclerotisation, which cross-links and stabilises the
74 cuticle by incorporating phenolic compounds, and (2) melanisation, which results in melanin
75 being distributed within the cuticle or deposited as granules (see Andersen (2010) for a
76 review). Central to both these processes is the production of 3,4-dihydroxyphenylalanine
77 (DOPA), which is produced by the hydroxylation of the semi-essential amino acid tyrosine
78 (Gorman et al., 2007; Moussian, 2010). As well as being important in cuticle production,
79 melanin is also involved in the encapsulation response against parasites within the body
80 cavity (Siva-Jothy et al., 2005). Consequently, tyrosinase POs are also found in haemocytes
81 and haemolymph as inactive pro-enzymes, and increase survival after infection with fungi and
82 Gram-positive bacteria (Binggeli et al., 2014), as well as being an important constitutive
83 immune effector in insects (Chase et al., 2000). Given the complex interactions between
84 internal physiology and cuticular melanisation and sclerotisation, the relationship between
85 cuticular colouration and immune defence might result from shared biochemical investment
86 via pleiotropic association (Silva et al., 2016). A comparatively little-studied feature of the
87 insect cuticle are the pore canals, the narrow ducts that perforate the cuticle perpendicular to
88 the surface, connecting the epidermis to the epicuticle (Locke, 1961; Zacharuk, 1976). Their
89 relationship with cuticular darkness and with cuticle-invading pathogens is unknown.
90 *Tenebrio molitor* has numerous pore canals in the exocuticle, running parallel to one another

91 (Delachambre, 1971; Wigglesworth, 1948). Renewed recent interest in pore canals in another
92 tenebrionid beetle, *Tribolium castaneum*, has shown that two cuticular proteins, TcCPR27 and
93 TcCPR18 are necessary for the correct formation of the pore canals in this species,
94 particularly in body regions where there is rigid cuticle such as the thoracic body wall (Noh et
95 al., 2014). Whilst a darker cuticle, which in *T. molitor* is fixed during early adulthood (e.g.
96 Thompson et al., 2002), confers an advantage in terms of parasite resistance (e.g. Barnes and
97 Siva-Jothy, 2000), it is likely to be coupled with costs in order for variation in cuticular colour
98 to be maintained within natural insect populations.

99

100 Resource constraints (e.g. protein and/or nitrogen acquisition) and competition for shared
101 resources by other physiological processes may result in trade-offs leading to phenotypic
102 variation in cuticular colour (Lee et al., 2008). Tyrosine is required for both immunity and
103 cuticular melanisation and sclerotisation, with the latter two processes occurring at each moult.
104 *Tenebrio molitor* larvae moult between 14 to 18 times (Morales Ramos et al., 2015; Park et
105 al., 2014), with loss of tyrosine *via* the exuvium at each moult (Andersen, 2004). Tyrosine can
106 be obtained from proteinaceous food, or synthesised directly from the essential amino acid
107 phenylalanine via phenylalanine hydroxylase (Chapman et al., 2013; Corrigan, 1970) but like
108 most animals, insects cannot synthesize the benzene ring in phenylalanine or tyrosine, thus
109 they must be sequestered via foraging. Given this, and the competing processes that require
110 tyrosine, it is likely to be a limiting resource, and so we hypothesise that it is an important
111 currency for trade-offs in insects.

112

113 In this study we first examine the relationship between cuticular architecture and colour in *T.*
114 *molitor*. We examine two aspects of cuticular architecture: exocuticular thickness and
115 cuticular ‘porosity’ (a compound-measure of pore-canal density and pore-canal lumen size).

116 We test whether these parameters are connected with variation in cuticle colour that is
117 generated both naturally (phenotypic response) and by selection (genetic response). We
118 predict that blacker (more resistant, see Barnes and Siva-Jothy, 2000) beetles will have a
119 thicker and/or less porous exocuticle compared to lighter beetles. Second, we address whether
120 dietary tyrosine influences cuticular colour and/or thickness. Given that tyrosine is likely to be
121 a limiting resource we predict that larval dietary supplementation will result in darker and
122 thicker adult cuticles. Finally, previous work found that non-immune challenged beetles
123 selected for a darker cuticle had a higher haemocyte load and increased PO levels (Armitage
124 and Siva-Jothy, 2005) we therefore also measured these two immune traits.

125

126

127 **Material and methods**

128 *1.1 Beetle culturing and colour selection*

129 Stock cultures were reared and maintained in an insectory at $26 \pm 2^\circ\text{C}$ in a LD 12:12h
130 photocycle. Their diet consisted of *ad libitum* access to water and rat diet (special
131 diets services: 77% cereal (wheat, maize, barley, wheatfeed), 15% vegetable proteins (soya
132 bean meal), 5% animal protein (fish meal) and 3% vitamins (major and trace) and amino
133 acids), with biweekly supplementation with apple chunks. Stock cultures contained 2000+
134 individuals at all stages of development kept in 25 x 30 x 50 cm tanks. Stock cultures were
135 initiated at least 8 generations (ca. 2 years) prior to the study by mixing individuals from four
136 cultures, and they were supplemented with adults from other stock cultures on a biannual
137 basis. Colour selected lines were produced as described in Armitage and Siva-Jothy (2005)
138 and were the result of selecting dorsal cuticular colour to be either dark (black) or light (tan).
139 To do this, virgin males and females from out-bred stock cultures with extremes of cuticular

140 phenotype (either black or tan) were allowed to mate monogamously to create the two
141 extremes of colour-selected lines (see supplementary materials for further details).

142

143 *1.2 Experiment 1: Is there a relationship between cuticle colour and cuticular architecture?*

144 To determine the relationship between cuticular colour and architecture, the thickness and
145 porosity of the cuticle of both stock and colour-selected beetles was correlated with its colour.

146 Beetles from the stock cultures and colour-selected lines were collected as pupae, their sex
147 determined, and kept in individual grid box cells. Only adult beetles that fell within a range of
148 0.100 – 0.110g were used. At between 7-10 days post imaginal eclosion cuticular colour was
149 assessed (cuticular colour ceases changing at 5.4 days after imaginal eclosion; Thompson et
150 al., 2002). All beetles were cold-anaesthetised on ice, weighed, and then a digital image of the
151 elytra was captured. From this image the elytra length was measured, and the degree of
152 cuticular darkness was analysed using Optimas 6® software to give a weighted average
153 luminance on a greyscale between 0 and 255 (0 = darkest, 255 = lightest; Thompson et al.,
154 2002).

155

156 *1.2.1 Measurement of exocuticular thickness*

157 Following colour determination, exocuticular thickness was determined for a subset of
158 beetles. Fifty-two beetles were taken from the stock culture (26 of each sex) as well as a
159 single female from each of 10 black-selected and 10 tan-selected lines, and a male from 9 of
160 the black-selected and 9 of the tan-selected lines (20 females, 18 males). While remaining
161 cold-anaesthetised on ice, the abdomen and head were excised with scissors to leave the
162 pronotum (Fig. S1a, 1). The cuticle from the dorsal side of the pronotum was cut away using
163 bow spring scissors and placed under a drop of water on a Petri dish and examined with a
164 stereo microscope (Leica MZ8). Fatty tissue was gently scraped away from the interior side of

165 the cuticle with a scalpel. The piece of cuticle was then sectioned into four (Fig. S1a, 2) and
166 three of the resulting quarters were further sectioned into wedge-shaped pieces (Fig. S1a, 3).
167 Three triangular wedges were placed on their sides and inserted into a piece of Blu-tack™ on
168 a glass slide, with the thin edge of the wedge of cuticle protruding (Fig. S1a, 4). The slide was
169 placed under a light microscope (Leitz Diaplan) and the thickness of the exocuticle (the
170 opaque, coloured layer including the epicuticle) at the thin edge of the wedge was drawn
171 using the camera lucida and measured using a pair of digital callipers (Mitutoyo, Digimatic).
172 This sampling method demonstrated repeatability at both the level of the individual using
173 three wedges of cuticle taken from the same beetle ($F_{51,104} = 7.32$, $P < 0.001$, $r = 0.69$), and at
174 the level of the drawing by making three repeated drawings of the same wedge of cuticle
175 ($F_{19,40} = 13.95$, $P < 0.001$, $r = 0.81$).

176

177 *1.2.2 Measurement of cuticular porosity*

178 Following colour determination, cuticular porosity was determined for a different subset of
179 beetles. Eleven male beetles were taken from the stock culture beetles as well as a single male
180 and female from each of 5 black-selected and 5 tan-selected lines (10 females, 10 males).
181 While remaining cold-anaesthetised on ice, the dorsal thoracic cuticle was dissected out and
182 stored in 2.5% glutaraldehyde in sodium cacodylate buffer (0.01M sodium cacodylate,
183 0.005M calcium chloride; pH 7.4). Each fixed cuticle slice was prepared for transmission
184 electron microscopy (TEM) by embedding it in resin. Ultrathin (70-90nm thick) sections were
185 taken transversely to the long axis of the pore canals (Reichert Ultracut E ultramicrotome),
186 and stained with 3% uranyl acetate and then Reynold's lead citrate. The sections were
187 mounted on copper grids and examined at 80Kv in a Transmission Electron Microscope
188 (Philips CM10). The cuticle was examined for pore canals: once identified (Wigglesworth,
189 1985; Reynolds, 2005, Pers. Comm.), three photomicrographs were taken of sections from

190 each beetle. The photomicrographs were taken approximately one full field of view away
191 from the edge of the section, at a magnification of 6600× (Fig. S1b). Pore canals were traced
192 onto an acetate sheet placed over the photomicrograph and the outline of each of the canals
193 was defined with a fine-tipped permanent marker (Fig. S1b). To ensure consistency in the
194 area drawn, another acetate with an outlined area 14 cm × 10 cm with markers denoting the
195 edges of the photomicrograph (16.5 cm × 21.5 cm), was sandwiched between the
196 photomicrograph and the acetate. The drawing was then assessed from a digitized image
197 obtained from a video camera (PULNiX TM/765) and analyzed using Optimas 6[®] software.
198 Analysis using Optimas 6[®] involved calibrating the drawing and setting a threshold
199 incorporating all of the pore canals; the pores could then be highlighted and data collected on
200 the number of pores and the total area of pores (porosity) in the region of interest. The total
201 area assessed in each photomicrograph was 74 μm². This sampling method demonstrated a
202 reasonable repeatability at both the level of the individual using photomicrographs of cuticle
203 from the same beetle (pore canal number: $F_{df=28,58} = 10.88$, $P < 0.001$, $r = 0.77$; total pore
204 canal area (porosity): $F_{df=28,58} = 8.37$, $P < 0.001$, $r = 0.71$), and at the level of the drawing
205 using the same photomicrograph (pore canal number: $F_{df=19,20} = 226.16$, $P < 0.001$, $r = 0.99$;
206 porosity: $F_{df=19,20} = 39.08$, $P < 0.001$, $r = 0.95$). Repeated temporal measures from an
207 individual could not be performed because the assay is destructive.

208

209 *1.3 Experiment 2: Is tyrosine a limiting resource?*

210 In a second experiment we tested whether dietary tyrosine influenced cuticular colour,
211 cuticular thickness, and immunity related parameters. We placed ten stock beetles (5 of each
212 sex) in a box with plain flour, with *ad libitum* access to apple and water, and allowed them to
213 mate. The stock beetles had a relatively dark cuticular colour that was closer to that of black-
214 selected lines than tan-selected lines (Thompson et al., 2002). 450 of the resulting larvae were

215 distributed between 45 boxes (10 larvae per box) and kept in an incubator at $26 \pm 1^\circ\text{C}$. The 45
216 boxes were distributed across three treatment groups: 15 boxes contained 60g white wheat
217 flour with a tyrosine supplement, 15 boxes contained 60g white wheat flour with a tryptophan
218 supplement, and 15 boxes contained 60g white wheat flour only. Both white wheat flour and
219 apple are low in tyrosine, phenylalanine and tryptophan (wheat flour (USDA Nutrient
220 Database Number: 20481): 0.312g, 0.520g, 0.127g per 100g respectively; apple (USDA
221 Nutrient Database Numbers: 9004/9014): 0.005g, 0.007g, 0.001g per 100g respectively) thus
222 allowing the experimental regulation of their levels in this study. Tryptophan was the control
223 treatment. It is a semi-essential amino acid involved in pigment production; it is converted to
224 ommochromes, a class of body and eye pigment widespread in arthropods and molluscs
225 (Corrigan, 1970), but it is not involved in melanisation or sclerotisation. 0.3g per 100g of each
226 of the two amino acids were supplemented to the flour at the start of the experiment; thus
227 roughly doubling the dose compared to the control. All boxes were examined weekly for
228 pupae. Collected pupae were sexed, weighed and kept in a grid box in individual cells. Adults
229 were not given access to food, so as not to confound feeding effects from the larval stage.
230 Eight days after imaginal eclosion the beetles were weighed again, elytra length was
231 measured using calipers (Digimatic, Mitutoyo) and cuticular colour was analysed as described
232 above (sample sizes: Control: Female (F) = 26, Male (M) = 16; Tryptophan: F = 29, M = 23;
233 Tyrosine: F = 21, M = 26). Beetles remained cold-anaesthetised on ice and a 1mm length of
234 sterile nylon monofilament (Sunline, fil classe I.G.F.A., Siglon V transparent; 0.128mm
235 diameter) was inserted through the pleural membrane between the 3rd and 4th sternite to
236 induce an immune challenge. Twenty-four hours after the immune challenge, beetles were
237 assayed for exocuticular thickness as described in the supplementary materials (sample sizes
238 = Control: F = 22, M = 15; Tryptophan: F = 23, M = 17; Tyrosine: F = 19, M = 19). In the

239 same beetles, haemocyte numbers were counted and PO measurements taken as described in
240 the supplementary materials.

241

242 *1.4 Statistical analyses*

243 All data was analysed using R statistical software (R Core Development Team, 2015). There
244 was a highly significant relationship between adult weight and elytra length across all three
245 types of beetle (stock, colour-selected, and experimentally-fed; Pearson's correlation: $t_{233} =$
246 10.8 , $P < 0.001$, $r = 0.578$): consequently, weight was used as an indicator of size, and
247 included as a covariate in any analyses where it significantly improved the model. All models
248 were checked visually for normality of the residuals and homogeneity of variances (Faraway,
249 2006) and P values were generated for the mixed models using the likelihood ratio test.

250

251 *1.4.1 Experiment 1:* The relationship between colour and exocuticular thickness in stock
252 beetles was analysed using a general linear model with a Gaussian error distribution and
253 identity link function. In the colour-selected beetles, the relationship between exocuticular
254 thickness and their selection colour (i.e. a fixed factor with two levels, black and tan), their
255 sex, and the interaction between the two, were analysed using a linear mixed effects models
256 implemented using the *lmer* function from the *lme4* package (Bates and Maechler, 2009),
257 with selection line included as the random effect to account for their shared rearing
258 environment. For both stock and colour-selected beetles, the relationship between colour and
259 all three measures of porosity (mean pore number, mean area per pore and mean total pore
260 area) were analysed using general linear models with a Gaussian error distribution and
261 identity link function. Again, for the colour-selected beetles, their selection colour as a fixed
262 factor, their sex, and the interaction between the two were assessed, whereas for the stock
263 beetles, only cuticular colour as a continuous variable was assessed because only males were

264 assessed for porosity in the stock beetles. Weight was not significant in any case so was
265 removed from the models.

266

267 *1.4.2 Experiment 2:* Each of the measured variables in the experimentally-fed beetles;
268 cuticular colour and thickness, PO level, and log transformed haemocyte counts, were
269 analysed using linear mixed effects models fitted with the *lmer* function as described above,
270 but here the fixed factors were experimental treatment and sex, as well as their interaction,
271 and the random effect was box number to account for the shared rearing environment. There
272 was a significant overall negative effect of adult weight on cuticle colour ($\chi^2_1 = 6.18, P =$
273 0.013) whereby heavier beetles had darker cuticles, and a positive effect on cuticle thickness
274 ($\chi^2_1 = 8.32, P = 0.004$) whereby heavier beetles had thicker cuticles, so in each of these
275 analyses weight was kept in the model. Additionally, the same models were run to check for
276 relationships between the measured PO level, and log transformed haemocyte counts as the
277 response variable and the beetles' cuticular colour and thickness as the predictor variables for
278 all of the experimentally fed beetles. Treatment was kept in each model as a covariate, and
279 adult weight was significant in each of the models so was retained. However, sex was not
280 significant for any model so was removed.

281

282

283 **Results**

284 *2.1 Experiment 1: the correlation between cuticular colour and architecture*

285 There was a significant relationship between cuticular colour and exocuticular thickness in
286 both stock and colour-selected beetles. In the stock beetles, there was a significant
287 relationship between colour and exocuticular thickness ($t_{48} = 3.27, P = 0.002$), with darker
288 beetles having thicker cuticles (Fig. 1a). There was no effect of sex ($t_{48} = 0.556, P = 0.581$),

289 and no interaction between colour and sex ($t_{48} = 0.559, P = 0.579$). In the colour-selected
290 beetles, there was a significant effect of selection colour on the exocuticular thickness of the
291 beetles ($\chi^2_1 = 19.8, P < 0.001$), with beetles from lines selected for black cuticular colour
292 having thicker cuticles than those from tan-selected lines (Fig. 1b). Although there was no
293 main effect of sex ($\chi^2_1 = 0.035, P = 0.851$) on exocuticular thickness, there was a significant
294 interaction between colour of line and sex ($\chi^2_1 = 5.03, P = 0.025$), suggesting that males
295 responded more strongly to colour selection in terms of their cuticular thickness (Fig. 1b).


296

297 There was a relationship between cuticular colour and porosity in both the stock beetles and
298 the colour-selected beetles. In stock beetles, there was a significant positive relationship
299 between cuticular porosity (mean total pore area) and cuticular colour ($t_9 = 3.24, P = 0.010$;
300 Fig. 1c), with lighter beetles, i.e. those with a tan cuticle, having more porous cuticle.

301 However, there was no relationship between cuticular colour and either pore number ($t_9 =$
302 $1.51, P = 0.166$) or mean area per pore ($t_9 = 0.069, P = 0.946$). Similarly, beetles selected for
303 a tan cuticle had a significantly higher porosity than black-selected beetles ($t_{18} = 3.31, P =$
304 0.004 ; fig 1d). Again, there was no main effect of sex upon porosity ($t_{17} = 1.74, P = 0.101$)
305 and no significant interaction between cuticle colour and sex ($t_{16} = 1.88, P = 0.078$). Neither
306 the mean pore number nor the mean area per pore differed according to selection colour (pore
307 number: $t_{18} = 0.553, P = 0.588$; area per pore: $t_{18} = 0.553, P = 0.588$). However, there was a
308 significant difference in mean area per pore between the sexes ($t_{17} = 2.24, P = 0.039$) with
309 males having a larger mean area per pore than females, but no difference in mean pore
310 number ($t_{17} = 1.27, P = 0.220$) between the sexes. In neither case was there an interaction
311 between colour and sex (pore number: $t_{16} = 1.09, P = 0.292$; area per pore: $t_{16} = 0.30, P =$
312 0.767).

313

314 **Figure 1.** The relationship between cuticular colour (note that lower numbers denote a darker
 315 cuticle) and a) exocuticular thickness for female and male beetles collected from the stock
 316 culture, b) mean (\pm s.e.) exocuticular thickness for males and females from black and tan
 317 colour-selected lines, c) porosity (mean total pore area) for male beetles collected from the
 318 stock culture, and d) mean (\pm s.e.) porosity for males and females from black and tan colour-
 319 selected lines.



320

321

322 2.2 Experiment 2: The effect of tyrosine on cuticular colour

323 There was a significant effect of experimental diet on cuticular colour ($\chi^2_1 = 7.55$, $P = 0.023$),


324 with beetles from the tyrosine supplemented diet having darker cuticles (Fig. 2a). There was

325 no effect of sex ($\chi^2_1 = 2.37, P = 0.123$), and no interaction between sex and treatment ($\chi^2_1 =$
326 $0.448, P = 0.799$). There was no effect of treatment on cuticular thickness ($\chi^2_1 = 4.67, P =$
327 0.097) but there was a significant effect of sex ($\chi^2_1 = 4.86, P = 0.028$), as well as a significant
328 interaction between sex and treatment ($\chi^2_1 = 15.3, P < 0.001$), suggesting females responded
329 more to tyrosine in terms of cuticular thickness than males did (Fig. 2b).

330

331 **Figure 2.** The effect of dietary tyrosine supplementation on a) mean (\pm s.e.) cuticular colour
332 (note that lower numbers denote a darker cuticle), and b) mean (\pm s.e.) exocuticular thickness
333 for males and females from the experimentally-fed beetles.

334



335

336

337 There were no effects of treatment or sex, or their interactions for either of the immune
 338 measures (i.e. average haemocyte count and PO level after nylon insertion; table S1).

339 However, when experimental treatment and weight were accounted for, there was a

340 significant positive relationship between cuticular thickness and PO activity ($\chi^2_1 = 5.09$, $P =$
341 0.024; Fig. 3); beetles with thicker cuticles had higher PO activity. Conversely, there was no
342 relationship between cuticular colour and PO activity ($\chi^2_1 = 0.41$, $P = 0.522$), nor was there a
343 relationship between cuticular colour ($\chi^2_1 = 2.86$, $P = 0.091$) or cuticular thickness ($\chi^2_1 =$
344 0.74, $P = 0.390$) and average haemocyte count.


345

346 **Figure 3.**

347 The relationship between exocuticular thickness and PO activity (dashed line), pooling
348 individuals across all feeding treatments, i.e., control (squares), tryptophan-supplemented
349 (circles) and tyrosine-supplemented (triangles) diets.

350

351



352

353

354

355 **Discussion**

356 Our results show a clear relationship between the colour and architecture of *T. molitor* cuticle,
357 and suggest there are phenotypic and genetic correlations between cuticular colour and
358 cuticular architecture. Both stock beetles and beetles selected for extremes of cuticular colour
359 had thicker less porous cuticles when they were darker, and thinner more porous cuticles
360 when they were lighter. There appeared to be a sex difference in selection for a darker cuticle;
361 the cuticles of males were thicker than of females in dark selected beetles. Supplementation of
362 the larval diet with tyrosine led to the development of a darker cuticle in adults, and again
363 there were sex specific effects whereby females had thicker cuticles than males after tyrosine
364 supplementation.

365

366 Black beetles had a thicker and a less porous exocuticle than tan beetles. This was true for
367 male and female beetles derived from stock cultures. It also applied to lines selected for
368 cuticular colour (although only male porosity was assessed here), suggesting that increasing
369 exocuticular thickness and porosity is a correlated genetic response to selection for darker
370 cuticular colour. Our results for cuticular thickness are corroborated by a recent study (Silva
371 et al., 2016), which demonstrated that all the cuticle layers, including the endocuticle, of *T.*
372 *molitor* cuticle were thicker in blacker adult beetles than in tan stock beetles; the authors
373 examined abdominal sternites, suggesting that our results can be generalised beyond the
374 pronotum. The consistency of both our results and those of Silva et al. (2016), despite
375 measurement of different layers also suggest that the exocuticle, measured in our study, is the
376 cuticular structure driving the relationship between colour and thickness. A similar
377 phenomenon has been found in larvae of the greater wax moth, *Galleria mellonella*
378 (Dubovskiy et al. 2013), whereby a melanic (dark) morph had a thicker cuticle than a non-

379 melanic morph. However, we are not aware of other studies where porosity has been
380 examined in relation to cuticular colour. In our study, males appeared to be more constrained
381 in reducing porosity after selection for a darker cuticle than females, perhaps reflecting an
382 evolutionary constraint on the reduction of pore canal size in males. Even though there was no
383 significant interaction between sex and mean pore area, males had a larger area per pore in the
384 dark-selected beetles compared to tan. *Tenebrio molitor* has numerous parallel pore canals in
385 the exocuticle (Delachambre 1971; Wigglesworth, 1948; 1985). Two proteins are important
386 for the proper formation of the pore canals and horizontal chitinous laminae in the cuticle of
387 another tenebrionid beetle, *Tribolium castaneum* (Noh et al., 2014); knockdown of the genes
388 led to beetles with less rigid elytra. However because the functional significance of pore
389 canals is not fully understood, it is not clear whether they can influence invasion by fungal
390 pathogens. Nonetheless it is intriguing that porosity co-varies with cuticular colour. Although
391 a direct test is currently lacking, we hypothesise that one explanation for the differential
392 resistance between dark and tan coloured beetles (Barnes and Siva-Jothy, 2000; Prokkola et
393 al., 2013) could be the correlated differences in cuticular architecture.

394

395 Despite the relationships between cuticular colour and architecture, our data does not allow
396 inference as to whether the patterns we found are driven by the process of melanisation, or
397 whether cuticular colour is a consequence of increased density/number of lamellae leading to
398 higher sclerotisation. On one hand, cuticular colour could depend upon the thickness of the
399 cuticle considering that beetles selected for darker cuticular colour showed a correlated
400 increase in cuticular thickness, suggesting that selection was on exocuticular thickness itself
401 rather than on the phenotypic expression of exocuticular thickness (i.e. colour). On the other
402 hand, the melanin content of sternites from black *T. molitor* has been shown to be higher than
403 that of tan beetles (Silva et al., 2016) suggesting that at least some of the colour difference in

404 our colour-selected lines and stock cultures could be due to melanin. Melanin does not seem
405 to be involved in sclerotisation of the elytra in another tenebrionid beetle species, *Tribolium*
406 *castaneum*, as its deposition does not increase resistance of the beetle's cuticle to puncture
407 (Roseland et al., 1987). Indeed, black mutants of *T. castaneum* and the cockroach *Blatella*
408 *germanica* divert more dopamine into the cuticular melanisation pathway, reducing the
409 substrate available for cross-linking and thus actually weakening the cuticle (Czapla et al.,
410 1990; Roseland et al., 1987). It would be interesting to see if these mutants were less resistant
411 to entomopathogens.

412

413 Dietary manipulation revealed tyrosine as an important factor determining cuticular colour;
414 tyrosine-fed beetles had significantly darker cuticles than both tryptophan- and control-fed
415 beetles. Interestingly, diet-induced darkening of the cuticle was accompanied by a thicker
416 cuticle only in female beetles. Male beetles had the same cuticular thickness, irrespective of
417 feeding treatment. Considering that diet-determined cuticular colour change was not sex-
418 dependent, but diet-determined cuticular thickness was, this suggests that females may
419 allocate their resources differently to males. This could also explain the differential response
420 between the sexes to colour selection, because under standard food conditions in the first
421 experiment, female cuticular thickness responded less to selection for cuticular darkness than
422 males, supporting a differential allocation of resources between the sexes here too. Although
423 the tyrosine supplement resulted in darker cuticles, darker cuticle did not correlate with
424 increased haemocyte density. Given the positive correlations that have been found between
425 the degree of cuticular melanisation and expression of physiological immune effectors (e.g.
426 Cotter et al., 2004; Armitage and Siva-Jothy, 2005; Reeson et al., 1998; Wilson et al., 2001),
427 it is perhaps surprising that there were no clear differences between the experimentally fed
428 treatments in the immune measures made here. In common with Armitage & Siva-Jothy

429 (2005), we found no difference in PO between the black and tan beetles after immune
430 challenge. Differences in the results between previous studies and our results could be
431 explained by methodology; for example, in this study haemocytes were counted after the
432 insertion of a nylon monofilament as an immune challenge, whereas other studies assayed
433 haemocytes without an immune challenge (e.g. Cotter et al., 2004). There could alternatively
434 be correlated immune responses to selection on cuticular colour that are not related to tyrosine
435 and that cuticular colour can be uncoupled from internal physiological defences. Interestingly,
436 there was a positive relationship between PO activity and cuticular thickness when all of the
437 experimentally fed beetles were pooled, supporting a relationship between humoral and
438 structural immunity. This positive relationship highlights the complex associations between
439 immune responsiveness, cuticular architecture, and the degree of cuticular darkness.

440 Pleiotropic interplay between cuticular melanisation and several other resistance traits have
441 been shown in *Galleria mellonella*, including a thickened cuticle and higher basal expression
442 of immunity (Dubovskiy et al., 2013), and these traits also correlate with higher resistance to
443 fungal entomopathogens. Tyrosine has the potential to be a limiting resource across other
444 taxa. For example, supplementing the diet of kittens with tyrosine resulted in black hair
445 compared to controls that had reddish-brown hair (Morris et al., 2002; see also Anderson et
446 al., 2002), and under a reduced phenylalanine and tyrosine dietary allowance, the black
447 melanin-containing feathers of the bib of juveniles of the male house sparrow, *Passer*
448 *domesticus*, which function as a sexual signal in adult males (Viega, 1993) were lighter in
449 colour (Poston et al., 2005).

450

451 Cuticular melanisation and sclerotisation share several of the same substrates and enzymes,
452 but are two distinct processes that may be independently regulated. Melanins interact with
453 other cuticular macromolecules, but appear to contribute little to cuticular sclerotisation

454 (Hopkins et al., 1984). During sclerotisation, tyrosine undergoes hydroxylation to form
455 dopamine derivatives, which are transported to the cuticular matrix, where inter-protein cross-
456 links and polymers form which thicken and strengthen the cuticle (Andersen, 2010).
457 However, if dopamine is present in the cuticle during the sclerotisation process, melanisation
458 should occur concurrent with sclerotisation (Andersen, 2010), leading to the formation of both
459 a thicker and a darker cuticle, i.e. an interactive process. Thus, cuticular colour appears likely
460 to be a consequence of cuticular thickness, and selection for increased resistance could act
461 directly upon cuticular thickness in *T. molitor*. How pore formation fits into this process is
462 less clear, and requires further study. Melanism has been associated with increased dopamine
463 levels in *Drosophila melanogaster*, and the metabolism of dopamine is a major source of
464 reactive oxygen species (ROS), which contributes to early senescence (Vermeulen et al.,
465 2006). There may also be resource-based costs of preventing damage caused by high levels of
466 PO/proPO, which are correlated with high levels of cuticular melanisation (e.g. serine
467 protease inhibitors). There are clearly complex trade-offs associated with insect cuticular
468 properties and its role in immunity. However, the results of this study provide a mechanism
469 for maintenance of cuticular colour variation in this species of beetle; darker cuticles are
470 thicker, but their production is potentially limited by resource constraints and differential
471 investments in resistance mechanisms between the sexes.

472

473

474 **Acknowledgements**

475 We thank John Proctor for preparing the samples for transmission electron microscopy, and
476 Jens Rolff for help during the preparation of a previous version of the manuscript. This work
477 was supported by the Leverhulme Trust (SEFE), the University of Sheffield (JDG, JJWT),
478 and NERC (MS-J, SAOA) [grant number GR3/12121].

479

480

481 **References**

482 1. Andersen, S.O. 2004. Chlorinated tyrosine derivatives in insect cuticle. *Insect*
483 *Biochemistry and Molecular Biology*, 34, 1079-1087.

484

485 2. Andersen, S.O. 2010. Insect cuticular sclerotization: a review. *Insect Biochemistry*
486 *and Molecular Biology*, 40, 166-178.

487

488 3. Anderson, P.J., Rogers, Q.R., Morris, J.G. 2002. Cats require more dietary
489 phenylalanine or tyrosine for melanin deposition in hair than for maximal growth. *The*
490 *Journal of Nutrition*, 132, 2037-2042.

491

492 4. Armitage, S.A.O., Siva-Jothy, M.T. 2005. Immune function responds to selection for
493 cuticular colour in *Tenebrio molitor*. *Heredity*, 94, 650–656.

494

495 5. Ashida, M., Brey, P.T. 1995. Role of the integument in insect defense: pro-phenol
496 oxidase cascade in the cuticular matrix. *Proceedings of the National Academy of*
497 *Sciences of the U.S.A.*, 92, 10698-10702.

498

- 499 6. Bailey, N. 2011. A test of the relationship between cuticular melanism and immune
500 function in wild-caught Mormon crickets. *Physiological Entomology*, 36, 155-164.
501
- 502 7. Barnes, A.I., Siva-Jothy, M.T. 2000. Density-dependent prophylaxis in the mealworm
503 beetle *Tenebrio molitor* L. (Coleoptera: Tenebrionidae): cuticular melanization is an
504 indicator of investment in immunity. *Proceedings of the Royal Society of London*
505 *Series B Biological Sciences*, 267, 177-182.
506
- 507 8. Bates, D., Maechler, M. 2010. lme4: linear mixed effects models using S4 classes.
508 Available at: <http://cran.r-project.org/web/packages/lme4/>
509
- 510 9. Binggeli O., Neyen C., Poidevin M., Lemaitre B. 2014. Prophenoloxidase activation is
511 required for survival to microbial infections in *Drosophila*. *PLoS Pathogens*, 10,
512 e1004067.
513
- 514 10. Boomsma, J.J., Jensen, A.B., Meyling, N.V., and Eilenberg, J. 2014. Evolutionary
515 interaction networks of insect pathogenic fungi. *Annual Review of Entomology*, 59,
516 467-485.
517
- 518 11. Chapman, R.F. (author), Simpson, S.J., Douglas, A.E. (Eds.) 2013. *The insects:*
519 *structure and function*. Cambridge University Press, New York.
520
- 521 12. Chase, M., Raina, K., Bruno, J., Sugumaran, M. 2000. Purification, characterization
522 and molecular cloning of prophenoloxidase from *Sarcophaga bullata*. *Insect*
523 *Biochemistry and Molecular Biology*, 30, 953–967.

- 524
- 525 13. Corrigan, J.J. 1970. Nitrogen metabolism in insects, in: Campbell, J.W. (Ed.)
526 Comparative Biochemistry of Nitrogen Metabolism. Academic Press, New York.
527
- 528 14. Cotter, S.C., Kruuk, L.E.B., Wilson, K. 2004. Costs of resistance: genetic correlations
529 and potential trade-offs in an insect immune System. Journal of Evolutionary Biology,
530 17, 421–429.
531
- 532 15. Czapla, T.H., Hopkins, T.L., Kramer, K.J. 1990. Cuticular strength and pigmentation
533 of five strains of adult *Blattella germanica* (L.) during sclerotization: correlations with
534 catecholamines, β -alanine and food deprivation. Journal of Insect Physiology, 36,
535 647–654.
536
- 537 16. Delachambre, J. 1971. La formation des canaux cuticulaires chez l'adult de *Tenebrio*
538 *molitor* L. étude ultrastructurale et remarques histochimiques. Tissue and Cell, 3, 499-
539 520.
540
- 541 17. Dubovskiy, I.M., Whitten, M.M.A., Yaroslavtseva, O.N., Grizanova, E.V., Greig, C.,
542 et al. 2013. More than a colour change: insect melanism, disease resistance and
543 fecundity. Proceedings of the Royal Society of London Series B Biological Sciences,
544 280, 20130584.
545
- 546 18. Faraway, J.J. 2006. Extending the Linear Model with R: Generalised Linear Mixed
547 Effects and Nonparametric Regression Models. Chapman and Hall/CRC, Boca Raton.
548

- 549 19. Golkar, L., LeBrun, R.A., Ohayon, H., Gounon, P., Papierok, B., Brey, P.T., 1993.
550 Variation of larval susceptibility to *Lagenidium giganteum* in three mosquito species.
551 Journal of Invertebrate Pathology, 62, 1–8.
552
- 553 20. Gorman, M.J., An, C., Kanost, M.R. 2007. Characterization of tyrosine hydroxylase
554 from *Manduca sexta*. Insect Biochemistry and Molecular Biology, 37, 1327–1337.
555
- 556 21. Hopkins, T.L., Morgan, T.D., Kramer, K.J., 1984. Catecholamines in haemolymph
557 and cuticle during larval, pupal and adult development of *Manduca sexta* (L.). Insect
558 Biochemistry, 14, 533-540.
559
- 560 22. Karl, I., Hoffmann, K.H., and Fischer, K. 2010. Cuticular melanisation and immune
561 response in a butterfly: local adaptation and lack of correlation. Ecological
562 Entomology, 35, 523-528.
563
- 564 23. Kuo, M.J., Alexander, M. 1967. Inhibition of the lysis of fungi by melanins. Journal of
565 Bacteriology, 94, 624-629.
566
- 567 24. Lee, K.P., Simpson, S.J., Wilson, K. 2008. Dietary protein-quality influences
568 melanization and immune function in an insect. Functional Ecology, 22, 1052-1061.
569
- 570 25. Locke, M. 1961. Pore canals and related structures in insect cuticle. The Journal of
571 Biophysical and Biochemical Cytology, 10, 589-618.
572

- 573 26. Lui, W., Xie, Y., Dong, J., Xue, J., Zhang, Y., Lu, Y., Wu, J. 2014. Pathogenicity of
574 three entomopathogenic fungi to *Matsucoccus matsumurae*. PLoS ONE, 9, e103350.
575
- 576 27. Mitsui, J., Kunimi, Y. 1988. Effect of larval phase on susceptibility of the armyworm,
577 *Pseudaletia separate* Walker (Lepidoptera, Noctuidae) to an entomogeneous
578 Deuteromycete, *Nomuraea rileyi*. Japanese Journal of Applied Entomology and
579 Zoology, 32, 129-134.
580
- 581 28. Morales Ramos, J.A., Kay, S., Rojas, M.G., Shapiro Ilan, D.I., Tedders, W.T. 2015.
582 Morphometric analysis of instar variation in *Tenebrio molitor* (Coleoptera:
583 Tenebrionidae). Annals of the Entomological Society of America, 108, 146-159.
584
- 585 29. Moret, Y., Moreau, J. 2012 The immune role of the arthropod exoskeleton.
586 Invertebrate Survival Journal, 9, 200-206.
587
- 588 30. Morris, J.G., Shiguang, Y., Rogers, Q.R. 2002. Red hair in black cats is reversed by
589 addition of tyrosine to the diet. The Journal of Nutrition, 132, 16465-16485.
590
- 591 31. Moussian, B. 2010. Recent advances in understanding mechanisms of insect cuticle
592 differentiation. Insect Biochemistry and Molecular Biology, 40, 363-375.
593
- 594 32. Noh, M.Y., Kramer, K.J., Muthukrishnan, S., Kanost, M.R., Beeman, R.W., Arakane,
595 Y. 2014. Two major cuticular proteins are required for assembly of horizontal laminae
596 and vertical pore canals in rigid cuticle of *Tribolium castaneum*. Insect Biochemistry
597 and Molecular Biology, 53, 22-29.

598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621

33. Park, J.B., Choi, W.H., Kim, S.H., Jin, H.J., Han, Y.S., Lee, Y.S., Kim, N.J. 2014. Developmental characteristics of *Tenebrio molitor* larvae (Coleoptera: Tenebrionidae) in different instars. *International Journal of Industrial Entomology*, 28, 5-9.
34. Pekrul, S., Grula, E.A. 1979. Mode of infection of the Corn Earworm (*Heliothis zea*) by *Beauveria bassiana*. *Journal of Invertebrate Pathology*, 70, 322-330.
35. Poston, J.P., Hasselquist, D., Stewart, I.R.K., Westneat, D.F. 2005. Dietary amino acids influence plumage traits and immune responses of male house sparrows, *Passer domesticus*, but not as expected. *Animal Behaviour*, 70, 1171–1181.
36. Prokkola, J., Roff, D., Kärkkäinen, T., Krams, I., Rantala, M.J. 2013. Genetic and phenotypic relationships between immune defense, melanism and life-history traits at different temperatures and sexes in *Tenebrio molitor*. *Heredity*, 111, 89-96.
37. R Core Development Team, 2015. R: a language and environment for statistical computing. R foundation for statistical computing, Vienna.
38. Reeson, A.F., Wilson, K., Gunn, A., Hails, R.S., Goulson, D. 1998. Baculovirus resistance in the noctuid *Spodoptera exempta* is phenotypically plastic and responds to population density. *Proceedings of the Royal Society of London Series B Biological Sciences*, 265, 1787–1791.

- 622 39. Rolff, J., Armitage, S.A.O., Coltman, D.W. 2005. Genetic constraints and sexual
623 dimorphism in immune defense. *Evolution*, 59, 1844-1850.
624
- 625 40. Roseland, C.R., Kramer, K.J., Hopkins, T.L. 1987. Cuticular strength and
626 pigmentation of rust-red and black strains of *Tribolium castaneum*: correlation with
627 catecholamine and β -alanine content. *Insect Biochemistry*, 17, 21-28.
628
- 629 41. Schabel, H.G. 1978. Percutaneous infection of *Hylobius pales* by *Metarhizium*
630 *anisopliae*. *Journal of Invertebrate Pathology*, 31, 180-187.
631
- 632 42. Silva, F.W.S., Araujo, L.S., Azevedo, D.O., Serrão, J.E., Elliot, S.L. 2016. Physical
633 and chemical properties of primary defences in *Tenebrio molitor*. *Physiological*
634 *Entomology*, 41, 121-126.
635
- 636 43. Siva-Jothy, M.T., Moret, Y., Rolff, J. 2005. Insect immunity: an evolutionary ecology
637 perspective. *Advances in Insect Physiology*, 32, 1-48.
638
- 639 44. Söderhäll, K., Ajaxon, R. 1982. Effect of quinones and melanin on mycelial growth of
640 *Aphanomyces* spp. and extracellular protease of *Aphanomyces astaci*, a parasite on
641 crayfish. *Journal of Invertebrate Pathology*, 39, 105-109.
642
- 643 45. St. Leger, R., Durrands, P.K., Charnley, K., Cooper, R. 1988. Role of extracellular
644 chymoelastase in the virulence of *Metarhizium anisopliae* for *Manduca sexta*. *Journal*
645 *of Invertebrate Pathology*, 52, 285-293.
646

- 647 46. Thompson, J.J.W., Armitage, S.A.O., Siva-Jothy, M.T. 2002. Cuticular melanisation
648 is time-constrained: blacker beetles darken faster. *Physiological Entomology*, 27, 136-
649 141.
- 650
- 651 47. Verhoog, M., Boven, A., Brakefield, P. 1996. Melanic moths and the ability to
652 encapsulate parasitoid eggs and larvae. *Proceedings of the Netherlands Entomological*
653 *Society*, 7, 127–133.
- 654
- 655 48. Vermeulen, C.J., Cremers, T.I., Westerink, B.H., Van De Zande, L., Bijlsma, R. 2006.
656 Changes in dopamine levels and locomotor activity in response to selection on virgin
657 lifespan in *Drosophila melanogaster*. *Mechanisms of Ageing and Development*, 127,
658 610-617.
- 659
- 660 49. Viega, J.P. 1993. Badge size, phenotypic quality, and reproductive success in the
661 house sparrow: a study on honest advertisement. *Evolution*, 47, 1161-1170.
- 662
- 663 50. Wigglesworth, V.B. 1948. The structure and deposition of the cuticle in the adult
664 mealworm, *Tenebrio molitor* L. (Coleoptera). *Quarterly Journal of Microscope*
665 *Science*, 89, 197-217.
- 666
- 667 51. Wigglesworth, V.B. 1985. The transfer of lipids in insects from the epidermal cells to
668 the cuticle. *Tissue Cell* 17, 249-265.
- 669
- 670 52. Wilson, K., Reeson, A.F. 1998. Density-dependent prophylaxis: evidence from
671 Lepidoptera–baculovirus interactions? *Ecological Entomology*, 23, 100-101.

672

673 53. Wilson, K.S., Cotter, C., Reeson, A.F., Pell, J.K. 2001. Melanism and disease
674 resistance in insects. *Ecology Letters*, 4, 637-649.

675

676 54. Wilson, K., Thomas, M.B., Blanford, S., Doggett, M., Simpson, S.J., Moore, S.L.
677 2002. Coping with crowds: density dependent disease resistance in desert locusts.
678 *Proceedings of the National Academy of Sciences of the U.S.A.*, 99, 5471–5475.

679

680 55. Zacharuk, R.Y. 1976. Structural changes of the cuticle associated with moulting. In:
681 *The insect integument* (ed. H.R. Hepburn). Elsevier, Amsterdam, 299-321.