

This is a repository copy of *Incidental retrieval of prior emotion mimicry*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/112568/>

Version: Accepted Version

Article:

Pawling, Ralph, Kirkham, Alexander, Hayes, Amy et al. (1 more author) (2017) Incidental retrieval of prior emotion mimicry. *Experimental Brain Research*. pp. 1173-1184. ISSN: 1432-1106

<https://doi.org/10.1007/s00221-017-4882-y>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

[Click here to view linked References](#)

Memory For Body States

Incidental retrieval of prior emotion mimicryRalph Pawling¹, Alexander J Kirkham², Amy E Hayes³ & Steven P Tipper²

Correspondence: Ralph Pawling, School of Natural Sciences and Psychology, Liverpool
John Moores University, Byrom Street, Liverpool, UK.

Telephone: 0151 9046315 Email: r.pawling@ljmu.ac.uk

Author Affiliations: ¹ School of Natural Sciences and Psychology, Liverpool John Moores
University, Liverpool, UK.

² Department of Psychology, University of York, York, UK.

³ School of Sport, Health and Exercise Sciences, Bangor University, Bangor, UK

Keywords: Embodied Cognition; Memory; Emotion; Psychophysiology

Character Count: 49,594

Acknowledgement. This work was supported by an Economic and Social Research
Council (ESRC) grant [ES/K000012/1] awarded to SPT and AEH. Data used in the
analysis of this study can be found at the following doi:

Abstract

When observing emotional expressions, similar sensorimotor states are activated in the observer, often resulting in physical mimicry. For example, when observing a smile, the zygomaticus muscles associated with smiling are activated in the observer; and when observing a frown, the corrugator brow muscles. We show that the consistency of an individual's facial emotion, whether they always frown or smile, can be encoded into memory. When the individuals are viewed at a later time expressing no emotion, muscle mimicry of the prior state can be detected, even when the emotion itself is task irrelevant. The results support simulation accounts of memory where prior embodiments of other's states during encoding are reactivated when re-encountering a person.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 As social animals humans must understand the current states of other's, predict
5 future actions and retrieve from memory information about a person that might
6 facilitate later interactions. Evidence supports the notion that such abilities may be
7 underpinned by processes of sensorimotor simulation. That is, representing the
8 observed actions (Avenanti, Candidi, & Urgesi, 2013) and internal states (Keyers, Kaas &
9 Gazzola, 2010) of others, involves the activation of similar visuomotor and
10 somatosensory states in the observer.

11 Research into sensorimotor simulation has focused on perception of motor
12 actions, after the discovery of cells in area F5 of the monkey, referred to as 'mirror
13 neurons' (Rizzolatti, Fogassi, and Gallese 2001). These cells are active when the animal
14 performs an action, but also when the animal observes the same or similar actions
15 performed by another actor (Di Pellegrino, Fadiga and Rizzolatti 1992).

16 Research studies in humans, utilising behavioural (e.g. Bach and Tipper 2007;
17 Griffiths and Tipper 2009), neuromodulatory (e.g. Fadiga, Fogassi, Pavesi and Rizzolatti
18 1995), and functional neuroimaging methods (e.g. Oosterhof, Wiggett, Diedrichsen,
19 Tipper and Downing 2010) provide evidence that observing actions indeed activates
20 similar motor states in the brain of the observer. Similar results have been shown for
21 observation of sensory experiences, such as viewing others receive noxious (Morrison,
22 Lloyd, di Pellegrino and Roberts 2004), and non-noxious touch (Keyers, Wicker,
23 Gazzola, Anton, Fogassi and Gallese 2004; Morrisson, Bjornsdotter and Olausson 2011).

24 Recently, interest in the role of simulation in understanding people's emotions
25 has increased. Facial expressions are a key observable aspect of emotion (Panksepp
26 2005), and motor simulation may be a route by which an observer understands the

1 meaning of these actions (Niedenthal, Mermillod, Maringer & Hess, 2010; Wood,
2 Rychlowska, Korb, & Niedenthal, 2016). Viewing facial expressions causes automatic
3 imitative responses in the facial muscles of the observer. For example, viewing another
4 person smile causes an almost immediate increase in activation in the zygomaticus
5 muscle of the cheek, whilst viewing a frown activates the corrugator muscle that draws
6 down the brow (e.g. Dimberg, Thunberg and Elmehed 2000; Dimberg, Thunberg and
7 Grunedal 2002). These changes in activity from the prior state of the musculature are
8 typically very small, non-visible, and only detectable through measurement of the
9 electrical activity at the muscle site, recorded via facial electromyography (EMG) (for
10 discussion of micro-expressions see Cacioppo, Bush and Tassinary 1992; Tassinary and
11 Cacioppo 1992). This ‘facial mimicry’ occurs rapidly, within a few hundred milliseconds
12 (Dimberg, 1997). It also appears automatic, occurring even when participants are
13 instructed not to move their faces (Dimberg et al. 2002), or when the emotion viewed is
14 irrelevant to the task in hand (Cannon, Hayes and Tipper 2009).

15 Recent models purport that facial mimicry represents a ‘spill over’ of
16 sensorimotor simulation (Wood et al. 2016). Vicarious activity in sensorimotor regions
17 elicited when viewing a facial expression (Sato, Kochiyama and Uono 2015; Schilbach,
18 Eickhoff, Mojzisch and Vogely 2008) is thought to trigger activation of other brain and
19 body states associated with the emotion being perceived, and this simulation aids the
20 recognition of the emotion. Whilst simulation may not always involve mimicry (Wood
21 et al. 2016), there is evidence that mimicry can augment or empower simulation of
22 other’s emotions (Lee et al. 2013). Blocking mimicry reduces accuracy in decoding the
23 meaning of facial expressions (Ipser and Cook 2015; Maringer, Krumhuber, Fischer and
24 Niedenthal 2011; Ponari, Conson, D’Amico, Grossi and Trojano 2012; Stel and
25 Knippenberg 2008), whilst amplifying mimicry can improve recognition. Participants

1 whose brow muscles had been paralysed via Botox injections performed worse than
2 controls in an emotion recognition task, whilst participants whose facial feedback was
3 amplified with proprioceptive taping performed better (Neal and Chartrand 2011). It's
4 believed that during facial mimicry afferent feedback from the facial muscles (Price and
5 Harmon-Jones 2015) adds to the overall simulation of the other's emotion already
6 occurring in sensorimotor regions of the brain (Niedenthal et al. 2010; Wood et al.
7 2016).

8 One issue yet to be addressed is whether such simulations might be reinstated as
9 part of long term memory for other's emotions. Reactivation accounts of episodic
10 memory suggest that retrieving a memory involves partial reactivation of the brain
11 states active during encoding, including those involved in sensory and motor processing
12 (Barsalou 1999; Buckner and Wheeler 2001; Glenberg 1979, for reviews see Buchanan
13 2007; Danker & Anderson 2010). Evidence for this comes from neuroimaging research
14 showing that retrieval of stimuli previously associated with differential sensory inputs
15 reactivated sensory specific cortical regions. For example, recalling an object encoded
16 in the form of a picture, as opposed to in word form, re-activated visual cortex (Vaidya,
17 Zhao, Desmond and Gabrieli 2002); remembering the sound associated with a
18 particular label reactivates areas of auditory cortex (Wheeler, Peterson and Buckner
19 2000); and remembering whether an object was previously presented in the presence
20 of an odor reactivates olfactory cortex (Gottfried, Smith, Rugg and Dolan 2004; for
21 limitations see Danker and Anderson 2010, pp6). Similarly, the retrieval from memory
22 of action words activated motor cortices when the words were encoded alongside
23 performance of the action, but not if the words were encoded without action
24 performance (Nyberg et al. 2001). Interestingly, simply *imagining* the enactment of the
25 action during encoding activated motor areas, which again became active during

1 retrieval. This latter finding is suggestive of the reactivation of prior simulative
2 processes.

3 In the same manner, might motor simulations of an individual's emotional state
4 be reactivated upon retrieval of that individual's identity? According to models such as
5 that of Wood et al. (2016), viewing another person's smile should activate the face
6 regions within the viewer's sensorimotor cortices, and might result in facial mimicry in
7 the zygomaticus muscle, with resultant afferent feedback, creating a simulation. Upon
8 meeting the same individual again, would this simulation then be reinstated as part of
9 implicit recall of the person's identity? And if so would a facial mimetic-like response
10 indicate such reinstatement – i.e could rapid and sub-perceptual reactivation of the
11 zygomaticus be detected? If this were to occur even if the individual being perceived
12 was now not smiling themselves, this would support the notion of a reactivation of prior
13 simulation.

14 The present study addresses this question. Participants were exposed to faces
15 that consistently smiled or frowned, whilst the activity of their own facial muscles was
16 recorded using facial EMG. Later the participants were re-exposed to the same faces
17 whilst completing a simple oddball task, and again facial EMG was recorded.
18 Importantly however, at re-exposure all the faces now had a neutral facial expression.
19 Covert facial mimicry responses, recorded during initial exposure to the emotional
20 versions of the faces would indicate the presence of sensorimotor simulation. However,
21 could reinstatement of these motor states be detected when the participants viewed the
22 neutral versions of the face? If so this would suggest the reactivation of the prior
23 simulation upon recalling the facial identity. A second issue was also addressed. Facial
24 mimicry appears to occur even when the facial expression being viewed is not relevant
25 to the task in hand. Similarly, reactivation of sensory or motor regions active during

encoding of episodic memories, occurs even when the sensory aspect of the memory is not being actively recalled (e.g. Vaidya et al. 2002). In the present study the question of whether task relevance would affect the reactivation of facial mimicry was addressed by having half the participants attend to the emotional aspect of the faces they viewed, whilst the other half ignored the emotion and attended to the identity of the faces.

Method

Participants

Participants were 36 female undergraduates from Bangor University, recruited in two cohorts of 18 participants. The first cohort completed the attend to identity condition (mean age = 21.9 years, SD = 2.1 years), and the second cohort the attend to emotion condition (mean age = 19.9 years, SD = 2.7 years). All gave informed consent. Women appear more responsive than men in their physiological reactions to emotional facial expressions (Dimberg and Lundquist 1990). Therefore a female cohort was selected as being more likely to demonstrate initial embodiment.

Design

Participants completed three tasks. First an initial learning task, during which face identities consistently expressed either smiling or frowning emotions and participants had to either identify oddball trials where the identity of the face changed mid-trial, or identified the emotion expressed by the face. Here it was predicted that participants would demonstrate facial mimicry even though this was never an explicit task goal.

Second, a few minutes after the first task, participants completed an implicit recall task where in the majority of trials they passively viewed static, neutral versions

of the faces seen during the learning phase. Their task was to monitor for occasional oddball targets, where either an identity change or an emotion change occurred, depending on the condition they had been assigned to for the initial encoding task. It was predicted that during the passive viewing trials, implicit retrieval of the prior mimicry state associated with a face identity would occur, if learning had taken place. Therefore, participants would show greater zygomaticus activity to those faces that had *previously* smiled than to those who *previously* frowned, and greater corrugator activity to those faces that *previously* frowned than to those who *previously* smiled.

Finally the participants completed an explicit recall task, viewing neutral versions of the previously seen faces, and making forced choice decisions about as to whether the face had previously smiled or frowned.

In a mixed design, participants were divided into two groups. One attended to the emotion expressed on the faces during both the encoding, and implicit retrieval tasks. The other ignored the emotion and attended to the identity of the face.

Stimuli

Morphing software was used to create dynamic facial expressions. Neutral, smiling and frowning versions of four female and four male faces were selected from the Nim Stim (<http://www.macbrain.org/mission.htm>) and KDEF (Lundqvist, Flykt and Ohman 1998) face sets. The eight identities used were selected to create four sex and attractiveness matched pairs, based on previously collected ratings of a larger set of faces, rated for attractiveness from 1-5. The male pairs were rated as 1.9 and 2.1, and 2.3 and 2.6. The female pairs as 2.7 and 2.9, and 3.7 and 3.9.

Dynamic morphs of each neutral face identity morphing into both expressions of every other face identity were also created, within sex. In total this created 24 identity change morphs, which were used as oddball trials in the initial learning task for participants tasked with attending to identity. For each participant eight morphs were randomly selected, with each face appearing once as the end face of an oddball morph. A further set of two neutral male and two neutral female faces were also selected as oddball trials for use in the subsequent implicit retrieval task for the participants who were tasked with attending to identity.

Procedure

After electrode placement, the participant completed the three tasks in order described above. The participants were told that the study was about sustained attention and a story of frontal lobe recording was used to disguise the EMG measures. The participants were not told that the same faces would be seen throughout the experiment, or that any recall would be required.

Initial learning/encoding task (mimicry). The procedure for the mimicry task can be seen in Figure 1. Participants pressed the spacebar to begin each trial. A fixation cross appeared for 2000ms. A face appeared, with a neutral expression lasting for 1500ms, before morphing to an emotional expression (the transition took 240ms), displaying a smile or a frown for 1500ms. After a blank screen of 2000ms the participant relaxed for 5000ms. All participants saw four identities (two male, two female) who *always* smiled, and four identities (two male, two female) who *always* frowned. Participants in the 'attend to emotion' condition (Figure 1a top panel) categorized the emotion they saw in each face display during the trial, pressing either

the Z or M key (counterbalanced). Participants in the 'attend to identity' condition (Figure 1b bottom panel) looked out for oddball trials, where a face would appear, then morph into the emotional face of another identity. Participants pressed the spacebar for oddballs, but did not respond to standard trials. Oddball trials always involved faces morphing into another face of the same emotion category. Face allocation to emotion was quasi-random for each participant, so that there were always two smiling females and two smiling males, and similarly two frowning males and females, and that the attractiveness balance was maintained. The task consisted of 64 standard trials, divided into four blocks of 16. In the attend to identity condition each block also contained two oddball trials, meaning a total of 72 trials.

Figure 1 here.

Implicit recall task (neutral faces). The procedure for the implicit recall task can be seen in Figure 2. Participants pressed the spacebar to initiate trials, after which they were viewed a fixation cross for 2000ms. A face appeared, with a neutral expression for 2000ms. After this the screen went blank for 2000ms and the participant was asked to relax for 5000ms. The task consisted of two blocks of 16 trials and each face identity appeared twice in each block. Each block contained two oddball trials. Oddballs in the attend to emotion condition (Figure 2a top) consisted of a face appearing for 1500ms, then morphing to a smile or frown; there were always two smiling and two frowning faces, selected from those used during encoding and balanced for sex. In the attend to identity condition (Figure 2b bottom) oddballs were brand new identities with neutral expressions. In both conditions participants responded to oddballs with a spacebar press and pressed nothing to standard trials. The selection of trial n's in both tasks was guided by the number of trials used in previous studies of

facial mimicry (Dimberg and Thunberg 1998; Hess and Philipott 1998; Vrana and Gross 2004). We used a larger number of trials during the initial encoding task than during implicit recall to try and ensure the participants learned the emotions associated with each face.

Figure 2 here

Explicit recall task. Finally, participants were shown the eight faces seen in the previous tasks and made a forced choice decision ('A' and 'L' keys assigned to condition, counterbalanced) as to whether the face had smiled or frowned during the encoding task. Faces were presented once, with neutral expressions, for 2000ms, and separated by a 2000ms fixation cross prior to presentation and blank screen afterwards.

EMG Recording

EMG data were collected from the zygomaticus (smile muscle) and corrugator (frown muscle), using a BioPac MP100 system (BioPac: Goleta, CA). Data were sampled at 2000Hz. Online filters were set to a bandpass of 1-5000Hz, and notch of 50Hz. Electrode placement and preparation were conducted as guided by Fridlund and Cacioppo, 1986. Recordings were made from the left side of the face (see Dimberg and Petterson 2000; Zhou and Hu 2004).

Offline the data were pass filtered with a 20-400Hz bandpass (as recommended in van Boxtel 2001), rectified, log10 transformed to reduce extreme values, and z-transformed within participant and muscle site. Change-scores were calculated trial-by-trial. For every trial the mean of the final 500ms of the fixation was used as a baseline and subtracted from each data point during the trial, transforming each point into a change from baseline score. The data were averaged over 200ms bins, using the

1 arithmetic mean. These bins were used as the level at which data were visualized. For
2 data analysis data were averaged using the arithmetic mean, over whole trial periods
3 (e.g. the period during which an emotional face was on screen, or the blank period after
4 this). Artifact trials were removed by eye before the data were matched to condition.
5 Error and post error trials were removed from the EMG analysis to account for error
6 related activity (Ekins-Brown, Saunders and Inzlicht 2016; Lindstrom, Mattsson-Marn,
7 Golkar and Olsson 2013) or post-error increases in effort (Van Boxtel and Jessuru
8 1993). The first trial of each task was removed upon observation of noisier data in
9 these trials. EMG data from oddball trials were not analysed. For the encoding task the
10 mean number of (non-oddball) trials removed per participant was: Attend to identity,
11 happy faces 4.0 trials (SD = 1.5), angry faces 4.7 trials (SD = 1.7); Attend to emotion,
12 happy faces 7.1 trials (SD = 2.2), angry faces 5.6 trials (SD = 1.4). For the implicit recall
13 task the mean number of (non-oddball) trials removed per participant were: Attend to
14 identity, previously happy faces 1.3 trials (SD = 1.0), previously angry faces 1.0 trials
15 (SD = 1.1); Attend to emotion, previously happy faces 2.1 trials (SD = 1.4), previously
16 angry faces 1.9 trials (SD = 1.1).

Results & Discussion

Behavioral Results for Learning/encoding and Implicit Recall Tasks

Accuracy rates were assessed for the encoding and implicit recall tasks. During
encoding, accuracy in the attend to identity condition referred to the percentage of
trials correctly categorized as oddballs, if they were such, or not responded to if they
were standard trials. In the attend to emotion condition accuracy referred to the
percentage of trials correctly identified as containing a happy or angry facial expression.
In the implicit recall tasks accuracy referred to the percentage of trials correctly

identified as oddballs if they were such (new faces in attend to identity / emotional faces in attend to emotion), or not responded to if they were standard trials. Accuracy rates were high across both the initial encoding and subsequent implicit recall tasks, suggesting the participants were attentive to the stimuli. During the learning tasks mean accuracies were above 98%, for both the attend to emotion and attend to identity conditions. During the implicit recall tasks accuracies were above 94% for both the attend to emotion and attend to identity conditions.

Mimicry during Learning/encoding Task

Facial mimicry was calculated using the EMG responses elicited by the emotion section of the morph videos, after the initial 1500ms neutral expression, when the face became expressive, smiling or frowning for 1500ms. This period also included the 2000ms blank screen after the offset of the face.

Corrugator (frown muscle). The activation states of the corrugator during and after the emotion expression face is seen are shown in Figure 3a. A three-way mixed analysis of variance was undertaken to analyse face emotion (happy / angry face), time (emotional face present / blank screen after face offset), and the between participants factor of task (attend to identity / attend to emotion).

Of primary interest are the significant results involving the face emotion factor, which are indications of emotional mimicry. Of most importance, there was a significant main effect of face emotion, where as predicted, the corrugator was more active when viewing angry than when viewing happy faces, $F(1,34) = 7.5, p = .01, \eta^2_p = .180$. Importantly, the mimicry of face emotion did not interact with whether participants were attending to the task relevant property of face emotion or irrelevant property of face identity, $F(1,34) = .015, p = .902, \eta^2_p < .001$. Interestingly there was also an

interaction between face emotion and time, $F(1,34) = 14.03, p = .001, \eta^2_p = .292$. Further analysis with uncorrected t-tests revealed that the effect of face emotion was detected while the emotional face was viewed, $t(35) = 4.6, p < .001$, but this effect was transient as there was no effect when the subsequent blank screen was viewed, $t(35) = .73, p > .1$.

The remaining significant effects did not involve the emotion factor, and thus indicate effects on the overall corrugator activation regardless of emotion. There was a significant effect of time, where corrugator activity increased from viewing faces to viewing the subsequent blank screen $F(1,34)=23.5, p < .001, \eta^2_p = .408$. The only significant between participants interaction effect was the interaction of task and time, $F(1,34) = 7.8, p = .009, \eta^2_p = .186$. Follow up uncorrected t-tests revealed that whilst overall activity in the corrugator during the face and blank periods differed significantly for those participants who attended to emotion, $t(17) = 5.1, p < .001$, the difference was not significant for those who attended to identity, $t(17) = 1.6, p > .1$.

Figure 3 here

Zygomaticus (smile muscle). The data are presented in Panel b of Figure 3. The data were entered into the same structure of analyses as the corrugator data. Again, significant results involving the face emotion factor are of primary interest. Of most importance, there was a significant main effect of face emotion $F(1,34) = 7.05, p = .012, \eta^2_p = .172$, where the zygomaticus was more active when viewing smiling faces as compared to viewing angry faces, and this did not interact with whether attention was focused on face emotion or identity $F(1,34) = 1.62, p = .289, \eta^2_p = .033$. Interestingly, there was no interaction in the zygomaticus between face emotion and time $F(1,34) = .88, p > .1, \eta^2_p = .055$, meaning that, in contrast to the corrugator

muscle, the mimicry effect remained stable into the blank screen period after the offset of the emotional face. Significant results not involving the face emotion factor were a main effect of time $F(1,34) = 7.09, p = .012, \eta^2_p = .173$, where activity increased across time from the period where the face was on screen into the blank screen after the offset of the face; and a main effect of task $F(1,34) = 12.4, p < .001, \eta^2_p = .267$, where the zygomaticus muscle was more active when attending to emotion than attending to identity.

Two key findings here are of note. First, even though participants were not asked to explicitly mimic the face emotion while attending to their particular task, the mimicry effect was still detected in both muscles. Secondly, this seems to be independent of the participant's task, as there is no interaction between face emotion and task of attending to emotion or attending to face identity. That is, whether directly attending to and making decisions about face emotion, or ignoring the emotion while reporting occasional identity changes, the mimicry effect is observed. This supports the notion that mimicry is evoked without intent (e.g., Cannon et al. 2009; Dimberg et al. 2000; Dimberg et al. 2002).

It is also worth noting that across both muscle sites the pattern of activity seen in the attend to emotion condition differed to that in the attend to identity condition. In the former it appeared that mimicry constituted a greater increase in activity from baseline levels, whereas in the latter it appeared as a decrease in activity from baseline to the incongruent expression. Finally it worth noting that mimicry seems to be longer lasting in the zygomaticus muscle, continuing in to the blank field after face offset; whereas in contrast, mimicry is only detected in the corrugator muscle while participants are actually viewing an emotion expressing face.

The next analysis examined the EMG data from the implicit recall section of the experiment, where the participants viewed the faces that had previously smiled or frowned, but this time showed neutral expressions. Here we predicted that there would be a reactivation of the facial mimicry responses detected in the encoding task.

Figure 4 here

Implicit Recall Task (Neutral Faces)

A three-way mixed analysis of variance was carried out on these data, within each muscle site, examining face emotion (previously appeared happy/previously appeared angry), time (neutral face present/ blank screen) and task (attend to identity/attend to emotion).

Corrugator: The data are presented in Figure 4, Panel a. Importantly, there was no main effect or interactions involving prior face emotion; this indicates that retrieval of prior emotional mimicry did not occur. The only significant main effects and interactions involved the factors of task and time. Participants showed more corrugator activity during the blank screen period than during the period where the face was onscreen, $F(1,34) = 18.7, p < .001, \eta^2_p = .355$, but this varied between those participants attending to emotion and those attending to identity, $F(1,34) = 18.6, p < .001, \eta^2_p = .353$. Uncorrected t-tests conducted to explore this interaction revealed that the increase in activity from face to blank period was only significant in those participants attending to emotion, $t(17) = 4.6, p < .001$. Overall, the participants attending to emotion showed greater corrugator activity as well, $F(1,34) = 8.3, p = .007, \eta^2_p = .195$.

Therefore the corrugator provides no evidence for retrieval of prior facial emotion body states. This might not be entirely unexpected given previous studies have

also observed less robust effects in corrugator than in zygomaticus muscles (e.g., Cannon, Hayes and Tipper 2010) and have reported less robust mimicry of the socially costly expression of anger, than on less costly expressions such as happiness and sadness (Bourgeois and Hess 2008; Hess and Bourgeois 2006; but see Kirkham, Pawling, Hayes, and Tipper 2015, for contrasting effects).

Zygomaticus: The data are presented in Figure 4, Panel b. Most importantly there was a main effect of prior face emotion, $F(1,34) = 7.7, p = .009, \eta^2_p = .184$, with increased zygomaticus activation when viewing a neutral face that had previously expressed a positive/smiling emotion as compared to a neutral face that had previously expressed anger and this did not interact with the face property attended (emotion or identity) $F(1,34) < .1, p = .9, \eta^2_p < .1$. The remaining significant effects did not involve the face emotion factor. There was a main effect of time, $F(1,34) = 23.7, p < .001, \eta^2_p = .410$, with activation increasing from the face to blank periods. Although the interaction of task and time was significant, $F(1,34) = 5.6, p = .02, \eta^2_p = .142$, follow up uncorrected t-tests revealed a significant increase in zygomaticus activity from the face to blank periods in both participants attending to identity, $t(17) = 2.9, p = .01$, and emotion, $t(17) = 4.0, p = .001$.

Explicit Recall Task

At the end of the study participants were presented with the faces and asked to recall whether the person had previously smiled or was angry. Data from two participants in the attend to identity condition were missing from this analysis due to experimenter error in saving the data, and a participant misunderstanding the key-presses. Mean percentage accuracy (correct identification of whether a face previously

smiled or frowned) was high amongst both those participants attending to emotion (mean = 95.1%, SD = 7.6%) and those attending to identity (mean = 82.03 %, SD = 20.4%). Both the attend to emotion ($p < .001$ for t-test against 50%) and attend to identity ($p < .001$) groups performed above chance on this task, but, perhaps predictably a comparison of the two groups' performance showed that the group who attended to emotion recalled significantly more faces' prior emotions correctly, $F(1,32) = 6.4, p = .016$. It is also worth noting that the attend to identity group also saw oddball trials at encoding, which might have interfered with learning.

Individual Differences in Mimicry

Finally, we make a tentative analysis concerning individual differences in mimicry processes. Although mimicry of another person's emotion is a robust effect observed in a number of previous studies, not all people produce detectable mimicry (see Manssuer, Pawling, Hayes and Tipper 2016, for other evidence for individual differences in evoked emotion and learning). This could have implications for later retrieval of prior body states, in that people who did not initially mimic while observing faces expressing emotion, cannot subsequently retrieve and reestablish a mimicry state.

Therefore we divided participants in to those who mimicked observed emotion in the first stage of the study and those who did not mimic. This was done by separating those participants who showed muscle activity trending in the expected direction in each muscle from those who did not. So for analysis of the zygomaticus we compared those participants who showed greater activity to smiles than frowns (mimickers) to those who showed the opposite (non-mimickers). And in the corrugator the opposite, with those who showed greater activity in response to frowns (mimickers) being compared to those who showed greater activity in response to smiles (non-mimickers).

1 This does not mean the participants in the mimicry group mimicked every face, only
2 that on average they showed an effect that suggested mimicry. We predicted that
3 retrieval of mimicry in the later retrieval stage can only be detected in the individuals
4 who originally mimicked, and while no significant effects were found in the corrugator
5 analysis, this was indeed the case for the zygomaticus muscle ($N = 25$; $F(1,24) = 9.7$, p
6 $= .005$, $\eta^2_p = .287$). In sharp contrast, those who did not mimic the observed emotion in
7 the initial stage of the study also did not produce any evidence of mimicry in the later
8 retrieval stage ($N = 11$, $F(1,10) = .011$, $p = .92$, $\eta^2_p = .001$). On the other hand, when
9 examining explicit recall of prior emotion the mimickers (mean accuracy = 90.2%, test
10 against chance $p < .001$) and non-mimickers (mean accuracy 86.4%, test against chance
11 $p < .001$) performed similarly. Hence explicit recall is not reliant on implicit retrieval of
12 prior body states (e.g., Toplinski 2011). Clearly the small sample size in the latter non-
13 mimicry group requires that we are cautious and tentative with our conclusions, but
14 retrieval of prior embodied states predicts just such a result, and hence it is worthy of
15 future investigation.

General Discussion

18 The current study provided a number of new observations that add to the
19 discussion surrounding the role of sensorimotor simulation in social cognition (Wood et
20 al., 2016). First, automatic mimicry of another person's emotion was again observed in
21 this study. Thus, even though participants were not asked to overtly mimic another
22 person's emotion, mimicry was nevertheless detected through the use of facial EMG.
23 This replicates previous research (Cannon et al. 2009; Dimberg et al. 2000; 2002), and
24 fits theory purporting that facial motor regions are vicariously activated when viewing
25 facial expressions (Wood et al. 2016).

Furthermore, the mimicry effect was observed both when emotion was relevant and irrelevant to the participant's goal, supporting the notion that facial mimicry is automatic, and can to be evoked even when another person's emotion is task irrelevant and not explicitly attended to (e.g., Cannon et al. 2009; Dimberg et al. 2000). It is worth noting that the patterns of muscle activity differed between the attention conditions. When attending to emotion, facial mimicry in both muscles appeared as a larger increase from baseline activity. However, in the attend to identity condition it took the form of a lesser reduction from baseline activity, for example, the zygomaticus activity decreased from baseline to a lesser degree for happy than angry faces. Similar results have been reported in other experiments (see Cannon et al. 2009; Dimberg et al. 2000). It might be that the pattern of activity in the attend to identity condition is indicative of a greater preparatory response in both muscles, prior to the presentation of the face – perhaps because the task required more effort. Or, as suggested in Cannon et al. (2009), it could be due a suppression of the task irrelevant emotion.

However, the core issue engaged here was the reactivation of simulation during later encounters. Embodied accounts of emotional memory processes (e.g., Niedenthal 2007) propose that perceptual-motor states are encoded during initial experience of a stimulus. In the current case this would be the activation of facial muscles while viewing another person express an emotion, which in accordance with simulation models, results from vicarious neural activity in facial motor regions. We predicted that such body states might be reactivated when encountering a person again, even if they were not expressing any emotion. We found that indeed, there is evidence for such reinstatement of prior processing, which we saw in the zygomaticus muscle. Here muscle activity reflecting mimicry during previous encoding of emotional faces can be detected when the faces are later viewed with neutral expressions. Interpreted in the

light of current theory this supports the notion that sensorimotor simulations occurring during encoding are reinstated during recall. This finding aligns with models of emotional memory and with evidence that during episodic memory retrieval, states of sensory and motor neural activity that were active at encoding are re-activated. Our data fit particularly well with evidence that neural simulations of actions encoded alongside verbal stimuli are reactivated upon later retrieval of the verbal information alone (Nyberg et al. 2001).

However there appear to be boundary conditions. First, the retrieval effects were only detected in the zygomaticus muscle and not the corrugator, despite initial mimicry in both muscles. It may be the case that the more subtle embodied responses are simply detected more easily in the zygomaticus (see Cannon et al. 2010). It is also perhaps noteworthy that the mimicry activity in the corrugator in the present study appeared fleeting, only differentiating between angry and happy faces when the faces were onscreen. Prior research has indicated that facial mimicry is affected by social context (for review see Hess and Fischer 2014), particularly the more socially costly mimicry of anger. It might be the case that reactivation of the corrugator inhibited either by potential social cost or because of reduced affiliation with faces who previously demonstrated consistently negative expressions (Bourgeois and Hess 2008).

In an exploratory analysis we compared reactivation effects in the zygomaticus between participants who did and did not demonstrate mimicry in this muscle. Interestingly, whilst mimickers and non-mimickers showed an equal ability to explicitly recall the face-emotion pairings, it was only amongst the mimickers that reactivation of mimicry was seen. The scope for conclusions on this result are limited by the small sample size and simplistic division of mimickers and non-mimickers. However, the results are none-the-less interesting and suggest that further exploration of individual

1 differences in the utilization, or outward expression of reactivated simulations is
2 needed. It is also worth noting the contrasting results of a similar study (Kirkham,
3 Pawling, Hayes and Tipper 2015). Here during encoding participants viewed emotional
4 faces who smiled and frowned in a manner that was either consistent, or inconsistent
5 with an emotional context, created by the simultaneous presentation of an emotional
6 image alongside the face. Mimicry was reduced for faces whose expressions were
7 inconsistent with the emotional context. However, in a later retrieval phase, where the
8 faces still smiled and frowned but no emotional context was presented, the prior
9 consistency or inconsistency of a face's expressions did not affect participant's mimicry
10 –i.e no retrieval of prior emotion consistency was found. However, unlike the current
11 study, which used neutral faces at retrieval, the facial mimicry evoked at retrieval may
12 have blocked any detection of prior mimicry states being reactivated.

13 Finally some limitations of the current study, and possible future directions
14 require discussion. Numerous experiments have explored the role of sensorimotor
15 simulations and facial mimicry in accurately decoding facial expressions (Ipser and
16 Cook 2015). The current study did not utilize a design that allowed for such an analysis
17 of the role that *reactivation* of simulations might play. For example, simulations might
18 be retrieved as part of predicting the other person's oncoming emotion (Heerey and
19 Crossley 2013), or might play a part in impression formation (Halberstadt, Winkielman,
20 Niedenthal and Dalle 2009). We did not record EMG activity during explicit recall,
21 where reactivation of facial mimicry might have predicted accurate retrieval of prior
22 emotion condition. We also used a relatively small number of faces, and an extension
23 using a larger face set might be better suited for exploring links between reactivated
24 facial mimicry and impression formation or accurate emotion recall.

1 It would also be interesting to investigate similar effects with non-facial stimuli
2 that elicit mimicry-like responses, such as emotional bodies (Tamietto et al. 2009), or
3 images (Dimberg 1986). It is difficult in the present study, but also in the literature as a
4 whole, to decipher whether facial mimicry responses constitute simulation, emotion
5 contagion or just emotional response (Hess and Fischer 2014). Perception deficits
6 specific to perceiving facial emotion when blocking facial mimicry, for example, support
7 the notion of mimicry as simulation (Ipser and Cook 2015; Neal and Chartrand 2011).
8 However, it would be enlightening to explore similar boundaries on the retrieval of
9 mimicry. For example, would blocking later reactivation of mimicry cause a decreased
10 accuracy in remembering prior emotions?

11 We sampled only female participants, and therefore our results can only be
12 generalized to the female population. Examining male participants might help cast light
13 on the individual differences that might underlie the apparent link seen in our
14 exploratory analysis between initial mimicry and later mimicry reactivation. We also
15 used faces that always smiled or frowned. Of course in the real world people hardly
16 ever express a single emotion consistently, hence a more ecologically valid follow up
17 could include identities that emote less predictably.

18 To conclude, we have confirmed that mimicry of another person's emotion takes
19 place without intention and even when attending to irrelevant properties of a face such
20 as identity. Of most importance, in support of simulation accounts of memory, we show
21 encoding of the relationship between a person's identity and their typical emotion, via
22 reactivation of prior motor simulation. That is, mimicry evoked while viewing a person
23 expressing emotion can be reactivated at a later time, even when they are no longer
24 expressing any emotion. And although tentative at this time, it appears that such

1 reactivations only occur in individuals who demonstrate mimicry responses in the first
2 instance.

References

- 3
4
5
6
7
8
9
10
11 Avenanti A, Candidi M, Urgesi, C (2013) Vicarious motor activation during action
12 perception: beyond correlational evidence. *Front Hum Neurosci.* doi:
13 10.3389/fnhum.2013.00185
14
15
16
17 Bach P, Tipper SP (2007) Implicit action encoding influences personal-trait judgments.
18
19 *Cognition.* doi: 10.1016/j.cognition.2005.11.003
20
21
22 Barsalou LW (1999) Perceptual symbol systems. *Behav Brain Sci.* doi:
23 10.1017/s0140525x99002149
24
25
26
27 Bourgeois P, Hess U (2008) The impact of social context on mimicry. *Biol Psychol.* doi:
28 10.1016/j.biopsycho.2007.11.008
29
30
31
32 Buchanan TW (2007) Retrieval of emotional memories. *Psychol Bull.*
33 doi:10.1037/0033-2909.133.5.761
34
35
36
37 Buckner RL, Wheeler ME (2001) The cognitive neuroscience of remembering. *Nat Rev*
38
39 *Neurosci* doi: 10.1080/17588928.2010.503602
40
41
42 Cacioppo JT, Bush LK, Tassinary LG (1992) Microexpressive Facial Actions as a Function
43
44 of Affective Stimuli: Replication and Extension. *Pers Soc Psychol B.*
45
46
47 doi:10.1177/0146167292185001
48
49
50
51
52 Cannon PR, Hayes AE, Tipper SP (2009) An electromyographic investigation of the
53
54 impact of task relevance on facial mimicry. *Cognition Emotion.* doi:
55 10.1080/02699930802234864
56
57
58
59 Cannon PR, Hayes AE, Tipper SP (2010) Sensorimotor fluency influences affect:
60
61 Evidence from electromyography. *Cognition Emotion.* doi:

- 10.1080/02699930902927698
- Danker JF, Anderson JR (2010) The ghosts of brain states past: Remembering reactivates the brain regions involved during encoding. *Psychol Bull.* doi: 10.1037/a0017937
- di Pellegrino G, Fadiga L, Fogassi VG, Rizzolatti G (1992) Understanding motor events: a neurophysiological study. *Exp Br Res.* doi: 10.1007/BF00230027
- Dimberg U (1986) Facial reactions to fear-relevant and fear-irrelevant stimuli. *Biol Psychol.* doi: 10.1016/0301-0511(86)90079-7
- Dimberg U (1997) Facial reactions: Rapidly evoked emotional responses. *J Psychophysiol.* doi: 10.1080/02699930143000356
- Dimberg U, Lundquist LO (1990) Gender differences in facial reactions to facial expressions. *Biol Psychol.* doi: 10.1016/0301-0511(90)90024-Q
- Dimberg U, Petterson M (2000) Facial reactions to happy and angry facial expressions: Evidence for right hemisphere dominance. *Psychophysiology.* doi: 10.1111/1469-8986.3750693
- Dimberg U, Thunberg M (1998) Rapid facial reactions to emotional facial expressions. *Scand J Psychol.* doi:10.1111/1467-9450.00054
- Dimberg U, Thunberg, M, Elmehed K (2000) Unconscious facial reactions to emotional facial expressions. *Psychol Sci.* doi: 10.1111/1467-9280.00221
- Dimberg U, Thunberg M, Grunedal S (2002) Facial reactions to emotional stimuli: Automatically controlled emotional responses. *Cognition Emotion.* doi: 10.1080/02699930143000356
- Elkins-Brown N, Saunders B, Inzlicht M (2016) Error-related electromyographic activity over the corrugator supercilii is associated with neural performance monitoring. *Psychophysiology.* doi: 10.1111/psyp.12556


- 1 Fadiga L, Fogassi L, Pavesi G, Rizzolatti G (1995) Motor facilitation during action
2 observation: a magnetic stimulation study. J Neurophysiol, 73: 2608-2611
3
- 4 Fridlund AJ, Cacioppo JT (1986) Guidelines for human electromyography research.
5 Psychophysiology. doi: 10.1111/j.1469-8986.1986.tb00676.x
6
- 7 Glenberg AM (1997) What memory is for? Beh Brain Sci. doi:
8 10.1017/S0140525X97000010
9
- 10 Gottfried JA, Smith APR, Rugg MD, Dolan RJ (2004) Remembrance of odors past: Human
11 olfactory cortex in cross-modal recognition memory. Neuron. doi:
12 10.1016/S0896-6273(04)00270-3
13
- 14 Griffiths D, Tipper SP (2009) Priming of reach trajectory when observing actions: Hand-
15 centred effects. Q J Exp Psychol. doi: 10.1080/17470210903103059
16
- 17 Halberstadt J, Winkielman P, Niedenthal P, Dalle N (2009) Emotional conception: How
18 embodied emotion concepts guide perception and facial action. Psychol Science.
19 doi: 10.1111/j.1467-9280.2009.02432.x
20
- 21 Heerey, EA, Crossley HM (2013) Predictive and reactive mechanisms in smile
22 reciprocity. Psychol Sci. doi:10.1177/0956797612472203
23
- 24 Hess U, Bourgeois P (2006) The social costs of mimicking: Why we should not both look
25 angry. Paper presented at the 7th Annual Meeting of the Society for Personality
26 and Social Psychology, January 26th-28th, Palm Springs, CA.
27
- 28 Hess U, Fischer A (2014) Emotional mimicry: Why and when we mimic emotions. Social
29 and Personality Psychology Compass. doi: 10.1111/spc3.12083
30
- 31 Hess U, Philippot P (1998) Facial reactions to emotional facial expressions : Affect or
32 cognition ? Cognition Emotion. doi: 10.1080/026999398379547
33
- 34 Ipser A, Cook R (2016) Inducing a Concurrent Motor Load Reduces Categorization
35 Precision for Facial Expressions. J Exp Psychol Human. doi:
36


- 1 10.1037/xhp0000177.
- 2 2 Keysers C, Kaas JH, Gazzola V. Somatosensation in social perception. Nat Reviews
- 3 Neurosci. doi: 10.1038/nrn2833
- 4 4 Keysers C, Wicker B, Gazzola V, Anton JL, Fogassi L, Gallese V (2004) A touching sight:
- 5 SII/PV activation during the observation and experience of touch. Neuron. doi:
- 6 10.1016/S0896-6273(04)00156-4
- 7 7 Kirkham AJ, Pawling R, Hayes AE, Tipper SP (2015) Facial mimicry and emotion
- 8 consistency: Influences of memory and context. PLoS ONE. doi:
- 9 10.1371/journal.pone.0145731
- 10 10 Lee IS, Yoon SS, Lee SH, Lee H, Park HJ, Wallraven C, Chae Y (2013) An amplification of
- 11 feedback from facial muscles strengthened sympathetic activations to emotional
- 12 facial cues. Auton Neurosci-Basic. doi: 10.1016/j.autneu.2013.06.009
- 13 13 Lindström BR, Mattsson-Mårn IB, Golkar A, Olsson A (2013) In your face: Risk of
- 14 punishment enhances cognitive control and error-related activity in the
- 15 corrugator supercilii muscle. PLoS ONE. doi: 10.1371/journal.pone.0065692
- 16 16 Lundqvist D, Flykt A, Ohman A (1998) The Karolinska Directed Emotional Faces (KDEF).
- 17 Stockholm: Department of Neurosciences, Karolinska Hospital.
- 18 18 Manssuer L, Pawling R, Hayes AE & Tipper SP (2015) The role of emotion in the
- 19 learning of trustworthiness from eye-gaze: Evidence from facial
- 20 electromyography. *Cognitive Neuroscience*, doi:
- 21 10.1080/17588928.2015.1085374
- 22 22 Maringer M, Krumhuber EG, Fischer AH, Niedenthal PM (2011) Beyond smile dynamics:
- 23 Mimicry and beliefs in judgments of smiles. Emotion. doi: 10.1037/a0022596
- 24 24 Morrison I, Björnsdotter M, Olausson H (2011) Vicarious responses to social touch in
- 25 posterior insular cortex are tuned to pleasant caressing speeds. J Neurosci. doi:


- 10.1523/JNEUROSCI.0397-11.2011
- Morrison I, Lloyd D, di Pellegrino G, Roberts N (2004) Vicarious responses to pain in anterior cingulate cortex: Is empathy a multisensory issue? Cogn Affect Behav Ne. doi: 10.3758/CABN.4.2.270
- Neal DT, Chartrand TL (2011) Embodied Emotion Perception: Amplifying and Dampening Facial Feedback Modulates Emotion Perception Accuracy. Soc Psycho Personal Sci. doi: 10.1177/1948550611406138
- Niedenthal PM (2007) Embodying Emotion. Science. doi: 10.1126/science.1136930
- Niedenthal PM, Mermillod M, Maringer M, Hess U (2010) The Simulation of Smiles (SIMS) model: Embodied simulation and the meaning of facial expression. Behav Brain Sci. doi: 10.1017/S0140525X10000865
- Nyberg L, Petersson KM, Nilsson LG, Sandblom J, Aberg C, Ingvar M. (2001) Reactivation of motor brain areas during explicit memory for actions. Neuroimage. doi: 10.1006/nimg.2001.0801
- Oosterhof NN, Wiggett AJ, Diedrichsen J, Tipper SP, Downing PE (2010) Surface-based information mapping reveals crossmodal vision-action representations in human parietal and occipitotemporal cortex. J Neurophysiol. doi: 10.1152/jn.00326.2010
- Overbeek TJM, van Boxtel A, Westerink JHDM (2012) Respiratory sinus arrhythmia responses to induced emotional states: Effects of RSA indices, emotion induction method, age, and sex. Biol Psychol doi: 10.1016/j.biopsycho.2012.05.011
- Panksepp J (2005) Affective consciousness: Core emotional feelings in animals and humans. Conscious Cogn. doi: 10.1016/j.concog.2004.10.004
- Ponari M, Conson M, D'Amico NP, Grossi D, Trojano L (2012) Mapping correspondence between facial mimicry and emotion recognition in healthy subjects. Emotion.

- doi: 10.1037/a0028588
- Price TF, Harmon-Jones E (2015) Embodied emotion: The influence of manipulated facial and bodily states on emotive responses. Wiley Interdiscip Rev Cog Sci. doi: 10.1002/wcs.1370
- Rizzolatti G, Fogassi L, Gallese V (2001) Neurophysiological mechanisms underlying the understanding and imitation of Action. Neuroscience. doi: 10.1038/35090060
- Sato W, Kochiyama T, Uono S (2015) Spatiotemporal neural network dynamics for the processing of dynamic facial expressions. Nature Publishing Group. doi: 10.1038/srep12432
- Schilbach L, Eickhoff SB, Mojzisch A, Vogeley K (2008) What's in a smile? Neural correlates of facial embodiment during social interaction. Soc Neurosci doi: 10.1080/17470910701563228
- Stel M, Knippenberg A (2008) The role of facial mimicry in the recognition of affect. Psychol Sci, 19. doi: 10.1111/j.1467-9280.2008.02188.x
- Tamietto M, Castelli L, Vighetti S, Perozzo P, Geminiani G, Weiskrantz L, de Gelder B (2009) Unseen facial and bodily expressions trigger fast emotional reactions. P Natl Acad Sci USA. 10.1073/pnas.0908994106
- Tassinari LG, Cacioppo JT (1992) Unobservable Facial Actions and Emotion. Psychol Sci. doi: 10.1111/j.1467-9280.1992.tb00252.x
- Topolinski S (2011) The sensorimotor contributions to implicit memory, familiarity and recollection. J Exp Psychol Gen. doi: 10.1037/a0025658
- Vaidya CJ, Zhao M, Desmond JE, Gabrieli JDE (2002) Evidence for cortical encoding specificity in episodic memory: Memory-induced re-activation of picture processing areas. Neuropsychologia. doi: 10.1016/S0028-3932(02)00053-2

- 1 Van Boxtel A (2001) Optimal signal bandwidth for the recording of surface EMG activity
2 of facial, jaw, oral and neck muscles. Psychophysiology. doi: 10.1111/1469-
3 8986.3810022
- 4 Van Boxtel A, Jessurun M (1993) Amplitude and bilateral coherency of facial and jaw-
5 elevator EMG activity as an index of effort during a two-choice serial reaction
6 task. Psychophysiology. doi:10.1111/j.1469-8986.1993.tb02085.x
- 7 Vrana SR, Gross D (2004) Reactions to facial expressions: Effects of social context and
8 speech anxiety on responses to neutral, anger, and joy expressions. Biol Psychol.
9 doi: 10.1016/j.biopsycho.2003.07.004
- 10 Wheeler ME, Petersen SE, Buckner RL (2000) Memory's echo : Vivid remembering
11 reactivates sensory-specific cortex. Procl Acad Natl Sci USA. doi:
12 10.1073/pnas.97.20.11125
- 13 Wood A, Rychlowska M, Korb S, Niedenthal P (2016) Fashioning the Face: Sensorimotor
14 Simulation Contributes to Facial Expression Recognition. Trends Cogn Sci. doi:
15 10.1016/j.tics.2015.12.010
- 16 Zhou R, Hu S (2004) Effects of viewing pleasant and unpleasant photographs on facial
17 EMG asymmetry. Percept Motor Skill. doi: 10.2466/pms.99.3f.1










Fig. 1. Time-course of trials from (a) attend to emotion condition and (b) attend to identity condition of the encoding task

Fig. 2. Time-course of trials from (a) attend to emotion condition and (b) attend to identity condition of the implicit recall task

Fig. 3. EMG responses during the encoding task from the (a) corrugator and (b) zygomaticus muscles when the participants viewed faces smiling (Happy) and frowning (Angry).

Changescorers are standardised units or change from baseline resting state. Dotted lines represent transition points in the timecourse of the trial, as illustrated above the graph: After fixation the face appeared with a neutral expression for 1500ms, transitioned to an emotional expression which remained on screen for 1500ms, before a blank screen appeared for 2000ms.

Fig. 4. EMG responses during the implicit retrieval task from the (a) corrugator and (b) zygomaticus muscles when participants viewed neutral faces. Changescorers are standardised units or change from baseline resting state. Dotted lines represent transition points in the timecourse of the trial, as illustrated above the graph: After fixation the face appeared with a neutral expression for 2000ms, before a blank screen appeared for 2000ms.