UNIVERSITY of York

This is a repository copy of Development of an efficient glucosinolate extraction method.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/113468/</u>

Version: Accepted Version

Article:

Doheny-Adams, Timothy, Redeker, Kelly Robert orcid.org/0000-0002-1903-2286, Kittipol, Varanya et al. (2 more authors) (2017) Development of an efficient glucosinolate extraction method. Plant Methods. 17. ISSN 1746-4811

https://doi.org/10.1186/s13007-017-0164-8

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

Plant Methods

Development of an efficient glucosinolate extraction method --Manuscript Draft--

Manuscript Number:	PLME-D-16-00100R1			
Full Title:	Development of an efficient glucosinolate extraction method			
Article Type:	Methodology			
Funding Information:	Biotechnology and Biological Sciences Research Council (BB/L002124/1)	Prof Ian Bancroft		
	Biotechnology and Biological Sciences Research Council (BB/K020463/1)	Prof Sue E Hartley		
Abstract:	Abstract Background			
	Glucosinolates, anionic sulfur rich secondary metabolites, have been ext studied because of their occurrence in the agriculturally important brassic their impact on human and animal health. There is also increasing interest biofumigant properties of toxic glucosinolate hydrolysis products as a me agricultural pests. Evaluating biofumigation potential requires rapid and a quantification of glucosinolates, but current commonly used methods of e to analysis involve a number of time consuming and hazardous steps; the to develop an improved method for glucosinolate extraction.			
	Three methods previously used to extract glucosinolates from brassicacae tissues, namely extraction in cold methanol, extraction in boiling methanol, and extraction in boiling water were compared across tissue type (root, stem leaf) and four brassicacae species (B. juncea, S. alba, R. sativus, and E. sativa). Cold methanol extraction was shown to perform as well or better than all other tested methods for extraction of glucosinolates with the exception of glucoraphasatin in R. sativus shoots. It was also demonstrated that lyophilisation methods, routinely used during extraction to allow tissue disruption, can reduce final glucosinolate concentrations and that extracting from frozen wet tissue samples in cold 80% methanol is more effective.			
	Conclusions			
	We present a simplified method for extractin does not require the use of a freeze drier or hazardous, and more time and cost effectiv to have comparable or improved glucosinol commonly used ISO method for major gluco studied: sinigrin and gluconasturtiin in B. jun gluconasturtiin in S. alba; glucoraphenin an glucosatavin, glucoerucin and glucoraphani	ng glucosinolates from plant tissues which boiling methanol, and is therefore less e. The presented method has been shown ate extraction efficiency relative to the osinolates in the Brassicaceae species ncea; sinalbin, glucotropaeolin, and d glucoraphasatin in R. sativus; and n in E. sativa.		
Corresponding Author:	Timothy William Doheny-Adams, Ph.D University of York UNITED KINGDOM			
Corresponding Author Secondary Information:				
Corresponding Author's Institution:	University of York			
Corresponding Author's Secondary Institution:				
First Author:	Timothy William Doheny-Adams, Ph.D			

First Author Secondary Information:	
Order of Authors:	Timothy William Doheny-Adams, Ph.D
	Kelly Redeker
	Varanya Kittipol
	Ian Bancroft
	Sue E Hartley
Order of Authors Secondary Information:	
Response to Reviewers:	We would like to thank the reviewers for their time and effort and their suggestions for improving the manuscript. We have addressed these suggestions to the best of our ability. Corrections have been highlighted in yellow in the document and the line numbers of those corrections are included below.
	REVIEWER #1: 1.The authors should provide the some chromatograms of their real smaples analysis as a supporting materials:
	Response: A supporting document with example chromatograms has been uploaded.
	2.The size of the column they used should be provided. For example, what is the size of "C18 column" (line 233). Response: the size of the column has been added (line 237)
	REVIEWER #2
	LN 103-104: A description of the desulfation process/mechanism and or reference is needed. Why is this a necessary step?
	Response: Now reads "Samples are subsequently desulfated by ion exchange chromatography on a DEAE sephadex column to remove impurities." Whether it is a necessary step or not and the steps involved are explored later in the manuscript with a direct comparison between desulfated and non-desulfated extractions.
	LN112: self-dimerisation via formation of a disulfide linkage during the extraction procedure. Response: Changed to: "In addition, prior to 2002 the major glucosinolate in leaves of E. sativa, 4-mercaptobutyl glucosinolate, was missed due to self-dimerisation via formation of disulfide linkages during extraction [22]"
	LN113-115: How many glucosinolates possess free thiol groups and the ability to self- react? The main difficulties lie in inhibiting myrosinase activity and degradation via heat or oxidation. Response:We appreciate the reviewers comment and agree that the vast majority of glucosinolates are not capable of self-dimerization. However, the previous sentence clearly demonstrates that at least one, important, glucosinolate has this potential. Therefore, we feel that this sentence remains valid as stated.
	LN156: Table m2 could be converted to text to reduce space. Response: We are willing to perform this action if the journal is concerned about space, otherwise we are content to let the table remain.
	LN160: What were average conditions for grinding prior to milling? Response: Lines 161-165 now read: "(i) Freeze dried plant tissue was homogenised to a roughly ground powder (approximately 0.1cm particle size) using a grinder (Lloytron, E5601BK) Homogenised ground samples were milled at a frequency of 20 Hz for 10 minutes (Retch, MM400) with 2 steel ball bearings to a fine powder (particle diameter <0.1mm). Samples were then sealed and stored at 20°C for up to 9 months."
	LN161: Frequency in Hz Response: corrected 2 instances (lines 161, 168) LN188: rev/min - rpm

Response: corrected (line 191 and 375) LN190: micron symbol µ Response: corrected (line 193) LN192-197: replace incubation with heated (one presumes this was not conducted in an incubation chamber. Response: "incubation" has been changed to "heated" (lines 197, 198)
LN195: magnetic stirrer hotplate corrected (line 198)
LN200: krmp? corrected to "8000 rpm" (line 203)
LN215: small letter p to represent para-nitrocatechol corrected (line 218)
LN215: check consistency re: sulfate vs sulphate Response: changed all instances of "sulphate" to "sulfate" (line 218)
LN218: Is DAE meant to be DEAE? Response: Yes, corrected (line 221)
LN223: For reduction of disulfide linkages, from dimerized please spell out TCEP. Response: changed and added. Now reads: " For the reduction of disulphide linkages, from dimerized desulfoglucosatavin in E. sativa extracts 3g TCEP (Tris(2- carboxyethyl)phosphine hydrochloride powder Sigma, C4706) was" (line 226 and 227)
LN228: High performance liquid chromatography Response: amended (line 228)
LN 492: 'krpm' vs rpm? Response: Changed "8 krpm" to "8000 rpm" for lines 494 and 496.
LN246: Provide a bold title for Mass Spectrometry Response: added (line 250)
LN257: First mention of TBA and ACN please spell out in first instance Response: amended (line 262 and 263)
LN273: Capital T used for time, check consistency earlier t is used Response: all "T=" have been changed to "t=" (lines 278, 636)
Acknowledgements have been amended to recognise the time and effort contributed by the reviewers (line 520).

Click here to view linked References

1 2 3	1	
4 5 6	2	Development of an efficient glucosinolate extraction method
7 8 0	3	Doheny-Adams T., Redeker K., Kittipol V., Bancroft I., Hartley S. E.
9 10 11	4	
12 13 14 15	5	
17 18 19	6	Author information
20 21	7	All Authors are employed at the University of York, in the United Kingdom
22 23 24	8	Postal address:
25 26	9	Department of Biology
27 28 29	10	University of York
30 31	11	Wentworth Way, York
32 33 34	12	YO10 5DD
35 35 36	13	United Kingdom
38 39	14	
40 41 42 43	15	Timothy Doheny-Adams: tim.doheny-adams@york.ac.uk
44 45 46	16	Kelly Redeker: kelly.redeker@york.ac.uk
47 48 49	17	Varanya Kittipol: varanya.kittipol@york.ac.uk
50 51 52	18	Ian Bancroft: ian.bancroft@york.ac.uk
53 54 55	19	Sue Hartley: sue.hartley@york.ac.uk
56 57 58 59 60	20	
62 63 64 65		1

22 Abstract

23 Background

Glucosinolates, anionic sulfur rich secondary metabolites, have been extensively studied because of
their occurrence in the agriculturally important brassicacae and their impact on human and animal
health. There is also increasing interest in the biofumigant properties of toxic glucosinolate
hydrolysis products as a method to control agricultural pests. Evaluating biofumigation potential
requires rapid and accurate quantification of glucosinolates, but current commonly used methods of
extraction prior to analysis involve a number of time consuming and hazardous steps; this study
aimed to develop an improved method for glucosinolate extraction.

31 Results

Three methods previously used to extract glucosinolates from brassicacae tissues, namely extraction in cold methanol, extraction in boiling methanol, and extraction in boiling water were compared across tissue type (root, stem leaf) and four brassicacae species (B. juncea, S. alba, R. sativus, and E. sativa). Cold methanol extraction was shown to perform as well or better than all other tested methods for extraction of glucosinolates with the exception of glucoraphasatin in R. sativus shoots. It was also demonstrated that lyophilisation methods, routinely used during extraction to allow tissue disruption, can reduce final glucosinolate concentrations and that extracting from frozen wet tissue samples in cold 80% methanol is more effective.

Conclusions

We present a simplified method for extracting glucosinolates from plant tissues which does not
require the use of a freeze drier or boiling methanol, and is therefore less hazardous, and more time
and cost effective. The presented method has been shown to have comparable or improved
glucosinolate extraction efficiency relative to the commonly used ISO method for major

glucosinolates in the Brassicaceae species studied: sinigrin and gluconasturtiin in B. juncea; sinalbin, glucotropaeolin, and gluconasturtiin in S. alba; glucoraphenin and glucoraphasatin in R. sativus; and glucosatavin, glucoerucin and glucoraphanin in E. sativa.

Background

Glucosinolates, B-thioglucoside N-hydroxysulfate derivatives, are secondary metabolites found in brassicaceae and related families [1]. Over 120 glucosinolates, which differ in variable aglycone side chains derived from an alpha-amino acid, have been identified and classified into aliphatic, aromatic and indole glucosinolates [2, 3]. Due to their prevalence in cultivated vegetables, spices, oils and animal feed, glucosinolates and their hydrolysis products have been much studied in the context of their effects on human and animal nutrition [4, 5]. Glucosinolates and their breakdown products have also been a focus of studies in dietary prevention of disorders linked to oxidative stress such as cancer and gastric ulcers [2, 6, 7] and more recently, potential undesirable dietary effects such as genotoxicity of glucosinolate breakdown products in broccoli [8] and Pak Choi [9]. The breakdown of glucosinolates has also been studied because of their potential use as agricultural pesticides in a technique known as biofumigation. In biofumigation a glucosinolate-rich crop is mulched into the field, releasing toxic secondary glucosinolate by-products, in order to reduce the incidence of pests, weeds and diseases in the following arable and horticultural crops [10, 11, 12, 13].

Evaluating biofumigation potential requires rapid and accurate quantification of glucosinolates, but current commonly used methods of extraction prior to analysis involve a number of time consuming and potentially hazardous steps. These steps are (i) lyophilisation, or freeze drying, and tissue disruption, (ii) extraction in water or methanol, (iii) purification of extract, typically by desulfation on DEAE sephadex, and (iv) separation and analysis of (desulfo)glucosinolates. These steps are outlined

in figure 1 and discussed in more depth below. This study aimed to improve glucosinolate extraction methods by finding alternatives to commonly used steps which are unnecessary or likely to introduce variability.

INSERT FIGURE 1 HERE

Myrosinase, an enzyme found in brassicacae and compartmentalised in cells in close proximity to glucosinolates, is responsible for hydrolysing glucosinolates upon plant tissue disruption. Accurate analysis of glucosinolates therefore requires inactivation of myrosinase prior to tissue disruption. This is achieved by first freezing then freeze drying the tissue which allows disruption by milling or grinding to occur in the absence of water (fig 1). Lyophilisation, or freeze drying, is used to remove water from glucosinolate-containing tissues while preventing myrosinase mediated glucosinolate hydrolysis through thermal inhibition. Publications on freeze drying plant tissue have focussed primarily on the production of heat or its implications in generating oxygen sensitive foodstuffs (e.g.-space, military or extreme-sport foodstuffs and instant coffee) [14]. To our knowledge, no study has yet examined the efficiency of freeze drying in maintaining glucosinolate concentrations. Freeze drying functions on the principle of sublimation: pressure is reduced below the triple point of water (6.12 mbar, 0.01°C) at which point sublimation of ice from the sample occurs. The cooling effect of sublimation should be high enough to ensure the sample remains below 0°C for the initial stage of freeze drying, thus minimizing enzyme-driven glucosinolate hydrolysis. Rapid sample loading and rapid initial pressure drop are also required to avoid sample defrosting before pressure is reduced below 6.12 mbar. Leaves have a high surface area to volume ratio and may defrost quickly, activating myrosinase and reducing final glucosinolate concentration. Despite the importance of the freeze drying process in glucosinolate extraction, many authors do not report details which are likely to affect final concentrations of glucosinolates (e.g.- how samples are transported, temperature of the room, whether a heating/cold plate is used and time taken for the pressure to drop).

The most commonly used methods for extraction of glucosinolates from plant material are based on the ISO 9167-1 method [15; highlighted in grey in figure 1], which was designed for extraction of glucosinolates from *B. napus* seed and has been adapted to suit the needs of researchers examining glucosinolate profiles of other plant species and tissue types. Although freeze drying is not explicitly detailed in the ISO 9167-1 method, it is an implicit requirement in order to avoid myrosinase mediated glucosinolate hydrolysis during disruption of leaf, stem or root tissues. Once the plant tissue is prepared, the ISO 9167-1 extraction is carried out at 75°C in 70% methanol for 10 minutes. Heating the sample is thought to be an essential step to denature myrosinase, thus preventing enzymatic hydrolysis of glucosinolates [16]. Samples are subsequently desulfated by ion exchange chromatography on a DEAE sephadex column to remove impurities. Desulfoglucosinolates are then separated and identified using HPLC with a reverse phase C18 column and a UV or MS detector. Hazards associated with boiling methanol [17] and the time required for extractions using this method have led researchers to seek alternatives. Replacing heated methanol with boiling water is reported to have comparable [18, 19], and in some cases better [20], extraction efficiencies. Although most glucosinolates are thermostable for the typical 10-30 minute heating period, indole glucosinolates such as 4-hydroxy-glucobrassicin and 4-methoxyglucobrassicin have been reported to degrade quickly at temperatures below 100°C [21]. In addition, prior to 2002 the major glucosinolate in leaves of *E. sativa*, 4-mercaptobutyl glucosinolate, was missed because it self-dimerises via formation of disulphide linkages during extraction [22]. A major challenge therefore to ensuring consistent and repeatable GSL analysis is to create extraction conditions in which myrosinase is inactive, and glucosinolates do not self-react or degrade. A single study, conducted exclusively on radish roots, has demonstrated that cold extraction in 80% methanol does not cause appreciable reduction in glucosinolate concentrations compared to more conventional heated extraction methods [23]. However, myrosinase activity can vary dramatically [24] and whether this method is suitable for extraction of glucosinolates from other glucosinolate containing plants has not previously been assessed.

A desulfation step is often carried out post extraction to purify desulfoglucosinolates and improve accuracy and identification from HPLC. However, the desulfation reaction of glucosinolates can be affected by feedback inhibition of the enzyme which causes incomplete desulfation of glucosinolates [25]. In addition, rhamnopyranosyloxy-benzyl glucosinolates extracted from M. oleifera have been shown to be completely converted and degraded by the desulfation purification step [26]. Due to these drawbacks, and the additional time and potential error extra steps can introduce, some authors have skipped the purification and desulfation steps entirely [19, 26, 27] (fig 1). We have tested each stage of glucosinolate analysis from the roots, stems and leaves of B. juncea, S. alba, R. sativus, E. sativa and B. napus and suggest a number of adjustments/improvements which can be made to reduce the costs, time and variability associated with glucosinolate analysis. Specifically, this study aims to address the following questions: 1) How do lyophilisation conditions affect glucosinolate concentrations? 2) Is lyophilisation a necessary step for glucosinolate extraction from green tissues? 3) Do extractions in hot methanol, cold methanol and boiling water yield comparable glucosinolate concentrations across a range of brassicacae species and tissue types? 4) How do desulfation time and enzyme concentration affect final glucosinolate concentrations? 5) Is desulfation a necessary step for glucosinolate extraction from green tissue? Materials and methods

Experiment	Fig	Species	Tissue	Freeze	Extraction	Desulfation	HPLC
				drying/tis			
				sue			
				disruption			
1-Effect of freeze drier	2	B. napus	Leaves	FD-A or	Cold	0.3 U/ml	ISO 9167-1
on GSL concentration				FD-B	methanol	for 24 H	method
				/mill			
2-Comparison of GSL	3	B. napus	Leaves	FD-A or	Cold	0.3 U/ml	ISO 9167-1
extraction from freeze				-20°C	methanol	for 24 H	method
dried tissue with				methanol			
extraction from wet							
tissue							
3-Comparison of	6, 7	R. sativus	Leaves,	FD-A	Hot	0.3 U/ml	ISO 9167-1
extraction methods		B. juncea	Stems,		methanol,	for 24 H	method
		S. alba	Roots		Cold		
		E. sativa			methanol,		
					Boiling		
					water		
4-Comparison of	8,9	R. sativus	Leaves,	FD-A	Cold	0.3 U/ml	ISO 9167-1
desulfation/purification		B. juncea	Stems,		methanol	for 12, 24,	Method for
methods		S. alba	Roots			48 H, and	desulfoGSL,
		E. sativa				5U/ml for	Herzallah
						16 H or	and Holly
						filtration	method for
							intact GSLs.

4 Table m1: summary of methods used

B. napus used in the freeze drying tests were grown in 1 L pots filled with Terra-green in a controlled temperature glasshouse (regulated from 17.6°C to 27.7°C). At 3-4 weeks post germination, leaves were removed and halved down the limits of the midrib, excluding the midrib from the final sample. Leaf halves were immediately frozen in liquid nitrogen and stored at -80°C for a maximum of 1 week. B. juncea (cv. ISCI99), R. sativus (cv. Bento), S. alba (cv. Ida Gold) and E. sativa (cv. Nemat) plants were grown by Barworth agriculture ltd. in a sandy loam soil dominated fields (coordinates: 53.000371, -0.290404) from 31/07/2014 to 25/09/2014. Total stem and total leaves were cut from flowering plants and immediately frozen in liquid nitrogen; root samples were gently washed and dried before freezing in liquid nitrogen. Samples were stored at -80°C for a maximum of 2 months.

Freeze drying

Samples wrapped loosely in aluminium foil were transported on dry ice and loaded into one of two freeze driers (table m2). Maximum loading time was 30 seconds.

Freeze drier	Room temp	Cooling	Time to	Lowest	Freezer	Model
	(°C)	plate	5 mbar	pressure	temperature	
			(s)	(mbar)	(°C)	
А	22	Yes	90	0.12	-45	Lyotrap, LTE
						scientific ltd.
						1 chamber.
В	28	No	65	0.16	-53	Thermo, Heto
						Powerdry LL3000.
						4-6 chambers.

Table m2: Freeze drier characteristics

Tissue disruption

(i) Freeze dried plant tissue was homogenised to a roughly ground powder (approximately 0.1cm particle size) using a grinder (Lloytron, E5601BK) Homogenised ground samples were milled at a frequency of 20 Hz for 10 minutes (Retch, MM400) with 2 steel ball bearings to a fine powder (particle diameter <0.1mm). Samples were then sealed and stored at 20°C for up to 9 months.

(ii) Frozen fresh B. napus leaf halves (experiment 2, table m1) were placed in 2ml eppendorf vials and stored at -20°C. 1.755ml of 80% methanol precooled at -20°C, 25µl of 5mM sinigrin and 2 small ball bearings were added. Samples were milled for 10 minutes at frequency 20 Hz (TissueLyser II, Qiagen). Final concentrations of methanol were estimated by incorporating average leaf moisture content of fresh B. napus leaves according to equation (1). Final concentration of methanol ranged from 79.3% to 79.9% and leaf moisture content accounted for less than 1% of final liquid volume.

(1)
$$C_{MeOHf} = \frac{c_{MeOHi} \times V_{MeOHi}}{m_{av} \times m_{dl} + V_{MeOHi}}$$

Where c_{MeOHf} is final methanol concentration (%)

c_{MeOHi} is initial methanol concentration (90%)

V_{MeOHi} is initial methanol volume (1.755 ml)

m_{av} is the average moisture content per dry weight (in this case 0.22 ml/g)

m_{dl} dry mass of leaf sample (g)

Glucosinolate extraction

Extractions were carried out in one of three ways (fig 1m). In each case 50µl of a 5mM

gluctropaeolin (for B. juncea samples) or 20 mM sinigrin (for all other samples) internal standard was

added:

Hot methanol extraction (based on the ISO 9167-1 method):

0.1g of plant material was preheated at 75°C for 3 minutes in a 20ml falcon tube. 4.95ml of 70:30 methanol:water, preheated to 75°C and the internal standard was added. The sample was incubated at 75°C for 10 minutes, and manually shaken every 2 minutes. The sample was then centrifuged at 4000 rpm (Jouan, model:B 3.11) for 10 minutes. Supernatent was stored at -20°C or desulfated directly.

Cold methanol extraction (Ishida et al. 2011, [23]):

5 ml of 80:20 methanol:water at 20°C was added to 0.1g plant tissue and the internal standard was added. The sample was shaken and left to stand for 30 minutes at room temperature. The sample was then mixed at 70 rpm with a platform rocker for a further 30 minutes (Bibby, STR6) before centrifugation at 4000 rpm (Jouan, model:B 3.11) for 10 minutes. Supernatent was then filtered through a 0.22 µm syringe filter (Millex GP) for direct injection on HPLC, or unfiltered if applied to sephadex column in a purification step.

Boiling water extraction (adapted from Herzallah and Holley, 2012, [19]):

25 ml of boiling water was added to 0.1g of freeze dried and milled plant tissue in a 150ml erlenmeyer flask and the internal standard was added. Sample was heated at 100°C and stirred with a magnetic stirrer hot plate for 10 minutes. Sample was heated for a further 4 H at 70°C before centrifugation at 4000 rpm (Jouan, model:B 3.11) for 10 minutes. Sample was topped up to 20ml with deionised water.

Purification and determination of activity of sulfatase:

Sulfatase from Helix pomatia type H-1 (Sigma, S9626) was purified according to Wathalet et al. (1999) [25]. 25 mg of sulfatase was added to 1ml 40% ethanol and centrifuged at 8000 rmp for 1 minute (eppendorf centrifuge, 54151). The supernatant was transferred to a fresh 2ml eppendorf tube, 1ml of pure ethanol was added to precipitate the sulfatase before being centrifuged at 8krmp

for 1 minute. The supernatant was discarded and the sulfatase pellet air dried and redissolved in 2ml of water.

Activity of sulfatase was determined based on the ISO 9167-1 method. 1 ml of buffered 0.15mM sinigrin solution (3ml of 5mM sinigrin, adjusted to 100ml with a solution containing 40ml 0.2% ethylene diamine, 73ml 0.2% acetic acid; adjusted to pH 5.8) in a quartz cuvette was placed in a UV spectrometer set to 229nm. At t=0, 25µl of diluted and undiluted purified sulfatase was added to the cuvette and measurements taken over the course of 4 hours. The tangent to t=0 was plotted and its gradient ($\Delta A/\Delta t$) measured. Activity was calculated using equation (2):

(2) Activity (U/ml) = $\frac{\Delta A \times 5.7}{\Delta t A_e}$

Where $\Delta A/\Delta t$ is the gradient at t=0 and A_e is the difference between absorbance at equilibrium and absorbance at t=0.

The activity for Sulfatase from Helix pomatia type H-1 (Sigma, S9626) given by the supplier is determined by desulfation of p-nitrocatechol sulfate and is an order of magnitude higher than the activity measured for desulfation of sinigrin using this method.

Desulfation of glucosinolates

As per the ISO 9067-1 method, columns were prepared with 0.5ml sephadex slurry (2g DEAE sephadex beads in 30 ml 2M acetic acid.) and activated with 2ml imizadole formate (6M). Columns were washed twice with 1ml water. The column was washed twice with 1ml 20mM sodium acetate (pH 4.0) and 75µl of purified sulfatase was added (5U/ml or 0.3U/ml). Columns were incubated at room temperature for either 12, 24 or 48 hours before elution of desulfoglucosinolates with two 1ml volumes of water. For the reduction of disulphide linkages, from dimerized desulfoglucosatavin in E. sativa extracts 3g TCEP (Tris(2-carboxyethyl)phosphine hydrochloride powder Sigma, C4706) was added to 1 ml of desulfated extract. Desulfoglucosinolates were stored at -20°C before high performance liquid chromatography analysis.

For the high sulfatase treatment, between 0.5 and 1 ml of sample was added due to insufficient sample volume remaining.

HPLC

A Waters 600E system controller attached to a Waters 717 autosampler, Waters 996 photodiode array detector and SphereClone 5µ ODS(2) column (Phenomonex) were used for separation and detection of desulfo and intact glucosinolates.

HPLC analysis of desulfoglucosinolates – adapted from ISO 9167-1

A reverse phase C18 column (Phenomonex, SphereClone 5µ ODS(2), 150mm x 4.6mm) was

equilibrated for 30 min with a mobile phase which consisted of 100% diH₂O. Flow rate was set to

1ml/min and samples separated according to programme for desulfoglucosinolates detailed in table

m3. Mobile phase solutions were degassed for 30 minutes in a sonicator (Decon, Sussex England).

Solution A: 100% diH₂O

Solution B: 70:30, diH₂O:acetonitrile

Time	% solution A	% solution B	Transition
0	100	0	
30	0	100	Linear gradient
35	0	100	
40	100	0	Linear gradient
50	100	0	

Table m3: mobile phase conditions for separation of desulfoglucosinolates.

Desulfoglucosinolates were quantified using 229nm wavelength within the UV spectrum. The HPLC PDA detector allowed a full spectrum analysis from 180 to 800nm, allowing comparative UV-Visible spectra analysis, which aided in identifying unknown glucosinolates. Through standard injections and HPLC-MS identification we were able to confirm the id's of these reported glucosinolates. Desulfated

1	248	purified standards: sinigrin (sigma aldrich), glucotropaeolin, glucoraphenin, glucoraphanin,
1 2 3 4	249	glucerucin, glucobrassicin, gluconasturtiin, sinalbin, progoitrin and glucoiberin (phytoplan).
5 6	250	Mass spectrometry
7 8 9	251	Major glucosinolates for which no commercial standard is available were identified using an MS
10 11	252	detector (Bruker maXis UHR-TOF) with the following settings:
12 13	253	Source: Standard electrospray (flow split 1/10 from LC)
14 15 16	254	Nebulizer: 2.0 bar
17 18	255	Dry gas: 6.0 L/min
19 20	256	Dry gas heater: 250C
21 22 23	257	Capillary voltage: 3500 V
24 25	258	Ion polarity: positive
26 27 20	259	Spectra rate: 1Hz
29 30 31	260	HPLC analysis of intact glucosinolates – adapted from Herzallah and Holly, 2012 [19]
32 33	261	A C18 column (Phenomonex, SphereClone 5u ODS(2)) was equilibrated for 3 h with a mobile phase
34 35 36	262	which consisted of 80 mL (0.02 M) TBA <mark>(tetrabutylammonium bromide)</mark> and 20 mL ACN
37 38	263	(acetonitrile) with detection at 229 nm. The flow rate was set at 1.0 ml/min and separated according
39 40	264	to programme for desulfoglucosinolates detailed in table m3.
41 42 43	265	Solution A: 100% TBA (0.02M)
44 45 46	266	Solution B: 70:30, TBA (0.02M):acetonitrile
40		
48 49 50	267	Glucosinolates were quantified using the chromatogram from 229nm and standard curves were
50 51 52	268	constructed using pure sinigrin (sigma aldrich), glucotropaeolin, glucoraphenin, glucoraphanin,
53 54 55	269	glucerucin, glucobrassicin, gluconasturtiin, sinalbin, progoitrin and glucoiberin (phytoplan).
56 57	270	In the case of glucoraphasatin in <i>R. sativus</i> leaves and glucotropaeolin in <i>B. juncea</i> minor alterations
58 59	271	were made to avoid peaks co-eluting. The mobile phase programme for <i>R. sativus</i> leaves was 100% A
60 61 62 63 64		13
h h		

for 5 minutes, followed by a 35 minute linear gradient to 66% B followed by a 5 minute linear gradient to 100% B followed by a 5 minute linear gradient to 100% A . For B. juncea leaves, an isocratic 85:15, TBA (0.02M):acetonitrile mobile phase for 70 minutes was used.

Determination of myrosinase activity

Activity of pure myrosinase was tested in water and 80% methanol solutions containing 0.25mM sinigrin and 0.1 mM ascorbic acid, a myrosinase cofactor (Burmeister et al. 2000). Myrosinase was added at t=0 and absorbance of sinigrin at 229nm was measured over the course of an hour. Activity was measured at room temperature (25°C).

Determination of glucosinolate thermostability

A 50µl of 10mM sinigrin, 10mM glucotropaeolin, 10mM glucobrassicin solution was added to 0.95ml water or 70% methanol preheated to 100°C or 75°C respectively and sealed in 1.5ml eppendorf tubes. Samples were maintained at either 100°C or 75°C for 5, 10, 30 and 60 minutes and intact glucosinolate concentrations analysed with HPLC following the adapted Herzallah and Holly (2012) method [19].

Calculation of glucosinolate content

Glucosinolate content, expressed in µmol/g were calculated according to the ISO 9067-1 method (equation (3)):

(3) Glucosinolate content =
$$\frac{A_g}{A_s} \times \frac{n}{m} \times K_g \times \frac{100}{100-w}$$

Where A_g is the peak area corresponding to desulfoglucosinolate;

As is the peak area corresponding to internal standard;

n is the quantity, in micromoles, of the internal standard;

m is the mass of the test portion;

K_g is the response factor of the desulfoglucosinolate relative to the internal standard;

w is the moisture and volatile matter content, expressed as a percentage by mass of the test sample.

296 Statistical analysis

Paired two tailed t-test analysis were carried out on total *B. napus* glucosinolate content per leaf half
in experiments 1 and 2 with Microsoft excel (table m1). For determination of significance of effect of
method on final glucosinolate content estimates in experiments 3 and 4 (table m1), repeat measure
ANOVA analyses were carried out for each glucosinolate with R statistical software package (version
3.3.1).

Results and discussion

1 Lyophilisation

Modifications to the ISO9167-1 method (specifically created for the extraction and analysis of glucosinolates from oil rape seed samples) are required for analysis of plant green tissues (leaves, stems and roots). A number of prior-to-analysis steps, such as sampling in the field, cleaning (if required), freezing, crushing, storage or/and shipping and reduction of sample amount have been discussed by Wathelet et al. (2004) and are not revisited here [28]. These preliminary steps are followed by lyophilisation, or freeze drying, to remove water from glucosinolate containing tissues while preventing myrosinase mediated glucosinolate hydrolysis through thermal inhibition. This process allows subsequent tissue disruption without risking glucosinolate degradation.

We tested reproducibility of glucosinolate concentrations extracted after lyophilisation in separate freeze driers (table 1). Fresh *B. napus* leaves were halved, loosely wrapped in foil, flash frozen in liquid nitrogen and transported in dry ice to be dried in separate freeze driers (table 1). Total glucosinolate concentrations were significantly higher in samples dried in freeze drier A than freeze drier B (fig 2a). In addition, samples dried in freeze drier B developed a darker hue and deformed more than samples in dried in freeze drier A (fig 2b). Plant tissue samples have been shown to deform during the freeze drying process when temperatures exceed the glass transition state and

melting point of water [29]. It is likely that samples placed into freeze drier B may have defrosted before the pressure had reduced below the 6.12 mbar required for sublimation due to higher temperatures and the lack of cooling plate. As a result, enzyme mediated hydrolysis of glucosinolates may have occurred at the initial stage. Additionally, as sublimation slows over time due to the remaining water vapour passing through a dry layer of increasing thickness and because water is increasingly more tissue bound, the sample temperature may have increased to above 0°C in freeze drier B, causing defrosting.

Freeze drier	Room temp	Cooling plate	Time to 5	Lowest	Freezer
	(°C)		mbar (s)	pressure	temperature
				(mbar)	(°C)
А	22	Yes	90	0.12	-45
В	28	No	65	0.16	-53

Table 1: Freeze drier characteristics

INSERT FIGURE 2 HERE

These results underline the need for a more substantive study to assess optimal conditions for

freeze drying plant tissues for glucosinolate analysis. It is clear that differences in freeze drying can

introduce significant variability in retained glucosinolate concentrations (fig 2a).

A cold methanol extraction method may be sufficient to 1) inactivate myrosinase and 2) efficiently

extract glucosinolates, precluding the need for the lyophilisation step altogether. We tested this by

comparing glucosinolates extracted from one half of a B. napus leaf in 80% methanol without freeze

drying against glucosinolates extracted from the other half, first dried in freeze drier A and then extracted using the cold methanol extraction method.

No significant difference in final glucosinolate concentration was found between the two methods (fig. 3). Freeze drying is an energy intensive and costly process requiring long drying times under continuous vacuum and the significant effect of freeze drier parameters on final glucosinolate concentrations (fig. 2a) highlights a potential source of variation between studies. If long term storage of plant tissue samples is not required, skipping the freeze drying step and extracting glucosinolates directly into cold methanol (-20°C) is cheaper, quicker and less hazardous.

INSERT FIGURE 3 HERE

2 Extraction

Some authors have highlighted that glucosinolates, specifically indole glucosinolates, are heat sensitive and are significantly degraded in temperatures ≥75°C in less than 10 minutes [21]. This has serious implications for accuracy and reliability of the ISO 9167-1 extraction method, which recommends extractions occur in boiling 70% methanol (75° C) for 10 minutes, as well as the less commonly used boiling water extraction (100° C). In order to first test whether thermal degradation of glucosinolates was likely to occur with these methods we measured the glucosinolate concentrations of pure sinigrin (aliphatic), glucotropaeolin (aromatic) and glucobrassicin (indole) in boiling water (fig 4) and boiling 70% methanol (data not shown). Sinigrin and glucotropaeolin did not significantly decrease over 60 minutes suggesting that extraction in boiling water or methanol is unlikely to affect the concentrations of these glucosinolates. However, glucobrassicin was thermally degraded at 100°C and data from extractions carried out at these temperatures or above (such as with microwave based methods) may underestimate the concentration of glucobrassicin and other indole glucosinolates. Boiling an extract in water for 10 minutes degrades glucobrassicin by an estimated 7%.

364	INSERT FIGURE 4 HERE						
365							
366							
367	Activity of pure myrosin	nase was tested at 25° C in wa	iter and 80% methanol s	solutions containing			
368	0.25mM sinigrin and 0.	1 mM ascorbic acid, a myrosir	nase cofactor [30]. Abso	rbance of sinigrin at			
369	229nm, at room tempe	rature (25°C), was measured	over the course of an ho	our after myrosinase			
370	addition. Myrosinase w	as inactive in 80% methanol (fig 5) suggesting that he	ating methanol at 75°C			
371	for 10 minutes in order	to inactivate myrosinase may	v be an unnecessary step	o for extracting			
372	glucosinolates from pla	nt tissue.					
373		INSERT FIGUE	RE 5 HERE				
374	Glucosinolates from B.	<i>juncea, S. alba, R. sativus</i> and	<i>E. sativa</i> leaves, stems a	and roots were			
375	extracted (i) in boiling v	vater for 10 minutes followed	by a 4h incubation at 7	0°C, (ii) in 70% methanol			
376	at 75°C, or (iii) in 80% n	nethanol at room temperatur	e (~20°C) for 30 minutes	standing followed by			
377	30 minutes shaking at 7	70 <mark>rpm</mark> . All extracts were cent	rifuged and desulfated v	with sulfatase according			
378	to the ISO 9167-1 method. Major glucosinolates from these species can be found in table 2.						
	Common name	Chemical name	Structure	Species, tissue type			
	Sinigrin	2-propenyl	Aliphatic	B. juncea L , S, R			
	Glucoraphenin	4-methylsulfinyl-3-butenyl	Aliphatic	R. sativus L, S, R			
	Glucoraphanin	4-methylsulfinylbutyl	Aliphatic	E. sativa L, S , R			
	Glucosatavin	mercaptobutyl	Aliphatic	E. sativa L , S, R			
	Glucoraphasatin	4-Methylthio-3-butenyl	Aliphatic	R. sativus L, S, R			
	or hydroxyglucoerucin						
	Glucoerucin	Methylthiobutyl	Aliphatic	E. sativa S, R			
				S. alba, R			

Sinalbin	4-Hydroxybenzyl	Aromatic	S. alba L, S, R
Glucotropaeolin	Benzyl	Aromatic	S. alba L, S, R
Gluconasturtiin	phenylethyl	Aromatic	B. juncea R
			S. alba R
Methoxyglucobrassicin	4-methoxy-3-indolylmethyl	Indole	S. alba R

Table 2: Glucosinolates examined in this study. L, S and R correspond to leaf, stem and root respectively. Letters in bold represent major glucosinolates of those tissues (>10 μmol/g dry weight).

Figure 6 compares glucosinolate concentrations obtained using the cold methanol method and boiling water method normalised against the ISO 9167-1 boiling methanol method. For most glucosinolates, across most tissue types and species, the three extraction methods yield similar results. We found that extraction with cold methanol produced a significantly higher estimated concentration of sinalbin in *S. alba* and sinigrin in *B. juncea* than the hot methanol extraction (fig 6). Surprisingly, given the sensitivity of glucobrassicin to thermal degradation (fig 4), extraction in boiling water did not significantly reduce the concentration of the indole glucosinolate: methoxyglucobrassicin relative to the other two methods. However, glucosatavin was extracted with lower efficiency from leaves of *E. sativa* using the boiling water method (fig 6). It seems unlikely that this glucosinolate is less thermostable than other glucosinolates and was therefore degraded by the extraction method since reduced extraction efficiencies are not observed for stem and root samples. There are no published explanations or hypotheses that might help to explain the observed lower extraction efficiencies for glucosatavin using the boiling water method. Glucoraphasatin extraction using cold methanol appears to be significantly less effective than the standard ISO method (fig 6), however this was driven by poor extraction efficiencies from R. sativus stems (fig 7). Ishida et al. reported a significant 5% increase in glucoraphasatin concentrations extracted from R. sativus roots using the cold methanol method [23]. In this study, extraction efficiencies of glucoraphenin in R.

sativus roots with a cold methanol method were comparable to extraction efficiencies using the boiling methanol method (fig 7).

INSERT FIGURE 6 HERE

INSERT FIGURE 7 HERE

No glucosinolates were detected in a subset of samples extracted in cold water indicating the presence of active myrosinase leading to their degradation (data not shown). However, the cold methanol extraction did not significantly affect the concentration of the internal standard relative to the boiling methanol method (data not shown), providing additional evidence that myrosinase is inactivated in 80% methanol without heating (fig 5).

These data demonstrate that 80% cold methanol can be used instead of boiling methanol to extract glucosinolates across a broad spectrum of brassicacae species and tissue types. With the exception of glucoraphasatin in R. sativus shoots, replacing hot 70% methanol with cold 80% methanol did not significantly reduce glucosinolate concentrations, yet marginally increased recovery of sinalbin in S. alba and sinigrin in B. juncea. It is advised, due to reduction in steps and hazard as well as improved or comparable glucosinolate recovery, that a cold methanol extraction is used instead of a boiling methanol extraction for most glucosinolate containing green tissues.

Purification

Purification of extract according to the ISO 9167-1 method is carried out by introducing 1ml of extract to a column containing 0.5ml of sephadex solution. The column is rinsed with a 20mM acetate buffer at pH 4.0 to avoid possible reduction of indole glucosinolates recovery [28]. 75µl of sulfatase solution with an activity above 0.05 U/ml is applied and left to act overnight. We tested the extraction efficiency of the ISO 9167-1 purification step at the described pH 4.0, at 20°C for 12, 24 and 48 hours. Complete desulfation of glucosinolates in rapeseed extract required a minimum of 11 hours in operating conditions of 30°C and pH 5.8 [25] so it was expected that an overnight 12 hour

desulfation period may be insufficient for complete desulfation of samples at room temperature. Figure 8 shows absorbance values for representative desulfoglucosinolate solutions from B. juncea, S. alba, R. sativus and E. sativa extracts treated with sulfatase solution for 12, 24 or 48 hours. In most cases, 12 and 24 hour incubation periods were insufficient for complete desulfation of glucosinolates. Glucoraphenin decreased in all R. sativus leaf samples tested, from 24 to 48 hours, while recovery of the internal standard increased, suggesting that specifically this desulfoglucosinolate is degraded during the purification process (fig 8).

INSERT FIGURE 8 HERE

Not all glucosinolates are desulfated on the column at the same rate [31], meaning that incomplete desulfation of extractions is likely to yield imprecise results: overestimating or underestimating the final concentration of glucosinolates which are desulfated guicker or slower respectively than the internal standard. In addition, relative and total concentrations of glucosinolates and degradation or rearrangement of glucosinolates during this process can also affect final concentrations [26, 31]. Use of higher sulfatase concentrations than outlined in the ISO method has been suggested for glucosinolate analysis in B. napus and B. oleracea [25, 31]. Figure 9 compares relative glucosinolate concentrations from B. juncea, S. alba, R. sativus and E. sativa purified with a low activity sulfatase solution (0.3U/ml) for 12H, 24H and 48H, a high activity sulfatase solution (5U/ml) and intact glucosinolates. All concentrations have been normalised to the intact glucosinolate values. Desulfated glucosinolates concentrations obtained with high concentration sulfatase compared well with intact glucosinolates (fig 9). However, both high sulfatase as well as low sulfatase treatments yielded lower glucoraphenin content estimates. Coupled with the reduction of the recovery of desulfoglucoraphenin from 24H to 48H (fig 8), these data suggest that glucoraphenin is degraded or transformed during the desulfation process.

INSERT FIGURE 9 HERE

Shorter desulfation times and lower sulfatase concentrations resulted in underestimation of the concentrations of glucoraphenin from R. sativus, glucoraphanin and glucosatavin from E. sativa, sinigrin from B. juncea, and sinalbin from S. alba and an overestimation of the concentrations of glucoraphasatin in *R. sativus* roots (fig 9). The overnight (12H-24H) incubation with 0.3 U/ml sulfatase solution yields inaccurate results for most major glucosinolates examined in this study. The ISO9167-1 method suggests that a diluted purified sulfatase solution with an activity exceeding 0.05U/ml should be used, which is shown to be insufficient for glucosinolate analysis from plant samples and conditions examined in this study (fig 9). Instead, if a desulfation step is carried out, use of a higher concentration of purified sulfatase (in this case, 5U/ml) is advised.

In all *E. sativa* leaf samples tested, recovery of monomeric desulfo-glucosatavin decreased and recovery of dimeric desulfo-glucosatavin increased between 24 and 48 hours. Bennet et al. (2002) previously hypothesised that dimeric glucosatavin is unlikely to be found in vivo and is probably an artefact of the extraction process [22]. We can confirm that glucosatavin forms dimers as a result of the desulfation step of the extraction and that without carrying this step out and instead quantifying intact glucosinolates, no dimeric glucosatavin was detected in these samples.

Given that glucoraphenin concentration estimates are lower from methods employing a desulfation step, and that this step is also responsible for the dimerization of glucosatavin, analysis of intact glucosinolates is preferable in most instances. It is out of the scope of this study to compare or improve separation and detection methods but it should be noted that major glucosinolates in this study were accurately measured by a HPLC-UV method adapted from Herzallah and Holley (2012) [19]. For examination of low abundance glucosinolates, and to avoid any potential inaccuracies due to contamination it is advised that an alternative HPLC method such as those suggested in Lee et al. (2013) or Forster et al. (2015) be used instead [26, 32].

471 Suggested method for glucosinolate extraction:

472 Tissue disruption

473 Depending on whether freeze drying is required:

474 1a – Freeze samples loosely wrapped in foil in liquid nitrogen and store at -80°C. Transport samples
475 to freeze drier in dry ice. Rapidly load samples onto a cool plate in freeze drier and ensure the
476 pressure drops to below 5 mbar in under 2 minutes. Mill samples once dried and store in airtight
477 containers in the dark.

479 or

1b – Freeze 50mg samples in liquid nitrogen in 2ml eppendorf tubes and store at -80°C (for larger
samples use larger tubes). Add a volume of 80% methanol precooled to -20°C ensuring that final
methanol concentration remains above 78% according to equation (1) in materials and methods.
Add an appropriate volume of internal standard sinigrin or glucotropaeolin (e.g. 100µM final
concentration). Disrupt tissue by adding 2 small ball bearings and agitating with a tissue lyser (e.g.
tissuelyserII, Qiagen) for 10 minutes at 20 rev/s. Alternatively use a plastic pestle to thoroughly grind
the sample taking care that to keep the media below 0°C. Continue directly to 2b.

Extraction

2a – For freeze dried tissue (1a). To 0.1g tissue, add 5ml of 80% methanol and 50μL of 20mM sinigrin
solution. Then

2b – shake sample once and leave to stand for 30 minutes. Shake sample for a further 30 minutes
(70 rev/s). Centrifuge at 4000 rpm and transfer supernatant to a fresh tube.

492 Desulfation

493 If desulfation is required, a high concentration sulfatase solution should be prepared by dissolving
494 15-25mg sulfatase in 1ml 40% ethanol and centrifuge at 8000 rmp for 1 minute. Transfer

495 supernatant to a fresh 2ml eppendorf tube and add 1ml of pure ethanol to precipitate the sulfatase

and centrifuge at 8000 rpm for 1 minute. Discard the supernatant and air dry the pellet before re-dissolving in 2ml of water. Proceed with desulfation according to ISO9167-1 method.

Conclusions

In this study we compared different methods for extracting and purifying glucosinolates from B. napus, B. junea, S. alba, E. sativa and R. sativus green tissues to highlight unnecessary or hazardous steps. We have presented a simplified method for extracting glucosinolates from plant tissues which does not require the use of a freeze drier or boiling methanol, and is therefore less hazardous, and more time and cost effective. The presented method has been shown to have comparable or improved glucosinolate extraction efficiency relative to the commonly used ISO method for major glucosinolates in the Brassicaceae species studied: sinigrin and gluconasturtiin in B. juncea; sinalbin, glucotropaeolin, and gluconasturtiin in S. alba; glucoraphenin and glucoraphasatin (roots but not shoots) in *R. sativus*; and glucosatavin, glucoerucin and glucoraphanin in *E. sativa*.

Declarations

Authors' contributions

TDA, KR, and SEH organized the project. VK carried out sample preparation and glucosinolate extractions on *B. napus* leaves. TDA performed all other experiments, analyzed the data, and wrote the paper; SEH, IB and KR reviewed and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We are grateful to Catherine Lilley, (University of Leeds), Peter Urwin (University of Leeds), Andy Barker (Barworth Agriculture ltd.) and Helen Barker (Barworth Agriculture ltd.) for collection of leaf, stem and root samples used in this study. We would also like to thank Thomas Hartley (University of

1	519	York) for statistical advice. We would also like to express our gratitude to the BBSRC for funding this
1 2 3	520	study. <mark>Finally, we would like to thank the reviewers for their time and effort in reviewing this</mark>
4 5 6	521	manuscript.
7 8 9	522	Competing interests
10 11 12	523	The authors declare that they have no competing interests.
13 14	524	Funding
15 16 17	525	This work was supported by UK Biotechnology and Biology Sciences Research Council
18 19 20	526	(BB/L002124/1) and (BB/K020463/1).
21 22 23	527	Ethics approval (NA)
24 25 26	528	Consent for publication (NA)
27 28 29	529	Availability of data and Material
30 31	530	Material used in this study is stored at the University of York and is available on request. Datasets
32 33 34	531	analysed in this study are available from the corresponding author on request.
36 37 38	532	
39 40	533	References
41 42 43	534	[1] Clarke DB. Glucosinolates, structures and analysis in food. Analytical Methods. 2010;2(4):310-25.
44 45	535	[2] Fahey JW, Zalcmann AT, Talalay P. The chemical diversity and distribution of glucosinolates and
48 47 48	536	isothiocyanates among plants. Phytochemistry. 2001 Jan 31;56(1):5-1.
49 50	537	[3] Mithen RF, Dekker M, Verkerk R, Rabot S, Johnson IT. The nutritional significance, biosynthesis
51 52 53	538	and bioavailability of glucosinolates in human foods. Journal of the Science of Food and Agriculture.
54 55	539	2000 May 15;80(7):967-84.
56 57 58	540	[4] VanEtten CH, Daxenbichler ME, Wolff IA. Natural glucosinolates (thioglucosides) in foods and
59 60 61	541	teeds. Journal of Agricultural and Food Chemistry. 1969 May;17(3):483-91.
62 63 64 65		25

[5] Cartea ME, Velasco P. Glucosinolates in Brassica foods: bioavailability in food and significance for human health. Phytochemistry reviews. 2008 Jul 1;7(2):213-29.

[6] Talalay P, Fahey JW. Phytochemicals from cruciferous plants protect against cancer by

modulating carcinogen metabolism. The Journal of Nutrition. 2001 Nov 1;131(11):3027S-33S.

[7] Shapiro TA, Fahey JW, Wade KL, Stephenson KK, Talalay P. Chemoprotective glucosinolates and

isothiocyanates of broccoli sprouts metabolism and excretion in humans. Cancer Epidemiology

Biomarkers & Prevention. 2001 May 1;10(5):501-8.

[8] Latté KP, Appel KE, Lampen A. Health benefits and possible risks of broccoli-an overview. Food and Chemical Toxicology. 2011 Dec 31;49(12):3287-309.

[9] Wiesner M, Schreiner M, Glatt H. High mutagenic activity of juice from pak choi (Brassica rapa

ssp. chinensis) sprouts due to its content of 1-methoxy-3-indolylmethyl glucosinolate, and its

enhancement by elicitation with methyl jasmonate. Food and Chemical Toxicology. 2014 May

31;67:10-6.

[10] Ngala BM, Haydock PP, Woods S, Back MA. Biofumigation with Brassica juncea, Raphanus

sativus and Eruca sativa for the management of field populations of the potato cyst nematode

Globodera pallida. Pest management science. 2015 May 1;71(5):759-69.

[11] Lord JS, Lazzeri L, Atkinson HJ, Urwin PE. Biofumigation for control of pale potato cyst

nematodes: activity of Brassica leaf extracts and green manures on Globodera pallida in vitro and in

soil. Journal of agricultural and food chemistry. 2011 Jun 30;59(14):7882-90.

[12] Mattner SW, Porter IJ, Gounder RK, Shanks AL, Wren DJ, Allen D. Factors that impact on the

ability of biofumigants to suppress fungal pathogens and weeds of strawberry. Crop Protection. 2008 Aug 31;27(8):1165-73.

[13] Bellostas N, Kachlicki P, Sørensen JC, Sørensen H. Glucosinolate profiling of seeds and sprouts of B. oleracea varieties used for food. Scientia Horticulturae. 2007 Nov 20;114(4):234-42.

[14] Ratti C. Hot air and freeze-drying of high-value foods: a review. Journal of food engineering.

	1	568	[15] ISO 9167-1, 1992 (NA 057-05-05 AA – Joint committee of DIN and DGF for the analysis of fats,
	1 2 3	569	oils and products thereof, related and primary products. (2012): Rapeseed – Determination of
	4 5	570	glucosinolate content – Part 1: Method using high-performance liquid chromatography (ISO 9167–
	6 7 0	571	1:1992/DAM 1:2012), German version EN ISO 9167-1:1995/prA1:2012.)
	。 9 10	572	[16] Ares AM, Nozal MJ, Bernal JL, Bernal J. Optimized extraction, separation and quantification of
-	11 12	573	twelve intact glucosinolates in broccoli leaves. Food chemistry. 2014 Jun 1;152:66-74.
-	13 14 15	574	[17] Church AS, Witting MD. Laboratory testing in ethanol, methanol, ethylene glycol, and
-	16 17	575	isopropanol toxicities. The Journal of emergency medicine. 1997 Oct 31;15(5):687-92.
-	18 19	576	[18] Rangkadilok N, Nicolas ME, Bennett RN, Premier RR, Eagling DR, Taylor PW. Determination of
4	20 21 22	577	sinigrin and glucoraphanin in Brassica species using a simple extraction method combined with ion
	23 24	578	pair HPLC analysis. Scientia Horticulturae. 2002 Dec 6;96(1):27-41.
	25 26	579	[19] Herzallah S, Holley R. Determination of sinigrin, sinalbin, allyl-and benzyl isothiocyanates by RI
	27 28 29	580	HPLC in mustard powder extracts. LWT-Food Science and Technology. 2012 Jul 31;47(2):293-9.
	30 31	581	[20] D. F. Stoin and R. D. Dogaru, Bull. USAMV-CN, 2007, 63, 77–82
	32 33	582	[21] Oerlemans K, Barrett DM, Suades CB, Verkerk R, Dekker M. Thermal degradation of
	34 35 36	583	glucosinolates in red cabbage. Food Chemistry. 2006 Mar 31;95(1):19-29.
	37 38	584	[22] Bennett RN, Mellon FA, Botting NP, Eagles J, Rosa EA, Williamson G. Identification of the majo
-	39 40 41	585	glucosinolate (4-mercaptobutyl glucosinolate) in leaves of Eruca sativa L.(salad rocket).
2	42 43	586	Phytochemistry. 2002 Sep 30;61(1):25-30.
2	44 45	587	[23] Ishida M, Kakizaki T, Ohara T, Morimitsu Y. Development of a simple and rapid extraction
2	46 47 48	588	method of glucosinolates from radish roots. Breeding science. 2011;61(2):208-11.
	49 50	589	[24] Piekarska A, Kusznierewicz B, Meller M, Dziedziul K, Namieśnik J, Bartoszek A. Myrosinase
	51 52	590	activity in different plant samples; optimisation of measurement conditions for spectrophotometri
	53 54 55	591	and pH-stat methods. Industrial Crops and Products. 2013 Oct 31;50:58-67.
	56 57	592	[25] Wathelet JP, Mabon N, Marlier M. Determination of glucosinolates in rapeseed improvement
1	58 59	593	the official HPLC ISO method (precision and speed). In Proceedings of the 10th International
(oU 61 62		27
(63 64		27
(65		

9	oils and products thereof, related and primary products. (2012): Rapeseed – Determination of
0	glucosinolate content – Part 1: Method using high-performance liquid chromatography (ISO 9167–
1	1:1992/DAM 1:2012), German version EN ISO 9167-1:1995/prA1:2012.)
2	[16] Ares AM, Nozal MJ, Bernal JL, Bernal J. Optimized extraction, separation and quantification of
3	twelve intact glucosinolates in broccoli leaves. Food chemistry. 2014 Jun 1;152:66-74.
4	[17] Church AS, Witting MD. Laboratory testing in ethanol, methanol, ethylene glycol, and
5	isopropanol toxicities. The Journal of emergency medicine. 1997 Oct 31;15(5):687-92.
6	[18] Rangkadilok N, Nicolas ME, Bennett RN, Premier RR, Eagling DR, Taylor PW. Determination of
7	sinigrin and glucoraphanin in Brassica species using a simple extraction method combined with ion-
8	pair HPLC analysis. Scientia Horticulturae. 2002 Dec 6;96(1):27-41.
9	[19] Herzallah S, Holley R. Determination of sinigrin, sinalbin, allyl-and benzyl isothiocyanates by RP-
0	HPLC in mustard powder extracts. LWT-Food Science and Technology. 2012 Jul 31;47(2):293-9.
1	[20] D. F. Stoin and R. D. Dogaru, Bull. USAMV-CN, 2007, 63, 77–82
2	[21] Oerlemans K, Barrett DM, Suades CB, Verkerk R, Dekker M. Thermal degradation of
3	glucosinolates in red cabbage. Food Chemistry. 2006 Mar 31;95(1):19-29.
4	[22] Bennett RN, Mellon FA, Botting NP, Eagles J, Rosa EA, Williamson G. Identification of the major
5	glucosinolate (4-mercaptobutyl glucosinolate) in leaves of Eruca sativa L.(salad rocket).
6	Phytochemistry. 2002 Sep 30;61(1):25-30.
7	[23] Ishida M, Kakizaki T, Ohara T, Morimitsu Y. Development of a simple and rapid extraction
8	method of glucosinolates from radish roots. Breeding science. 2011;61(2):208-11.
9	[24] Piekarska A, Kusznierewicz B, Meller M, Dziedziul K, Namieśnik J, Bartoszek A. Myrosinase
0	activity in different plant samples; optimisation of measurement conditions for spectrophotometric
1	and pH-stat methods. Industrial Crops and Products. 2013 Oct 31;50:58-67.
2	[25] Wathelet JP, Mabon N, Marlier M. Determination of glucosinolates in rapeseed improvement of
3	the official HPLC ISO method (precision and speed). In Proceedings of the 10th International
	27

594	Rapeseed Congress 1999 (p. 185). Gosford, NSW, Australia: The Regional Institute Ltd
595	[26] Förster N, Ulrichs C, Schreiner M, Müller CT, Mewis I. Development of a reliable extraction and
596	quantification method for glucosinolates in Moringa oleifera. Food chemistry. 2015 Jan 1;166:456-
597	64.
598	[27] Troyer JK, Stephenson KK, Fahey JW. Analysis of glucosinolates from broccoli and other
599	cruciferous vegetables by hydrophilic interaction liquid chromatography. Journal of Chromatography
600	A. 2001 Jun 15;919(2):299-304.
601	[28] Wathelet JP, Iori R, Leoni O, Rollin P, Quinsac A, Palmieri S. Guidelines for glucosinolate analysis
602	in green tissues used for biofumigation. Agroindustria. 2004;3(3):257-66.
603	[29] Karathanos VT, Anglea SA, Karel M. Structural collapse of plant materials during freeze-drying.
604	Journal of Thermal Analysis and Calorimetry. 1996 Nov 1;47(5):1451-61.
605	[30] Burmeister WP, Cottaz S, Rollin P, Vasella A, Henrissat B. High resolution X-ray crystallography
606	shows that ascorbate is a cofactor for myrosinase and substitutes for the function of the catalytic
607	base. Journal of Biological Chemistry. 2000 Dec 15;275(50):39385-93.
608	[31] Hennig K, Verkerk R, Bonnema G, Dekker M. Pitfalls in the desulphation of glucosinolates in a
609	high-throughput assay. Food chemistry. 2012 Oct 15;134(4):2355-61.
610	[32] Lee JG, Bonnema G, Zhang N, Kwak JH, de Vos RC, Beekwilder J. Evaluation of glucosinolate
611	variation in a collection of turnip (Brassica rapa) germplasm by the analysis of intact and desulfo
612	glucosinolates. Journal of agricultural and food chemistry. 2013 Apr 10;61(16):3984-93.
613	
614	
615	
616	
617	
	28

618	
619	
620	Figure legends
621	
622	Fig 1: A broad outline of common extraction methods used for glucosinolate analysis. Highlighted in grey is the
623	ISO 9167-1 method which was originally intended for glucosinolate extraction from <i>B. napus</i> seed but is
624	commonly used for glucosinolate extraction and analysis in all glucosinolate containing plant tissues.
625	
626	Fig 2: (a) total glucosinolate concentration of <i>B. napus</i> leaf halves dried in freeze drier B are significantly lower
627	(paired t-test, p=0.009) than leaf halves dried in freeze drier A; (b) <i>B. napus</i> leaf tissue dried with freeze drier B
628	is deformed and darker Error bars represent standard error.
629	
630	Fig 3: There is no difference in final glucosinolate concentrations between freeze drying or direct extraction in -
631	20°C methanol. B. napus leaves were cut in half and frozen. One half was freeze dried prior to glucosinolate
632	extraction, the other half was extracted directly into -20°C methanol (n=12; paired t-test, p=0.15; R squared =
633	0.96). The dashed line represents equivalence of x and y.
634	
635	Fig 4: Concentrations of representative aliphatic (sinigrin) and aromatic (glucotropaeolin) glucosinolates were
636	not reduced over the course of an hour at 100°C. The representative indole glucosinolate (glucobrassicin) is
637	degraded at 100°C. Asterisks represent significant difference from concentration at t=0 (paired t-test, p<0.05).
638	
639	Fig 5: Spectrophotometric analysis of sinigrin hydrolysis kinetics in water and 80% methanol (n=3) by purified
640	myrosinase (0.05mg/ml) at room temperature (25°C).
	29

Fig 6: Extraction of glucosinolates ($\geq 1 \mu mol/g$) in plant tissues across the three extraction methods. Glucosinolate concentrations from the cold methanol and boiling water extraction methods are normalised to the glucosinolate concentrations obtained from the ISO9167-1 (75°C methanol) method (n=4-12). Error bars represent standard error. Asterisks represent a significant effect of extraction method on glucosinolate concentration (repeat measure ANOVA, p<0.05). Fig 7: The cold extraction method yields less glucoraphasatin in R. sativus stems relative to the ISO 9167-1 (boiling methanol) extraction method (n=4). Values normalised to the ISO method results. Error bars represent standard error. Asterisks represent significant difference from the ISO 9167-1 (boiling methanol) method (paired t test, p<0.05). Fig 8: Absorbance values for representative desulfoglucosinolate extracts from B.juncea, S. alba, R. sativus and E. sativa extracts treated with sulfatase solution for 12, 24 or 48 hours. These values are reflective of desulfoglucosinolate recovery and not the initial glucosinolate concentration. Fig 9: Desulfoglucosinolate content extracted from B. juncea, E. sativa and R. sativus tissue incubated with 75µl low concentration sulfatase (0.3 U/ml) over 12H, 24H or 48H, and with a high concentration sulfatase (5U/ml) over 24H, normalised to glucosinolate content of the same samples prior to sulfatase treatment. E. sativa leaf samples were treated with TCEP post desulfation to undimerise didesulfoglucosatavin. Asterisks indicate a significant effect of purification method on glucosinolate concentration (repeat measure ANOVA, p<0.05). 'd' indicates the purification method yields a significantly different glucosinolate concentration relative to the intact glucosinolates (paired t-test, p<0.05).







Click here to download Figure Figure 2.pptx 👱

















Example chromatograms

Click here to access/download Supplementary Material Example Chromatograms.docx