



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/114157/>

Version: Accepted Version

Article:

Shanklin, V.E., Snowden, J.A. and Greenfield, D.M. (2018) Late treatment effects following bone marrow transplant: Efficacy of implementing international guidelines. *European Journal of Cancer Care*, 27 (2). e12623. ISSN: 0961-5423

<https://doi.org/10.1111/ecc.12623>

This is the peer reviewed version of the following article: Shanklin, V. E. et al (2016), Late treatment effects following bone marrow transplant: Efficacy of implementing international guidelines, *European Journal of Cancer Care*, which has been published in final form at <https://doi.org/10.1111/ecc.12623>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

**Late treatment effects following bone marrow transplant:
efficacy of implementing international guidelines**

Journal:	<i>European Journal of Cancer Care</i>
Manuscript ID	Draft
Manuscript Type:	Original Article
Keywords:	Haematopoietic cell transplantation, Allogeneic, Autologous, Late effects, Screening, Prevention

SCHOLARONE™
Manuscripts

Review

1
2
3
4 **Title: Late treatment effects following bone marrow transplant: efficacy of implementing**
5 **international guidelines**
6

7
8 **Concise title: Efficacy of late effects screening**
9

10 **Abstract**
11

12
13 An increasing cohort of haematopoietic stem cell transplantation (HSCT) survivors has raised
14 awareness of long-term and late effects. Updated recommendations for HSCT late effects screening
15 were published in 2012 (Majhail et al, 2012). We aimed to assess the clinical efficacy of a dedicated
16 screening clinic to identify problems in HSCT survivors using the international guidelines. Clinic
17 letters and test results of the first 59 consecutive patients attending the screening clinic were
18 evaluated. 30 females and 29 males (mean age of 49 years, range 22-74) were included. The mean
19 time since transplant was 6 years (0.5-18). 49/65 transplants were allogeneic. Primary indications for
20 HSCT were myeloid (56%), lymphoid (37%), solid tumour (5%) and auto-immune diseases (2%). 134
21 complications were reported (mean 2, range 0-8), with 114 documented further actions/referrals.
22 The most commonly reported concerns were pain 18/59(31%), fatigue 14/59(24%), sexual function
23 14/59(24%) and sleep disturbance 11/59(19%). Second primary malignancies were recorded in 5
24 cases. Implementation and audit of the international late effect screening guidelines confirm the
25 need for systematic long-term physical and psychological screening and care, thus ensuring timely
26 and efficient identification of problems and the opportunity to minimise morbidity effects and
27 optimise health.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 **Keywords:** Haematopoietic cell transplantation, Allogeneic, Autologous, Late effects, Screening,
50 Prevention
51
52
53
54
55
56
57
58
59
60

Introduction

Autologous or allogeneic HSCT survivors have higher risks of developing medical complications compared with the general population or other cancer survivors. Life expectancy of HSCT patients is lower than that of an age, gender and nationality-matched general population. This lag in life expectancy decreases but continues to exist until at least 15-20 years after HSCT (Majhail & Rizzo, 2013). Disease relapse, however, is the dominant cause of mortality within two years post HSCT (Majhail & Rizzo, 2013; Wingard et al, 2011). Significant early treatment complications include the risk of severe infection and graft versus host disease (Syrjala et al, 2012). Non relapse and infection complications include secondary cancer, endocrine, pulmonary and sexual dysfunction, bone degeneration and cardiovascular complications (Tichelli et al, 2013). Other complications including xerostomia and dry eyes, may not directly impact mortality but could significantly reduce quality of life. Indeed any system can be affected with severity ranging from life-threatening to mild and reversible (Trotti et al, 2003).

Awareness of long term effects of HSCT and implication of long term follow-up ensures identification and treatment of complications thereby optimising survival rates (Sun et al, 2013). However, mortality rates are still increased even after 10-30 years post transplantation being 4-9 fold higher than reported in an age-matched general population (Tichelli et al, 2013). A systematic approach to screening guidance and recommendations, alongside specific and common late effects of HSCT, was published in 2012 (Majhail et al, 2012) and adapted for clinical use in our transplant centre with systematic screening coordinated through a nurse-led late effects clinic. Thus the aims of our clinical audit were:

1. To evaluate the clinical efficacy of the screening clinic to identify late effects following HSCT
2. To evaluate the number and nature of subsequent actions or referrals.

Subjects and Methods

Our centre implemented the internationally agreed guidelines in 2013 using a locally developed Standardised Operating Policy (SOP) through a nurse-led late effects screening clinic. 59 consecutive patients who attended the screening clinic for the first time during 1st Sept 2013 to 31st Aug 2014 were included in the audit. Suitable patients were referred into the clinic by the transplant physicians in the same institution. Patients were in remission, with no active GvHD and without infection. The late effects screening clinic was held in an out patients facility and run in parallel to the consultant led transplant clinics. Audit approval confirmation was given by the Clinical Effectiveness Unit of our hospital trust.

Data was extracted from the clinic letter annotated from the consultation and from results of the requested investigations as per the SOP. Presence or absence of each variable and investigation was coded and current results and actions were recorded in a spreadsheet. The accuracy of requesting investigations as per the departmental SOP was reviewed to check whether these results were deranged or within normal limits.

Results

Demographics

Thirty patients were female and twenty nine male. Fifty four patients received one transplant, 4 had two transplants and 1 had three transplants. 49/65(75%) of transplants were allogeneic. Primary indications for HSCT for audit participants are given in Figure 1. Mean age of patients at assessment was 49 years (range 22-74), mean years from initial diagnosis was 8 (range 1-28) and mean years since transplant was 6 (0.5-18).

1
2
3
4 Forty four of 59 (75%) had chemotherapy treatment prior to transplant with 12 of these having
5 further high dose treatments. The most widely used chemotherapy regimen was FLAG-Ida (n=8).
6
7
8 Thirteen patients had TBI radiotherapy as part of their conditioning. Fifty one patients were given
9 pre-transplant conditioning regimens with 8 patients not recorded as having conditioning. Thirteen
10 of 59 received biological therapy, most receiving interferon (n=5) or rituximab (n=5). The most
11 commonly identified health problems are given in Table 1.
12
13
14
15

16 ***Health problems by system***

17 **Skin**

18
19
20
21
22
23 Nine patients reported skin concerns including dryness or rash. Seven had documented referrals for
24 further review by the consultant in charge of the screening clinic or to the GP for monitoring of skin
25 lesions.
26
27
28

29 **Head, eyes, ears nose and throat**

30
31
32
33 Ten patients suffered ocular complications mainly with dry eyes or 'gritty eyes'. Ocular action was
34 documented for 16 people i.e. were recommended an optician appointment for routine eye testing
35 or referred to an ophthalmologist for specialist review.
36
37
38

39
40 Thirteen patients reported oral complications mainly of xerostomia (dry mouth). Action or referral
41 was documented in 11, 1 being advised a routine dental appointment, 2 needing no further action
42 and 1 who had no action documented.
43
44
45

46 **Respiratory**

47
48
49
50 Seven patients reported respiratory problems, principally chest infections and coughs, and five
51 further action documented. Fifty seven had recorded pulmonary function tests (PFTs) requested as
52 part of routine post HSCT screening. Reported results indicated 21 had low TLCO, 3 low FEV1 and 4
53 with both low FEV1 and TLCO.
54
55
56
57
58
59
60

Cardiovascular

Thirteen cases of elevated blood pressure were recorded with 8 cases identified in clinic and 5 pre-existing. Three patients were found to have cardiac complications and all were referred to a cardiologist for specialist review. Fifty six had recorded echocardiogram reports, of which 49 were reported to have good systolic function with 22 reports of mild problems such as aortic regurgitation.

Thirty out of 47 patients who had lipids evaluated had deranged results, with 1 report of further action. Of these, 23 had elevated triglycerides but no recorded further actions.

GI

Two patients reported upper GI complications such as suspected silent reflux and excess gastric acid. Action was recorded for both patients. Six patients suffered lower GI complications, predominantly diarrhoea and urgent defecation- of these 5 had documented action. Forty one had liver function tests performed, 11 results were deranged, but no further action documented.

Genital

Fourteen patients reported problems with sexual function with 13 being referred for further investigation/specialist advice. Reports of 4 patients with early menopause as a result of HSCT were documented, 3 had documented action or referral. One man identified with testosterone deficiency, was referred to an endocrinologist.

Neurological

Six patients prioritised memory loss as a major concern, however no specific actions were documented. Fourteen patients reported persistent fatigue. Seven had documentation of further action such as referral to a specialist. All patients would normally have repeat bloods to exclude organic causes. Low energy, tiredness and sleep problems were the main concern of patients when

1
2
3
4 asked to prioritise concerns; 20 experienced difficulties with sleepiness and fluctuating energy levels,
5
6 11 reported problems sleeping at night of these- 7/11 had documentation of action.
7

8
9 Eighteen patients reported pain. The plan of action for pain management was documented although
10
11 2 patients had no further action documented.
12

13 **Musculo-skeletal**

14
15
16
17 DEXA scans results were available in 17/59 patients. Four patients were found to have low bone
18
19 mineral density with further action being to evaluate specific problems and consider
20
21 supplementation of Vitamin D. For others scanned, recommendations were given for time interval
22
23 before repeat DEXA and lifestyle advice e.g. reducing alcohol consumption. Twelve patients had
24
25 Vitamin D levels tested with 4 being below normal. 2 had actions reported of instigation of Vitamin D
26
27 supplementation.
28

29 **Haematological**

30
31
32
33 Eighteen out of twenty four patients were found to have high ferritin levels. Two had reported
34
35 evidence of further action (venesections) being taken. Twenty had Vitamin B12 levels tested, 2 of
36
37 these were deranged but no action was documented.
38

39 **Endocrine**

40
41
42
43 TSH was measured in 48 patients, resulting in 12 deranged results and 1 documented referral/action.
44

45 **Immunology**

46
47
48
49 Thirty letters mentioned vaccination programmes with evidence of 23/30 having a reported plan
50
51 such as recommending an annual flu jab or requests for GP reinstatement of vaccine programmes
52
53 which may have been paused to manage GVHD. Five patients experienced complications from a
54
55 compromised immune system- predominantly GVHD, recurrent chest infections and multiple
56
57
58
59
60

1
2
3
4 infections varying from sinusitis to UTI. All actions were documented- often the problem was already
5
6 managed by other specialists but those with recurrent infections required further investigations.
7
8

9 **Psychological**

10
11 Six patients were referred to a psychologist with an additional one patient who was recommended to
12
13 see a psychologist and 1 referred to a psychiatrist.
14
15

16 **Malignancies**

17
18 Six malignancies were recorded post transplantation of which 5 were recent onset primary skin
19
20 lesions and the other new malignancy was liver metastases from previous colon cancer.
21
22

23
24 Figure 2 gives frequency of complications and frequency of documented action.
25
26
27
28
29

30 **Discussion**

31
32 To our knowledge this is the first time the new international late effect screening guidelines have
33
34 been formally audited (Majhail et al, 2012). One hundred and thirty four new problems or concerns
35
36 were identified in 59 consecutive transplant patients with 114 documented further action or
37
38 referrals. The most frequently concerns were pain, fatigue, sexual function and sleep patterns.
39
40 Second malignancy was reported in 5 cases.
41
42
43

44 Pain is a well-recognised unmet need of cancer patients (Vogelzang et al, 1997) and also evident in
45
46 HSCT patients (Niscola et al, 2008). Pain is often associated with fatigue with 30-42% of HSCT
47
48 survivors troubled by fatigue in contrast to <20% prevalence of pain (Mosher et al, 2011).
49
50 Interestingly, Vogelzang found that whilst oncologists believed that pain affected their patients to a
51
52 higher degree than fatigue, patients believed that fatigue affected their lives more daily than pain
53
54 (61% v 19%) (Vogelzang et al, 1997). The reported sources of pain are mainly due to the mucosal
55
56
57
58
59
60

1
2
3
4 tissue injury induced by the conditioning regimen but also include pain from infection, from acute or
5
6 chronic GVHD as well as adverse drug reactions to GVHD drug prophylaxis. Furthermore, some late
7
8 complications of the actual transplant may be associated with pain (Niscola et al, 2008). The
9
10 management strategy of pain that is recommended is using the analgesic ladder- NSAIDs or
11
12 paracetamol for mild pain, partial mixed opiate receptor agonists and mild opioids e.g. oxycodone for
13
14 moderate pain relief and finally, the pure μ -opiate receptor agonists or strong opioids, e.g. morphine
15
16 and fentanyl, are chosen for severe pain relief. With one third of our cohort reporting pain as an
17
18 issue, there is a clear need for regular assessment and monitoring with implications for palliative and
19
20 supportive care in addition to further investigation for cause.
21
22

23
24 Further detail as to whether physiological causes of fatigue such as anaemia or vitamin D deficiency
25
26 were excluded or that it was confirmed as idiopathic with no identifiable cause. It is important to
27
28 routinely screen for anaemia in this population of HSCT survivors.
29
30

31 Sexual function was also a highly prevalent concern in our cohort with 14 (7 male, 7 female)
32
33 reporting problems- such as early menopause and testosterone deficiency. Psychosexual functioning
34
35 has been found to be worse in HSCT patients compared with patients treated with chemotherapy
36
37 alone and significantly worse compared with healthy subjects (Thygesen et al, 2012). The effect is
38
39 long-lasting with survivors at increased risk of sexual function problems even 5-10 years after
40
41 transplantation (Frey Tirri et al, 2015). The late effects guidelines (Majhail et al, 2012) suggest clinical
42
43 and endocrinologic gonadal assessment at 1 year post HSCT in all women post-pubertal at
44
45 transplantation with the frequency of subsequent assessments guided by clinical need. Women
46
47 should have annual gynaecologic evaluation as part of general health screening to rule out for
48
49 example atrophic vaginitis, a common problem after HSCT at which time HRT may be considered in
50
51 those who are postmenopausal (Frey Tirri et al, 2015). In men, gonadal function (particularly FSH, LH
52
53 and testosterone) should be assessed if symptomatic eg lack of libido or erectile dysfunction. It is
54
55
56
57
58
59
60

1
2
3
4 also important to rule out GvHD of the genital tract (Frey Tirri et al, 2015). Once again systematic
5
6 enquiry is indicated, even for intimate topics to allow patients permission to open a conversation and
7
8 be reassured, itself considered therapeutic (Yi & Syrjala, 2009).
9

10
11 HSCT survivors, particularly allogeneic patients, are at risk of developing secondary malignancies of
12
13 the skin. In our cohort, 5 out of 6 of the malignancies reported post transplantation, were skin
14
15 lesions. In terms of risk and prevention of secondary cancers, clinic letters gave detailed advice of the
16
17 risks of UV exposure, smoking, alcohol and the importance of a healthy lifestyle. This identified that
18
19 the post HSCT review is also a useful mechanism for health promotion. The late effects guidelines do
20
21 have a section titled 'healthy lifestyle recommendations for all patients' which covers this.
22
23

24
25 We have demonstrated here that any system can be affected thus a complete systems review is
26
27 indicated. Screening and taking a history of patient centred concerns such as pain, fatigue, sexual
28
29 function and sleep are as important as ruling out physiological disorders to maximised health and
30
31 recovery after HSCT.
32
33

34
35 This evaluation was limited by the decision to use conveniently available source data including the
36
37 clinic letter, most recent blood test results and last ECHO, DEXA and PFT in the audit. This included
38
39 DEXA scan reports which were only available if done within the previous 18 months gives scan
40
41 reports within the last 18 months. Furthermore, data regarding initial diagnoses and treatment was
42
43 inconsistently documented. Additional searching was required to retrieve more accurate
44
45 information.
46
47
48
49

50 **Conclusion**

51
52
53 In summary, this study described the population of 59 new patients to the late effects screening
54
55 clinic with a mean 6 years since transplant. We demonstrated a high prevalence of systematic
56
57
58
59
60

1
2
3
4 complications and evidence for further actions to assess these complications. This new nurse-led
5
6 screening clinics demonstrates clinical efficacy in identifying late complications using holistic
7
8 screening and confirms the need for long-term follow-up post-transplantation. Ultimately, screening
9
10 clinics such as this aim to pick up complications earlier to prevent/reduce morbidity therefore playing
11
12 a central role in optimising quality of life for HSCT survivors. Future evaluation should more
13
14 accurately assess the severity of each concern using a scoring system such as the Common Toxicity
15
16 Criteria (CTC) system (Trotti et al, 2003).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conflict of Interest

There are no conflicts of interest.

Acknowledgements

We are grateful to both the Bone Marrow Transplant Team and Clinical Effectiveness and Audit department for their help and support with this project.

For Peer Review

References

Frey Tirri B, Häusermann P, Bertz H, Greinix H, Lawitschka A, Schwarze C-P *et al.* (2015) Clinical guidelines for gynecologic care after hematopoietic SCT. Report from the international consensus project on clinical practice in chronic GVHD. *Bone Marrow Transplant*, 50(1), 3–9.

Majhail NS, Rizzo JD, Lee SJ, Aljurf M, Atsuta Y, Bonfim C *et al.* (2012) Recommended screening and preventive practices for long-term survivors after hematopoietic cell transplantation. *Biology of Blood and Marrow Transplantation*, 18(3), 348–71.

Majhail NS, Rizzo JD. (2013) Surviving the cure: long term followup of hematopoietic cell transplant recipients. *Bone Marrow Transplant*, 48(9), 1145–51.

Mosher CE, DuHamel KN, Rini C, Corner G, Lam J, Redd WH. (2011) Quality of life concerns and depression among hematopoietic stem cell transplant survivors. *Support Care Cancer*, 19(9), 1357–65.

Nicola P, Romani C, Scaramucci L, Dentamaro T, Cupelli L, Tendas A *et al.* (2008) Pain syndromes in the setting of haematopoietic stem cell transplantation for haematological malignancies. *Bone Marrow Transplant*, 41(9), 757–64.

Sun CL, Kersey JH, Francisco L, Armenian SH, Baker KS, Weisdorf DJ *et al.* (2013) Burden of morbidity in 10+ year survivors of hematopoietic cell transplantation: report from the bone marrow transplantation survivor study. *Biology of Blood and Marrow Transplantation*, 19(7), 1073–80.

1
2
3
4 Syrjala KL, Martin PJ, Lee SJ. (2012) Delivering care to long-term adult survivors of hematopoietic cell
5
6 transplantation. *Journal of Clinical Oncology*, 30, 3746-51.
7

8
9
10 Thygesen KH, Schjødt I, Jarden M. (2012) The impact of hematopoietic stem cell transplantation on
11
12 sexuality: a systematic review of the literature. *Bone Marrow Transplant*, 47(5), 716–24.
13

14
15
16 Tichelli A, Rovó A. (2013) A clever person solves a problem; a wise person avoids it. *Biology of Blood
17
18 and Marrow Transplantation*, 19(7), 1004-5.
19

20
21
22 Trotti A, Colevas AD, Setser A, Rusch V, Jaques D, Budach V *et al.* (2003) CTCAE v3.0: development of
23
24 a comprehensive grading system for the adverse effects of cancer treatment. *Seminars in Radiation
25
26 Oncology*, 13(3), 176–81.
27
28

29
30
31 Wingard JR, Majhail NS, Brazauskas R, Wang Z, Sobocinski KA, Jacobsohn D *et al.* (2011) Long-Term
32
33 Survival and Late Deaths after Allogeneic Bone Marrow Transplantation. *Journal of Clinical Oncology*,
34
35 29(16), 2230-2239.
36
37

38
39
40 Vogelzang NJ, Breitbart W, Cella D, Curt GA, Groopman JE, Horning SJ *et al.* (1997) Patient, caregiver,
41
42 and oncologist perceptions of cancer-related fatigue: results of a tripart assessment survey. The
43
44 Fatigue Coalition. *Seminars in Hematology*, 34(3), 4–12.
45
46

47
48
49 Yi JC, Syrjala KL. Sexuality After Hematopoietic Stem Cell Transplantation. (2009) *The Cancer Journal*,
50
51 15(1), 57–64.
52
53
54
55
56
57
58
59
60

1
2
3
4 **Figure legends**
5

6
7 Figure 1:

8
9 Primary indications for HSCT for audit participants
10


11
12 Figure 2:

13
14 Frequency of identified problems/concerns and documented actions/referrals
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1



Peer Review

Figure 2



Peer Review