

This is a repository copy of *Long-term care provision, hospital bed blocking, and discharge destination for hip fracture and stroke patients*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/114270/>

Version: Accepted Version

---

**Article:**

Gaughan, James Michael [orcid.org/0000-0002-8409-140X](https://orcid.org/0000-0002-8409-140X), Gravelle, Hugh Stanley Emrys [orcid.org/0000-0002-7753-4233](https://orcid.org/0000-0002-7753-4233), Santos, Rita [orcid.org/0000-0001-7953-1960](https://orcid.org/0000-0001-7953-1960) et al. (1 more author) (2017) Long-term care provision, hospital bed blocking, and discharge destination for hip fracture and stroke patients. *International Journal of Health Economics and Management*. ISSN 2199-9031

<https://doi.org/10.1007/s10754-017-9214-z>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

**International Journal of Health Economics and Management**  
**Long-term care provision, hospital bed blocking, and discharge destination for hip fracture and stroke patients**  
 --Manuscript Draft--

<b>Manuscript Number:</b>	IHFE-D-13-00133R2
<b>Full Title:</b>	Long-term care provision, hospital bed blocking, and discharge destination for hip fracture and stroke patients
<b>Article Type:</b>	Manuscript
<b>Keywords:</b>	hospitals; Length of stay; long-term care; care homes; bed blocking; substitution
<b>Corresponding Author:</b>	James Gaughan Centre for Health Economics, University of York York, UNITED KINGDOM
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	Centre for Health Economics, University of York
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	James Gaughan
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	James Gaughan
	Hugh Gravelle
	Rita Santos
	Luigi Siciliani
<b>Order of Authors Secondary Information:</b>	

## Response to Reviewer

*We thank the reviewer for their comments. We set out how we have responded to each of these below.*

### Reviewer #2: Minor Comments

Page 10, line 42: There is a missing reference.

*We apologise for this incorrect formatting, which has now been corrected. The reference is to equation (1).*

Tables: The tables are hard to read. Ideally they should be self-explanatory. Currently, we have a long list of variables where it is difficult to know which one are the important one. I recommend to group the variables with appropriate headings following the classification of Sections 2.3, 2.4, and 2.5.

*We thank the reviewer for this suggestion. We have added titles for groups of variables in Tables as suggested. We have also highlighted in bold the key variables for the analysis and reordered tables so that all variables discussed in each subsection 2.3, 2.4 and 2.5 are grouped together under the same subheading.*

# Long-term care provision, hospital bed blocking, and discharge destination for hip fracture and stroke patients

**Authors:** James Gaughan<sup>1</sup>, Hugh Gravelle<sup>1</sup>, Rita Santos<sup>1</sup>, Luigi Siciliani<sup>1,2</sup>

**Institutions:** <sup>1</sup>Centre for Health Economics, University of York, York, UK

<sup>2</sup>Department of Economics and Related Studies, University of York, York, UK

**Correspondence author:** James Gaughan, Centre for Health Economics, University of York, York; email: [james.gaughan@york.ac.uk](mailto:james.gaughan@york.ac.uk) tel: 01904 321423 fax: 01904 321402

**Conflict of interest:** None

**Acknowledgements** The research reported in this paper was funded by a grant from the Department of Health to the Policy Research Unit in the Economics of Health and Social Care Systems. The views expressed are those of the authors and not necessarily those of the funders. We are grateful for comments and suggestions from Henning Øien and participants in the 3rd Norwegian Workshop in Health Economics (Geilo, Norway, 20-21 March 2013), and in the Second Workshop on the Economics of Long Term Care held at Toulouse School of Economics (Toulouse, France, 22 November 2012).

## Abstract

We examine the relationship between long-term care supply (care-home beds and prices) and (i) the probability of being discharged to a care home and (ii) length of stay in hospital for patients admitted to hospital for hip fracture or stroke. Using patient level data from all English hospitals and allowing for a rich set of demographic and clinical factors, we find no association between discharge destination and long term care beds supply or prices. We do, however, find evidence of bed blocking: hospital length of stay for hip fracture patients discharged to a care home is shorter in areas with more long-term care beds and lower prices. Length of stay is over 30% shorter in areas in the highest quintile of care home beds supply compared to those in the lowest quintile.

Words: 132

**Keywords:** hospitals; length of stay; long-term care; care homes; bed blocking; substitution

JEL: I11

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Long-term care provision, hospital bed blocking, and discharge  
destination for hip fracture and stroke patients**

## Abstract

We examine the relationship between long-term care supply (care-home beds and prices) and (i) the probability of being discharged to a care home and (ii) length of stay in hospital for patients admitted to hospital for hip fracture or stroke. Using patient level data from all English hospitals and allowing for a rich set of demographic and clinical factors, we find no association between discharge destination and long term care beds supply or prices. We do, however, find evidence of bed blocking: hospital length of stay for hip fracture patients discharged to a care home is shorter in areas with more long-term care beds and lower prices. Length of stay is over 30% shorter in areas in the highest quintile of care home beds supply compared to those in the lowest quintile.

Words: 132

# 1. Introduction

The provision of health care and long term care for the elderly is a consistent focus of policy makers in the U.K. and other OECD countries (Department of Health 2001; Department of Health 2011; Glendinning 2003; OECD 2011; Wanless 2006). Around 10% of individuals over 75 years old used both health and long-term care in 2006/7 in England (Bardsley et al 2012). Long term care has costs and outcome consequences on health care and vice versa (Fernandez and Forder 2008; Forder 2009; Vetter 2003). In England, acute hospital care and long term care are organised and funded separately and differently.

There is long standing concern over coordination for patients requiring health and long-term care, in particular the delayed discharge of patients from hospital (Baumann et al 2007; Bryan et al 2006; House of Commons 2003; National Audit Office 2000). To improve integration policy makers need information about the effects of provision of one type of care on the other. In this paper we examine two questions where there is little quantitative evidence: the extent to which accessibility of long-term care affects the length of stay in hospital and the probability of a patient being discharged back to their homes rather than to a care home.

We focus on patients who suffer a hip fracture or stroke. The conditions are selected for their high incidence, impairing effects on patients, and the consequent need for long-term care for the elderly (Kasteridis et al 2015; de Meijer et al 2011). The British Orthopaedics association reports there were around 70,000 hip fractures in the U.K. in 2007, including many older patients with complex clinical needs. The incidence is expected to rise (British Orthopaedic Association 2007). Further, 10-20% of hip fracture patients admitted to hospital from their own home ultimately utilise some institutional care (National Clinical Guidelines Centre 2011). The national stroke strategy published by the Department of Health in 2007 reports



1 that there are approximately 110,000 strokes each year in England and that 75% of these  
2 occur among people aged 65 and over. Stroke is the single largest cause of disability, with a  
3 third of people suffering from a stroke having a long-term disability (Department of Health  
4 2007; National Audit Office 2005). Estimates of the proportion of care home residents who  
5 have had a stroke vary between 25% and 45% (CQC 2011). Hip fracture and stroke patients  
6 are thus a policy concern (Department of Health 2001) and have been the focus of past  
7 research in England (Bond et al 2000). Both require immediate hospital care and longer term  
8 rehabilitation. Such rehabilitation could take place in hospital but also at the patient's home  
9 or in a long-term care facility.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 Higher care home bed supply, at given prices, implies a shorter waiting time for a bed to  
25 become available and may thus increase the probability of discharge to a care home. It may  
26 also imply a shorter hospital length of stay since patients will have shorter waits for a place in  
27 a care home. Higher prices will reduce probability of the patient opting for a care home as  
28 opposed to returning to their own home and may induce patients to search for longer, thereby  
29 increasing hospital length of stay.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 We examine two questions. First, we investigate whether access to long-term care in nursing  
42 and residential homes (as measured by beds and prices) influences the probability that  
43 patients who are admitted to hospital from their home are discharged to a care home. Second,  
44 we investigate the bed blocking hypothesis that the supply of long-term care influences  
45 length of stay in hospital.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1.1. Institutional Background

Emergency hospital care in England is predominantly provided by the National Health Service (NHS) through 166 acute public hospitals, known as Trusts. NHS care is funded by general taxation, with 152 local health authorities (Primary Care Trusts, PCTs) receiving capitated budgets from the Department of Health from which they pay for hospital care provided to their populations. Patients do not pay for hospital care provided by NHS hospitals.

Long-term care is provided by over 18,000 care homes<sup>1</sup> (Laing and Buisson 2010). In 2014 about 74% of care home beds were provided by for-profit firms, 17% by voluntary organisations and 8% by Local Authorities (LAs) (Jarrett 2016).<sup>2</sup> Providers are generally small and the supply and price of care home beds is largely driven by local demand, the cost of provision, and by competition (Forder and Fernandez 2012; Allan and Forder 2015).

Around 60% of care home users (Forder 2007) are self-funders and pay at least in part for their long-term care. Those with low wealth are subsidised by their LA. The price charged for subsidised patients is lower than for self-funders because of the bargaining power of the LAs (OFT 2005). This gives a financial incentive to care homes to give priority to self-funders, within the constraints of any locally negotiated contract with LA commissioners. Despite potentially considerable out-of-pocket expenses associated with long-term care, the

---

<sup>1</sup> “Care home” is the usual term in England for a residential institution providing long-term personal care such as help with bathing and dressing. Some care homes also provide nursing care and are referred to as nursing homes (Age UK 2015). A care home which does not provide nursing care has, in the past, been sometimes referred to as a residential home. Other countries use a different terminology. For example, in the U.S. care homes might be referred to as Assisted Living, Personal Care Home or Residential Care facilities with precise definitions determined at the state level. Skilled Nursing Facilities or Nursing Homes perform the same general function as nursing homes in the U.K. (Pioneer Network 2011). As this study uses English data, we use the English terminology definitions and so care homes include nursing and residential homes.

<sup>2</sup> The 152 Local Authorities (LAs) with responsibility for social care are elected political bodies funded by local taxes and central government grants. The 166 local acute health trusts are part of the NHS and are appointed by the Department of Health.

1 proportion of individuals covered by private long term care insurance in England is very  
2 small, with just under 22,000 people holding such insurance in 2008 (Comas-Herrera et al  
3  
4  
5 2009).  
6

7  
8  
9 The majority of residents in care homes are long stay. Care homes also provide some  
10 rehabilitation services. Steventon and Roberts (2012) found that 39% of LA funded  
11  
12 admissions to care homes in three English LAs in 2005/06 were short stay but that the median  
13  
14 length of stay was 18 months.  
15  
16  
17  
18  
19  
20  
21  
22

### 23 1.2. *Previous Literature*

24

25 Previous studies investigating discharge destination of hospital patients find that age, gender  
26  
27 and living arrangements are key drivers of discharge to a care home as opposed to their own  
28  
29 home. Other factors include comorbidities (Aharonoff et al 2004; Gilbert et al 2010),  
30  
31 ethnicity (Aharonoff et al 2004; Ellis and Trent 2001), urbanisation (Gilbert et al 2010) and  
32  
33 income deprivation (Gilbert et al 2010; Picone et al 2003). Patients discharged to a care home  
34  
35 are also likely to have a longer stay in hospital (Castelli et al 2015; Wong et al 2010).  
36  
37  
38  
39  
40  
41

42 Picone et al (2003) investigates the determinants of hospital length of stay and discharge  
43  
44 destination of 4608 U.S. Medicare patients following hip fracture, stroke or heart attack.  
45  
46 They show that potential supply of informal care (being married and number of children)  
47  
48 increases the likelihood of being discharged home. They also find that supply of institutional  
49  
50 care (concentration of skilled nursing home or rehabilitation hospital beds) is associated with  
51  
52 shorter length of stay. The only English study of discharge destination is Bond et al (2000)  
53  
54 who examine the discharge destination of 440 stroke and 572 hip fracture patients in six  
55  
56 English NHS hospitals. They find that the probability of being discharged to a care home  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 increases with the supply of care home beds. In line with these studies, we find a negative  
2 association between care home beds supply and length of stay in hospitals, and a positive but  
3 insignificant association between care home beds supply and the probability of being  
4 discharged to a care home.  
5  
6  
7  
8  
9

10  
11 Although there is an extensive literature on the substitution between informal and formal  
12 long-term care (Bolin et al 2008; Bonsang 2009; Grabowski et al 2012; Van Houtven and  
13 Norton 2004), there is limited evidence on the effect of care homes supply on health care  
14 utilisation, i.e. on the substitution between long-term social care and hospital care. Fernandez  
15 and Forder (2008) find that LAs with more home help hours and nursing and residential care  
16 beds have a lower rate of hospital delayed discharges and lower emergency readmission rates.  
17 Forder (2009) uses small area data on 8,000 census areas in England and finds that a £1  
18 increase in spending on care homes reduces hospital expenditure by £0.35. Gaughan et al  
19 (2015) use LA level data on all types of hospital patients and find that greater supply of care  
20 home beds in the LA is associated with a reduction in delayed discharge from hospital.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Holmas et al (2010) investigate the effect of fining owners of long-term care institutions  
40 which prolong length of stay at hospitals in Norway. Surprisingly, hospital length of stay is  
41 longer when the fines are used, which they interpret as an example of monetary incentives  
42 crowding out intrinsic motivation. Holmas et al (2013) find that greater expenditure on long  
43 term care by Norwegian local authorities reduces both overall length of hospital stay and stay  
44 in hospital when medically ready for discharge.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

56 Our study contributes to the literature on the relationship between long-term care and health  
57 care by using a large and rich individual patient level data set for two disabling and high  
58  
59  
60  
61  
62  
63  
64  
65

1 incidence conditions so that we can control more precisely for patient diagnoses, socio-  
2 economic characteristics and hospital policies than previous English area level studies.  
3  
4  
5  
6

## 7 **2. Data and Methods**

### 8 *2.1. Sample*

9  
10 A detailed account of the data set construction is in the Data Appendix. We use cross-section  
11 administrative data from Hospital Episodes Statistics (HES). Our sample is all patients aged  
12 65 or over, resident in England, treated in NHS hospitals, admitted from home as an  
13 emergency with a primary diagnosis of hip fracture or stroke, and discharged in the financial  
14 year April 2008 to March 2009. We exclude patients who are admitted from a care home  
15 since they are very likely to return to the same care home. We analyse the two samples (hip  
16 fracture and stroke) separately. While coordination between acute and long-term care services  
17 is important for patients with either condition, the characteristics and needs of the two patient  
18 groups differ. As well as being treated in different hospital departments, which may have  
19 different policies about discharge, stroke is a potentially more severe condition and with a  
20 more varied prognosis.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 Patients who die in hospital, are discharged to a penal institution or to a secure psychiatric  
42 unit, have incomplete spells, or for whom final discharge destination is not known, are  
43 excluded from the analysis. We also exclude patients at Hospital Trusts with less than 10 hip  
44 fracture and stroke patients per year. The estimation samples are 21,959 hip fracture patients  
45 and 33,101 stroke patients.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2.2. *Dependent variables*

Our dependent variables are the patient's hospital length of stay and whether the patient is discharged to a long-term care institution as opposed to returning home following hospitalisation. Patient length of stay is the number of days between admission and discharge from hospital at the end of their spell, allowing for patients to be transferred between hospitals during their spell.

## 2.3. *Patient characteristics*

For each patient we have age, gender, number of diagnoses and procedures, day of discharge and whether the patient is transferred to a different provider during their hospital spell. HES diagnostic fields are used to construct three co-morbidity dummy variables based on the Charlson index (Charlson et al 1987). We distinguish between (i) no Charlson co-morbidities, (ii) a single non-severe co-morbidity, (iii) at least one severe or at least two non-severe co-morbidities. Since stroke is one of the Charlson co-morbidities we exclude it when constructing these variables for stroke patients.

We also distinguish between different types of hip fracture and stroke. The major types of hip fracture relate to the location of the fracture. The categories are femoral neck, peritrochanteric, and subtrochanteric, where the trochanters refer to protrusions of bone just below the ball of the hip. Prognosis and length of stay varies by type of fracture (Butler et al 2009; Clague et al 2002). The most critical distinction for stroke patients is between strokes due to an infarction, where a vessel supplying blood to the brain is blocked and those due to a haemorrhage, where there is bleeding in or around the brain. Although patient outcome is driven by stroke severity as well as by type of stroke (Jørgensen et al 1995) haemorrhagic strokes generally lead to worse outcomes in terms of disability and mortality.

#### 2.4. *Formal long-term care*

We have data on the number of registered beds and prices of different types of rooms for all providers in England whose main client group is patients aged 65 or over or with dementia (Laing and Buisson 2010). We include these two categories of provider since they match with the age restrictions of our sample and it is these groups of patients most likely to require care in a care home following hip fracture or stroke. Care homes specialising in dementia are likely to cater to patients in our group given that its prevalence increases rapidly with age. Moreover, having a stroke might cause dementia and conversely a patient suffering with dementia might have a hip fracture from a fall.

While stroke is a more severe condition, the Stroke Association suggests that a patient can be accommodated in a residential home depending on need (Stroke Association 2013). There are almost no specialist stroke care homes and at least 25% of patients in care homes have had a stroke (CQC 2011). There is a small amount of specialist rehabilitation in the care home sector but these providers do not cater specifically for care following stroke or hip fracture.

We compute the number of beds and average prices in residential and nursing homes within a 10km radius of the centroid of each patient's small area of residence (Lower Super Output Area (LSOA)). LSOAs are small geographic areas with an average population of 1,500 in 2001. There are 32,482 LSOAs in England served by 166 acute hospital trusts. We also have a measure of average care home quality based on ratings by the sector regulator (the Care Quality Commission (CQC)).

1 To allow for differences in population we use the mid-2008 population of retirement age (60  
2 years and over for women; 65 years and over for men) living in LSOAs whose centroids are  
3 within 10km of each patient's LSOA centroid.  
4  
5  
6

## 7 8 9 *2.5. Socioeconomic characteristics of patient areas of residence*

10 Information on socioeconomic characteristics is not available at the individual level in HES.  
11  
12 We attribute socioeconomic variables from the 2001 Census and the 2004 Index of Multiple  
13 Deprivation to patients by their LSOA of residence. The variables include the proportions of  
14 non-white residents, households with a single pensioner, and those reporting self-assessed  
15 health as "not good" from the three categories (good, fairly good and not good). The  
16 proportion of single pensioner households provides some adjustment for availability of  
17 informal care as much of this care is provided by a spouse or other relative living with the  
18 patient. We measure income deprivation as the proportion of the LSOA's population aged  
19 over 60 who are claiming income support, income related job seekers allowance, or pension  
20 credit guarantee. LSOAs are classified as urban, town or village. We include a dummy  
21 variable for patients resident in a London LSOA to allow for peculiarities of health and long-  
22 term care provision in the capital. Because of lack of information on acute care in Scotland  
23 and Wales, we include dummy variables indicating whether a patient's LSOA is within 10km  
24 of the English border with Scotland or Wales.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

## 48 *2.6. Methods*

49 We use models with the same explanatory for both patient discharge destination and patient  
50 length of stay except for using day of discharge in modelling length of stay. We estimate  
51 linear probability models for a patient being discharged to a care home as opposed to the  
52 patient's home, separately for hip fracture and stroke. The model is  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



$$y_{ij} = \beta_1 m_{ij} + \beta_2 x_{ij} + \beta_3 B_{ij} + \beta_4 P_{ij} + h_j + \varepsilon_{ij} \quad (1)$$

where  $y_{ij}$  is an indicator variable equal to one if the patients  $i$  in hospital  $j$  is discharged to a care home and zero if discharged to their own home.  $m_{ij}$  is a vector of patient morbidity variables and  $x_{ij}$  is a vector of patient demographic and socio-economic characteristics.  $B_{ij}$  is care home beds supply in the 10km area around the centroid of the patient's LSOA of residence, and  $P_{ij}$  is average care home prices in the same area. Notice that care home beds and prices vary across patients treated within the same hospital since each hospital will draw patients from many different LSOAs. We allow for non-linear effects by measuring price and beds supply as indicators for the quintiles of the national distribution of these variables across all LSOAs.  $h_j$  is a hospital effect. Except where otherwise stated, we estimate all models with cluster (hospital) robust standard errors.

For the investigation of patient length of stay we use the natural logarithm of length of stay as the dependent variable in (1). We estimate separate models for patients discharged to a care home and for those discharged to their home.

To choose between fixed or random hospital effects we estimate auxiliary regressions with the same explanatory variables as (1) but with hospital effects replaced by the hospital level means of the explanatories (Mundlak 1978; Wooldridge 2010, p. 332). For all models we find that a joint Wald test rejects the null hypothesis that the coefficients on the hospital level means are zero. We therefore use the fixed effects specifications. Thus the reported coefficients on beds or prices show whether, within a given hospital, patients residing in LSOAs with a higher long-term care supply have greater length of stay and or probability of being discharged to a care home.

1 Care home beds and prices are measured for each patient as the sum (beds) and means  
2 (prices) for care homes within 10km of the patient's LSOA of residence. We argue that these  
3  
4 are not correlated with unobservable factors affecting individual patient LoS and discharge  
5 destination. First, we control for hospital fixed effects and these will pick any variations in  
6  
7 hospital policies which might affect length of stay or discharge destination for all patients  
8  
9 treated in the hospital and which might be correlated with care home supply and prices if care  
10  
11 homes locate near hospitals which have a tendency to keep patients in longer or to discharge  
12  
13 them to care homes. We identify the effect of care home beds and prices from their variation  
14  
15 across patients treated within the same hospital but living in different LSOAs and thus facing  
16  
17 different beds supply and prices. Second, we have good data on individual patient morbidity  
18  
19 which might affect length of stay and discharge destination. We also include measures of  
20  
21 LSOA level demographics, socioeconomic characteristics, morbidity, retirement, and single  
22  
23 pensioner households which might affect area level demand for long term care and hence  
24  
25 supply and prices. Third, our dependent variables are for patients with two specific conditions  
26  
27 whilst care homes cater for a much wider set of patients so it is unlikely that care home beds  
28  
29 supply and prices will be driven by demand from stroke and hip fracture patients.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 Our models make the simplifying assumption that decisions on discharge destination and  
42  
43 length of stay are independent. As a robustness check, we model the decisions jointly with a  
44  
45 two-step selection model (Heckman 1979) with a probit model for discharge destination and  
46  
47 adding the inverse Mills ratio derived from it to the linear length of stay models in the second  
48  
49 stage. Using hospital fixed effects in the first stage probit model would yield biased  
50  
51 estimates. Instead, we replace each explanatory variable with its hospital level mean and with  
52  
53 patient level deviations from the mean. The hospital level means will pick up unobserved  
54  
55 hospital effects. As an exclusion condition, we include the rate of discharge to care at the  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 hospital level in the first stage but not in the second, arguing that the hospital rate of  
2 discharge to care will capture internal policies which affect patient probabilities of discharge  
3 to care but will not directly affect length of stay. We also run the model without this variable,  
4  
5 relying on non-linearity for identification.  
6  
7  
8  
9

### 10 11 **3. Results**

#### 12 13 *3.1. Summary statistics*

14  
15 Table 1 provides descriptive statistics. Similar proportions of hip (14.5%) and stroke (13.5%)  
16 patients are discharged to a care home. The average hospital length of stay is 22 days for hip  
17 fracture and 29 days for stroke. Length of stay is shorter for patients returning to their home  
18 (20 days for hip fracture and 23 for stroke) than for those discharged to a care home (33 days  
19 and 62 days). There are on average 2,300-2,400 care-home beds within 10km with an average  
20 price around £524 per week. The average care home quality rating is ‘good’ (a score of 3).  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

#### 32 33 34 *3.2. Discharge destination: hip fracture*

35  
36 Table 2 reports the results from linear probability discharge destination models. The  
37 probability of being discharged to a care home is greater for patients who are older, female,  
38 and have more diagnoses. Compared to patients aged 65-74 years old, patients who are 75-84  
39 and older than 84 years have 6.2 and 11.4 percentage points higher probabilities of being  
40 discharged to a care home. Men have a 1.6 percentage points smaller probability. An  
41 additional diagnosis or procedure increases the probability by 1.1 and 0.5 percentage points.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

52  
53  
54 There is some, though weak, association of the probability of discharge to a care home with  
55 long term care beds. Compared to patients in the lowest quintile of beds supply, the  
56 estimated probability of being discharged to care for patients in LSOAs in the second, third,  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

fourth and fifth beds quintiles are 0.9, 2.1, 2.5 and 4.1 percentage points greater. However, none of the coefficients achieves 5% statistical significance

### 3.3. *Discharge destination: stroke*

The effects of covariates on the discharge destination of stroke patients are similar to those for hip fracture patients. Patients who are 75-84 years old and older than 84 years have 5.4 and 12.5 percentage points higher probability of being discharged to a care home, compared to those aged 65-74. Men have a smaller probability (by 4.3 percentage points) of being discharged to a care home.

An additional diagnosis increases the probability by 1.8 percentage points. Transferred patients have 2.9 percentage points higher probability. Compared to patients with cerebral infarction, the probability of discharge to care is 1.4 percentage points higher if stroke is haemorrhagic.

The probability of stroke patients being discharged to care is not associated with the supply of care home beds, or with their price.

### 3.4. *Length of stay: hip fracture*

Since we use the logarithm of the length of stay as the dependent variable, the coefficients in Table 3 are the proportionate changes in length of stay in days from a one unit increase in the explanatory variable. Older patients have a longer length of stay. Among patients discharged to care, those who are older than 84 years have 12% longer stays than those aged 65-74. For patients discharged home, those aged 75-84 stay 21% and those aged over 84 stay 32% longer.

1  
2 Patients discharged to care stay 6.7% longer if male, 8.9% longer with an additional  
3  
4 diagnosis, and 7.7% with an additional procedure. Patients transferred to a different hospital  
5  
6 have more than 80% longer stays. Surprisingly, having Charlson comorbidities reduces  
7  
8 length of stay. Patients discharged to care who live in villages and sparsely populated areas  
9  
10 have 11% shorter stays than those living in urban areas. The fifth most income deprived  
11  
12 quintile of the population have 8% longer length of stay than those in less deprived areas if  
13  
14 discharged home. Patients discharged on Monday have longer stays compared with those  
15  
16 discharged on other days of the week, perhaps because some could not be discharged during  
17  
18 the weekend as relevant staff were not available.  
19  
20  
21  
22  
23  
24  
25

26  
27 The accessibility of long-term care beds is associated with shorter lengths of stay for patients  
28  
29 discharged to a care home: patients in LSOAs in higher quintiles of long-term care beds have  
30  
31 shorter hospital stay than those in lower quintiles. Those in the top two quintiles have lengths  
32  
33 of stay which are 22% and 32% shorter than those in the bottom quintile, a difference which  
34  
35 is quantitatively large and statistically significant at 5%. There is no association between beds  
36  
37 supply and length of stay for hip fracture patients discharged home.  
38  
39  
40  
41  
42

43  
44 There is some indication that patients in areas with higher care-home prices stay longer  
45  
46 before being discharged to a care home. Patients in the fourth and fifth quintile of the price  
47  
48 distribution have stays which are 16% ( $p = 0.030$ ) and 18% ( $p = 0.050$ ) longer than those in  
49  
50 the bottom quintile. There is no association between prices and length of stay for hip fracture  
51  
52 patients discharged to their home.  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 3.5. *Length of stay: stroke*

1  
2 Table 4 suggests that among patients discharged home, those who are 75-84 years old and  
3  
4 older than 84 years have greater length of stay (respectively by 16% and 33%). Older patients  
5  
6 discharged to care have shorter stays (by 7% and 23%). Men have 15% shorter stays if  
7  
8 discharged home. An additional procedure increases length of stay by 5% if discharged to  
9  
10 care and 10% if discharged home. An additional diagnosis increases length of stay by 7% and  
11  
12 13%. Length of stay for patients transferred to another hospital is 51% greater if discharged  
13  
14 home and 91% greater if discharged into long term care. Patients with Charlson comorbidities  
15  
16 have a shorter length of stay whether discharged to their own home or to care. Patients living  
17  
18 in areas in the fifth most income deprived quintile have 4% longer length of stay if  
19  
20 discharged home. The type of stroke also affects length of stay: compared to patients whose  
21  
22 stroke is caused by cerebral infarction, length of stay is shorter by 50% and 67% for patients  
23  
24 discharged to care and those discharged to home whose cause of stroke is unspecified.  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 Greater supply of long term care beds is not associated with the length of stay of patients  
35  
36 discharged to a care home. In contrast, more long-term care beds reduces length of stay for  
37  
38 patients discharged home (by 21% in the highest quintile) and the association is significant at  
39  
40 5% in the fourth and fifth quintiles. This is surprising since we would not expect beds supply  
41  
42 to affect the hospital length of stay of patients discharged to their own home. Length of stay  
43  
44 is not associated with the care home prices whether patients are discharged home or to care  
45  
46  
47 home.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 3.6. *Robustness checks*

We tested the sensitivity of our results to different estimation methods and specifications. We interacted income deprivation with the supply and price variables and found that their effect did not vary with deprivation. We measured long-term care supply using patient areas of 20km and 30km radii instead of 10km. We estimated models with beds per capita, rather than entering beds and population separately. We estimated Cox proportional hazard models for length of stay. The key results were not affected and are reported in Gaughan et al (2013).

Table 5 has results from the selection correction model which allows for the interdependence of decisions on length of stay discharge destination. They are very similar to those from the simpler separate linear models and indicate that accounting for selection does not alter the main findings.

## **4. Discussion**

### *4.1. Discharge destination*

As expected, and in line with earlier findings (Bond et al 2000) older patients and those with greater morbidity are more likely to be discharged to a care home when they leave hospital. Other things equal, men are less likely to enter a care home on discharge. This may be because they are more likely to have a spouse or a partner who can provide informal care at home since women have longer life expectancy and tend to be younger than their partners (Wilson and Smallwood 2008).

There is only very weak evidence in our results that accessibility of long term care affects the discharge destination for hip fracture or stroke patients. We find that hip fracture patients who live in areas with higher care home beds supply are more likely to be discharged to a care home. The magnitude of the effects are quite large (4.1% percentage points in the

1 highest quintile compared with an unconditional mean probability of 14.5%) but is not  
2 statistically significant at the conventional (5%) level.  
3  
4  
5  
6

7 There is no association between beds supply and discharge destination for stroke patients,  
8 who are arguably more severely impaired than hip fracture patients and therefore less able to  
9 substitute formal care with informal care at home.  
10  
11  
12  
13  
14  
15

16 Prices are not associated with discharge destination for either hip fracture or stroke patients.  
17 This suggests that the demand for a care home is at most determined by non-monetary costs,  
18 such as the waiting time.  
19  
20  
21  
22  
23  
24  
25

26 In summary, our results suggest that discharge destination is mostly driven by patient  
27 characteristics such as severity, age and comorbidities rather than by care home prices and  
28 bed supply. Stroke patients discharged to a care home have on average been diagnosed with  
29 two additional conditions, received 0.3 more procedures, are more likely to have the more  
30 serious haemorrhagic stroke, and to have more comorbidities. Hip fracture patients  
31 discharged to care are diagnosed with around one additional condition, receive 0.2 more  
32 procedures and have a rate of comorbidity five or six per cent higher than those discharged  
33 home.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

#### 48 *4.2. Length of stay*

49  
50

51 Patient severity is also a key determinant of length of stay. More secondary diagnoses and  
52 procedures are associated with significantly longer stays. The primary diagnosis is also  
53 important. The effect of age is similar for hip fracture and stroke patients discharged to their  
54 own homes: length of stay is over 30% greater for patients over 85. Older hip fracture  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 patients also have longer stays if discharged to care homes, though the effects of age are  
2 around one third as large as for patients discharged to their own homes. Counter-intuitively,  
3  
4 older stroke patients discharged to a care home have shorter hospital stays. These most  
5  
6 complex patients may have the highest priority for nursing home care and have least access to  
7  
8 potential substitutes with care at home, a hypothesis supported in part by the significantly  
9  
10 longer stay for older patients who are discharged home.  
11  
12  
13  
14  
15  
16

17 Patients in areas with greater income deprivation who are discharged to their own home stay  
18  
19 longer in hospitals. Cookson and Laudicella (2011) also find that English elective hip-  
20  
21 replacement patients in poorer areas have greater length of stay. Since poorer individuals are  
22  
23 generally in worse health, income deprivation may also be a proxy of poor health and  
24  
25 hospitals may want patients discharged home to be in better health than those discharged to a  
26  
27 care home where there will be more support.  
28  
29  
30  
31  
32  
33

34 Within a given hospital, hip fracture patients who are discharged to a care home have a  
35  
36 shorter hospital stay if they live in an area with a greater supply of long-term care beds. The  
37  
38 results are consistent with bed blocking: once patients are medically fit for discharge their  
39  
40 length of stay is determined by factors outside the control of the hospital, such as local care  
41  
42 home supply.  
43  
44  
45  
46  
47  
48

49 We also find that hip fracture patients living in areas with higher long-term care prices have  
50  
51 longer hospital stays. This suggests that patients intending to enter a care home take longer  
52  
53 to search for a place when bed prices are higher.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 The difference between the length of stay of patients discharged home and to a care home is  
2 particularly marked for stroke patients. This is expected as stroke is in general a more severe  
3 condition. Most critically, the prognosis following a stroke ranges widely from a very short  
4 stay of a few days to a protracted stay extending into months. The latter case, involving  
5 extensive rehabilitation in a hospital setting, is where patients ultimately discharged to care  
6 are concentrated. Patients discharged to their own home quickly are likely to have been less  
7 severe and have been treated most promptly as these are important factors in determining  
8 outcomes. Stroke patients discharged to a care home are much more likely to have a severe  
9 comorbidity (28% compared to 21%) and also more likely to have had a stroke caused by  
10 haemorrhage (16% compared to 13%).  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 The length of stay of stroke patients discharged to a care home is not affected by the  
27 availability of beds, their price or the quality of care homes in their area. The differences in  
28 results between stroke and hip fracture may be explained by stroke being a more impairing  
29 condition for which patients require much more intensive post hospital care. The demand for  
30 long-term care for stroke patients may therefore be less affected by beds supply and prices.  
31 Note however that the coefficients in the top quintiles are negative and for the top quintile the  
32 coefficient is quantitatively large (a reduction in length of stay by 20%) though not  
33 statistically significant ( $p = 0.17$ ).  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 In contrast to hip fracture patients, we find that for stroke patients, greater availability of beds  
50 reduces length of stay of those discharged to their own home: patients in the highest quintile  
51 of beds supply have a 21% shorter stay than those in the lowest quintile. This may be due to  
52 care homes beds supply being positively correlated with provision of assistance for stroke  
53 patients in their homes.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 As in the case of discharge destination decision, patient severity is the key determinant of  
3  
4 length of stay. However, for hip fracture patients the availability of formal long term care  
5  
6 also appears to have an important impact for those discharged to institutional care. For  
7  
8 patients discharged to their own home the availability of informal care from relatives is also  
9  
10 likely to affect length of stay. We do not have patient-level data on the home circumstances  
11  
12 of hospital patients but do use a small area measure of the proportion of pensioners living  
13  
14 alone. We find, in line with Picone et al (2003), that hip fracture patients discharged to their  
15  
16 home have longer hospital stays in small areas with a higher proportion of single pensioners.  
17  
18  
19  
20  
21  
22  
23

#### 24 *4.3. Concluding remarks*

25  
26 Our results suggest that accessibility of long term care matters for hip fracture patients.  
27  
28 Greater long-term care supply and lower prices are associated with shorter hospital length of  
29  
30 stay for patients discharged to care homes. The effect can be quantitatively large with 20-  
31  
32 30% shorter length of stay for patients with most availability (at the highest bed quintiles).  
33  
34  
35  
36  
37  
38

39 The results are substantially different for stroke patients. Hospital length of stay is not  
40  
41 associated with price (in contrast to hip fracture patients). Counter-intuitively, those  
42  
43 discharged home have a shorter length of stay if availability of long term care beds is high.  
44  
45 The differences between stroke and hip fracture may be that stroke results in more severe and  
46  
47 longer lasting impairment.  
48  
49  
50  
51  
52

53 Overall, for hip fracture patients we find evidence consistent with the ‘bed-blocking’  
54  
55 hypothesis that availability of long-term care affects the length of stay of patients who no  
56  
57 longer need to be in acute hospital and are ready to be discharged. Caring for such patients in  
58  
59  
60  
61  
62  
63  
64  
65

1 hospital is more costly than long-term care. Our results suggest that for hip fracture patients  
2 an expansion of the long-term care sector can reduce hospital length of stay and reduce the  
3 total cost of caring for these patients.  
4  
5  
6  
7  
8  
9

## 10 **Acknowledgements**

11  
12  
13 The research reported in this paper was funded by a grant from the Department of Health to  
14 the Policy Research Unit in the Economics of Health and Social Care Systems. The views  
15 expressed are those of the authors and not necessarily those of the funders. Hospital Episode  
16 Statistics are ©2006/7-2010/11 and re-used with the permission of The Health & Social Care  
17 Information Centre. We are grateful for comments and suggestions from Henning Øien and  
18 participants in the 3rd Norwegian Workshop in Health Economics (Geilo, Norway, 20-21  
19 March 2013), and in the Second Workshop on the Economics of Long Term Care held at  
20 Toulouse School of Economics (Toulouse, France, 22 November 2012).  
21  
22  
23  
24  
25

## 26 **References**

27  
28 Age UK: Care homes: Finding the right carehome, London, (2015) at  
29 [http://www.ageuk.org.uk/Documents/EN-GB/Information](http://www.ageuk.org.uk/Documents/EN-GB/Information_guides/AgeUKIG06_Care_homes_inf.pdf?dtrk=true)  
30 [guides/AgeUKIG06\\_Care\\_homes\\_inf.pdf?dtrk=true](http://www.ageuk.org.uk/Documents/EN-GB/Information_guides/AgeUKIG06_Care_homes_inf.pdf?dtrk=true) on 04.03.16  
31  
32  
33  
34  
35

36 Aharonoff G., Barsky A., Hieber R., Zuckerman J. and Koval K.: Predictors of discharge to a  
37 skilled nursing facility following hip fracture surgery in New York State. *Gerontology*, 50,  
38 298-302 (2004)  
39  
40  
41  
42

43 Allan S. and Forder J.: The determinants of care home closure. *Health Economics*,  
44 24(Supplement 1), 132-145 (2015)  
45  
46  
47  
48

49 Bardsley M., Georghiou T., Chassin L., Lewis G., Steventon A. and Dixon J.: Overlap of  
50 hospital use and social care in older people in England. *Journal of Health Services Research*  
51 *and Policy*, 17, 3, 133-139 (2012)  
52  
53  
54  
55

56 Baumann M., Evans S., Perkins M., Curtis L., Netten A., Fernandez J. and Huxley P.:  
57 Organisation and features of hospital, intermediate care and social services in English sites  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

with low rates of delayed discharge. *Health and Social Care in the Community*, 15, 4, 295-305 (2007)

Bolin K., Lindgren B.I. and Lundborg P.: Informal and formal care among single-living elderly in Europe. *Health Economics*, 17, 3, 393–409 (2008)

Bond J., Gregson B., Smith M., Lecouturier J., Rousseau N. and Rodgers H.: Predicting place of discharge from hospital for patients with a stroke or hip fracture on admission. *Journal of Health Services Research and Policy*, 5, 133-9 (2000)

Bonsang E.: Does informal care from children to their elderly parents substitute for formal care in Europe? *Journal of Health Economics*, 28, 143-154 (2009)

British Orthopaedic Association: The care of patients with fragility fracture (2007)

Bryan K., Gage H. and Gilbert K.: Delayed transfer of older patients from hospital: Causes and policy implications. *Health Policy*, 76, 194-201 (2006)

Butler M., Forte M., Kane R.L., Joglekar S., Duval S.J., Swiontkowski M. and Wilt T.: Treatment of common hip fractures. Evidence Report/Technology Assessment Number 184, Rockville M.D.: Agency for healthcare research and quality (2009)

Care Quality Commission: *Supporting Life After Stroke*. (2011).  
<http://www.cqc.org.uk/content/services-people-who-have-had-stroke-and-their-carers>

Castelli A., Daidone S., Jacobs R., Kasteridis P. and Street A.: The determinants of costs and length of stay for hip fracture patients. *PLoS ONE*, 10, 7, 1-14 (2015)

Castelli A., Laudicella M. and Street A.: Measuring NHS Output Growth. *CHE Research Papers*, 43 (2008)

Charlson M.E., Pompei P., Ales K.L. and Mackenzie C.R.: A new method of classifying prognostic co-morbidity in longitudinal-studies - development and validation. *Journal of Chronic Disease*, 40, 5, 373-383 (1987)

1 Clague J.E., Craddock E., Andrew G., Horan M.A. and Pendleton N.: Predictors of outcome  
2 following hip fracture: Admission time predicts length of stay and in-hospital mortality.  
3 *Injury, International Journal of Care for the Injured*, 33, 1-6 (2002)  
4  
5  
6

7  
8  
9 Cookson R. and Laudicella M.: Do the poor still cost more? The relationship between small  
10 area income deprivation and length of stay for elective hip replacement in the English NHS  
11 from 2001 to 2008. *Social Science and Medicine*, 72, 173-184 (2011)  
12  
13  
14

15  
16 Comas-Herrera A., Wittenberg R. and Pickard L.: From commission to commission:  
17 Financing long-term care in England. Financing long-term care in Europe: Institutions,  
18 markets and models (eds. Costa-Font J. and Courbage C.), Palgrave Macmillan (2009)  
19  
20  
21

22  
23 Department of Health: National Service Framework for Older People, London (2001)  
24  
25

26  
27 Department of Health: National stroke strategy, London (2007)  
28  
29

30  
31 Department of Health: NHS Support for Social Care 2010/11-2012/13, London (2011)  
32  
33

34  
35 Ellis A.A. and Trent R.B. Hospitalized fall injuries and race in California. *Injury Prevention*,  
36 7, 316-20 (2001)  
37  
38

39  
40 Fernandez J. and Forder J.: Consequences of local variations in social care on the  
41 performance of the acute health care sector. *Applied Economics*, 40, 12, 1503-1518 (2008)  
42  
43  
44

45  
46 Forder J.: Self-funded social care for older people: An analysis of eligibility, variations and  
47 future projections. *PSRRU DP 2507* (2007) <http://www.pssru.ac.uk/pdf/dp2505.pdf>  
48  
49

50  
51 Forder J.: Long-term care and hospital utilisation by older people: An analysis of substitution  
52 rates. *Health Economic*, 18, 1322-1338 (2009)  
53  
54

55  
56 Forder J. and Fernandez J-L.: Geographical differences in the provision of care home services  
57 in England. *PSSRU Discussion Paper 2824(revised)* (2012)  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Gaughan, J., Gravelle, H., Santos, R., Siciliani, L.: Long term care provision, hospital length of stay and discharge destination for hip fracture and stroke patients. *CHE Research Paper* 86 (2013)

Gaughan, J., Gravelle, H., Siciliani, L.: Testing the bed-blocking hypothesis: Does higher supply of nursing and care homes reduce delayed hospital discharges? *Health Economics*, 24(S1), 32–44 (2015)

Gilbert R., Todd C., May M., Yardley L. and Ben-Shlomo Y.: Socio-demographic factors predict the likelihood of not returning home after hospital admission following a fall. *Journal of Public Health*, 32, 117-124 (2010)

Glendinning C.: Breaking down barriers: integrating health and care services for older people in England. *Health Policy*, 65, 139-151 (2003)

Grabowski D., Norton E. and Van Houtven C. Informal care. In: Jones A. (Eds), *The Elgar Companion in Health Economics*, Second edition, Edward Elgar Publishing, Cheltenham, p. 307-317 (2012)

Heckman J.: Sample selection bias as a specification error. *Econometrica*. 47, 153-161 (1979)

Holmås T.H., Islam M.K. and Kjerstad E.: Between two beds: Inappropriately delayed discharges from hospitals. *International Journal of Health Care Finance and Economics*, 13, 3-4, 201-217 (2013)

Holmås T.H., Kjerstad E., Lurås H. and Straume O.R.: Does monetary punishment crowd out pro-social behaviour? A natural experiment on hospital length of stay, *Journal of Economic Behavior & Organization*, 75, 261-267 (2010)

House of Commons, Committee of Public Accounts: Ensuring the Effective Discharge of Older Patients from NHS Acute Hospitals: Thirty-Third Report of Session 2002–2003, HC 459 Stationery Office, London (2003)

1 Jarman B., Aylin P. and Bottle A.: Discharge destination and length of stay: Differences  
2 between US and English hospitals for people aged 65 and over. *British Medical Journal*, 328,  
3 605 (2004)  
4  
5

6  
7 Jarret T.: Social care: The state of the care home market (England). Briefing Paper, House of  
8 Commons Library (2016)  
9

10  
11  
12 Jørgensen H.S., Nakayama H., Raaschou H.O. and Olsen T.S.: Intracerebral haemorrhage  
13 versus infarction: Stroke severity, risk factors and prognosis. *Annals of Neurolog*, 38, 1, 45-  
14 50 (1995)  
15  
16

17  
18  
19  
20 Kasteridis, P., Street, A., Dolman, M., Gallier, L., Hudson, K., Martin, J., and Wyer, I.: Who  
21 would benefit most from improved integrated care? Implementing an analytical strategy in  
22 South Somerset. *Journal of Integrated Care*, 15, 1-11 (2015)  
23  
24

25  
26  
27 Laing and Buisson: Care homes complete dataset, Laing and Buisson (2010)  
28

29  
30  
31 de Maijer, C., Koopmanschap, M., d’Uva, T. B. and van Doorslaer, E.: Determinants of long-  
32 term care spending: Age, time to death or disability?. *Journal of Health Economics*, 30, 425-  
33 438 (2011)  
34  
35

36  
37  
38 Mundlak Y.: On the pooling of time series and cross section data. *Econometrica*, 46, 69-85  
39 (1978)  
40  
41

42  
43  
44 National Audit Office: Inpatient Admissions and Bed Management in NHS Acute Hospitals,  
45 London: Stationary Office (2000)  
46

47  
48  
49 National Audit Office: *Reducing brain damage: Faster access to better stroke care*, London  
50 (2005)  
51

52  
53  
54  
55 National Clinical Guidelines Centre: The management of hip fracture in adults, London,  
56 National Clinical Guidelines Centre (2011)  
57

58  
59  
60 OECD: Help wanted? Providing and paying for long-term care, *OECD publications* (2011)  
61



1 OFT: Care homes for older people in the UK: A market study (2005)  
2  
3  
4

5 Picone G., Wilson R.M. and Chou S-Y.: Analysis of hospital length of stay and discharge  
6 destination using hazard functions with unmeasured heterogeneity. *Health Economics*, 12,  
7 1021-1034 (2003)  
8  
9

10  
11  
12 Pioneer Network: Long-term care and culture change glossary: Definitions of common terms  
13 used in long-term care and culture change (2011) at  
14 <http://www.pioneernetwork.net/Consumers/PickerGlossary/> accessed on 04.03.2016  
15  
16  
17

18  
19  
20 Steventon A. and Roberts A.: Estimating length of stay in publicly-funded residential and  
21 nursing care homes: A retrospective analysis using linked administrative data sets. *BMC*  
22 *Health Services Research*, 12, 377-385 (2012)  
23  
24

25  
26  
27 Stroke Association: Accommodation after stroke (2013) at  
28 [https://www.stroke.org.uk/sites/default/files/accommodation\\_after\\_stroke.pdf](https://www.stroke.org.uk/sites/default/files/accommodation_after_stroke.pdf) accessed on  
29 12.04.2016  
30  
31  
32

33  
34  
35 Van Houtven C.H., Norton E.C.: Informal care and health care use of older adults. *Journal of*  
36 *Health Economic*,. 23, 6, 1159-1180 (2004)  
37  
38

39  
40 Vetter N.: Inappropriately delayed discharge from hospital: What do we know? *British*  
41 *Medical Journal*,. 326, 7395, 927-928 (2003)  
42  
43

44  
45  
46 Wanless D.: Securing good care for elderly people: Taking a long-term view. Kings Fund,  
47 London (2006)  
48  
49

50  
51 Wilson B. and Smallwood S.: Age differences at marriage and divorce. *Population Trends*.  
52 132, Summer, 17-25 (2008)  
53  
54

55  
56  
57 Wong A., Elderkamp-de Groot R., Polder J. and van Exel J.: Predictors of long-term care  
58 utilization by Dutch hospital patients aged 65+. *BMC Health Services Research*, 10, 110  
59 (2010)<http://www.biomedcentral.com/1472-6963/10/110>  
60  
61  
62

1  
2 Wooldridge J. *Econometric Analysis of Cross Section and Panel Data*. second edition.  
3  
4 (2010). M.I.T Press.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 1. Summary statistics**

	Hip Fracture				Stroke			
	mean	sd	min	max	mean	sd	min	Max
<i>Dependent variables</i>								
Discharged to care home	0.145	0.352	0	1	0.135	0.342	0	1
Length of stay if discharged to care home	32.68	27.98	2	167	62.07	44.16	1	460
Length of stay if discharged home	19.95	17.90	2	168	23.38	30.13	1	394
<i>Patient characteristics</i>								
Age group: 65-74	0.166	0.372	0	1	0.299	0.458	0	1
Age group: 75-84	0.409	0.492	0	1	0.429	0.495	0	1
Age group: 85plus	0.425	0.494	0	1	0.271	0.445	0	1
Male patient	0.223	0.416	0	1	0.466	0.499	0	1
Total diagnoses	5.713	2.919	1	39	6.234	3.511	1	32
Total procedures	2.818	1.546	0	24	2.671	1.786	0	22
Patient transferred in CIPS	0.049	0.217	0	1	0.137	0.343	0	1
Pertrochanteric fracture	0.228	0.420	0	1				
Subtrochanteric fracture	0.029	0.167	0	1				
Unspecified hip fracture	0.743	0.437	0	1				
Stroke caused by a hemorrhage					0.137	0.344	0	1
Stroke caused by an infarction					0.615	0.487	0	1
Stroke not hemorrhage or infarction					0.213	0.410	0	1
Occluded cerebral vessels no infarction					0.003	0.051	0	1
Other stroke					0.032	0.176	0	1
No Charlson comorbidities	0.493	0.500	0	1	0.515	0.500	0	1
1 minor Charlson comorbidity	0.334	0.472	0	1	0.266	0.442	0	1
≥2 minor or ≥1 major Charlson comorbidity	0.172	0.378	0	1	0.218	0.413	0	1
Discharged on Monday	0.152	0.359	0	1	0.174	0.379	0	1
Discharged on Tuesday	0.191	0.393	0	1	0.186	0.389	0	1
Discharged on Wednesday	0.190	0.392	0	1	0.185	0.388	0	1
Discharged on Thursday	0.180	0.385	0	1	0.180	0.385	0	1
Discharged on Friday	0.209	0.406	0	1	0.216	0.411	0	1
Discharged on Saturday	0.052	0.223	0	1	0.044	0.204	0	1
Discharged on Sunday	0.017	0.131	0	1	0.015	0.122	0	1
<i>Socioeconomic characteristics of patient areas of residence</i>								
% LSOA 60+ pop on income based benefit	19.65	11.81	1.0	95.0	19.81	12.31	1.0	95.0
% LSOA pop who are non-white	6.21	11.20	0	90.45	7.09	13.00	0	94.8
% LSOA pop with good SAH	67.13	6.51	37.3	87.6	67.22	6.34	37.3	87.0
% LSOA pop with fairly good SAH	23.03	3.47	10.4	36.1	23.07	3.41	10.7	37.3
% LSOA pop with not good SAH	9.84	3.62	1.7	31.0	9.71	3.51	1.7	31.0
% single pensioner households in LSOA	16.09	6.01	0.5	51.0	15.92	6.00	0.0	51.0
Patient resident in London	0.089	0.284	0	1	0.103	0.304	0	1
LSOA within 10km of Scottish boarder	0.001	0.026	0	1	0.001	0.029	0	1
LSOA within 10km of Welsh boarder	0.012	0.109	0	1	0.015	0.121	0	1
Urban > 10k people	0.791	0.407	0	1	0.788	0.409	0	1
Town and fringe	0.111	0.314	0	1	0.114	0.318	0	1
Village or hamlet and isolated dwellings	0.098	0.297	0	1	0.098	0.297	0	1
Total retired population within 10km (000s)	67.2	62.1	0.5	328.7	70.9	67.1	348	328.2
<i>Formal long-term care</i>								
Care home beds within 10km (000s)	2.31	1.79	0	7.81	2.41	1.92	0	7.82
Beds within 10km/retired population	0.037	0.010	0	0.116	0.037	0.010	0	0.116
Average price within 10km	523.21	93.05	232	971	525.25	91.49	232	961
Average care home rating within 10km	3.03	0.17	1	4	3.03	0.17	1	4
Number of patients	21959				33101			

Notes: CIPS: continuous in-patient spell. LSOA: lower super output area. SAH: self-assessed health.

**Table 2. Determinants of discharge to care home**

	Hip Fracture		Stroke	
	b	p	b	p
<i>Patient characteristics</i>				
Age 75-84	0.062	0.000	0.054	0.000
Age 85plus	0.114	0.000	0.125	0.000
Male	-0.016	0.005	-0.043	0.000
Number diagnoses	0.011	0.000	0.018	0.000
Number procedures	0.005	0.021	0.002	0.110
Patient transferred	0.005	0.793	0.029	0.025
Pertrochanteric fracture	0.000	0.938		
Subtrochanteric fracture	0.004	0.766		
Stroke caused by a hemorrhage			0.014	0.035
Stroke hemorrhage or infarction			-0.019	0.000
Occluded cerebral no infarction			0.009	0.776
Other stroke			-0.074	0.000
1 minor Charlson comorbidity	0.031	0.000	0.002	0.655
≥2 minor/≥1 major Charlson comorbidity	0.019	0.047	-0.001	0.926
<i>Socioeconomic characteristics of patient areas of residence</i>				
LSOA 5 <sup>th</sup> income deprivation quintile	-0.013	0.168	-0.012	0.050
% LSOA pop non white	0.000	0.615	0.000	0.381
% LSOA not good SAH	0.000	0.788	0.004	0.002
% LSOA single pensioner household	0.001	0.278	-0.001	0.080
London LSOA	-0.031	0.364	-0.027	0.371
LSOA 10km of Scottish boarder	-0.012	0.897	-0.069	0.252
LSOA 10km of Welsh boarder	-0.003	0.923	0.024	0.238
Town and fringe	-0.007	0.340	-0.003	0.560
Village, hamlet, isolated dwellings	-0.011	0.204	-0.004	0.604
Population within 10km (100000s)	0.013	0.578	0.011	0.449
<i>Formal long-term care</i>				
<b>Beds within 10km second quintile</b>	0.009	0.396	0.002	0.790
<b>Beds within 10km third quintile</b>	0.021	0.088	-0.012	0.238
<b>Beds within 10km fourth quintile</b>	0.025	0.263	0.018	0.188
<b>Beds within 10km top quintile</b>	0.041	0.197	0.014	0.512
<b>Price within 10km second quintile</b>	-0.010	0.339	0.002	0.855
<b>Price within 10km third quintile</b>	-0.012	0.326	0.011	0.267
<b>Price within 10km fourth quintile</b>	0.006	0.739	0.019	0.127
<b>Price within 10km top quintile</b>	-0.004	0.817	0.020	0.175
<b>Care home ratings 10km mean</b>	-0.005	0.834	0.001	0.935
Constant	-0.021	0.788	-0.063	0.239
Hospital effects	FE		FE	
R <sup>2</sup>	0.0327		0.074	
Observations	21959		33101	

Notes. Fixed effect panel data linear probability model of discharge to care home versus discharge to own home with cluster robust standard errors. The  $\chi^2$  statistic for the auxiliary regression test (see section 2.6) for the consistency of the random error specification is  $\chi^2(29)= 28.15$ ,  $p = 0.5097$  for hip fracture and  $\chi^2(31)= 125.85$ ,  $p = 0.000$  for stroke.

**Table 3. Determinants of length of stay: hip fracture**

	Discharged to care		Discharged to home	
	b	p	b	p
<i>Patient characteristics</i>				
Age 75-84	0.065	0.128	0.209	0.000
Age 85plus	0.120	0.007	0.315	0.000
Male	0.067	0.026	0.008	0.464
Number diagnoses	0.089	0.000	0.081	0.000
Number procedures	0.077	0.000	0.080	0.000
Patient transferred	0.855	0.000	0.870	0.000
Pertrochanteric fracture	-0.036	0.272	-0.005	0.651
Subtrochanteric fracture	0.035	0.631	0.117	0.000
1 minor Charlson comorbidity	0.009	0.769	-0.039	0.001
≥2 minor/ ≥1 major Charlson comorbidity	-0.136	0.000	-0.074	0.000
Discharged on Tuesday	-0.087	0.037	-0.076	0.000
Discharged on Wednesday	-0.175	0.000	-0.082	0.000
Discharged on Thursday	-0.110	0.003	-0.086	0.000
Discharged on Friday	-0.141	0.000	-0.116	0.000
Discharged on Saturday	-0.154	0.006	-0.189	0.000
Discharged on Sunday	-0.197	0.057	-0.135	0.000
<i>Socioeconomic characteristics of patient areas of residence</i>				
LSOA 5 <sup>th</sup> income deprivation quintile	0.023	0.547	0.075	0.000
% LSOA pop non white	0.000	0.844	0.001	0.281
% LSOA not good SAH	-0.006	0.365	-0.008	0.002
% LSOA households single pensioner	-0.002	0.303	0.005	0.000
London LSOA	-0.185	0.468	-0.018	0.784
LSOA 10km of Scottish boarder	0.448	0.000	0.099	0.079
LSOA 10km of Welsh boarder	-0.224	0.084	-0.140	0.012
Town and fringe	0.009	0.849	0.000	0.989
Village, hamlet isolated dwellings	-0.105	0.045	-0.048	0.014
Population within 10km (100000s)	0.230	0.040	0.048	0.095
<i>Formal long-term care</i>				
<b>Beds within 10km second quintile</b>	-0.049	0.336	-0.012	0.466
<b>Beds within 10km third quintile</b>	-0.064	0.391	0.011	0.598
<b>Beds within 10km fourth quintile</b>	-0.216	0.036	0.007	0.835
<b>Beds within 10km top quintile</b>	-0.319	0.020	-0.022	0.641
<b>Price within 10km second quintile</b>	-0.014	0.769	-0.030	0.178
<b>Price within 10km third quintile</b>	0.006	0.925	-0.044	0.116
<b>Price within 10km fourth quintile</b>	0.162	0.030	-0.031	0.342
<b>Price within 10km top quintile</b>	0.175	0.050	-0.010	0.792
<b>Care home ratings 10km mean</b>	0.127	0.182	0.030	0.428
Constant	1.977	0.000	1.782	0.000
Hospital effects	FE		FE	
R <sup>2</sup>	0.305		0.311	
Observations	3175		18784	

Notes. Dependent variable: natural logarithm of length of stay. Coefficients are the proportionate change in length of stay in days from a one unit increase in the explanatory variable. Fixed effects panel data models with cluster robust standard errors. The  $\chi^2$  statistic for the auxiliary regression test (see section 2.6) for the consistency of the random error specification is  $\chi^2(35)= 115.56$ ,  $p= 0.000$  for patients discharged to care and  $\chi^2(35)= 100.87$ ,  $p = 0.000$  for those discharged to their own home.

**Table 4. Determinants of length of stay: stroke**

	Discharged to care		Discharged to home	
	b	p	b	p
<i>Patient characteristics</i>				
Age 75-84	-0.074	0.035	0.160	0.000
Age 85plus	-0.225	0.000	0.327	0.000
Male	0.027	0.282	-0.146	0.000
Number diagnoses	0.069	0.000	0.129	0.000
Number procedures	0.052	0.000	0.099	0.000
Patient transferred	0.509	0.000	0.909	0.000
Stroke caused by a hemorrhage	-0.080	0.010	0.026	0.293
Stroke hemorrhage or infarction	-0.087	0.005	-0.202	0.000
Occluded cerebral vessels no infarction	0.083	0.569	-0.192	0.106
Other stroke	-0.497	0.000	-0.672	0.000
1 minor Charlson comorbidity	-0.128	0.000	-0.077	0.000
≥2 minor or ≥1 major Charlson comorbidity	-0.207	0.000	-0.045	0.041
Discharged on Tuesday	-0.075	0.024	-0.156	0.000
Discharged on Wednesday	-0.080	0.019	-0.170	0.000
Discharged on Thursday	-0.102	0.008	-0.164	0.000
Discharged on Friday	-0.099	0.008	-0.313	0.000
Discharged on Saturday	-0.223	0.000	-0.423	0.000
Discharged on Sunday	-0.203	0.117	-0.675	0.000
<i>Socioeconomic characteristics of patient areas of residence</i>				
LSOA fifth income deprivation quintile	0.022	0.530	0.043	0.038
% LSOA pop non white	0.001	0.461	0.001	0.211
% LSOA not good SAH	-0.013	0.020	0.003	0.352
% LSOA households single pensioner	0.004	0.103	-0.001	0.334
London LSOA	0.179	0.112	-0.126	0.103
LSOA within 10km of Scottish boarder	0.860	0.000	-0.008	0.899
LSOA within 10km of Welsh boarder	0.022	0.765	0.002	0.981
Town and fringe	0.040	0.259	-0.039	0.051
Village or hamlet and isolated dwellings	0.007	0.879	-0.005	0.833
Population within 10km (100000s)	0.044	0.667	0.184	0.000
<i>Formal long-term care</i>				
<b>Beds within 10km second quintile</b>	0.020	0.655	-0.026	0.289
<b>Beds within 10km third quintile</b>	-0.052	0.411	-0.050	0.153
<b>Beds within 10km fourth quintile</b>	-0.061	0.522	-0.111	0.035
<b>Beds within 10km top quintile</b>	-0.196	0.167	-0.209	0.003
<b>Price within 10km second quintile</b>	-0.013	0.786	-0.024	0.449
<b>Price within 10km third quintile</b>	-0.060	0.247	0.018	0.617
<b>Price within 10km fourth quintile</b>	-0.046	0.530	-0.016	0.678
<b>Price within 10km top quintile</b>	-0.051	0.624	-0.032	0.477
<b>Care home ratings within 10km mean</b>	0.122	0.206	-0.073	0.144
Constant	3.069	0.000	1.678	0.000
Hospital effects	FE		FE	
R <sup>2</sup>	0.253		0.337	
Observations	4465		28636	

Notes. Dependent variable: natural logarithm of length of stay. Coefficients are the proportionate change in length of stay in days from a one unit increase in the explanatory variable. Fixed effects panel data models with cluster robust standard errors. The  $\chi^2$  statistic for the auxiliary regression test (see section 2.6) for the consistency of the random error specification is  $\chi^2(37)= 114.35$ ,  $p = 0.000$  for patients discharged to care and  $\chi^2(37)= 450.81$ ,  $p = 0.000$  for those discharged to their own home.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 5. Comparison of selection correction and separate linear models**

	Discharge Destination				Ln Length of Stay if Discharged to Care				Ln Length of Stay if Discharged Home			
Hip Fracture Models												
	Linear FE		Selection correction First Stage		Linear FE		Selection correction 2nd Stage		Linear FE		Selection correction 2nd Stage	
	b	p	B	p	b	p	b	P	b	p	b	p
<b>Beds within 10km 2nd quintile</b>	0.009	0.396	0.042	0.335	-0.049	0.336	-0.036	0.445	-0.012	0.466	-0.013	0.426
<b>Beds within 10km 3rd quintile</b>	0.021	0.088	0.105	0.070	-0.064	0.391	-0.028	0.669	0.011	0.598	0.012	0.602
<b>Beds within 10km 4th quintile</b>	0.025	0.263	0.114	0.159	-0.216	0.036	-0.181	0.046	0.007	0.835	0.008	0.799
<b>Beds within 10km top quintile</b>	0.041	0.197	0.176	0.141	-0.319	0.020	-0.246	0.061	-0.022	0.641	-0.022	0.635
<b>Price within 10km 2nd quintile</b>	-0.010	0.339	-0.065	0.173	-0.014	0.769	-0.024	0.643	-0.030	0.178	-0.031	0.086
<b>Price within 10km 3rd quintile</b>	-0.012	0.326	-0.086	0.143	0.006	0.925	-0.010	0.883	-0.044	0.116	-0.044	0.043
<b>Price within 10km 4th quintile</b>	0.006	0.739	0.009	0.906	0.162	0.030	0.203	0.015	-0.031	0.342	-0.030	0.270
<b>Price within 10km top quintile</b>	-0.004	0.817	-0.035	0.703	0.175	0.050	0.211	0.040	-0.010	0.792	-0.009	0.789
Inverse Mills ratio							0.233	0.000			-0.324	0.035
Stroke Models												
	b	p	B	p	b	p	b	P	b	p	b	p
<b>Beds within 10km 2nd quintile</b>	0.002	0.790	0.019	0.598	0.020	0.655	0.023	0.596	-0.026	0.289	-0.022	0.346
<b>Beds within 10km 3rd quintile</b>	-0.012	0.238	-0.061	0.204	-0.052	0.411	-0.075	0.213	-0.050	0.153	-0.065	0.045
<b>Beds within 10km 4th quintile</b>	0.018	0.188	0.091	0.162	-0.061	0.522	-0.048	0.551	-0.111	0.035	-0.092	0.036
<b>Beds within 10km top quintile</b>	0.014	0.512	0.079	0.410	-0.196	0.167	-0.174	0.143	-0.209	0.003	-0.199	0.002
<b>Price within 10km 2nd quintile</b>	0.002	0.855	0.004	0.929	-0.013	0.786	-0.004	0.932	-0.024	0.449	-0.025	0.349
<b>Price within 10km 3rd quintile</b>	0.011	0.267	0.059	0.206	-0.060	0.247	-0.032	0.572	0.018	0.617	0.030	0.335
<b>Price within 10km 4th quintile</b>	0.019	0.127	0.092	0.116	-0.046	0.530	-0.028	0.696	-0.016	0.678	0.003	0.936
<b>Price within 10km top quintile</b>	0.020	0.175	0.088	0.235	-0.051	0.624	-0.037	0.698	-0.032	0.477	-0.014	0.770
Inverse Mills ratio							0.334	0.076			-0.778	0.082

Notes. There are 21,959 hip fracture observations and 33,101 stroke observations. Linear FE model coefficients are from the models in Tables 2-4. Selection correction models include hospital means of patient level variables and deviations from means. Coefficients are those on the deviations

## Data Appendix

1  
2 A patient is admitted from home if their admission code is “usual place of residence,  
3 including no fixed abode” or “temporary place of residence when usually resident elsewhere,  
4 for example, hotels and residential educational establishments” (admisorc = 19, 29). Care  
5 homes are not considered usual residence and are therefore excluded (see more below). Each  
6 HES record covers a single finished consultant episode (FCE) during which the patient is  
7 continuously under a single consultant (senior hospital doctor).  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

19 We link FCEs into continuous inpatient stays (CIPS) to allow for changes of consultant,  
20 including transfers to other hospitals. We combine FCEs into CIPS using the methodology in  
21 Castelli et al. (2008) and Cookson and Laudicella (2011). We include patients whose CIPS  
22 finish in the financial year 1 April 2008 to 31 March 2009 and start between 1 April 2007 and  
23 31 March 2009.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 Patients are coded as being discharged to their home if their HES discharge destination field  
35 indicates usual or temporary residence (disdest 19, 29), to a long-term care facility if their  
36 destination is an NHS-run nursing home, a residential home or group home (disdest 54), a  
37 local authority care home (disdest 69) or non-NHS (other than local authority) residential  
38 care home (disdest 85).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 Hip fracture patients have a primary diagnosis ICD10 code of S72.0 (fracture of neck of  
50 femur or unspecified femur fracture), S72.1 (pertrochanteric fracture) or S72.2  
51 (subtrochanteric fracture) (Jarman et al 2004). Stroke patients have primary ICD10 code I60-  
52 2 (intracerebral haemorrhage), I63 (cerebral infarction), I64 (unspecified stroke), I66 or I67.2,  
53 I69.8 or R47.0 (other form of stroke).  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2 There are 33,082 hip fracture and 59,316 stroke emergency admissions in our study period.  
3  
4 We exclude 11,113 hip fracture and 26,211 stroke patients from the analysis because: the  
5  
6 patient dies in-hospital (4,253 for hip fracture and 15,501 for stroke), the hospital spell is  
7  
8 incomplete (2,080 for hip fracture and 2,518 for stroke), the patient is discharged elsewhere  
9  
10 than to usual residence or care home (1,595 for hip fracture and 3,211 for stroke), is admitted  
11  
12 from elsewhere than usual residence where “usual residence” excludes a care home (1,910 for  
13  
14 hip fracture and 2,437 for stroke), has a repeat emergency admission (376 for hip fracture and  
15  
16 1,194 for stroke), is treated in a hospital with 10 or fewer cases in 2008/9 (46 for hip fracture  
17  
18 and 60 for stroke). We exclude cases with very long length of stay, and the logarithm of the  
19  
20 length of stay is more than three standard deviations above the mean (205 for hip fracture),  
21  
22 and cases with missing data (658 for hip fracture and 1294 for stroke).  
23  
24  
25  
26  
27  
28  
29  
30

31 HES records the patient’s Lower Super Output Area (LSOA) of residence. There are 32,482  
32  
33 LSOAs in England with an average population of 1500 in 2001. We compute the number of  
34  
35 beds in care homes within 10km of the centroid of the patient’s LSOA of residence. For each  
36  
37 provider, we have the minimum and maximum price by type of room (single, shared) and  
38  
39 type of care (nursing, non-nursing). We compute the average price for care homes within  
40  
41 10km of each LSOA centroid. 1682 care homes (14%) do not report any price and we impute  
42  
43 the price of these from the average price for providers in the same quintile of beds supply.  
44  
45 We measure the average quality of care homes within 10km by assigning numerical values 1  
46  
47 to 4 to the CQC quality ratings (poor, adequate, good, and excellent). We use the same  
48  
49 strategy as for missing price data to impute quality rating for 1953 providers without  
50  
51 information (16.5%).  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65