



This is a repository copy of *Connexin43 intercellular communication drives the early differentiation of human bone marrow stromal cells into osteoblasts.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/114525/>

Version: Accepted Version

---

**Article:**

Talbot, J., Brion, R., Lamora, A. et al. (4 more authors) (2017) Connexin43 intercellular communication drives the early differentiation of human bone marrow stromal cells into osteoblasts. *Journal of Cellular Physiology*. ISSN 0021-9541

<https://doi.org/10.1002/jcp.25938>

---

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**Connexin43 intercellular communication drives the early differentiation of human bone marrow stromal cells into osteoblasts**

Julie Talbot<sup>1</sup>, Régis Brion<sup>1,2,3,4</sup>, Audrey Lamora<sup>1,2,3,4</sup>, Mathilde Mullard<sup>1,2,3</sup>, Sarah Morice<sup>1,2,3</sup>,  
Dominique Heymann<sup>1,4,5</sup> and Franck Verrecchia<sup>1,2,3,\*</sup>

<sup>1</sup>INSERM, UMR 957, Nantes, France

<sup>2</sup>INSERM, UMR 1238, Nantes, France

<sup>3</sup>Université Bretagne Loire, Nantes, France

<sup>4</sup>CHU, Nantes, France

<sup>5</sup>Department of Oncology and metabolism, INSERM European Associated Laboratory, University of Sheffield, Sheffield, UK.

**\*Address correspondence to:** Franck Verrecchia, INSERM UMR1238, Faculté de Médecine, 1 rue Gaston Veil, 44000 Nantes, France, Phone: +33 240412842, Fax: +33 240412860, E-mail: [franck.verrecchia@inserm.fr](mailto:franck.verrecchia@inserm.fr)

**Running head:** Role of Cx43 in hBMSCs differentiation

**Keywords :** cell differentiation, connexin43, gap junction, human bone marrow stromal cells, intercellular communication, osteoblast

**Total number of figures :** 5

**Total number of tables :** 1

Conflict of interest : NO

## Abstract

Although it has been demonstrated that human bone marrow stromal cells (hBMSCs) express the ubiquitous connexin43 (Cx43) and form functional gap junctions, their role in the early differentiation of hBMSCs into osteoblasts remains poorly documented. Using *in vitro* assays, we show that Cx43 expression and gap junctional intercellular communication (GJIC) are increased during the differentiation of hBMSCs into osteoblasts, both at the protein and mRNA levels. Two independent procedures to reduce GJIC, a pharmacological approach with GJIC inhibitors (18 $\alpha$ -glycyrrhetic acid and Gap27 peptide) and a molecular approach using small interfering RNA against Cx43, demonstrated that the presence of Cx43 and functional junctional channels are essential to the ability of hBMSCs to differentiate into osteoblasts *in vitro*. In addition, a reduced GJIC decreases the expression of *Runx2*, the major transcription factor implicated in the control of osteoblast commitment and early differentiation of hBMSCs into osteoblasts, suggesting that GJIC mediated by Cx43 is implicated in this process. Together our results demonstrate that GJIC mediated by the Cx43 channels plays a central role throughout the differentiation of hBMSC into osteoblasts, from the early stages to the process of mineralization.

## Introduction

Bone formation is a regulated process that takes place during embryonic development, growth, remodeling and repair. It begins with the commitment of mesenchymal stem cells (MSC) into the osteoblastic differentiation program, continues with the early differentiation of stem cells into osteoblast progenitors and the subsequent differentiation of these osteoblast progenitors into immature osteoblasts, and finally the late differentiation of these immature osteoblast into mature osteoblasts (Augello 2010; Vaananen 2005; Neve et al., 2011). These MSC, located mainly in the bone marrow stroma, can also differentiate into other mesenchymal cell lineages such as fibroblasts, chondrocytes and bone marrow stromal cells including adipocytes, depending on the activated signaling transcription pathways (Friedenstein et al., 1987; Owen 1988; Marie 2008). At the end of the osteogenic differentiation process, mature osteoblasts produce many markers such as alkaline phosphatase (ALP), collagen type I (COL1), bone sialoprotein (BSP), osteopontin and osteocalcin (OCN) (Wennberg et al., 2000). This process results in the formation of an extracellular organic matrix which will be mineralized.

Connexins, the structural proteins of gap junctions, are a family of transmembrane proteins that oligomerize into hemichannels containing six connexin subunits (Sosinsky 2005). Upon reaching the cell surface, two hemichannels pair to complete an intercellular gap junction channel, which directly links the cytoplasm of neighboring cells and mediates the exchange of low-molecular-mass molecules (<1000 Da), including cAMP, inositol trisphosphate, and  $Ca^{2+}$  (Sohl and Willecke, 2004; Nielsen et al., 2012).

Multiple lines of evidence have established that Cx43, the most abundant connexin in bone cells, and subsequent intercellular channels play a major role in skeletal development and for the function of osteoblasts (Stains and Civitelli 2016; Plotkin and Stains, 2015). Cx43 gene mutations have thus been directly associated with abnormalities of skeletal development observed for example in oculodentodigital and cranometaphysal dysplasias (Paznekas et al., 2003; Laird, 2014). Specifically, it has been demonstrated that Cx43 expression is required for osteoblastic gene expression and function in primary and cultured osteoblasts (Lecanda et al., 1998; Lecanda et al., 2000). For example, it has been shown that gap junctional communication is required for the maturation process of osteoblastic cells in culture and for the stimulation of osteoblast

mineralization in response to parathyroid hormone (Schiller et al., 1992; Schiller et al., 2001a; Schiller et al., 2001b). In addition, *in vivo* experiments using either Cx43<sup>-/-</sup> or Cx43<sup>fl/-</sup>;Col1a.1-2.3 kb-Cre mice have demonstrated that Cx43 play a crucial role in osteoblastic differentiation and function (Plotkin and Stains, 2015).

However, although it was shown that human MSC (hMSC) express Cx40, Cx43 and Cx45 (Valiunas et al., 2004, Talbot et al., 2013), and form functional gap junctions, the role of connexins in the *in vitro* differentiation of human MSCs into osteoblast progenitors remains poorly documented. In this context, the aim of the present work was to determine the specific role of Cx43 in the early differentiation of hMSCs into osteoblasts and throughout the process of osteoblast differentiation by using small interfering RNA against Cx43 as well as via a pharmacological approach with GJIC inhibitors (18 $\alpha$ -glycyrrhetic acid and Gap27 peptide).

## Materials and methods

### Ethics statement

Bone marrow aspirates were obtained from healthy volunteers (age=46±12; range=36-67) during orthopaedic surgical procedures after exposure of the iliac crest in the orthopaedic department of the University Hospital of Tours (France). The written consent of informed patients was obtained for the use of tissue samples for anonymized research, in accordance with French law (Art. L. 1245-2 du code de la santé publique, Loi n° 2004-800 du 6 août 2004 Journal Officiel du 7 août 2004). The donors had no significant medical history. The study followed the ethical guidelines of the University Hospital of Tours and was approved by the ethics committee "Comite de protection des personnes" (Tours - Region Centre Ouest).

### Cell culture

Human bone marrow stromal cells (hBMSCs) were isolated by Ficoll density gradient separation as previously described (Chang et al., 2009). Non-adherent cells were removed by replacing the culture medium and hBMSCs were then cultured in medium composed of Dulbecco's Modified Eagle Medium (Lonza, Belgium), 10% fetal bovine serum (Hyclone Perbio, France) with 1 ng/ml basic fibroblast growth factor (bFGF; R&D systems, UK). Adherent cells were characterized by flow cytometry (CD45<sup>-</sup>, CD34<sup>-</sup>, CD105<sup>+</sup>, CD73<sup>+</sup>, CD90<sup>+</sup> and CD146<sup>+</sup>, purity ≥ 99%) prior to further experiments. The expression of each marker was not changed in the presence of bFGF.

### Osteoblast differentiation

hBMSCs were seeded at  $1 \times 10^4$  cells/cm<sup>2</sup> in 24-well plates in culture medium without bFGF. After 3 days, when the cells were confluent, the medium was replaced by fresh medium supplemented with vitamin D3 ( $10^{-8}$  M; Hoffmann-La Roche, Switzerland) and dexamethasone ( $10^{-8}$  M; Sigma, France), referred to as "osteogenic medium" in the text. Three days later, ascorbic acid (50 µg/ml; Sigma) and β-glycerophosphate ( $10^{-7}$  M; Sigma) was added to allow mineralization. Alizarin red staining was used to detect the mineralized nodules formed, as previously described (Chipoy et al., 2004). Briefly, around day 15, cells were fixed in ice-cold 70% ethanol for 1h and incubated with Alizarin red S (40 mM, pH 7.4; Sigma) for 10 min at room temperature. After washing, images were

captured using a stereo microscope (Stemi 2000-C; Zeiss, Germany), and mineralized surfaces were quantified using the Qwin software (Leica, Germany). This software quantifies the red pixel intensity of the whole well surface and expresses these values in arbitrary units. Two gap junction inhibitors were used: the Cx43 mimetic peptide Gap27 (250  $\mu$ M; GeneCust, France) and the chemical agent 18 $\alpha$ -glycyrrheticin (50  $\mu$ M; Sigma). Medium in presence or absence of GJIC inhibitors was replaced by fresh medium every 24h. Control cultures were treated with the solvent (DMSO, Sigma) only.

### **Western Blot**

Cells were lysed in a lysis buffer (1% SDS, 10mM Tris pH 7.4, 1 mM Sodium orthovanadate) and protein concentration was determined by BCA kit (Sigma). 10  $\mu$ g of total protein extracts in Laemmli buffer (62.5 mM Tris-HCl, pH 6.8, 2% SDS, 10% glycerol, 5% 2-mercaptoethanol, 0.001% bromophenol blue) were separated by SDS-polyacrylamide gel electrophoresis, and transferred to Immobilon-P membranes (Millipore, France). Membranes were immunoblotted with mouse monoclonal anti-connexin43 (dilution 1/1000, Sigma) or mouse monoclonal anti-GAPDH (dilution 1/2000, Abcam, France). Antibody binding was visualized with the enhanced chemiluminescence system (SuperSignal West Pico Chemiluminescent Substrate, ThermoScientific, France). For quantification, luminescence was detected with a Charge Couple Device (CCD) camera and analyzed using the GeneTools program (Syngene, UK).

### **Real-time polymerase chain reaction**

Total RNA was extracted using NucleoSpin<sup>®</sup>RNAII (Macherey Nagel, France). 1  $\mu$ g of total RNA was used for first strand cDNA synthesis using ThermoScript RT-PCR System (Invitrogen, France). DNase I treatment (25 units, 10 min) of total RNA was performed to eliminate genomic contamination. Real-time PCR was performed with a Chromo4 instrument (Biorad, France) using SYBR Green Supermix reagents (Biorad). Primer sequences are provided in Table 1. Calibration curves with different amounts of cDNA were used to validate the primers according the MIQE guidelines (Bustin et al., 2009). Cyc1 (cytochrome c-1) and Hprt 1 (Hypoxanthine guanine phosphoribosyl transferase 1) were used as internal controls to normalize the data to determine the relative expression of the target genes using the  $2^{-\Delta\Delta Ct}$  method.

### **Plasmid constructs, transient cell transfections and reporter assays**

The Cx43 promoter/gene reporter construct -2400Cx43-lux was a gift from GI Fishman, The Leon H. Charney Division of Cardiology, New York University School of Medicine, New York, NY and has been described previously (De Leon et al., 1994). Transient cell transfections were performed with jetPEI™ (Polyplus-transfection, France). The phRLMLP-*Renilla* luciferase expression vector was cotransfected in every experiment to monitor transfection efficiencies. Luciferase activity was determined with the Dual-Luciferase reporter assay system (Promega, France).

### **RNA interference**

Modified hBMSCs cells were obtained by lentiviral cell transduction as previously described (Trichet et al., 2006). Oligonucleotides were designed and cloned into pSUPER to produce the short hairpin RNA shRNACx43 directed against the human Cx43 gene. The targeted sequence was TGCTGCGAACCTACATCAT. The cloned oligonucleotides were controlled by sequencing (Genome Express) before subcloning them with the upstream H1 promoter into the vector pFG12. As a control, a vector pFG12 was developed to produce shRNAs targeting the LacZ gene. The resulting constructs were used for lentivirus production following the protocols provided with the ViraPower Lentiviral Expression System (Invitrogen) to transduce hBMSCs.

### **Parachute assay**

Gap junctional intercellular communication (GJIC) was determined by “parachute” assay, as previously described (Ziambaras et al., 1998, Talbot et al., 2013; Tellez-Gabriel et al., 2017). Cell cultures were divided into “acceptor” and “donor” cell groups. Adherent acceptor cells were loaded with Dil (Sigma; 5  $\mu$ M), a nontransferable membrane dye, for 1h, and confluent donor cells were dyed with calcein-acetoxymethylester (calcein-AM; Sigma; 1  $\mu$ M), a gap junction permeant dye, for 45 min. After loading, cells were washed in PBS to remove the excess dye. Then, the calcein-AM-dyed cells were trypsinized and parachuted on the Dil-dyed acceptor cell layer at a cell ratio of 1:10 and incubated at 37°C for 2h to allow calcein transfer via gap junctions. Dye coupling was analyzed using a FACS FC500 cytometer (Beckman Coulter). For the analysis of results, gates were established on the day of each experiment using non-dyed cells, cells dyed with calcein-AM,



and cells dyed with Dil.

#### **Cell proliferation assay (test XTT)**

Cell viability was determined using a 2,3- bis(2 methoxy-4 nitro-5-sulfohenyl)-2H-tetrazolium-5-carboxanilide (XTT) reagent assay kit (Roche Molecular Biomedicals, Germany). Two thousand cells per well were plated into 96-well plates. At confluence, cells were cultured for 17 days in culture medium in the presence or absence of GJIC inhibitors.

#### **Statistical analysis**

Analyses were performed using GraphPad Prism 6.0 software (GraphPad Software). The unpaired t-test was used for the comparison of two conditions. An ANOVA test was used for the multi-group comparisons of more than two samples. A two-way ANOVA test was used for multi-group comparisons of data that compare two variables. The results are given as mean  $\pm$  SD. Results with  $p < 0.05$  were considered significant.

## Results

### ***Increased production of Cx43 during the differentiation of hBMSCs into osteoblasts in vitro***

We first analyzed the *in vitro* differentiation of hBMSC into osteoblasts under our experimental conditions (see materials and methods). As shown in Figure 1A, incubation of hBMSCs in monolayer cultures with osteogenic medium over 2 weeks resulted in osteoblastic differentiation process as demonstrated by a marked increase in mineral deposition revealed by Alizarin red staining. Concurrently, there was an increase in the expression of genes associated with osteoblast differentiation (Figure 1B). The expression of *Runx2*, the main transcriptional factor involved in the osteoblast commitment and the early differentiation of hBMSCs into osteoblast progenitors (Marie, 2008), was increased quickly after stimulation of hBMSCs with osteogenic medium. Indeed, quantitative real-time PCR demonstrated that *Runx2* mRNA levels increased from day 1 until day 3 and then decreased. Quantitative real-time PCR demonstrated that the levels of transcripts encoding *Alkaline Phosphatase (ALP)*, a gene associated with the differentiation of osteoblast progenitors into immature osteoblasts, increased from day 3 to day 7, and then slowly returned to basal level. The transcript levels of *Bone Sialoprotein (BSP)* and *Osteocalcin (OCN)*, two genes associated with the late differentiation of immature osteoblasts into mature osteoblasts, gradually increased starting at day 13 and day 7, respectively, until the end of the experiment. As expected, the expression of *Sox9*, a transcription factor implicated in chondrocyte differentiation was not modified after stimulation of hBMSCs with osteogenic medium (data not shown).

We secondly analyzed the production of Cx43 during hBMSC differentiation into osteoblasts under our experimental conditions. Interestingly, Western-Blot analysis showed an increase in Cx43 protein level after treatment of hBMSCs with osteogenic medium for 9 days, which remained stable until day 15 (Figure 2A). To determine whether the induction of Cx43 production during hBMSC differentiation occurred via modulation of the corresponding gene, the Cx43 mRNA steady-state levels following treatment with osteogenic medium were measured by real-time PCR. As shown in Figure 2B, induction of Cx43 mRNA was observed in response to osteogenic medium after 9 days of treatment, and Cx43 levels remained elevated until day 17. By contrast,

no change in the expression of Cx45 was observed during this process of hBMSC differentiation (Figure 2B). Although the presence of Cx40 has been demonstrated in hMSC (Valiunas et al., 2004), its expression level remains very low (under the limit of capacity of detection by PCR in our experimental conditions) throughout the differentiation process (data not shown). To further explore whether the effect of osteogenic medium on Cx43 gene expression takes place at the transcriptional level, transient cell transfections were performed with the Cx43 promoter/gene reporter construct -2400Cx43-lux. As shown in Figure 2C, osteogenic medium induced a 6.5-fold transactivation of the Cx43 promoter in hBMSCs. To determine whether the increase in Cx43 gene expression in hBMSCs during osteogenic differentiation is associated with gap junctional intercellular communication changes, GJIC was analyzed by FACS using the parachute assay. GJIC was significantly enhanced (26-fold increase) after 17 days of incubation of hBMSCs with osteogenic medium (Figure 2D).

Together, these results demonstrated an increase in Cx43 production and subsequent cell-cell communication during the differentiation of hBMSCs into osteoblasts.

***Small interfering RNA against Cx43 and treatment with 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide efficiently reduce GJIC between hBMSCs***

To investigate the role of Cx43 and GJIC during the early differentiation of hBMSCs into osteoblast progenitors and throughout the process of hBMSC differentiation into mature osteoblasts, Cx43 was stably knocked down in hBMSCs by transduction with a lentiviral vector or the cells were treated with the GJIC inhibitor 18 $\alpha$ -glycyrrhetic acid or with the peptide Gap27.

As shown in Figure 3A, shCx43 cells were characterized by a significant (70%) reduction in Cx43 mRNA levels compared to shLacZ or parental cells used as control. This decrease was confirmed at the protein level (Figure 3B). The knocked down of Cx43 does not change the Cx45 gene expression in hBMSCs (Figure 3A). In addition, GJIC analysis using the parachute assay demonstrated that GJIC was significantly decreased (by 50%) in shCx43 cells compared to shLacZ cells or parental hBMSCs (Figure 3C). Of note, the Cx43 knock-down does not affected the proliferation rate of the cells (data not shown). As shown in Figure 3D both, 18 $\alpha$ -glycyrrhetic (50  $\mu$ M) acid and Gap27 peptide (250  $\mu$ M) significantly reduce GJIC between hBMSCs by respectively 70% and 60% after 24h incubation of the cells. Of note, in the experimental

conditions used, the viability of cells was not affected in the presence of 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide throughout the experiment (Figure 3E).

Together these results demonstrated that small interfering RNA against Cx43 or treatment with two GJIC inhibitors efficiently reduce GJIC between hBMSCs

### ***Cx43 knocked down blocks the differentiation of hBMSCs into osteoblasts***

We next investigated the effect of the Cx43 knocked down on the osteogenic differentiation of hBMSCs. As shown in Figure 4A, an early decrease in *Runx2* expression was observed in shCx43 cells in response to osteogenic medium compared to that measured in shLacZ cells or in parental cells. Indeed, *Runx2* mRNA steady-state levels were increased by approximately 3-fold in parental and shLacZ cells after treatment of hBMSCs with osteogenic medium for 1 day, but only by 1.7-fold in shCx43 cells. Furthermore, a significant decrease in *ALP*, *BSP* and *OCN* expression levels was observed in shCx43 cells in response to osteogenic medium compared to those measured in shLacZ or in parental cells. *ALP* mRNA steady-state levels were increased by approximately 9-fold in parental and shLacZ cells, but only by 2-fold in shCx43 cells after 7 days of incubation with osteogenic medium (Figure 4A). *OCN* and *BSP* mRNA steady-state levels were increased by approximately 180- and 16-fold, respectively, in parental and shLacZ cells, and only by 120- and 8-fold in shCx43 cells after treatment of hBMSCs with osteogenic medium for 17 days (Figure 4A). Interestingly, after 17 days of treatment with osteogenic medium, a marked decrease (60%) in mineral deposition was measured in shCx43 monolayer cultures compared to that measured in shLacZ or in parental cultures (Figure 4B). Finally, the expression level of *Cx45* was evaluated by quantitative real-time PCR throughout the process of differentiation. As shown in Figure 4C, the expression level of *Cx45* is not modified by the knocked down of Cx43 in hBMSC and remains unchanged throughout the process of differentiation. In contrast and as expected, the expression level of *Cx43* was decreased after hBMSC infection with the shRNA against Cx43 and remains very low throughout the process of differentiation.

Together, these results indicate a crucial role of Cx43 during the early differentiation of hBMSCs into osteoblasts and throughout the process of osteoblast differentiation.

### ***Treatment with GJIC inhibitors inhibits the differentiation of hBMSCs into osteoblasts***

To investigate the role GJIC, and not only the role of Cx43 expression, during the early differentiation of hBMSCs into osteoblast progenitors and throughout the process of hBMSC differentiation into mature osteoblasts, the cells were treated with 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide.

As shown in Figure 5A, a significant decrease in *Runx2* expression level in response to osteogenic medium was observed in the presence of 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide compared to hBMSCs cultured without inhibitor. Treatment of hBMSCs with osteogenic medium for 1 day induced a 3.6-fold increase in *Runx2* mRNA steady-state levels. By contrast, *Runx2* level increased by only 1.4-fold and 1.6-fold in the presence of 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide. A significant decrease in *ALP*, *BSP* and *OCN* expression levels in response to osteogenic medium was also observed in the presence of 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide compared to hBMSC controls (Figure 5A). *ALP* mRNA steady-state levels were increased by 8.6-fold in the absence of a GJIC inhibitor, while treatment with 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide induced only a 2.2-fold and 3.2-fold increase in mRNA levels, respectively, after 7 days of incubation of hBMSCs in osteogenic medium. Treatment of hBMSCs with osteogenic medium for 17 days induced an increase in *OC* and *BSP* mRNA steady-state levels of 137- and 10-fold, respectively. By contrast, *OCN* and *BSP* levels increased by only 30- and 2.3-fold in the presence of 18 $\alpha$ -glycyrrhetic acid, and by 63- and 4.7-fold in the presence of Gap27 peptide. In agreement with these data, a significant decrease in mineral deposition was measured in the presence of 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide compared to cells cultured in the absence of GJIC inhibitors. After 17 days of treatment with osteogenic medium, hBMSCs treated with 18 $\alpha$ -glycyrrhetic acid or Gap27 peptide showed significantly less Alizarin red S staining (95% and 60%, respectively) compared to cells cultured without GJIC inhibitor (Figure 5B).

Together, these results indicate an important role GJIC during the early differentiation of hBMSCs into osteoblasts and throughout the process of osteoblast differentiation.

## Discussion

Previous *in vitro* studies have demonstrated the importance of Cx43 in osteoblast function and during the late stages of osteoblast differentiation specifically during the differentiation of immature osteoblasts into mature osteoblasts able to mineralize (Plotkin and Bellido, 2013; Stains and Civitelli, 2016). *In vivo* experiments using either Cx43<sup>-/-</sup> or Cx43<sup>fl/-</sup>;Col1a.1-2.3 kb-Cre mice have demonstrated that Cx43 play a crucial role in osteoblastic differentiation (Plotkin and Stains, 2015). However, the role of Cx43 in the early differentiation of hBMSCs into osteoblasts *in vitro* remains poorly documented.

In the present study, we first demonstrate that Cx43 expression is increased at the transcriptional level during the late differentiation of hBMSCs into mature osteoblasts. Many transcription factors control osteoblastogenesis. Among them, activator protein-1 (AP-1), a dimeric transcription factor formed by Fos (c-Fos, Fra-1, Fra-2, FosB) and Jun proteins (c-Jun, JunB, JunD) is required for bone formation and remodeling (Eferl and Wagner 2003). For example, most Fos proteins are implicated in the proliferation and differentiation of osteoblasts (Grigoriadis et al., 1993). Interestingly, one AP-1-binding site has been identified in the human Cx43 proximal promoter P1 (approximately 150 nucleotides up- and downstream of the transcription initiation site), suggesting that this AP-1 response element could be involved in the increased expression of Cx43 during hBMSC differentiation into osteoblasts.

Additionally, we demonstrate that Cx43 and subsequent GJIC play an important role throughout the differentiation of hBMSCs into osteoblasts from the early stages to the process of mineralization. It is well recognized that gap junctional communication plays an important role in the control of differentiation into different cell types and tissues (Duval et al., 2002). Specifically, it has been shown that Cx43 and GJIC are crucial in the regulation of self-renewal and maintenance of pluripotency in embryonic stem cells (Todorova et al., 2008). With regard to hBMSC differentiation, osteoblastogenesis may be schematically defined by three phases (Marie, 2008): 1) the osteoblast commitment of hBMSCs and the early differentiation of hBMSCs into osteoblast progenitors, 2) the differentiation of these osteoblast progenitors into immature osteoblasts, and 3) the differentiation of immature osteoblasts into mature osteoblasts able to mineralize. All these three phases are characterized by sequentially expressed genes. The first stage is characterized

by the expression of *Runx2*, the earliest and the most specific transcription factor of mesenchymal stem cell commitment toward osteoblastic lineage (Wagner and Karsenty, 2001). *Runx2* is indeed expressed in all mesenchymal condensations before osteoblast differentiation during early murine skeleton development at E10.5-E12.5 (Ducy et al., 1997). Our *in vitro* results showed that *Runx2* is induced very early, 1 day after stimulation of hBMSCs with osteogenic medium. This early increase in *Runx2* expression precedes the increased expression of ALP (expression peak at day 7), a gene specific to the differentiation of osteoblast progenitors into immature osteoblasts (the second stage of the differentiation process). In this work, we specifically demonstrated that the early increase in *Runx2* expression observed after stimulation of hBMSCs with differentiation medium is inhibited both by the use of an shRNA against Cx43 or by GJIC inhibitors, suggesting that Cx43 as well as GJIC play a role during the commitment of hBMSCs toward osteoblastic lineage or at least during the early stages of hBMSC differentiation into osteoblast progenitors. In addition, using hBMSCs rather than osteoblast progenitors used in previous studies, our results also demonstrated that Cx43 plays a central role during the differentiation of osteoblast progenitors into immature osteoblasts and later during the differentiation of these immature osteoblasts into mature osteoblasts.

Regarding the convergence of Cx43 and *Runx2* expressions, it has been previously demonstrated that the overexpression or knockdown of Cx43 is able to regulate the expression and/or the transcriptional activity of *Runx2* in several cells such as MC3T3-E1, mouse calvaria osteoblastic, human fetal osteoblastic or longitudinal ligament cells (Stains et al., 2003; Li et al., 2006; Thi et al., 2010; Niger et al., 2011; Niger et al., 2012; Yang et al., 2016). The studies on the molecular mechanism underlying this process have identified two effectors, the extracellular signal-regulated kinase (ERK) and the protein kinase C delta (PKC), responsible for Cx43-regulation of *Runx2* expression and/or *Runx2* transcriptional activity (Lima et al., 2009; Niger et al., 2013; Niger et al., 2012; Li et al., 2015; Tu et al., 2016).

Gap junctions can mediate the passage of nucleotides, amino acids, ions and second messengers between coupled cells allowing ionic and metabolic signaling communication (Harris, 2007). Among these messengers able to cross through gap junctional channels, there is evidence that the spatial and temporal patterns of calcium are crucial in the regulation of the differentiation process. Both the stable increase in  $\text{Ca}^{2+}$  intracellular concentration ( $[\text{Ca}^{2+}]$ ), and the oscillations

of  $[Ca^{2+}]$  are a mode of signaling in both excitable and non-excitable cells. In bone marrow-derived stem cells, spontaneous  $[Ca^{2+}]$  oscillations were shown to depend on Inositol P3-mediated  $Ca^{2+}$  release followed by  $Ca^{2+}$  influx from the extracellular compartment (Kawano et al., 2002). In this context, it has been shown that  $[Ca^{2+}]$  oscillations can be regulated by intercellular coupling through gap junctions (Dupont et al., 2007). We can thus hypothesize that Cx43-mediated gap junctional intercellular communication drives the differentiation of hBMSCs into osteoblasts via its ability to propagate  $[Ca^{2+}]$  oscillations. However, future studies are required to establish the nature of the signals,  $Ca^{2+}$  or others, which drive the hBMSC differentiation process.



## Legends of Figures

### **Figure 1:** *In vitro* differentiation of hBMSCs toward osteoblasts

hBMSCs were cultured in the presence or absence of osteogenic medium up to 17 days. (A) At each time point, cultures were fixed and stained with Alizarin red S. Results show one representative experiment of three independent experiments. (B) *Runx2*, *Alkaline Phosphatase* (ALP), *Bone Sialoprotein* (BSP) and *Osteocalcin* (OCN) mRNA steady-state levels were determined by quantitative RT-PCR, and are expressed relative to the control culture in absence of osteogenic medium. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate.

### **Figure 2:** *Increase of Cx43 production during the differentiation of hBMSCs into osteoblasts.*

hBMSCs were cultured in the presence or absence of osteogenic medium (Diff) up to 17 days. (A) Cx43 production was detected by Western Blot analysis. Result shown is representative of three independent experiments. (B) *Cx43* and *Cx45* mRNA steady-state levels were determined by quantitative RT-PCR, and expressed relative to the control culture in absence of osteogenic medium. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate. (C) hBMSC cells were incubated in the presence or absence of osteogenic medium for 9 days. After incubation, cell cultures were transfected with the -2400Cx43-lux construct, and incubation continued for another 48h. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate. (\*\*p<0.01) (D) FACS analysis of calcein transfer in hBMSCs treated or not with osteogenic medium during 17 days. Histogram plots (left panel) represent the mean of fluorescence recorded in FL1 channel, which corresponds to calcein-positive cells. Control peak corresponds to cells dyed with Dil only, and parachute peak to cells double-labeled. Histograms (right panel) indicate mean of the relative transfer-ratio of calcein between cells, which represents a direct measure of the degree of dye coupling. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate. (\*\*p<0.01)

### **Figure 3:** *Small interfering RNA against Cx43 and GJIC inhibitors efficiency block GJIC in hBMSCs*

(A) *Cx43* and *Cx45* mRNA steady-state levels were determined by quantitative RT-PCR in parental (P), shLacZ and shC*x43* cells as indicated. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate. (\*\* $p < 0.05$ ) (B) *Cx43* production was detected by Western Blot analysis in parental, shLacZ and shC*x43* cells as indicated. Result shown is representative of three independent experiments. (C) FACS analysis of calcein transfer in parental, shLacZ and shC*x43* hBMSCs treated or not with osteogenic medium (Diff) for 17 days. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate. (\*\*\* $p < 0.01$ ). (D) FACS analysis of calcein transfer in hBMSCs treated or not with 18 $\alpha$ -glycyrrhetic acid (50  $\mu$ M) or Gap27 peptide (250  $\mu$ M) for 24h. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate. (\*\*\* $p < 0.01$ ). (E) hBMSCs viability, treated or not with 18 $\alpha$ -glycyrrhetic acid (50  $\mu$ M) or Gap27 peptide (250  $\mu$ M), was determined using a 2,3- bis(2 methoxy-4 nitro-5-sulfophenyl)-2H-tetrazolium-5-carboxanilide (XTT) reagent assay kit. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate.

**Figure 4:** *Small interfering RNA against Cx43 inhibits the differentiation of hBMSCs into osteoblasts.*

(A) Parental, shLacZ and shC*x43* cells were cultured in the presence or absence of osteogenic medium. The expression of *Runx2* was determined by quantitative RT-PCR after incubation of hBMSCs with osteogenic medium for 1 day. The expression of *ALP* was determined by quantitative RT-PCR after incubation of hBMSCs with osteogenic medium for 7 days. The expression of *BSP* and *OCN* mRNA steady-state levels were measured by quantitative RT-PCR after incubation of hBMSCs with osteogenic medium for 17 day. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate. (\*\* $p < 0.05$ , \*\*\* $p < 0.01$ ). (B) Parental, shLacZ and shC*x43* hBMSCs were cultured in the presence or absence of osteogenic medium. 17 days after induction of osteogenesis, cultures were fixed and stained with Alizarin red S. Results show one representative experiment of three independent experiments. (C) Parental, shLacZ and shC*x43* cells were cultured in the presence or absence of osteogenic medium. The expression of *Cx43* and *Cx45* were determined by quantitative RT-PCR throughout the differentiation process. Results indicate mean  $\pm$  SD of three independent experiments performed in duplicate.

**Figure 5:** *GJIC inhibitors reduce the differentiation of hBMSCs into osteoblasts.*

(A) hBMSCs were cultured in the presence of GJIC inhibitors ( $\alpha$ -glycyrrhetic acid or Gap27) as indicated. The expression of *Runx2* was determined by quantitative RT-PCR after treatment with osteogenic medium for 1 day. The expression of *ALP* was determined by quantitative RT-PCR after treatment with osteogenic medium for 7 days. 17 days after treatment of the cells with osteogenic medium (Diff), *BSP* and *OCN* mRNA steady-state levels were determined by quantitative RT-PCR. Bars indicate mean  $\pm$  SD of three independent experiments performed in duplicate. (\*\* $p < 0.05$ ; \*\*\* $p < 0.01$ ). (B) Cultures were fixed and stained with Alizarin red S 17 days after induction of osteogenesis in the presence or absence of 18 $\alpha$ -glycyrrhetic acid (50  $\mu$ M) or Gap27 (250  $\mu$ M). Results show one representative experiment of three independent experiments.

**Acknowledgments** : Jerome Amiaud for its technical help

## Literature cited

- Augello A, De Bari C. 2010. The regulation of differentiation in mesenchymal stem cells. *Hum Gene Ther* 21:1226-1238.
- Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. 2009. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 55:611-622.
- Chang Y, Hsieh P-H, Chao CC-K. 2009. The efficiency of Percoll and Ficoll density gradient media in the isolation of marrow derived human mesenchymal stem cells with osteogenic potential. *Chang Gung Med J* 32:264-275.
- De Leon JR, Buttrick PM, Fishman GI. 1994. Functional analysis of the connexin43 gene promoter in vivo and in vitro. *J Mol Cell Cardiol* 26:379-389.
- Ducy P, Zhang R, Geoffroy V, Ridall AL, Karsenty G. 1997. *Osf2/Cbfa1*: a transcriptional activator of osteoblast differentiation. *Cell* 89:747-54.
- Dupont G, Combettes L, Leybaert L. 2007. Calcium dynamics: spatio-temporal organization from the subcellular to the organ level. *Int Rev Cytol* 261:193-245.
- Duval N, Gomès D, Calaora V, Calabrese A, Meda P, Bruzzone R. 2002. Cell coupling and Cx43 expression in embryonic mouse neural progenitor cells. *J Cell Sci* 115:3241-3251.
- Eferl R, Wagner EF. 2003. AP-1: a double-edged sword in tumorigenesis. *Nat Rev Cancer* 3:859-68.
- Friedenstein AJ, Chailakhyan RK, Gerasimov UV. 1987. Bone marrow osteogenic stem cells: in vitro cultivation and transplantation in diffusion chambers. *Cell Tissue Kinet* 20:263-272.
- Grigoriadis AE, Schellander K, Wang ZQ, Wagner EF. 1993. Osteoblasts are target cells for transformation in c-fos transgenic mice. *J Cell Biol* 122:685-701.
- Harris AL. 2007. Connexin channel permeability to cytoplasmic molecules. *Prog Biophys Mol Biol* 94:120-143.
- Kawano S, Shoji S, Ichinose S, Yamagata K, Tagami M, Hiraoka M. 2002. Characterization of Ca(2+) signaling pathways in human mesenchymal stem cells. *Cell calcium* 32:165-174.
- Laird DW. 2014. Syndromic and non-syndromic disease-linked Cx43 mutations. *FEBS Lett* 588:1339-48.
- Lecanda F, Towler DA, Ziambaras K, Cheng SL, Koval M, Steinberg TH, Civitelli R. 1998. Gap junctional communication modulates gene expression in osteoblastic cells. *Mol Biol Cell* 9:2249-2258.
- Lecanda F, Warlow PM, Sheikh S, Furlan F, Steinberg TH, Civitelli R. 2000. Connexin43 deficiency causes delayed ossification, craniofacial abnormalities, and osteoblast dysfunction. *J Cell Biol* 151:931-944.
- Li Z, Zhou Z, Saunders MM, Donahue HJ. 2006. Modulation of connexin43 alters expression of osteoblastic differentiation markers. *Am J Physiol Cell Physiol* 290:C1248-55.
- Li S, Zhang H, Li S, Yang Y, Huo B, Zhang D. 2015. Connexin 43 and ERK regulate tension-induced signal transduction in human periodontal ligament fibroblasts. *J Orthop Res* 33:1008-14.
- Lima F, Niger C, Hebert C, Stains JP. 2009. Connexin43 potentiates osteoblast responsiveness to fibroblast growth factor 2 via a protein kinase C-delta/Runx2-dependent mechanism. *Mol Biol*

Cell 20:2697-708.

- Marie PJ. 2008. Transcription factors controlling osteoblastogenesis. *Arch Biochem Biophys* 473: 98-105.
- Neve A, Corrado A, Cantatore FP. 2011. Osteoblast physiology in normal and pathological conditions. *Cell Tissue Res* 343:289-302.
- Nielsen MS, Axelsen LN, Sorgen PL, Verma V, Delmar M, Holstein-Rathlou NH. 2012. Gap junctions. *Compr Physiol* 2:1981-2035.
- Niger C, Lima F, Yoo DJ, Gupta RR, Buo AM, Hebert C, Stains JP. 2011. The transcriptional activity of osterix requires the recruitment of Sp1 to the osteocalcin proximal promoter. *Bone* 49:683-692.
- Niger C, Buo AM, Hebert C, Duggan BT, Williams MS, Stains JP. 2012. ERK Acts in Parallel to PKC delta to Mediate the Connexin43-dependent Potentiation of Runx2 Activity by FGF2 in MC3T3 Osteoblasts. *Am J Physiol Cell Physiol* 302:C1035-44.
- Niger C, Luciotti MA, Buo AM, Hebert C, Ma V, Stains JP. 2013. The regulation of runt-related transcription factor 2 by fibroblast growth factor-2 and connexin43 requires the inositol polyphosphate/protein kinase C $\delta$  cascade. *J Bone Miner Res* 28:1468-77.
- Owen M. 1988. Marrow stromal stem cells. *J Cell Sci Suppl* 10:63-76.
- Paznekas WA, Karczeski B, Vermeer S, Lowry RB, Delatycki M, Laurence F, Koivisto PA, Van Maldergem L, Boyadjiev SA, Bodurtha JN, Jabs EW. 2009. GJA1 mutations, variants, and connexin 43 dysfunction as it relates to the oculodentodigital dysplasia phenotype. *Hum Mutat* 30:724-33.
- Plotkin LI, Bellido T. 2013. Beyond gap junctions: Connexin43 and bone cell signaling. *Bone* 52:157-66.
- Plotkin LI, Stains JP. 2015. Connexins and pannexins in the skeleton: gap junctions, hemichannels and more. *Cell Mol Life Sci* 72:2853-67.
- Schiller PC, Mehta PP, Roos BA, Howard GA. 1992. Hormonal regulation of intercellular communication: parathyroid hormone increases connexin 43 gene expression and gap-junctional communication in osteoblastic cells. *Mol Endocrinol* 6:1433-40.
- Schiller PC, D'Ippolito G, Balkan W, Roos BA, Howard GA. 2001a. Gap-junctional communication is required for the maturation process of osteoblastic cells in culture. *Bone* 28:362-9.
- Schiller PC, D'Ippolito G, Balkan W, Roos BA, Howard GA. 2001b. Gap-junctional communication mediates parathyroid hormone stimulation of mineralization in osteoblastic cultures. *Bone* 28:38-44.
- Söhl G, Willecke K. 2004. Gap junctions and the connexin protein family. *Cardiovasc Res* 62:228-32.
- Sosinsky GE, Nicholson BJ. 2005. Structural organization of gap junction channels. *Biochim Biophys Acta* 1711:99-125.
- Stains JP, Civitelli R. 2016. Connexins in the skeleton. *Semin Cell Dev Biol* 50:31-9.
- Stains JP, Lecanda F, Screen J, Towler DA, Civitelli R. 2003. Gap junctional communication modulates gene transcription by altering the recruitment of Sp1 and Sp3 to connexin-response elements in osteoblast promoters. *J Biol Chem* 278:24377-24387.

- Talbot J, Brion R, Picarda G, Amiaud J, Chesneau J, Bougras G, Stresing V, Tirode F, Heymann D, Redini F, Verrecchia F. 2013. Loss of connexin43 expression in Ewing's sarcoma cells favors the development of the primary tumor and the associated bone osteolysis. *Biochim Biophys Acta* 1832:553-564.
- Tellez-Gabriel M, Charrier C, Brounais-Le Royer B, Mullard M, Brown HK, Verrecchia F, Heymann D. 2017. Analysis of gap junctional intercellular communications using a dielectrophoresis-based microchip. *Eur J Cell Biol* 96:110-118.
- Thi MM, Urban-Maldonado M, Spray DC, Suadicani SO. 2010. Characterization of hTERT-immortalized osteoblast cell lines generated from wild-type and connexin43-null mouse calvaria. *Am J Physiol Cell Physiol* 299:C994-C1006.
- Todorova MG, Soria B, Quesada I. 2008. Gap junctional intercellular communication is required to maintain embryonic stem cells in a non-differentiated and proliferative state. *J Cell Physiol* 214:354-362.
- Trichet V, Benezech C, Dousset C, Gesnel MC, Bonneville M, Breathnach R. 2006. Complex interplay of activating and inhibitory signals received by Vgamma9Vdelta2 T cells revealed by target cell beta2-microglobulin knockdown. *J Immunol* 177:6129-6136.
- Tu B, Liu S, Liu G, Li Z, Sun Y, Fan C. 2016. Inhibition of connexin 43 prevents trauma-induced heterotopic ossification. *Sci Rep* 16;6:37184.
- Väänänen HK. 2005. Mesenchymal stem cells. *Ann Med* 37:469-479.
- Valiunas V, Doronin S, Valiuniene L, Potapova I, Zuckerman J, Walcott B, Robinson RB, Rosen MR, Brink PR, Cohen IS. 2004. Human mesenchymal stem cells make cardiac connexins and form functional gap junctions. *J Physiol (Lond)* 555:617-626.
- Wagner EF, Karsenty G. 2001. Genetic control of skeletal development. *Curr Opin Genet Dev* 11:527-32.
- Wennberg C, Hessle L, Lundberg P, Mauro S, Narisawa S, Lerner UH, Millán JL. 2000. Functional characterization of osteoblasts and osteoclasts from alkaline phosphatase knockout mice. *J Bone Miner Res* 15:1879-1888.
- Yang H, Shi L, Shi G, Guo Y, Chen D, Chen D, Shi J. 2016. Connexin 43 Affects Osteogenic Differentiation of the Posterior Longitudinal Ligament Cells via Regulation of ERK Activity by Stabilizing Runx2 in Ossification. *Cell Physiol Biochem* 38:237-47.
- Ziambaras K, Lecanda F, Steinberg TH, Civitelli R. 1998. Cyclic stretch enhances gap junctional communication between osteoblastic cells. *J Bone Miner Res* 13:218-228.




Figure 1






Figure 2




Figure 3




Figure 4





Figure 5

**Table 1.** Primer sequence for quantitative RT-PCR

| Gene   | Gene (full name)                                     | Accession number | Forward primer        | Reverse primer          |
|--------|--|------------------|-----------------------|-------------------------|
| Cx43   | Connexin43   | NM_000165.3      | gcctgaactgcctttcat    | ctccagtcacccatgttgc     |
| OC     | Osteocalcin  | NM_199173.2      | ggcgctacctgtatcaatgg  | tcagccaactcgtcacagtc    |
| BSP    | Bone sialoprotein                                    | NM_004967        | cgaatacacggggtcaatg   | gtagctgtactcatctcataggc |
| ALP    | Alkaline phosphatase                                 | NM_000478.3      | aacaccaccagggaac      | ggcacaatgccacagatt      |
| Runx2  | Runt-related transcription factor 2                  | NM_004348.3      | gcctaggcgattcaga      | gctcttactgagagtgaagg    |
| SOX9   | SRY (sex determining region Y)-<br>box9              | NM_000346.3      | gtaccgcactgcacaac     | tcgctcgttcagaagtctc     |
| Cx45   | Connexin 45  | NM_005497.3      | tgggtaaccgaagtctggaca | gaacagtgagccagatcttccc  |
| Hprt 1 | Hypoxanthine guanine<br>phosphoribosyl transferase 1 | NM_000194.1      | tgacctgattatgtcatacc  | cgagcaagacgttcagctct    |
| Cyc1   | Cytochrome c-1                                       | NM_001916        | gcatgggtgaggactacg    | ggccaggaaagtaggggttg    |