



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/116046/>

Version: Accepted Version

Article:

Ostergaard, M, Eshed, I, Althoff, C et al. (2017) Whole-body MRI in inflammatory arthritis - Systematic literature review and first steps towards standardization and an OMERACT scoring system. *Journal of Rheumatology*, 44 (11). pp. 1699-1705. ISSN: 0315-162X

<https://doi.org/10.3899/jrheum.161114>

© 2017 The Journal of Rheumatology. This is a pre-copy-editing, author-produced PDF of an article accepted for publication in *The Journal of Rheumatology* following peer review. The definitive publisher-authenticated version, Mikkel Østergaard, Iris Eshed, Christian E. Althoff, Rene P. Poggenborg, Torsten Diekhoff, Simon Krabbe, Sabine Weckbach, Robert G.W. Lambert, Susanne J. Pedersen, Walter P. Maksymowych, Charles G. Peterfy, Jane Freeston, Paul Bird, Philip G. Conaghan and Kay-Geert A. Hermann *The Journal of Rheumatology* June 2017, jrheum.161114; DOI: <https://doi.org/10.3899/jrheum.161114>, is available online at: <https://doi.org/10.3899/jrheum.161114>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Whole-body MRI in inflammatory arthritis - Systematic literature review and first steps towards standardization and an OMERACT scoring system

Journal:	<i>The Journal of Rheumatology</i>
Manuscript ID	2016-1114.R1
Manuscript Type:	OMERACT
Date Submitted by the Author:	n/a
Complete List of Authors:	Ostergaard, Mikkel; Rigshospitalet, Copenhagen Center for Arthritis Research, Center for Rheumatology and Spine Diseases; University of Copenhagen, Department of Clinical Medicine Eshed, Iris; Sheba Medical Center, Diagnostic Imaging Althoff, Christian; Charité, Campus Mitte, Department of Radiology Poggenborg, René; Rigshospitalet Glostrup, Copenhagen Center for Arthritis Research, Center for Rheumatology and Spine Diseases Diekhoff, Torsten; Charité Medical School, Department of Radiology Krabbe, Simon; Rigshospitalet Glostrup, Copenhagen Center for Arthritis Research, Center for Rheumatology and Spine Diseases Weckbach, Sabine Lambert, Robert; University of Alberta, Radiology & Diagnostic Imaging Pedersen, Susanne; Rigshospitalet Glostrup, Copenhagen Center for Arthritis Research, Center for Rheumatology and Spine Diseases Maksymowych, Walter; University of Alberta, Medicine Peterfy, Charles; Spire Sciences inc Freeston, Jane; Chapel Allerton Hospital, Academic Unit of Musculoskeletal Disease Bird, Paul; Combined Rheumatology Practice, Conaghan, Philip; University of Leeds, Leeds Institute of Rheumatic and Musculoskeletal Medicine Hermann, Kay; Charité Medical School, Dept. of Radiology
Keywords:	Magnetic resonance imaging, Arthritis, OMERACT, Synovitis, enthesitis

Whole-body MRI in inflammatory arthritis – Systematic literature review and first steps towards standardization and an OMERACT scoring system

Mikkel Østergaard, Iris Eshed, Christian Althoff, Rene P Poggenborg, Torsten Diekhoff, Simon Krabbe, Sabine Weckbach, Robert Lambert, Susanne J Pedersen, Walter P Maksymowych, Charles G Peterfy, Jane Freeston, Paul Bird, Philip G Conaghan, Kay-Geert A. Hermann

Key indexing terms

MRI, OMERACT, arthritis, synovitis, enthesitis,

Initials, surnames, appointments, academic degrees and e-mails of all authors, and njejeame of Departments and Institutions to which the work should be attributed

M Østergaard, MD, PhD, DMSc, Professor, Copenhagen Center for Arthritis Research (COPECARE), Center for Rheumatology and Spine Diseases, Rigshospitalet, Glostrup, Denmark, and the Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark.
mo@dadlnet.dk

I Eshed, MD, Associate professor, Department of Diagnostic imaging, The Sheba Medical Center, affiliated with the Sackler School of Medicine, Tel Aviv University, Israel
iriseshed@gmail.com

C Althoff, MD, Senior Consultant, Arthritis Imaging Research Group, Dept. of Radiology, Charité Medical School, Berlin, Germany
Christian.Alothoff@charite.de

RP Poggenborg, MD, PhD, Copenhagen Center for Arthritis Research (COPECARE), Center for Rheumatology and Spine Diseases, Rigshospitalet, Glostrup, Denmark,
poggenborg@dadlnet.dk

T Diekhoff, MD, Consultant, Arthritis Imaging Research Group, Dept. of Radiology, Charité Medical School, Berlin, Germany
torsten.diekhoff@gmail.com

S Krabbe, MD, PhD, Copenhagen Center for Arthritis Research (COPECARE), Center for Rheumatology and Spine Diseases, Rigshospitalet, Glostrup, Denmark.
simonkrabbe@gmail.com

S Weckbach, MD, Professor of Radiology, Diagnostic and Interventional Radiology, University Hospital Heidelberg, Heidelberg, Germany
Sabine.Weckbach@med.uni-heidelberg.de

R Lambert, MB BCh, FRCR, FRCPC, Professor, Department of Radiology & Diagnostic Imaging, University of Alberta, Edmonton, Alberta, Canada.
rlambert@ualberta.ca

SJ Pedersen, MD, PhD, Copenhagen Center for Arthritis Research (COPECARE), Center for Rheumatology and Spine Diseases, Rigshospitalet, Glostrup, Denmark.
susanne_juhl_ped@dadlnet.dk

WP Maksymowych, MB ChB, FRCPC(C), Professor, FACP Division of Rheumatology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Canada.
walter.maksymowych@ualberta.ca

CG Peterfy, MD, PhD, FRCPC, Chief Executive Officer, Spire Sciences inc., Boca Raton, FL, USA
Charles.Peterfy@spiresciences.com

J Freeston, MD, PhD, Consultant Rheumatologist and Honorary Clinical Associate Professor, St James' University and Chapel Allerton Hospitals, Leeds
J.E.Freeston@leeds.ac.uk

Paul Bird, BMed (Hons), FRACP, PhD, Grad Dip MRI, Associate Professor, University of NSW, Sydney, Australia
PBird@optimusresearch.com.au

Philip G Conaghan, MB, BS, PhD, FRACP, FRCP, Professor of Musculoskeletal Medicine, Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, & NIHR Leeds Musculoskeletal Biomedical Research Unit, UK
p.conaghan@leeds.ac.uk

Kay-Geert Hermann, MD, PhD, Senior Consultant, Arthritis Imaging Research Group, Dept. of Radiology, Charité Medical School, Berlin, Germany
kghermann@gmail.com

Sources of support in the form of grants or industrial support

None.

Address for correspondence

Mikkel Østergaard,

COPECARE, Center for Rheumatology and Spine Diseases, Rigshospitalet,

Nordre Ringvej 57, DK-2600 Glostrup, Denmark

Tel: +45-38633014 (secretary) or +45-21603865 (mobile), Fax: +45-38634192

E-mail: mo@dadlnet.dk

Running footline

Whole-body MRI in arthritis

ABSTRACT

Aim: Whole-body MRI (WBMRI) is a relatively new technique which potentially enables assessment of the overall inflammatory status of people with arthritis, but standards for image acquisition, definitions of key pathologies and a quantification system are required. Our aim was to perform a systematic literature review (SLR) and to develop consensus definitions of key pathologies, anatomical locations for assessment, a set of MRI sequences and imaging planes for the different body regions and a preliminary scoring system for WBMRI in inflammatory arthritis.

Methods: A SLR was initially performed, searching for whole-body MRI studies in arthritis/osteoarthritis/spondyloarthritis/enthesitis. These results were presented to a meeting of the MRI in Arthritis Working Group together with an MR-image review. Following this, preliminary standards for WBMRI in inflammatory arthritides were developed with further iteration at the Working Group meetings at OMERACT 13.

Results: The SLR identified 10 relevant original articles (7 cross-sectional/3 longitudinal, mostly focusing on synovitis and/or enthesitis in spondyloarthritis, 4 with reproducibility data). The Working Group decided on inflammation in peripheral joints and entheses as primary focus area, and then developed consensus MRI definitions for these pathologies, selected anatomical locations for assessment, agreed on a core set of MRI sequences and imaging planes for the different regions, and proposed a preliminary scoring system. It was decided to test and further develop the system by iterative multi-reader exercises.

Conclusion: These first steps in developing an OMERACT WBMRI scoring system for use in inflammatory arthritides offer a framework for further testing and refinement.

INTRODUCTION

Magnetic resonance imaging (MRI) is now frequently used as an outcome measure in rheumatology clinical trials. By objectively assessing both disease activity and structural damage MRI has provided new insights into disease pathogenesis and treatment response. The Outcome Measures in Rheumatology (OMERACT) MRI Working Group has been instrumental in advancing the utility of MRI in clinical research, and the OMERACT rheumatoid arthritis (RA) MRI score (RAMRIS)(1-6) for evaluating bone erosion, osteitis (bone marrow edema) and synovitis in RA, is now the standard method used in clinical trials. Further, supplementary RAMRIS joint space narrowing and tenosynovitis scores have recently been developed and validated(7-10). The group has also developed and validated a psoriatic arthritis (PsA) MRI scoring method (PsAMRIS)(11-13), and the Assessment of SpondyloArthritis International Society (ASAS)/OMERACT working group has validated scoring methods for assessing inflammation in sacroiliac joints(14) and spine(15) in patients with ankylosing spondylitis (AS).

A disadvantage of conventional MRI is the limited anatomical area that is assessed in a typical examination. Whole-body MRI (WBMRI) is a relatively new technique currently used as a screening tool for evaluating multifocal bone lesions in diseases such as multiple myeloma. WBMRI allows assessment of the entire body in one examination in less than an hour, and thereby can potentially provide a global assessment of the inflammatory status of an arthritis patient(16-25). This may improve the utility of MRI in AS, RA, and particularly PsA, which presents with varying patterns of arthritis, enthesitis, spondylitis and/or dactylitis. However, standards for image acquisition and definitions of key pathologies need to be established, and a system for quantification needs to be developed and validated.

In order to develop this tool according to OMERACT Filter 2 as an applicable measurement instrument for the relevant pathophysiological domain of inflammation (26), we performed a systematic literature review to establish the current status of WBMRI in imaging of peripheral joints. Thereafter based on published data and review of MR-images, we decided on consensus definitions of key pathologies, anatomical locations for assessment, a set of MRI sequences and imaging planes for the different regions of the body and a preliminary scoring system for WBMRI in inflammatory arthritis. This new work, which has not been published before, provides the first international consensus report on WBMRI and a useful novel framework for further development of WBMRI as an outcome measure in inflammatory arthritides.

METHODS.

Literature review

A systematic literature review was undertaken. The population of interest was patients with arthritis and/or enthesitis, the intervention whole-body MRI, a control group was not mandatory, the outcomes included lesions observed and intra/interreader agreement, and the study design should either be cross-

sectional or longitudinal original studies (PICOS). A literature search was done by one author (KGH) on January 19, 2016, using Medline searching for “whole body MRI” AND (“arthritis” OR “osteoarthritis” OR “spondyloarthritis” OR “enthesitis”)” yielded 43 results. One article in press and not yet indexed in Pubmed was added. A flow diagram, made in accordance with the PRISMA guidelines (www.equator-network.org/reporting-guidelines/prisma/), is provided in Figure 1. After manually excluding non-english articles, review papers and case reports, 22 articles remained. Manual exclusion of papers not truly applying the WBMRI technique left 18 articles. Finally, 8 articles focusing only on the axial skeleton were excluded(27-34), leaving 10 articles which used WBMRI for assessment of the extremities (Table 1)(16-25). The QUADAS-2 tool was used to assess risk of bias and concerns for applicability (Supplementary Figure 1)(www.quadas.org).

Consensus process

Members of the OMERACT MRI in Arthritis Working Group and other researchers who had previously worked with WBMRI participated in a one-day meeting in Berlin, January 21, 2016. At this meeting, the literature search was presented and discussed, followed by presentations by all groups who had previously published WBMRI data from patients with inflammatory arthritides. This was followed by a discussion of challenges in developing and applying WBMRI as an outcome measure, and of the following predefined topics: initial focus area, selection of key pathologies to assess, MRI definitions of key pathologies, selection of anatomical locations for assessment, core MRI sequences and imaging planes, and development of a preliminary assessment system. These issues were further discussed and refined during email communications and meetings at the OMERACT conference in Whistler, Canada, May 2016.

RESULTS

Literature review

Characteristics of the 10 publications that used the WBMRI technique to assess the extremities, with or without additional examination of the axial skeleton are provided in Table 1(16-25). The manuscripts described 7 cross-sectional and 3 longitudinal studies, mainly in SpA/PsA. They reported WBMRI-visualization of peripheral synovitis, effusion, osteitis, enthesitis and, to a limited extent, bone erosions. Four studies included Reproducibility data (Table 1).

Initial focus area

It was decided initially to focus on assessment of inflammation, as opposed to damage, in the extremities. Inflammation was chosen because total inflammatory load was considered to be clinically most important, and because the requirements for spatial resolution were believed to be less challenging than those for assessing bone erosion. Much standardization has been done in axial

SpA/AS(14;15;35-37), and the extremities were prioritized, because consensus scoring systems exist only for a limited number of regions (mainly hand and wrist joints).

Selection of key pathologies

It was decided to focus on joints and entheses and that these should both be assessed separately for inflammation in the soft tissues and inflammation in the bone (Appendix).

Definition of key pathologies

With previously published OMERACT MRI definitions used as a starting point, definitions of the joint and enthesal pathologies were developed(Appendix), taking into account the MRI sequences available.

Selection of anatomical locations for assessment

It was agreed that no peripheral joints should be excluded before reader exercises, with exception of the elbow joints, which based on the experience of the group are always located outside the field of view, because of peripheral positioning in the MRI-bore.

Since there is a large number of entheses in the body, a choice of which entheses to examine had to be made, for feasibility. The selection was based on existing clinical enthesitis indices and the ability of MRI to visualize those specific anatomical locations. A questionnaire was circulated to all participating groups, and from this a preliminary set of entheses to be assessed in reading exercises were chosen by consensus (Appendix). The most informative imaging plane for the specific region was also taken into account, since generally only one plane could be selected per region for feasibility reasons.

MRI sequences and planes

For evaluating inflammation, it was considered crucial to have either a short tau inversion recovery (STIR) or T2-weighted fat-suppressed (T2FS) sequences, and/or a fat suppressed T1-weighted sequence after intravenous gadolinium-contrast injection (T1-postGd). A T1-weighted sequence before contrast injection (T1-preGd) was not considered mandatory for assessing inflammation, but due to its high anatomical resolution, availability of a T1-preGd facilitates exact anatomical localization of imaging pathologies. If structural damage is to be assessed, it is crucial to include T1-pre-Gd.

The recommended imaging plane depends on the anatomical region (Appendix). The planes were selected with the aim of optimally presenting the most common and important pathologies in the individual regions.

Assessment (scoring) system

It was decided that all assessed pathologies in all selected joints and entheses would be scored 0-2, as follows: 0: no inflammation, 1: mild/moderate inflammation, 2: severe inflammation.

Total scores would be calculated for each of the following: a. Joints – synovitis; b. joints – osteitis; c. entheses – soft tissue inflammation; d. entheses – osteitis. It was also decided to calculate composite scores (joint inflammation index: a+b, enthesal inflammation index: c+d, and total peripheral inflammation index: a+b+c+d).

DISCUSSION

This report describes the first international consensus effort regarding the use of WBMRI in different arthritides. A literature review, an MR-image review, and discussion among physicians experienced in WBMRI and/or developing MRI scoring systems led to consensus on important pathologies and locations for assessment, MRI-definitions of these, core MRI sequences and imaging planes, and a preliminary scoring system. Future data are likely to modify these decisions, and the preliminary nature of the decisions was fully acknowledged by the group. Nevertheless, the group felt that the present work formed a useful framework for further development of WBMRI as an outcome measure in inflammatory arthritides.

Joints and entheses were selected as the key pathologies for scoring. Pericapsular inflammation was discussed as a relevant manifestation in some joints, particularly in SpA, including PsA. However, the group felt that this disease feature was not suitable for follow-up in clinical trials, and not assessable without contrast injection and higher WBMRI-image quality than currently available. The group therefore decided to exclude it. Similarly, bursitis, tenosynovitis, tendonitis, and dactylitis were excluded, although some of these may be considered in the future, depending on technical developments and new information about the individual importance of such pathologies in the different diseases. It was also decided not to distinguish large joints from small joints, until more information about this becomes available and data-driven conclusions can be made.

The use of WBMRI is currently challenged by the examination time, which limits the image quality and spatial resolution attainable to significantly less than what can be achieved with conventional, single-location MRI. It also limits the number of imaging planes and pulse sequences. However, imaging speed is constantly improving(22), and probably this limitation will be less significant in the future.

The proposed assessment system is not designed for one specific disease, but rather is meant to be tested in iterative exercises in different inflammatory arthritides. After such exercises, separate scoring systems specifically designed for individual diseases, such as RA or PsA, may be developed.

The planned next step of the group will be an initial multireader exercise to test the feasibility and reproducibility of the assessment system, followed by data-driven modifications and improvements.

A strength of the present initiative was that the most groups identified by the literature search as experienced in WBMRI in arthritis were represented in the consensus discussions. Limitations include that only one investigator searched one database, and extracted data. Searching in more databases and a broader search strategy could have diminished the risk of missed articles.

In conclusion, WBMRI offers significant potential as a measure of the total inflammatory burden in patients with arthritides. The present manuscript describes the first steps in developing an OMERACT WBMRI scoring system, and provides a useful framework for the further development of WBMRI as an outcome measure in inflammatory arthritides.

For Peer Review

Text boxes:

Text box 1: New work presented in this manuscript:

- A systematic literature review of WBMRI examination in arthritides
- Selection of key pathologies for assessment by WBMRI in arthritides
- Development of preliminary definitions of the key pathologies
- Selection of anatomical locations for assessment
- Selection of recommended MRI sequences and planes for WBMRI
- Development of a preliminary assessment (scoring) system for WBMRI

For Peer Review

References

Reference List

- (1) Østergaard M, Peterfy C, Conaghan P, McQueen F, Bird P, Ejbjerg B, et al. OMERACT Rheumatoid Arthritis Magnetic Resonance Imaging Studies. Core set of MRI acquisitions, joint pathology definitions, and the OMERACT RA-MRI scoring system. *J Rheumatol* 2003;30:1385-6.
- (2) Østergaard M, Edmonds J, McQueen F, Peterfy C, Lassere M, Ejbjerg B, et al. An introduction to the EULAR-OMERACT rheumatoid arthritis MRI reference image atlas. *Ann Rheum Dis* 2005;64 Suppl 1:i3-i7.
- (3) Bird P, Conaghan P, Ejbjerg B, McQueen F, Lassere M, Peterfy C, et al. The development of the EULAR-OMERACT rheumatoid arthritis MRI reference image atlas. *Ann Rheum Dis* 2005;64 Suppl 1:i8-i10.
- (4) Conaghan P, Bird P, Ejbjerg B, O'Connor P, Peterfy C, McQueen F, et al. The EULAR-OMERACT rheumatoid arthritis MRI reference image atlas: the metacarpophalangeal joints. *Ann Rheum Dis* 2005;64 Suppl 1:i11-i21.
- (5) Ejbjerg B, McQueen F, Lassere M, Haavardsholm E, Conaghan P, O'Connor P, et al. The EULAR-OMERACT rheumatoid arthritis MRI reference image atlas: the wrist joint. *Ann Rheum Dis* 2005;64 Suppl 1:i23-i47.
- (6) McQueen F, Østergaard M, Peterfy C, Lassere M, Ejbjerg B, Bird P, et al. Pitfalls in scoring MR images of rheumatoid arthritis wrist and metacarpophalangeal joints. *Ann Rheum Dis* 2005;64 Suppl 1:i48-i55.
- (7) Østergaard M, Bøyesen P, Eshed I, Gandjbakhch F, Lillegraven S, Bird P, et al. Development and preliminary validation of a magnetic resonance imaging joint space narrowing score for use in rheumatoid arthritis: potential adjunct to the OMERACT RA MRI scoring system. *J Rheumatol* 2011;38:2045-50.
- (8) Døhn UM, Conaghan PG, Eshed I, Boonen A, Bøyesen P, Peterfy CG, et al. The OMERACT-RAMRIS Rheumatoid Arthritis Magnetic Resonance Imaging Joint Space Narrowing Score: Intrareader and Interreader Reliability and Agreement with Computed Tomography and Conventional Radiography. *J Rheumatol* 2014;41:392-7.
- (9) Glinatsi D, Lillegraven S, Haavardsholm EA, Eshed I, Conaghan PG, Peterfy C, et al. Validation of the OMERACT Magnetic Resonance Imaging Joint Space Narrowing Score for the Wrist in a Multireader Longitudinal Trial. *J Rheumatol* 2015;42:2480-5.
- (10) Glinatsi D, Bird P, Gandjbakhch F, Haavardsholm EA, Peterfy CG, Vital EM, et al. Development and validation of the OMERACT rheumatoid arthritis magnetic resonance (RAMRIS) tenosynovitis scoring system in a multi-reader exercise. *J Rheumatol* (OMERACT submitted) 2016.
- (11) Østergaard M, McQueen F, Wiell C, Bird P, Boyesen P, Ejbjerg B, et al. The OMERACT psoriatic arthritis magnetic resonance imaging scoring system

- (PsAMRIS): definitions of key pathologies, suggested MRI sequences, and preliminary scoring system for PsA Hands. *J Rheumatol* 2009;36:1816-24.
- (12) Bøyesen P, McQueen FM, Gandjbakhch F, Lillegraven S, Coates L, Wiell C, et al. The OMERACT Psoriatic Arthritis Magnetic Resonance Imaging Score (PsAMRIS) is reliable and sensitive to change: results from an OMERACT workshop. *J Rheumatol* 2011;38:2034-8.
 - (13) Glinatsi D, Bird P, Gandjbakhch F, Mease PJ, Boyesen P, Peterfy CG, et al. Validation of the OMERACT Psoriatic Arthritis Magnetic Resonance Imaging Score (PsAMRIS) for the Hand and Foot in a Randomized Placebo-controlled Trial. *J Rheumatol* 2015;42:2473-9.
 - (14) Landewe RB, Hermann KG, van der Heijde DM, Baraliakos X, Jurik AG, Lambert RG, et al. Scoring sacroiliac joints by magnetic resonance imaging. A multiple-reader reliability experiment. *J Rheumatol* 2005;32:2050-5.
 - (15) Lukas C, Braun J, van der Heijde D, Hermann KG, Rudwaleit M, Østergaard M, et al. Scoring inflammatory activity of the spine by magnetic resonance imaging in ankylosing spondylitis: a multireader experiment. *J Rheumatol* 2007;34:862-70.
 - (16) Song IH, Hermann K, Haibel H, Althoff CE, Listing J, Burmester G, et al. Effects of etanercept versus sulfasalazine in early axial spondyloarthritis on active inflammatory lesions as detected by whole-body MRI (ESTHER): a 48-week randomised controlled trial. *Ann Rheum Dis* 2011;70:590-6.
 - (17) Weckbach S, Schewe S, Michaely HJ, Steffinger D, Reiser MF, Glaser C. Whole-body MR imaging in psoriatic arthritis: additional value for therapeutic decision making. *Eur J Radiol* 2011;77:149-55.
 - (18) Althoff CE, Sieper J, Song IH, Haibel H, Weiss A, Diekhoff T, et al. Active inflammation and structural change in early active axial spondyloarthritis as detected by whole-body MRI. *Ann Rheum Dis* 2013;72:967-73.
 - (19) Karpitschka M, Godau-Kellner P, Kellner H, Horng A, Theisen D, Glaser C, et al. Assessment of therapeutic response in ankylosing spondylitis patients undergoing anti-tumour necrosis factor therapy by whole-body magnetic resonance imaging. *Eur Radiol* 2013.
 - (20) Schanz S, Henes J, Ulmer A, Kotter I, Fierlbeck G, Claussen CD, et al. Magnetic resonance imaging findings in patients with systemic scleroderma and musculoskeletal symptoms. *Eur Radiol* 2013;23:212-21.
 - (21) Axelsen MB, Eshed I, Duer-Jensen A, Moller JM, Pedersen SJ, Østergaard M. Whole-body MRI assessment of disease activity and structural damage in rheumatoid arthritis: first step towards an MRI joint count. *Rheumatology (Oxford)* 2014;53:845-53.
 - (22) Mackie SL, Pease CT, Fukuba E, Harris E, Emery P, Hodgson R, et al. Whole-body MRI of patients with polymyalgia rheumatica identifies a distinct subset with complete patient-reported response to glucocorticoids. *Ann Rheum Dis* 2015;74:2188-92.

- (23) Poggendorf RP, Eshed I, Østergaard M, Sorensen IJ, Moller JM, Madsen OR, et al. Enthesitis in patients with psoriatic arthritis, axial spondyloarthritis and healthy subjects assessed by 'head-to-toe' whole-body MRI and clinical examination. *Ann Rheum Dis* 2015;74:823-9.
- (24) Poggendorf RP, Pedersen SJ, Eshed I, Sorensen IJ, Moller JM, Madsen OR, et al. Head-to-toe whole-body MRI in psoriatic arthritis, axial spondyloarthritis and healthy subjects: first steps towards global inflammation and damage scores of peripheral and axial joints. *Rheumatology (Oxford)* 2015;54:1039-49.
- (25) Althoff CE, Sieper J, Song IH, Weiss A, Diekhoff T, Haibel H, et al. Comparison of Clinical Examination versus Whole-body Magnetic Resonance Imaging of Enthesitis in Patients with Early Axial Spondyloarthritis during 3 Years of Continuous Etanercept Treatment. *J Rheumatol* 2016;43:618-24.
- (26) Boers M, Kirwan JR, Gossec L, Conaghan PG, d'Agostino MA, Bingham CO, III, et al. How to choose core outcome measurement sets for clinical trials: OMERACT 11 approves filter 2.0. *J Rheumatol* 2014;41:1025-30.
- (27) Weber U, Hodler J, Kubik RA, Rufibach K, Lambert RG, Kissling RO, et al. Sensitivity and specificity of spinal inflammatory lesions assessed by whole-body magnetic resonance imaging in patients with ankylosing spondylitis or recent-onset inflammatory back pain. *Arthritis Rheum* 2009;61:900-8.
- (28) Weber U, Maksymowych WP, Jurik AG, Pfirmann CW, Rufibach K, Kissling RO, et al. Validation of whole-body against conventional magnetic resonance imaging for scoring acute inflammatory lesions in the sacroiliac joints of patients with spondylarthritis. *Arthritis Rheum* 2009;61:893-9.
- (29) Weber U, Hodler J, Jurik AG, Pfirmann CW, Rufibach K, Kissling RO, et al. Assessment of active spinal inflammatory changes in patients with axial spondyloarthritis: validation of whole body MRI against conventional MRI. *Ann Rheum Dis* 2010;69:648-53.
- (30) Song IH, Hermann KG, Haibel H, Althoff CE, Poddubnyy D, Listing J, et al. Relationship between active inflammatory lesions in the spine and sacroiliac joints and new development of chronic lesions on whole-body MRI in early axial spondyloarthritis: results of the ESTHER trial at week 48. *Ann Rheum Dis* 2011;70:1257-63.
- (31) Jurik AG, Zejden A, Lambert RG, Rufibach K, Hodler J, Maksymowych WP, et al. Pitfalls in MR morphology of the sterno-costoclavicular region using whole-body MRI. *Clin Radiol* 2013;68:785-91.
- (32) Song IH, Hermann KG, Haibel H, Althoff CE, Poddubnyy D, Listing J, et al. Prevention of new osteitis on magnetic resonance imaging in patients with early axial spondyloarthritis during 3 years of continuous treatment with etanercept: data of the ESTHER trial. *Rheumatology (Oxford)* 2015;54:257-61.
- (33) Song IH, Hermann KG, Haibel H, Althoff CE, Poddubnyy D, Listing J, et al. Inflammatory and fatty lesions in the spine and sacroiliac joints on whole-body MRI in early axial spondyloarthritis--3-Year data of the ESTHER trial. *Semin Arthritis Rheum* 2016;45:404-10.

- (34) Weiss BG, Bachmann LM, Pfirrmann CW, Kissling RO, Zubler V. Whole Body Magnetic Resonance Imaging Features in Diffuse Idiopathic Skeletal Hyperostosis in Conjunction with Clinical Variables to Whole Body MRI and Clinical Variables in Ankylosing Spondylitis. *J Rheumatol* 2016;43:335-42.
- (35) Rudwaleit M, Jurik AG, Hermann KG, Landewe R, van der Heijde D, Baraliakos X, et al. Defining active sacroiliitis on magnetic resonance imaging (MRI) for classification of axial spondyloarthritis: a consensual approach by the ASAS/OMERACT MRI group. *Ann Rheum Dis* 2009;68:1520-7.
- (36) Hermann KG, Baraliakos X, van der Heijde DM, Jurik AG, Landewe R, Marzo-Ortega H, et al. Descriptions of spinal MRI lesions and definition of a positive MRI of the spine in axial spondyloarthritis: a consensual approach by the ASAS/OMERACT MRI study group. *Ann Rheum Dis* 2012;71:1278-88.
- (37) Lambert RG, Bakker PA, van der Heijde D, Weber U, Rudwaleit M, Hermann KG, et al. Defining active sacroiliitis on MRI for classification of axial spondyloarthritis: update by the ASAS MRI working group. *Ann Rheum Dis* (Published online) 2016.

Funding statement:

There was no funding for this systematic literature review and consensus process. PGC is supported in part by the National Institute for Health Research (NIHR) Leeds Musculoskeletal Biomedical Research Unit. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Table 1: Characteristics of publications on WBMRI of the extremities, retrieved by the literature search

First author	Year	Journal	Disease	Patients	Controls	Sites	Inflammatory lesions	Structural lesions	Manufacturer	Field strength	Gd	Sequences	Planes	Duration	TP (week)	Readers	Read setup	Reproducibility
Song IH	2011	Ann Rheum Dis	axSpA	76	None	27	enthesitis	ND	Siemens	1.5T	No	T1, STIR	cor, sag	65 mins.	0, 24 & 48	2	consensus	ND
Weckbach S	2011	Eur J Radiol	PsA	30	None	15	osteitis, effusion, synovitis, enthesitis	erosions, destruction	Siemens	1.5T	yes	STIR, dynT1*, VIBE	Cor	45 mins.	0	2	consensus	ND
Althoff CE	2013	Ann Rheum Dis	axSpA	75	None	J:12; E:30	effusion/synovitis, enthesitis	erosions	Siemens	1.5T	No	T1, STIR	cor, sag	65 mins.	0	2	consensus	ND
Karpitscka M	2013	Eur Radiol	AS	10	None	NA	synovitis, enthesitis	ND	Siemens	1.5T	yes	T1, STIR, T1fs/Gd	cor	-	0, 26 & 52	2	independent	NA
Schanz S	2013	Eur Radiol	SSc	18	None	NA	fasciitis, myositis, tenosynovitis, BME	ND	Siemens	1.5T	yes	STIR, T1**, T1/Gd**	cor	30 mins.	0	2	independent	inter-reader, kappa: 0.8–1.0
Axelsen MB	2014	Rheumatology	RA	20	None	J:76; E:30	Synovitis, enthesitis	erosions	Philips	3T	yes	T1, STIR, T1/Gd	cor, ax	60 mins.	0	1	ND	intra-reader agreement rate 85–100%

Mackie SL	2015	Ann Rheum Dis	PMR	22	16 (RA)	10	extracapsular PMR pattern	ND	Siemens	3T	yes	T2fs, Dixon-VIBE	ax, 3D-Reco	19 mins.	0	2	consensus	ND
Poggenborg RP	2015	Ann Rheum Dis	PsA, axSpA	36	12	35	enthesitis	ND	Philips	3T	yes	T1, STIR, T1fs/Gd	cor, ax	61 mins.	0	1	ND	intra-reader ICC: 0.58/0.85
Poggenborg RP	2015	Rheumatology	PsA, axSpA	36	12	76	synovitis, BME	erosions	Philips	3T	yes	T1, STIR, T1fs/Gd	cor, ax	61 mins.	Baseline	1	ND	intra-reader ICC: 0.31-1.0
Althoff CE	2016	J Rheumatol	axSpA	41	None	21	enthesitis	ND	Siemens	1.5T	No	T1, STIR	cor, sag	65 mins.	baseline, year 2 & 3	2	consensus	ND

AS: ankylosing spondylitis; ax: axial; AxSpa: axial spondyloarthritis; BME: bone marrow edema; cor: coronal; dyn: dynamic; E: entheses; Gd: gadolinium contrast, FS: fat suppression ICC: intraclass correlation coefficient, J: joints; NA: not available, ND: not done; PMR: polymyalgia rheumatic; PsA: psoriatic arthritis, RA: rheumatoid arthritis; sag: sagittal; STIR: short tau inversion recovery, T: tesla; TP: time point for MRI examination; VIBE: volumetric interpolated breath-hold examination, WBMRI: whole-body magnetic resonance imaging

APPENDIX: Preliminary OMERACT WBMRI assessment system of inflammation in inflammatory arthritides.**Initial Focus Area**

Assessing inflammation in the extremities

Selection of Key Pathologies

Inflammation in joints:

- Soft tissues (synovitis)
- Bone (osteitis)

Inflammation at entheses (enthesitis):

- Soft tissue (soft tissue inflammation)
- Bone (osteitis)

Definitions of Key Pathologies

Joints: Synovitis

Procedure: If T1-postGd images are available, synovitis should be assessed according to option a. If only STIR/T2FS images are available: Synovitis/effusion should be assessed according to option b.

Definitions:

- Option a. Definition of synovitis, based on T1-postGd images: An area in the synovial compartment that shows above-normal post-gadolinium enhancement on T1-weighted images, of a thickness greater than the width of the normal joint capsule
- Option b. Definition of synovitis/effusion, based on STIR/T2FS images: (to be used if STIR/T2FS images, but not T1-postGd images, are available): An area in the synovial compartment that shows high signal intensity on T2-weighted fat-saturated or STIR images, of a thickness greater than the width of the normal joint capsule and joint fluid.¹

Joints: Osteitis

Procedure: If STIR/T2FS images are available, assess bone edema according to option a. If only T1-postGd images are available: Assess intraosseous post-Gd enhancement according to option b.

Definitions:

- Option a. Definition of osteitis, based on STIR/T2FS images: A lesion within the trabecular bone, with ill-defined margins and high signal intensity on T2-weighted fat-saturated and STIR images (“bone marrow edema”)
- Option b. Definition of osteitis, based on T1-postGd images: A lesion within the trabecular bone marrow, with ill-defined margins, which shows above-normal enhancement (signal intensity increase) on T1-weighted after iv. Gadolinium contrast injection (“bone marrow post-contrast enhancement”)

Entheses: Enteseal soft tissue inflammation

Procedure: If T1-postGd images are available, enteseal soft tissues should be assessed according to option a. If only STIR/T2FS images are available, enteseal soft tissues should be assessed according to option b.

Definitions:

- Option a. Definition of enteseal soft tissue inflammation, based on T1-postGd images: Above-normal post-gadolinium enhancement of enteseal soft tissues on T1-weighted images.
- Option b. Definition of enteseal soft tissue inflammation, based on STIR/T2FS images:

High signal intensity of the enteseal soft tissues on T2-weighted fat-saturated or STIR images.

Entheses: osteitis

Procedure: If STIR/T2FS images are available, assess bone edema according to option a. If only T1-postGd images are available: Assess intraosseous post-Gd enhancement according to option b.

Definitions

- Option a. Definition of osteitis, based on STIR/T2FS images:

A lesion within the enteseal bone marrow, with ill-defined margins and high signal intensity on T2-weighted fat-saturated and STIR images (“bone marrow edema”).

- Option b. Definition of osteitis, based on T1-postGd images:

A lesion within the enteseal bone marrow, with ill-defined margins, which shows above-normal enhancement (signal intensity increase) on T1-weighted after iv. Gadolinium contrast injection (“bone marrow post-contrast enhancement”).

Anatomical Locations for Assessment

Joints:

- All peripheral joints, except the elbow.
- Joints of the chest wall: sternoclavicular joint, costosternal joints, manubriosternal joint

Entheses:

- Upper extremity: Insertion of supraspinatus tendon into humerus,
- Pelvis: Anterior superior iliac spine, posterior superior iliac spine, iliac crest (excluding the anterior and posterior superior iliac spines), ischial tuberosity, pubic symphysis.
- Lower extremities: greater trochanter of femur, medial femoral condyle, lateral femoral condyle, insertion of the quadriceps femoris tendon into patella, insertion of the patellar ligament into patella, insertion of the patellar ligament into the tibial tuberosity, insertion of the calcaneal (Achilles) tendon into calcaneus, insertion of the plantar aponeurosis into calcaneus.

MRI Sequences and Imaging Planes:

Recommended imaging planes:

- Spine: Sagittal
- Shoulder/anterior chest wall: Coronal
- Sacroiliac joints: Coronal oblique
- Wrist and hand: Coronal
- Pelvis: Axial²
- Knee: Axial (+ if possible sagittal²)
- Ankle: Sagittal
- Feet: Axial (will provide coronal view of foot)

Recommended MRI sequences:

- T1-PostGd or alternatively STIR/T2FS,
- T1 without contrast (not mandatory if only inflammation is assessed)

Scoring System

- For each selected joint, synovitis (a) and osteitis (b) are scored separately (0-2)³

- For each selected entheses, soft tissue inflammation (c) and osteitis (d) are scored separately (0-2)

- Individual scores: 0: None, 1: Mild-moderate, 2: severe⁴

- Sum scores: joint inflammation index: a+b, enteseal inflammation index: c+d, total peripheral inflammation index: a+b+c+d.

¹Enhancing synovitis on T1-postGD may appear with low signal on STIR/T2FS, presumably because of high collagen content.

²Additional sagittal plane needed for adequate assessment of patellar and quadriceps tendon insertions. Additional coronal plane improves hip joint assessment

³When only STIR/T2wFS is available, the assessment will cover synovitis and effusion.

⁴A 0-2 score was agreed by the group at the Berlin meeting, since it was considered difficult to reliably score with more steps with the current image quality. Definitions of mild/moderate/severe are as described for RAMRIS/PsAMRIS (1, 11). At the OMERACT meeting participants in the Special Interest Group suggested using a 0-3 score (none, mild, moderate, severe), and this approach may also be tested.

Figure 1: Flow Diagram for the Systematic Literature Review, in accordance with the PRISMA statement.

