

This is a repository copy of *The dynamics of response as measured by multiple composite* outcome tools in the Tight COntrol of inflammation in early Psoriatic Arthritis (TICOPA) trial.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/116102/

Version: Supplemental Material

Article:

Coates, LC, Mahmood, F, Emery, P et al. (2 more authors) (2017) The dynamics of response as measured by multiple composite outcome tools in the TIght COntrol of inflammation in early Psoriatic Arthritis (TICOPA) trial. Annals of the Rheumatic Diseases, 76 (10). ISSN 0003-4967

https://doi.org/10.1136/annrheumdis-2017-211137

© The Authors 2017. Published by BMJ Publishing Group Ltd (& EULAR). This is an author produced version of a paper published in Annals of the Rheumatic Diseases. Uploaded in accordance with the publisher's self-archiving policy. No commercial use is permitted unless otherwise expressly granted.

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Supplementary table S4. Domain elements of the (a) PASDAS (b) GRACE and (c) CPDAI for the group with oligoarthritis.

(a) PASDAS (VAS: visual analogue scale. SF36-PCS: estimated physical component score of the Short Form 36).

Values are mean, median and range.

Domain	Tight control n = 21	Standard Care n = 23
Patient global VAS (mm)	32.1, 28, 1 - 77	29.0, 22, 5 - 82
Physician global VAS (mm)	28.0, 26, 5 - 77	30.3, 28, 2 - 52
Tender joint count (0-68)	3.0, 2, 0 - 12	4.2, 3, 1 - 28
Swollen joint count (0-66)	2.1, 2, 0 - 6	2.0, 2, 0 - 4
Enthesitis (0-6)	1.0, 0.6, 0 – 4.7	1.0, 0.3, 0 - 6
Dactylitis (0-20)	0.3, 0, 0 - 3	0.4, 0, 0 - 3
CRP (mg/L)	24.6, 6.9, 5 - 153	12.1, 6.2, 5 - 49
Sf36-PCS (0-100)	39.5, 38.8, 24 - 51	40.3, 41.6, 20 - 50

(b) Modified GRACE (VAS: visual analogue scale. PASI: psoriasis area and severity index. PsAQoL: psoriatic arthritis quality of life. HAQ: Health assessment questionnaire)

Values are mean, median and range.

Domain	Tight control n = 26	Standard Care n = 28
Patient global VAS (0 – 100)	30.0, 27.5, 0 - 77	29.0, 24.5, 5 - 82
Pain VAS (0 – 100)	36.3, 35, 2 - 94	38.1, 29.5, 8 - 79
Tender joint count (0-68)	3.0, 2.5, 0 - 12	3.7, 3, 0 - 28
Swollen joint count (0-66)	2.1, 2, 0 - 6	2.0, 2, 0 - 4
HAQ (0 – 3)	0.5, 0.43, 0 - 2	0.4, 0.25, 0 – 1.75
PASI (0 – 72)	1.8, 1.5, 0 – 5.7	2.6, 1.8, 0 – 9.5
PsAQoL (0 – 20)	5.5, 4.5, 0 - 17	4.5, 2, 0 - 19

(c) CPDAI.

Values are mean, median and range.

Domain	Tight control n = 22	Standard Care n = 24
Tender joint count (0-68)	3.0, 2, 0 - 12	4.2, 3, 0 - 28
Swollen joint count (0-66)	2.1, 2, 0 - 6	2.0, 2, 0 - 4
HAQ (0 – 3)	0.5, 0.6, 0 - 2	0.4, 0.3, 0 – 1.75
PASI (0 – 72)	1.8, 1.6, 0 – 5.7	2.5, 2.0, 0 – 9.5
Est DLQI	3.2, 3.0, 1.3 – 8.0	3.2, 3.3, 1.3 – 8.6
Enthesitis (0-6)	1.0, 0.6, 0 - 6	1.0, 0.3, 0 - 6
Dactylitis (0-20)	0.3, 0, 0 - 3	0.4, 0, 0 - 3
BASDAI (0 – 10)	4.0, 4.6, 0.6 – 7.0	3.9, 3.9, 0.7 – 8.0
PsAQoL (0 – 20)	5.8, 5.5, 0 – 17	4.5, 2.0, 0 – 19