



UNIVERSITY OF LEEDS

This is a repository copy of *The dynamics of response as measured by multiple composite outcome tools in the Tight Control of inflammation in early Psoriatic Arthritis (TICOPA) trial*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/116102/>

Version: Supplemental Material

Article:

Coates, LC, Mahmood, F, Emery, P et al. (2 more authors) (2017) The dynamics of response as measured by multiple composite outcome tools in the Tight Control of inflammation in early Psoriatic Arthritis (TICOPA) trial. *Annals of the Rheumatic Diseases*, 76 (10). ISSN 0003-4967

<https://doi.org/10.1136/annrheumdis-2017-211137>

© The Authors 2017. Published by BMJ Publishing Group Ltd (& EULAR). This is an author produced version of a paper published in *Annals of the Rheumatic Diseases*. Uploaded in accordance with the publisher's self-archiving policy. No commercial use is permitted unless otherwise expressly granted.

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Supplementary table S1. The grid required to assess the categories of the CPDAI

	None (0)	Mild (1)	Moderate (2)	Severe (3)
Peripheral Arthritis		≤ 4 joints; normal function (HAQ <0.5)	≤ 4 joints but function impaired; or > 4 joints, normal function	> 4 joints <u>and</u> function impaired
Skin Disease		PASI ≤ 10 and est DLQI ≤ 10	PASI ≤ 10 but est DLQI >10; or PASI > 10 but est DLQI ≤ 10	PASI > 10 <u>and</u> est DLQI > 10
Enthesitis		≤ 3 sites; normal function (HAQ <0.5)	≤ 3 sites but function impaired; or >3 sites but normal function	>3 sites <u>and</u> function impaired
Dactylitis		≤ 3 digits; normal function (HAQ <0.5)	≤ 3 digits but function impaired; or >3 digits but normal function	>3 digits <u>and</u> has function impaired
Spinal Disease		BASDAI <4; normal function (PsAQoL < 6)	BASDAI >4 but normal function; BASDAI <4 but function impaired (PsAQoL ≥ 6)	BASDAI >4 <u>and</u> function impaired