



This is a repository copy of *Random walk attachment graphs*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/116490/>

Version: Published Version

Article:

Cannings, C. and Jordan, J. (2013) Random walk attachment graphs. ELECTRONIC COMMUNICATIONS IN PROBABILITY, 18 (77). ISSN 1083-589X

<https://doi.org/10.1214/ECP.v18-2518>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Random walk attachment graphs

Chris Cannings* Jonathan Jordan*

Abstract

We consider the random walk attachment graph introduced by Saramäki and Kaski and proposed as a mechanism to explain how behaviour similar to preferential attachment may appear requiring only local knowledge. We show that if the length of the random walk is fixed then the resulting graphs can have properties significantly different from those of preferential attachment graphs, and in particular that in the case where the random walks are of length 1 and each new vertex attaches to a single existing vertex the proportion of vertices which have degree 1 tends to 1, in contrast to preferential attachment models.

Keywords: random graphs; preferential attachment; random walk.

AMS MSC 2010: 05C82.

Submitted to ECP on December 21, 2012, final version accepted on September 3, 2013.

There is currently great interest in the preferential attachment model of network growth, usually called the Barabási-Albert [2, 1] model, though it dates back at least to Yule [11], and was discussed also by Simon [10]. In the simplest version of this an existing graph is incremented at each stage by adding a single new vertex which then attaches to a single pre-existing vertex; this latter is chosen from amongst those of the pre-existing graph with probability proportional to the degree of that vertex. In the Barabási-Albert model the new vertex will connect to m vertices, where m is fixed and is a parameter of the model, but here we only consider the case $m = 1$. One of the best known properties of the model is that it produces a power law degree distribution, as shown rigorously by Bollobás et al [3].

One weakness of this model and its generalisations is that this implicitly requires a calculation across all the existing vertices, or at least a knowledge of the total degree (sum of the vertex degrees) of the graph. This requirement then destroys the potential for this model to have emergent properties from local behaviour.

A possible solution to this was proposed by Saramäki and Kaski [9]. In their model the new vertex simply chooses a single vertex from the graph and then executes a random walk of length ℓ step initiated from that vertex. Saramäki and Kaski [9] and Evans and Saramäki [6] claim that this reproduces the Barabási-Albert degree distribution, even when $\ell = 1$. It is clear that this is the case if the random walk is run for long enough to have converged to its stationary distribution. However we will prove that in the particular case $\ell = 1$ the degree sequence does not converge to a power law distribution, but rather to a degenerate limiting distribution in which almost every vertex has degree 1.

*University of Sheffield. E-mail: c.cannings@shef.ac.uk, jonathan.jordan@shef.ac.uk

1 The Model

Let G_0 be an arbitrary (perhaps connected) graph, with v_0 vertices and e_0 edges. Form G_{n+1} from G_n by adding a single vertex. This vertex chooses a single vertex (i.e. this corresponds to $m = 1$ in the Barabási-Albert model) to connect to by picking a vertex uniformly at random in G_n and then, conditional on the vertex chosen, performing a simple random walk of length ℓ on G_n , starting from the randomly chosen vertex, and then choosing to connect to the destination vertex. Most of the time we will assume that ℓ is deterministic, but we will also consider a particular case where ℓ is replaced by a random variable.

2 Number of leaves

We first consider the number of leaves in the graph. Let $p_d^{(n)}$ be the proportion of vertices in G_n with degree d , and let $L_n = p_1^{(n)}$, i.e. the proportion of leaves. The number of edges in G_n will be $n + e_0$, the total degree will thus be $2(n + e_0)$, and the number of vertices will be $n + v_0$. Let V_n be the vertex initially chosen at random at step n , and let W_n be the vertex selected by the random walk, so the new vertex connects to W_n . We now prove the main result, which applies to the case where $\ell = 1$.

Theorem 2.1. *When $\ell = 1$, as $n \rightarrow \infty$, $L_n \rightarrow 1$, almost surely.*

Proof. We assume that G_0 is not a star. If G_0 is a star, then it is clear that, with probability 1, G_n will eventually not be a star, so we can just wait until this happens and re-label the first non-star graph as G_0 . If G_n is not a star each vertex has at least one neighbour which is not a leaf, and in particular no leaves have a leaf as their neighbour. If V_n is a leaf, which has probability L_n , then W_n will be one of its neighbours, which will not be a leaf, so in this case the number of leaves increases by 1. Hence, considering the conditional expectation of the number of leaves in G_{n+1} ,

$$\mathbb{E}((n + v_0 + 1)L_{n+1}|G_n) \geq (n + v_0)L_n + L_n = (n + v_0 + 1)L_n, \quad (2.1)$$

and so $\mathbb{E}(L_{n+1}|G_n) \geq L_n$ and so $(L_n)_{n \in \mathbb{N}}$ is a submartingale taking values in $[0, 1]$, and thus converges almost surely and in \mathcal{L}^2 to a limit, which we call L_∞ .

To show that $L_\infty = 1$ almost surely, note that conditional on V_n having degree d the probability of W_n not being a leaf is at least $1/d$, so we can make (2.1) sharper, getting

$$\mathbb{E}(L_{n+1}|G_n) \geq L_n + \sum_{d=2}^{\infty} \frac{p_d^{(n)}}{(n + v_0 + 1)d}. \quad (2.2)$$

The total degree of non-leaves in G_n is $2(n + e_0) - L_n(n + v_0) = (2 - L_n)(n + v_0) + 2(e_0 - v_0)$, and the number of non-leaves is $(1 - L_n)(n + v_0)$, so the average degree of non-leaves is $\frac{2 - L_n}{1 - L_n} + \frac{2(e_0 - v_0)}{(n + v_0)(1 - L_n)}$. Hence at least half the non-leaves have degree at most $2 \left(\frac{2 - L_n}{1 - L_n} + \frac{2(e_0 - v_0)}{(n + v_0)(1 - L_n)} \right)$ and so

$$\mathbb{E}(L_{n+1}|G_n) \geq L_n + \frac{1 - L_n}{2(n + 1)} \left(2 \left(\frac{2 - L_n}{1 - L_n} + \frac{2(e_0 - v_0)}{(n + v_0)(1 - L_n)} \right) \right)^{-1} \quad (2.3)$$

and so

$$\mathbb{E}(L_{n+1}) \geq \mathbb{E}(L_n) + \frac{1}{2(n + 1)} \mathbb{E} \left(\frac{1 - L_n}{2} \left(\frac{2 - L_n}{1 - L_n} + \frac{2(e_0 - v_0)}{(n + v_0)(1 - L_n)} \right)^{-1} \right). \quad (2.4)$$

If $\mathbb{E}(L_\infty) = \lim_{n \rightarrow \infty} \mathbb{E}(L_n) < 1$, then for some fixed $c < 1$ we must have $L_n \leq c$ with positive probability. The expectation on the right of (2.4) is then bounded away

from zero for large n , giving a contradiction and showing that $\mathbb{E}(L_\infty) = 1$ and thus that $L_\infty = 1$ almost surely. \square

It should be noted that the argument for Theorem 2.1 is dependent on the walk length being fixed at 1. For example, define a sequence of random variables $(X_n)_{n \in \mathbb{N}}$ which are independent and identically distributed with $P(X_n = 0) = p$ and $P(X_n = 1) = 1 - p$, and let the walk length from V_n to W_n be X_n , rather than a fixed ℓ as previously.

Then, by the same argument as before

$$\mathbb{E}(L_{n+1} - L_n | G_n, X_{n+1} = 1) \geq \frac{1 - L_n}{2} \frac{1 - L_n}{2(n + v_0 + 1)(2 - L_n)} + O(n^{-2}).$$

As there can be at most one more leaf in G_{n+1} than in G_n , we also have

$$\mathbb{E}(L_{n+1} - L_n | G_n, X_{n+1} = 1) \leq \frac{1 - L_n}{n + v_0 + 1} + O(n^{-2}).$$

Also, if there are no random walk steps from the initially chosen vertex the probability that the new vertex connects to a leaf is simply L_n , so

$$\mathbb{E}((n + v_0 + 1)L_{n+1} | G_n, X_{n+1} = 0) = (n + v_0)L_n + 1 - L_n,$$

and hence

$$\mathbb{E}(L_{n+1} - L_n | G_n, X_{n+1} = 0) = \frac{1}{n + v_0 + 1}(1 - 2L_n).$$

So, if we have $X_n = 0$ with probability p and 1 with probability $1 - p$ for all n independently of each other

$$\mathbb{E}(L_{n+1} - L_n | G_n) \geq \frac{1}{n + v_0 + 1} \left[p(1 - 2\lambda) + (1 - p) \frac{(1 - \lambda)^2}{4(2 - \lambda)} \right] + O(n^{-2}). \quad (2.5)$$

Similarly,

$$\mathbb{E}(L_{n+1} - L_n | G_n) \leq \frac{1}{n + v_0 + 1} [1 - \lambda(1 + p)] + O(n^{-2}). \quad (2.6)$$

The right hand side of (2.5) is negative if

$$L_n < \frac{1 + 9p - 2\sqrt{8p^2 + p}}{1 + 7p}$$

and n is sufficiently large and the right hand side of (2.6) is negative if $L_n > \frac{1}{1+p}$ and n is sufficiently large.

Note that

$$\frac{1 + 9p - 2\sqrt{8p^2 + p}}{1 + 7p} - \frac{1}{1 + p} \geq 0$$

for $p \in [0, 1]$ with equality only at $p = 0$ and $p = 1$, and that

$$\frac{1 + 9p - 2\sqrt{8p^2 + p}}{1 + 7p} \leq 1,$$

with equality only if $p = 0$.

A version of the argument of Lemma 2.6 of [8] now shows that, almost surely,

$$\liminf_{n \rightarrow \infty} L_n \geq \frac{1}{1 + p}$$

and

$$\limsup_{n \rightarrow \infty} L_n \leq \frac{1 + 9p - 2\sqrt{8p^2 + p}}{1 + 7p}.$$

So we do not get a similar result to Theorem 2.1 in this setting.

3 G_0 Bipartite

We now consider a special case which demonstrates that, for all odd ℓ , the random walk model of [9] differs fundamentally from that of the Barabási-Albert model.

Assume that G_0 is a bipartite graph, with the two parts coloured as red and blue. Then, in both models, for all n the graph G_n will be bipartite, and the parts can be coloured red and blue consistently for each n . Let the proportion of red vertices in G_n be R_n . We begin with the random walk model.

Theorem 3.1. *We have R_∞ such that R_n converges almost surely to R_∞ . If ℓ is even, then $R_\infty = \frac{1}{2}$, almost surely, while if ℓ is odd R_∞ is a random variable with a Beta distribution.*

Proof. Conditional on G_n , V_n will be red with probability R_n . If ℓ is odd W_n will be of opposite colour to V_n , which implies that the new vertex (which connects to W_n) will be of the same colour as V_n , and thus, conditional on G_n , will be red with probability R_n and blue with probability $1 - R_n$. Hence in this case the colours of vertices are equivalent to the colours of the balls in a standard Pólya urn (where when a ball is drawn two of the same colour are returned), and so by classical results on the Pólya urn (see, for example, Theorem 2.1 in [8]) R_n converges almost surely to R_∞ where R_∞ has a Beta distribution whose parameters depend on G_0 .

If ℓ is even then W_n is of the same colour as V_n and so the new vertex is of opposite colour to V_n . Hence this case corresponds to a two-colour generalised Pólya urn where a ball is selected and a ball of the opposite colour is added, namely a Friedman urn with $\alpha = 0$ and $\beta = 1$. In this case $R_n \rightarrow \frac{1}{2}$ almost surely; see for example Freedman [7], and Theorem 2.2 in [8]. \square

Theorem 3.2. *In the Barabási-Albert model $R_\infty = \frac{1}{2}$ almost surely.*

Proof. In this model it is possible to associate the selection of a vertex with an urn model by considering half-edges, and giving each half-edge the colour of its associated vertex, i.e. each edge is split into a red half and a blue half. The selection of a vertex with probability proportional to its degree is then equivalent to selecting a half-edge uniformly at random and then selecting the associated vertex. As the new edge added in G_{n+1} will always consist of a blue half and a red half, the proportion of red half-edges must converge to $\frac{1}{2}$, and as a red vertex is added if and only if a blue vertex is selected, the proportion of red vertices will converge to $\frac{1}{2}$, almost surely. \square

So in this respect the behaviour of the random walk model is different from the Barabási-Albert model when ℓ is odd, regardless of the size of ℓ .

4 Discussion

We have demonstrated that the model of Saramäki and Kaski is fundamentally different from that of Barabási and Albert, unless we allow an indefinite length for the random walk component. It does have the advantage of not requiring a global calculation, retaining the local behaviour characteristic which is desirable in models of emergent behaviour. An alternate approach might be to imagine that the addition of edges is affected by the vertices in G_n , rather than by the new vertex. Thus each vertex in G_n could link to a new vertex as it arises with probability proportional to its degree, independently of all other vertices, as in the variant of preferential attachment studied by Dereich and Mörters [4, 5]. This, of course, destroys one of the usual assumptions of the preferential attachment model that the number of new links is some fixed value m , though we could substitute the condition that the average number added was fixed.

The urn model approach is interesting particularly since there is much known about these (see for example the survey paper by Pemantle [8]). We might generalise the model to consider directed graphs where there are k colours c_i ; $i = 0, k - 1$, with directed edges only between a vertex of colour c_i and one of colour $c_{(i+1) \pmod k}$. When a new vertex is added it links at random to a vertex and then takes ℓ random steps along directed edges, its colour then being determined. The case $\ell \neq 0 \pmod k$ will have the proportions of each colour converging to $1/k$, whereas for $\ell = 0 \pmod k$ there will be a Dirichlet distribution with parameters depending on G_0 .

References

- [1] R. Albert, A.-L. Barabási, and H. Jeong. Mean-field theory for scale-free random networks. *Physica A*, 272:173–187, 1999.
- [2] A.-L. Barabási and R. Albert. Emergence of scaling in random networks. *Science*, 286:509–512, 1999. MR-2091634
- [3] B. Bollobás, O. Riordan, J. Spencer, and G. Tusnády. The degree sequence of a scale-free random graph process. *Random Structures and Algorithms*, 18:279–290, 2001. MR-1824277
- [4] S. Dereich and P. Mörters. Random networks with sublinear preferential attachment: Degree evolutions. *Electronic Journal of Probability*, 14:1222–1267, 2009. MR-2511283
- [5] S. Dereich and P. Mörters. Random networks with concave preferential attachment rule. *Jahresberichte der Deutschen Mathematiker Vereinigung*, 113:21–40, 2011. MR-2760002
- [6] T. Evans and J. Saramäki. Scale free networks from self-organisation. *Physical Review E*, 72:026138, 2005. MR-2177389
- [7] D.A. Freedman. Bernard Friedman’s urn. *Ann. Math. Statist.*, 36:956–970, 1965. MR-0177432
- [8] R. Pemantle. A survey of random processes with reinforcement. *Probability Surveys*, 4:1–79, 2007. MR-2282181
- [9] J. Saramäki and K. Kaski. Scale-free networks generated by random walkers. *Physica A*, 341:80–86, 2004. MR-2092677
- [10] H.A. Simon. On a class of skew distributions. *Biometrika*, 42:425–440, 1955. MR-0073085
- [11] G. U. Yule. A mathematical theory of evolution, based on the conclusions of Dr. J. C. Willis, F.R.S. *Philosophical Transactions of the Royal Society of London, B*, 213:21–87, 1925.

Acknowledgments. The first author acknowledges support from the European Union through funding under FP7-ICT-2011-8 project HIERATIC (316705).

Electronic Journal of Probability

Electronic Communications in Probability

Advantages of publishing in EJP-ECP

- Very high standards
- Free for authors, free for readers
- Quick publication (no backlog)

Economical model of EJP-ECP

- Low cost, based on free software (OJS¹)
- Non profit, sponsored by IMS², BS³, PKP⁴
- Purely electronic and secure (LOCKSS⁵)

Help keep the journal free and vigorous

- Donate to the IMS open access fund⁶ (click here to donate!)
- Submit your best articles to EJP-ECP
- Choose EJP-ECP over for-profit journals

¹OJS: Open Journal Systems <http://pkp.sfu.ca/ojs/>

²IMS: Institute of Mathematical Statistics <http://www.imstat.org/>

³BS: Bernoulli Society <http://www.bernoulli-society.org/>

⁴PK: Public Knowledge Project <http://pkp.sfu.ca/>

⁵LOCKSS: Lots of Copies Keep Stuff Safe <http://www.lockss.org/>

⁶IMS Open Access Fund: <http://www.imstat.org/publications/open.htm>