

This is a repository copy of *Pulse generation in the quorum machinery of Pseudomonas aeruginosa*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/116745/>

Version: Accepted Version

Article:

Alfiniyah, Cicik Alfiniyah, Bees, Martin Alan and Wood, Andrew James orcid.org/0000-0002-6119-852X (2017) Pulse generation in the quorum machinery of *Pseudomonas aeruginosa*. *Bulletin of Mathematical Biology*. pp. 1360-1389. ISSN 1522-9602

<https://doi.org/10.1007/s11538-017-0288-z>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Pulse generation in the quorum machinery of *Pseudomonas aeruginosa*

Cicik Alfiniyah · Martin A. Bees ·
A. Jamie Wood

Received: date / Accepted: date

Abstract *Pseudomonas aeruginosa* is a Gram-negative bacterium that is responsible for a wide range of infections in humans. Colonies employ quorum sensing (QS) to coordinate gene expression, including for virulence factors, swarming motility and complex social traits. The QS signalling system of *P. aeruginosa* is known to involve multiple control components, notably the *las*, *rhl* and *pqs* systems. In this paper, we examine the *las* system and, in particular, the repressive interaction of *rsaL*, an embedded small regulative protein, employing recent biochemical information to aid model construction. Using analytic methods we show how this feature can give rise to excitable pulse generation in this subsystem with important downstream consequences for rhamnolipid production. We adopt a symmetric competitive inhibition to capture the binding in the *lasI-rsaL* intergenic region and show our results are not dependent on the exact choice of this functional form. Furthermore, we examine the coupling of *lasR* to the *rhl* system, the impact of the predicted capacity for pulse generation and the biophysical consequences of this behaviour. We hypothesise that the interaction between the *las* and *rhl* systems may provide

C. Alfiniyah
Department of Mathematics,
University of York, York YO10 5DD, UK
Department of Mathematics,
Universitas Airlangga, Surabaya 60115, Indonesia
E-mail: caa520@york.ac.uk

M. A. Bees
Department of Mathematics,
University of York, York YO10 5DD, UK

A. J. Wood
Department of Mathematics,
University of York, York YO10 5DD, UK
Department of Biology,
University of York, York YO10 5YW, UK

a quorum memory to enable cells to trigger rhamnolipid production only when they are at the edge of an established aggregation.

Keywords *Pseudomonas aeruginosa* · Quorum sensing · Excitable behaviour · Bifurcation analysis

1 Introduction

Pseudomonas aeruginosa is a common Gram-negative bacterium responsible for a wide range of infections, including those of the urinary and gastrointestinal tract, the skin, and, most prominently, the respiratory system in immunocompromised hosts and sufferers of Cystic Fibrosis (CF). *P. aeruginosa* is a well-studied opportunistic pathogen in many contexts; it is well-known for its ability to form biofilms (O’Loughlin et al 2013; Singh et al 2015), its swarming behaviour (Daniels et al 2004; Shrout et al 2006), its rapid acquisition of resistance to antibiotics (Shih and Huang 2002) and its quorum sensing (QS) behaviour (Fuqua et al 2001). QS in *P. aeruginosa* is of particular interest because the mechanism is more complex than the originally-discovered, prototypical Lux homolog positive-feedback loop (e.g., James et al 2000; Shadel and Baldwin 1991) and the number of genes regulated by QS is large (Sitnikov et al 1995), especially those associated with virulence (O’Loughlin et al 2013). Mathematical models of QS in *Pseudomonas aeruginosa* have received a lot of attention. They provide the formalism to summarize current understanding as well as the means to explore mechanisms and evaluate emergent solution behaviour. Here, we develop a model description, employing recent genomic information and bioinformatic techniques, and explore mechanisms for the generation of pulses and memory effects for downstream rhamnolipid production.

In *P. aeruginosa* quorum sensing is governed by a hierarchical LuxI/luxR system, which consists of two homolog pairs: LasI/LasR and RhII/RhlR (Miller and Bassler 2001). Under this process, formation of the HSL autoinducers *N*-(3-oxododecanoyl)-HSL and *N*-(butyryl)-HSL are synthesised by LasI and RhII, respectively (see Fig. 1). It should be noted, however, that signalling systems of *las* and *rhl* are specific in their activation of autoinducers, i.e. *N*-(3-oxododecanoyl)-HSL is unable to activate RhlR and, similarly, LasR cannot be activated by *N*-(butyryl)-HSL (Latifi et al 1995; Pearson et al 1997). Although biochemically independent, the *las* system is able to exert control of the *rhl* system through the transcriptional promotion of the RhlR gene by LasR/*N*-(3-oxododecanoyl)-HSL (Pesci et al 1997; Latifi et al 1996). As well as gene regulation effects, the *rhl* system has an important function of modulating rhamnolipid production via *rhlAB*. Rhamnolipids are particularly important in swarming motility where they are postulated to lower surface tension and allow expansion of the colony through their surfactant and wetting properties, driving the bacteria to swarm on surfaces (Glick et al 2010; Kohler et al 2000). In addition, a quinolone system (Dubern and Diggle 2008) may also modulate

these interconnecting feedback loops (for simplicity, we do not model this aspect of QS here).

The first models of QS in *P. aeruginosa* were of the Lux (James et al 2000) and the Las systems (Dockery and Keener 2001). Both descriptions, and subsequent models, highlight the existence of a fold bifurcation structure for the concentration of the response regulator in response to bulk cell concentration. The seminal paper by Dockery and Keener (2001) provides the foundation for the emergence of QS based on formal mass action arguments. However, there have been significant increases in biochemical knowledge of this system in the last 15 years. Fagerlind et al (2003) constructed a large mass action model of the coupled Las and Rhl systems, including the effects of both RsaL and Vfr. This work also emphasises the existence of the classic two-fold bifurcation diagram for the activation level of the system (typically for the levels of liganded LasR) with respect to the external concentration of HSL. Subsequent work Fagerlind et al (2005) then explored how anti-virulence drugs (followed by Skindersoe et al (2008)) are able to quorum quench this system. The qualitative model of Viretta and Fussenegger (2004) does not include the effect of the *rsaL* negative loop and lacks the capacity to deal with the non-linear effects predicted here. The production of rhamnolipid was modelled and tested empirically by Chen et al (2004). The exhaustive rule-based approach of Schaadt et al (2013) includes the effect of *rsaL* but does not include the kinetic possibilities that emerge from non-linear interactions. There are many other studies on this system but, increasingly, these are based on the perspective of computational rather than mathematical modelling (e.g., Dockery and Keener 2001; Fagerlind et al 2005; Schaadt et al 2013), typically with a non-mechanistic emphasis.

In this article we focus in detail on the Las system and its internal regulation and modulation for individual cell. The *las* system is composed of LasI, autoinducer *N*-(3-oxododecanoyl)-HSL, LasR and RsaL (Pesci et al 1997). Importantly, biochemical evidence has now firmly established that LasR exists as a dimer in solution, with each monomer liganded by a single HSL (*N*-(3-oxododecanoyl)-HSL) molecule with additional evidence supporting higher multimers upon DNA binding (Schuster and Greenberg 2006). This is in contrast to assumptions of Dockery and Keener (2001) who assume a simpler mechanism for binding. Mathematically, the biochemical evidence is consistent with a Hill number of at least two and possibly much higher. In contrast, the RsaL transcriptional repressor is a helix-turn-helix protein that binds the promoter of *lasI* (De Kievit et al 2002) and exists as a monomer in the cell (Rampioni et al 2007a), leading to a Hill number of one. The transcription of both genes is promoted and regulated via binding of the two proteins to the same intergenic region between the *lasI* and *rsaL* operons, so except for rates the functional form for the transcription is likely to be identical (but we discuss variations of this in the appendix). This improvement in biochemical knowledge offers clear guidance for expected mathematical forms in the equations.




Fig. 1: The quorum sensing signalling system in *Pseudomonas aeruginosa* is composed of *las* and *rhl* systems. Arrows and barred arrows indicate activating (positive) and inhibiting (negative) regulatory interactions, respectively. Shapes on the diagram depict autoregulation terminology. Letters associated with each arrow reflect the associated time scale (ms = millisecond, s = second, and min = minute). Symbols associated with each shape are detailed in Table 1.

We begin with an investigation of the behaviour of gene regulation by constructing a mathematical model of a single cell. In order to focus on a biologically plausible region of parameter space we conduct an extensive search of published data, noting deviations from modelling choices in the literature. The dynamical system exhibits a range of interesting solution behaviour. We find that bistable solutions and oscillations are possible and explore the bifurcation structure. Of significant interest is the potential for excitability in the system, in the presence of both single and multiple steady states, such that a modest perturbation from a low level steady state triggers a large-amplitude excursion around phase space that eventually returns the system to low steady state. The extracellular concentration of HSL can have a significant impact on the dynamics. As signal molecules can accumulate and diffuse in the external environment we demonstrate how pulse generation of the *las* system can be propagated in space.

This paper is organised as follows. In section 2, we describe in greater detail the biological system as well as the mathematical approach. In Section 3, we conduct a numerical exploration of our model, highlighting the key parameters in determining LasI and RsaL equilibria. We then employ extracellular concentration as a driving factor in the dynamical behaviour of the system. We conclude by discussing our findings, identify challenges and suggest future work in Section 4.

2 Methods

We construct the governing equations using mass action kinetics, except where noted, guided by the literature (e.g., Fagerlind et al 2003). First, consider the LasR regulator (R_L), its binding activator, the autoinducer 3O-C12-HSL (H_L), and the complex LasR/3O-C12-HSL (R_{LH}). If LasR associates at rate k_L^+ , dissociates at rate k_L^- , is produced at a rate β_0 and degrades or dilutes at rate γ_L , then we may write

$$\frac{dR_L}{dt} = -k_L^+ R_L H_L + k_L^- R_{LH} + \beta_0 - \gamma_L R_L, \quad (1)$$

In a similar manner, we formulate an equation for the complex, with degradation or dilution rate γ_{RL} , such that

$$\frac{dR_{LH}}{dt} = k_L^+ R_L H_L - k_L^- R_{LH} - \gamma_{RL} R_{LH}. \quad (2)$$

where $\beta_0, \gamma_L, \gamma_{RL}$ are positive constants.

New biochemical data have indicated approximately steady cellular levels (Ishihama et al 2014) of many transcription factors and, therefore, we hypothesise it is the change in activation rather than production *per se* that dictates the dominant dynamics; unlike previous studies that consider the regulated production of LasR. Therefore, we shall assume the association and dissociation processes occur over a sufficiently fast time scale that the production and

Table 1. Description of dimensional variables

Variable	Description	Unit
R_L	LasR	nM
H_L	3O-C12-HSL	nM
R_{LH}	LasR/3O-C12-HSL complex	nM
I_L	LasI	nM
\hat{I}_L	lasI mRNA	nM
S_L	RsaL	nM
\hat{S}_L	rsaL mRNA	nM
R_R	RhlR	nM
H_R	C4-HSL	nM
R_{RH}	RhlR/C4-HSL complex	nM
\hat{R}_R	rhlR mRNA	nM
I_R	RhlI	nM
\hat{I}_R	rhlI mRNA	nM

loss terms can be safely neglected: we disregard terms multiplied by β_0 , γ_L and γ_{RL} .

The autoinducer 3O-C12-HSL (H_L) is created in the system via the activity of the LasI synthase (I_L), which we take to be at rate β_{HL} , and is naturally lost from the system at rate γ_{HL} . The most significant loss of the autoinducer from the cell is via diffusion through the cell membrane, a process we account for separately. Taking a simplified description of diffusion we can express the diffusive term as being proportional to the concentration difference across the membrane of H_L , where we take the extracellular concentration to be H_{ex} . Therefore, D_{HL} represents an additional loss rate, which is multiplied by the concentration difference, $H_L - H_{ex}$, yielding

$$\frac{dH_L}{dt} = \beta_{HL}I_L - \gamma_{HL}H_L - D_{HL}(H_L - H_{ex}). \quad (3)$$

The enzyme LasI (I_L) is produced by the lasI gene through a transcription and translation process of lasI-mRNA (\hat{I}_L) at rate α_L and degrades at rate γ_{IL} , such that

$$\frac{dI_L}{dt} = \alpha_L\hat{I}_L - \gamma_{IL}I_L. \quad (4)$$

In a similar fashion, the inhibitor RsaL (S_L) is produced by rsaL genes through transcription and translation of rsaL-mRNA (\hat{S}_L) at rate α_S and degrades at rate γ_S , providing

$$\frac{dS_L}{dt} = \alpha_S\hat{S}_L - \gamma_S S_L. \quad (5)$$

Transcription at the lasI promoter site (\hat{I}_L) is activated by the LasR/3O-C12-HSL complex (R_{LH}). The production process is assumed to follow a Hill form with a Hill number p . Recent biochemical evidence strongly suggests a Hill number > 1 in contrast to the arguments of Dockery and Keener (2001); the activated form of lasR is at least dimeric (Schuster and Greenberg 2006) and it is possible that it forms a tetramer on the DNA. For mathematical simplicity we adopt the smaller potential value, $p = 2$, in the analysis that follows. In addition, the RsaL transcriptional repressor has been shown to bind to the

Table 2. Parameters employed in the model

Par	Description	Standard value	Unit	Value / Range	Comments (Based on) / Ref
k_L^+	rate of binding reaction between LasR and 3O-C12-HSL		$\text{nM}^{-1}\text{min}^{-1}$		ratio $\frac{k_L^-}{k_L^+} \sim 1000 - 2000 \text{ nM}$
k_L^-	dissociation reaction rate of LasR/3O-C12-HSL		min^{-1}		ratio $\frac{k_L^-}{k_L^+} \sim 1000 - 2000 \text{ nM}$, (Welch et al 2000)
α_L	rate at which LasI produced by lasI mRNA	0.5	min^{-1}	0.5	2 min to translate protein, (Alon 2006)
α_S	rate at which RsaL produced by rsaL mRNA	0.5	min^{-1}	0.5	2 min to translate protein, (Alon 2006)
β_{HL}	rate at which 3O-C12-HSL produced by LasI	8×10^2	min^{-1}	8×10^2	(Raychaudhuri et al 2005)
β_L	max. production rate of LasI when lasI mRNA is activated by LasR/3O-C12-HSL	1	nM min^{-1}	1	estimate
β_S	max. production rate of RsaL when rsaL mRNA is activated by LasR/3O-C12-HSL	1	nM min^{-1}	1	estimate
β_{L0}	basal production rate of lasI mRNA	0.1	nM min^{-1}	0.1	basal transcription rate of a protein, (Alon 2006)
β_{S0}	basal production rate of rsaL mRNA	0.1	nM min^{-1}	0.1	basal transcription rate of a protein, (Alon 2006)
K_L	affinity constant between LasR/3O-C12-HSL and lasI mRNA	116	nM	1-1000	(Alon 2006)
K_S	affinity constant between LasR/3O-C12-HSL and rsaL mRNA	116	nM	1-1000	(Alon 2006)
K_{SL}	dissociation constant of inhibitor RsaL to lasI mRNA	185	nM	1-1000	(Alon 2006)
γ_L	degradation rate of LasR	0.01	min^{-1}	0.01	(Alon 2006)
γ_{IL}	degradation rate of LasI	0.01	min^{-1}	0.01	(Alon 2006)
γ_S	degradation rate of RsaL	0.0025	min^{-1}	0.01	(Alon 2006)
γ_{HL}	degradation rate of 3O-C12-HSL	0.01	min^{-1}	0.01	(Alon 2006)
γ_{RL}	degradation rate of LasR/3O-C12-HSL	0.01	min^{-1}	0.01	(Alon 2006)
γ_{mL}	degradation rate of lasI mRNA	0.2	min^{-1}	0.2	5 min halflife of RNA, (Alon 2006)
γ_{mS}	degradation rate of rsaL mRNA	0.2	min^{-1}	0.2	5 min halflife of RNA, (Alon 2006)
D_{HL}	diffusion constant of 3O-C12-HSL	60	min^{-1}	$0 - 10^4$	(Pai and You 2009)
H_{ex}	concentration of extracellular 3O-C12-HSL	10	nM	10-100	estimate
R_{L0}	total concentration of LasR and LasR/3O-C12-HSL	200	nM	200	QseB in <i>E. coli</i> , (Ishihama et al 2014)

lasI-rsaL intergenic region but, unlike LasR, RsaL is a helix-turn-helix protein that exists as a monomer in the cell (Rampioni et al 2007a), and thus we will adopt Hill number $q = 1$ for the inhibition factor for production of RsaL (\hat{S}_L), mathematically equivalent to a Michaelis-Menten equation. The transcription of both genes is promoted and regulated via binding of the two proteins to the same intergenic region between the lasI and rsaL operons, so the functional form for the transcription is identical, with the exception of the numerical values of the transcription and loss rates. Note that this implies a negative feedback relation between the RsaL protein and its own production, which has not been historically represented on graphical depictions of the *las* system, but strongly implied by analysis of Rampioni et al (2007a). Biochemically there is insufficient evidence to determine whether the binding in the intergenic region results in competitive, uncompetitive or non-competitive binding system, or indeed whether there is a symmetry in the expression rates in each direction with all configurations of binding at the intergenic region. In the analysis below we adopt a symmetric competitive binding form (we choose $K_L = K_S$ in the analysis below), but show in Appendix I that qualitatively the results are not dependent on this functional choice. With basal expression of β_{L0} and a loss rate of γ_{mL} this leads to the following expression for lasI:

$$\frac{d\hat{I}_L}{dt} = \beta_L \frac{R_{LH}^p}{K_L^p (1 + S_L^q / K_{SL}^q)^p + R_{LH}^p} - \gamma_{mL} \hat{I}_L + \beta_{L0}. \quad (6)$$

Similarly, with basal expression of β_{S0} and a loss rate of γ_S for rsaL we obtain

$$\frac{d\hat{S}_L}{dt} = \beta_S \frac{R_{LH}^p}{K_S^p (1 + S_L^q / K_{SL}^q)^p + R_{LH}^p} - \gamma_{mS} \hat{S}_L + \beta_{S0}. \quad (7)$$

The numerical values of the temporal processes are such that we are able to make a number of assumptions regarding the timescales to simplify the system. We assume that the dominant, slowest processes are protein production from mRNA via translation and folding. Therefore, other processes, namely the liganding of regulators, DNA binding, synthetase operation, and the transcription of DNA are much faster and we can assume that four of our differential equations are at a quasi-steady state, such that

$$\hat{I}_L = \frac{\beta_L R_{LH}^p}{\gamma_{mL} \left(K_L^p \left(1 + \frac{S_L^q}{K_{SL}^q} \right)^p + R_{LH}^p \right)} + \frac{\beta_{L0}}{\gamma_{mL}}, \quad (8)$$

and

$$\hat{S}_L = \frac{\beta_S R_{LH}^q}{\gamma_{mS} \left(K_S^p \left(1 + \frac{S_L^q}{K_{SL}^q} \right)^p + R_{LH}^p \right)} + \frac{\beta_{L0}}{\gamma_{mS}}. \quad (9)$$

From equation (1) and (2), we have a *Moiety conservation* equation

$$\frac{dR_{LH}}{dt} + \frac{dR_L}{dt} = 0 \Rightarrow R_{LH} + R_L = R_{L0}, \text{ where } R_{L0} \text{ is a constant.} \quad (10)$$

Initially, we make the simplifying assumption that HSL diffusion is rapid and, therefore, that the equation for HSL, H_L can also be written in a quasi-steady state, providing

$$H_L = \frac{\beta_{HL} I_L}{D_{HL}} + H_{ex}. \quad (11)$$

Later, we relax this constraint. With the simplifications above, the system of equations for the *las* system, constructed for the LasI and RsaL loops, become just two differential equations. By assuming there is negligible basal production of *lasI* and *rsaL* genes ($\beta_{L0} = 0$ and $\beta_{S0} = 0$) the governing equations become

$$\frac{dI_L}{dt} = \frac{\alpha_L \beta_L}{\gamma_{mL}} \frac{R_{L0}^2 H_L^2}{K_L^2 \left(1 + \frac{S_L}{K_{SL}}\right)^2 \left(H_L + \frac{k_L^-}{k_L^+}\right)^2 + R_{L0}^2 H_L^2} - \gamma_{IL} I_L, \quad (12)$$

and

$$\frac{dS_L}{dt} = \frac{\alpha_S \beta_S}{\gamma_{mS}} \frac{R_{L0}^2 H_L^2}{K_S^2 \left(1 + \frac{S_L}{K_{SL}}\right)^2 \left(H_L + \frac{k_L^-}{k_L^+}\right)^2 + R_{L0}^2 H_L^2} - \gamma_S S_L, \quad (13)$$

with H_L as in equation (11).

We nondimensionalize this model by writing

$$\eta = \frac{\beta_{HL} k_L^+}{D_{HL} k_L^-} I_L, \quad \xi = \frac{S_L}{K_{SL}}, \quad \text{and} \quad \tau = \frac{\alpha_L \beta_L \beta_{HL} k_L^+}{\gamma_{mL} D_{HL} k_L^-} t, \quad (14)$$

so that (12) and (13) become

$$\frac{d\eta}{d\tau} = \frac{(\eta + a_1)^2}{a_2 (1 + \xi)^2 (\eta + a_1 + 1)^2 + (\eta + a_1)^2} - a_3 \eta, \quad (15)$$

$$\frac{d\xi}{d\tau} = \frac{a_4 (\eta + a_1)^2}{a_2 (1 + \xi)^2 (\eta + a_1 + 1)^2 + (\eta + a_1)^2} - a_5 \xi, \quad (16)$$

where

$$a_1 = \frac{H_{ex} k_L^+}{k_L^-}, \quad a_2 = \left(\frac{K_L}{R_{L0}}\right)^2, \quad (17)$$

$$a_3 = \frac{\gamma_{IL} \gamma_{mL} D_{HL} k_L^-}{\alpha_L \beta_L \beta_{HL} k_L^+}, \quad a_4 = \frac{\alpha_S \beta_S}{\gamma_{IL} \gamma_{mS} K_{SL}} a_3,$$

$$a_5 = \frac{\gamma_S}{\gamma_{IL}} a_3.$$

Here, a_1 , a_2 , a_3 , a_4 , and a_5 are positive constants. The biological interpretation of this model is that ξ inhibits the expression of both η and ξ , which describes negative feedback from competitive inhibition by RsaL to the expression of both *lasI* and *rsaL* genes. The binding of RsaL to the bidirectional *lasI-rsaL* counters positive feedback, thereby balancing levels of HSL. In addition, both η and ξ degrade exponentially. The model thus constructed is a type of incoherent feed forward motif (Alon 2006) as can be seen from Fig. 1.

Table 3. Non-dimensional Parameters involved in the model

Name	Description	Standard value	Range
a_1	the effect of extracellular signal molecules	0.01	$5 \times 10^{-3} - 10^{-1}$
a_2	a squared ratio of concentration of Las components	0.3	$25 \times 10^{-6} - 25$
a_3	the degradation of LasI relative to signal molecule production	0.3	0 – 100
a_4	the control of binding of RsaL to LasI	0.4	0 – 25×10^3
a_5	the degradation of RsaL relative to signal molecule production	0.075	0 – 25

3 Results

The non-dimensional set of differential equations (15) and (16) have been investigated analytically with the assistance of Mathematica (10; Wolfram) and solved numerically using Matlab (R2016a; MathWorks). We note that the fixed points of this system lie upon the straight line $a_3 a_4 \eta = a_5 \zeta$ but substitution of this expression into either nullcline leads to a quintic equation for the fixed points that does not provide sufficient further simplification to yield general expressions for stability bounds.

3.1 Parameter ranges in the system

Table 2 lists parameters that have either been adopted from the literature based on experimental evidence or estimated, as stated. Moreover, some parameters are chosen for the following reasons. The basal production rate of genes-mRNA can be considered as similar to the basal transcription rate of a protein. This is because the transcriptional regulator protein activates genes-mRNA in a very fast process before encoding the protein. We take typical values of total concentration of LasR and LasR/3O-C12-HSL to be 200 nM, as for the concentration of QseB in *E. coli* (Ishihama et al 2014). As described in the previous section, we simplify the governing equations by assuming quasi-steady states for the fast reactions. This results in just 5 non-dimensional parameters (see Table 3).

The dimensionless variable a_1 is proportional to H_{ex} , the extracellular HSL concentration, an important factor in controlling both the intracellular HSL production and cell-cell communication. We shall see that a_1 influences the location of the key bifurcation.

Parameter a_2 depends on K_L and represents the relative binding strength of the LasR dimer, and a_4 assigns binding control of RsaL to LasI through parameter K_{SL} , consequently it gives the relative production of RsaL to LasI. Both parameters a_2 and a_4 have a wide potential range, $25 \times 10^{-6} - 25$ and $0 - 25 \times 10^3$, respectively. This is because the affinity constant between transcriptional regulator protein to genes-mRNA is largely unmeasured and depends on the chain structure of the signal molecule (short or long chain; 1 – 1000 nM) (Alon 2006).

Parameter a_3 is inversely proportional to β_{HL} , the HSL production rate. It is related to a_5 , which describes the degradation of RsaL relative to HSL production. It is important to note that we set $a_5 < a_3$ due to the transcription factor being more stable than the synthase. We shall see that the parameters a_3 and a_1 allow for fold and Hopf bifurcations in the system, leading to bistability and oscillations, respectively, as well as excitable dynamics, and provide suitable control parameters that can be varied in experiments.

3.2 Excitable dynamics in the LasI and RsaL phase plane




Fig. 2: Qualitative dynamical behaviour of the *Pseudomonas aeruginosa* quorum sensing system. (a) Four phase portraits of interest resulting from intersection between LasI and RsaL (η and ξ , respectively) nullclines. As the parameter a_3 varies there can be one intersection point with no excitable dynamics (I and IV; for $a_3 = 0.1$ and 0.6 , respectively), three intersection points (II; for $a_3 = 0.2$), or one intersection point with excitable dynamics (III; for $a_3 = 0.3$). (b) Excitable dynamics in the LasI and RsaL phase plane. Sufficiently large perturbations result in an excursion trajectory around the phase plane. (c,d) Time variation of η and ξ corresponding to the excitable trajectory of LasI and RsaL, respectively. All other parameters are as in Table 3.

The simplified system (15) and (16) involves just two variables, which we investigate with phase plane analysis. By varying a_3 , four phase portraits of interest can be identified, as depicted in Fig. 2a. We plot the two nullclines $\dot{\eta} = 0$ and $\dot{\xi} = 0$, where η and ξ represent the dimensionless of LasI and RsaL concentration, respectively. Fixed points for the system lie at their intersection.

A qualitative description of the model behaviour is now possible, with reference to classical excitable systems such as the Fitzhugh-Nagumo model (Murray 1993). Cases I and IV (see Fig. 2a), which correspond to small and large values of the bifurcation parameter $a_3 = 0.1$ and 0.6 , respectively, possess a single stable fixed point with the nullclines positioned so that there is no possibility of excitable behaviour (see below). For case II (Fig. 2a) there are two further intersections, which yield the central zone of the well-known *S*-shaped bifurcation diagram with three fixed points in a stable-unstable-stable configuration. This phenomenon has been reported in many experiments on autoregulation of genes (e.g., Alves and Dilo 2005; Angeli et al 2004; Pognard 2014); more detail is presented in Fig. 3. The feedback loop that consists of both positive and negative autoregulation creates two possible LasI production states, “on” and “off” (at the large and small stable steady states, respectively), affecting signal molecule concentration. This is a familiar pattern that represents quorum sensing; the cells have a low steady state (off) from which it is possible to jump past an intermediate unstable state – either via stochastic fluctuations or changes in the external parameters, in particular the background level of HSL represented here by the parameter a_1 – to a stable high steady state (on). Crucially, the return to the low steady state typically is not reversible and the system must trace a hysteretic loop to return to the low steady state via changes in the external parameters.

In Case III there is a single fixed point and the nullcline $\dot{\eta} = 0$ has a positive local maximum (which distinguishes it from cases II and IV); see Fig. 2a). This case allows for excitability in the LasI and RsaL phase plane caused by the $\dot{\eta}$ nullcline having cubic-like shape with a positive local maximum ($a_3 < 0.55$). By considering the behaviour of a perturbation to the steady state solution (see Fig. 2b), if the production of the (*N*-(3-oxododecanoyl)-HSL) molecule is sufficiently small, then global stability of the single fixed point ensures that the trajectory returns to the steady state. The variable ξ will engage, rapidly deplete and return the system to the small fixed point with a short loop. However, interesting behaviour occurs when there is a greater perturbation of (*N*-(3-oxododecanoyl)-HSL). For then the variable η pushes the system to the right in the phase plane, so that there is an excitable pulse of (*N*-(3-oxododecanoyl)-HSL). Consequently, there is no short route back to the stable fixed point and the trajectory undergoes a large excursion before returning. A small region of parameter space in Case II can also result in excitability ($0.27 \leq a_3 \leq 0.28$). Here there are three intersections between η and ξ nullclines, which are low stable fixed point, an intermediate saddle and a high unstable fixed points. In this region of parameter space, the saddle-unstable fixed point region is not accessible to trajectories that initiate in a zone close to the stable fixed point (see Appendix II, particularly Fig. 8).

The initial condition $\eta(0) = \eta_0$ and concentration of extracellular HSL have a large effect on the amplitude and duration of the excited pulse, as can be seen in Fig. 2. The results in Fig. 2b are plotted as a function of time in Fig. 2c and 2d, where the purple dashed-line and red dashed-line represent either excitable or non-excitable trajectories for large and small perturbations, respectively (for $a_1 = 0.01$ or, equivalently, $H_{ex} = 10nM$). By increasing external HSL concentration in the system via the variable a_1 (equivalent to increasing H_{ex} from $10nM$ to $= 300nM$), we find solutions tending to the single high steady state irrespective of the initial perturbation (shown in Fig. 2c and 2d by blue solid and black dotted lines; discussed later).

3.3 Bifurcation structure

The QS circuitry of *P. aeruginosa* is complex, with the *las* system itself consisting of both positive and negative feedback loops. The *lasI* gene is activated by LasR/3O-C12-HSL leading to increased (*N*-(3-oxododecanoyl)-HSL) production, positive feedback. However, LasR/3O-C12-HSL also activates the *rsaL* gene and induces expression of RsaL. RsaL inhibits expression of *lasI* and *rsaL* genes, and this process constitutes negative feedback. In general, this negative feedback loop is employed to maintain a homeostatic balance in the system. However, the interaction of both positive and negative feedback can yield bistability (and associated *hysteresis*; (Pfeuty and Kaneko 2009)). There are many other studies on the dynamical behaviour of gene regulatory networks involving positive and negative feedback loops. Song et al (2007) demonstrated that interlocked positive and negative feedback loops play essential roles in bistability and oscillations. More complex bifurcations of co-dimension one or two have also been explored (Hat et al 2016). Varying $a_1 > 0$, the effect of extracellular signal molecule, alters the qualitative dynamical behaviour of LasI and RsaL. Thus initially we investigated the dynamics of our system with a one-parameter bifurcation analysis of LasI versus a_1 . We used continuation methods to track the evolution of solutions for η versus a_1 .

Fig. 3, a-e, depicts five qualitatively different dynamical behaviours, each at a different fixed value of a_3 , the degradation of LasI relative to signal molecule production. As shown in Fig. 3a-d, there are at least four different bifurcation diagrams. Bifurcations of steady states do not appear in Fig. 3e. Fig. 3a, for $a_3 = 0.26$, depicts the potential for *irreversible* bistability. The left-hand saddle-node bifurcation can cross the η -axis for sufficiently small a_3 , preventing a drop from the upper to the lower branch. In addition, in Fig. 3b-d, *reversible* bistability of LasI is evident in the range of a_1 bounded by the two limit points, SN_1 and SN_2 , which denote saddle-node bifurcations. In the low steady state branch, the concentration of LasI is low until a_1 level exceeds the critical value SN_1 ($a_1 = 0.029, 0.032$, and 0.0329 , for Fig. 3b, c, and d, respectively), at which point the concentration of LasI increases abruptly to a high value. In similar manner, starting with a_1 high, the concentration of LasI does not drop significantly until a_1 reduces below a critical value, either SN_2 ($a_1 =$






Fig. 3: Bifurcation diagrams for autoregulation of the *las* system with respect to extracellular signal molecule concentration a_1 at five values of a_3 : sub-figures a, b, c, d, and e are for $a_3 = 0.26, 0.3, 0.32, 0.323, \text{ and } 0.34$, respectively. Solid lines depict stable steady states (no eigenvalues with positive real part); blue and green-dashed lines depict unstable steady states with one or two eigenvalues with positive real part, respectively; open (solid) circles denote maximum and minimum values of $\eta(\text{LasI})$ on unstable (stable) limit-cycles. The red stars indicate values of parameter a_1 and coordinate $\eta(\text{LasI})$ at bifurcations. Codimension-1 singular points marked as SN indicate a saddle-node point (or limit point); *sub*-Hopf indicates a subcritical Hopf bifurcation point; *sup*-Hopf indicates a supercritical Hopf bifurcation point. All other parameters are as in Table 3.



a.



b.

Fig. 4: Two-dimensional bifurcation diagram for (a_1, a_3) . The bifurcation lines divide the parameter domain into six regions $\mathcal{R}_1, \dots, \mathcal{R}_6$. Each of these regions is explained in the main text. The blue lines depict the saddle-node lines, defined by the locus of saddle-node bifurcation points (or limit points), with subscripts 1 or 2 as in Figure (3). The Hopf lines are constructed by subcritical-Hopf and supercritical-Hopf bifurcation points, which are presented by brown lines. The bistable region, \mathcal{R}_3 , consists of *reversible* (above of gray-dash lines) and *irreversible* (below of gray-dash lines). The red stars mark particular values of parameters a_1 and a_3 indicating a cusp, Bogdanov-Takens point (BT) or generalized-Hopf point (GH).

0.023, 0.0318, and 0.0328, for Fig. 3b, c, and d, respectively) or an earlier bifurcation such as a sub-critical Hopf bifurcation. With $a_3 = 0.3$ (Fig. 3b), subcritical Hopf (sub-Hopf) bifurcation points exist for the high fixed point in the bistability region, at $a_1 = 0.024$. The upper steady-state solution is unstable between SN_2 and sub-Hopf point, and stable beyond the sub-Hopf point where $a_1 > 0.24$. For $a_3 = 0.32$ (Fig. 3c), the two saddle-nodes (SN_1 and SN_2) move close together. Consequently, the bistability region is narrow. Here, the Hopf bifurcation becomes supercritical (sup-Hopf; at $a_1 = 0.032$). For $a_3 = 0.323$ (Fig. 3d), the sup-Hopf point disappears and the saddle-node bifurcations collide in Fig. 3e for $a_3 \geq 0.325$; the bistable regime ceases to exist.

Fig. 4 provides a two-dimensional bifurcation diagram in the (a_1, a_3) -plane. In general, for a value of a_1 smaller than at the cusp point, decreasing a_3 from a large value we move through a region with one relatively small stable fixed point for η to one small stable fixed point with excitable dynamics. Decreasing a_3 further provides a bistable regime before eventually reaching a region with one large stable fixed point for sufficiently small a_3 . The bistability region collapses through the collision of the two saddle-node lines at the cusp ($a_1 = 0.033$, $a_3 = 0.324$), a possibility that might be inferred from Fig. 2a. For $a_1 > 0.033$, the bistable regime ceases to exist.

Three codimension-2 singular points, a cusp point, a Bogdanov-Takens bifurcation point (BT), and a generalized-Hopf bifurcation point (GH), are identified in Fig. 4. The Bogdanov-Takens bifurcation, at $(a_1, a_3) = (0.032, 0.321)$, is caused by coalescence of a saddle-node point (SN_2) and a Hopf bifurcation point (sup-Hopf). A sup-Hopf line extends to the lower right of the saddle-node lines close to the cusp in the bifurcation diagram. The generalized-Hopf bifurcation is at $(a_1, a_3) = (0.317, 0.002)$, a transition point from sup-Hopf to sub-Hopf. It is clear that the bifurcation structure is complex, especially close to the cusp point, although the solution space is dominated by the fold bifurcation and excitability.

The response diagram (Fig. 4) illustrates that the bifurcation lines divide the (a_1, a_3) parameter plane into six regions, as depicted by region $\mathcal{R}_1, \dots, \mathcal{R}_6$. Detailed phase diagrams associated with each region are given in Appendix II. In summary, \mathcal{R}_1 has either a monostable fixed point (one stable steady state) with low concentrations of LasI and RsaL for $a_3 \geq 0.55$ (Fig. 8a), or excitable solutions, as discussed in the previous section (Fig. 8b). In the narrow region \mathcal{R}_2 (see Fig. 4b for detail figure), bounded by SN_2 and sub-Hopf, three steady states arise, but only the lower steady state is stable (Fig. 8c and Fig. 8d). Excitable solutions are also possible. Region \mathcal{R}_3 is divided into four domains with different behaviour. Three steady states arise in which the upper and lower steady states are stable, but the middle state is unstable. In the first domain, the upper state in \mathcal{R}_3 is surrounded by an unstable limit cycle (Fig. 8e). A homoclinic bifurcation arises to give the second domain (Fig. 8f). Upon decreasing a_3 further in this region, a high stable spiral exists with a large domain of attraction (Fig. 8g and Fig. 8h). Region \mathcal{R}_4 is monostable with high concentrations of LasI and RsaL (Fig. 8i). In the area approaching the

cusp point \mathcal{R}_5 and \mathcal{R}_6 , two small distinct regions arise (see Fig. 4b). Region \mathcal{R}_5 provides stable oscillations and \mathcal{R}_6 bistability. In summary, the bifurcation lines divide the (a_1, a_3) -plane into regions of distinct solution behaviour: monostability (\mathcal{R}_1 and \mathcal{R}_4), bistability or excitation (\mathcal{R}_2 , \mathcal{R}_3 , and \mathcal{R}_6), and oscillatory \mathcal{R}_5 .

3.4 Travelling wave of a pulse in a linear chain of cells

We demonstrated the potential for the *las* system to act as a pulse generator for a single cell in the previous section. To illustrate how this effect may translate into cell-cell communication within a colony we consider a simplified linear chain of cells. The goal is to provide proof of principle that it is possible to set up a pulse train when the individual cells are coupled to the dynamics of their neighbours. For simplicity, we assume that diffusion across each cell membrane is such that the intracellular concentration of HSL is at a kinetic equilibrium (i.e. $\frac{dH_L}{dt} = 0$, as before) and, additionally, assume that there is a neighbourhood surrounding each cell where the extracellular concentration can be modelled simply by

$$\frac{dH_{ex}}{dt} = D_{HL}(H_L - H_{ex}) - \Delta H_{ex}, \quad (18)$$

where Δ is the isotropic loss of H_{ex} from this loosely defined neighbourhood. Writing in a dimensionless form using equations (11) and (14) we obtain

$$\frac{d\zeta}{d\tau} = \eta - D\zeta, \quad (19)$$

where

$$\zeta = H_{ex} \frac{\alpha_L \beta_L \beta_{HL}}{\gamma_{mL} D_{HL}^2 K_D^2}, \quad D = \Delta \frac{\gamma_{mL} D_{HL} K_D}{\alpha_L \beta_L \beta_{HL}}, \quad K_D = \frac{k_L^-}{k_L^+}. \quad (20)$$

Additionally, we construct a chain of otherwise identical systems, (η_i, ζ_i) , for $i = 1, 2, \dots$, coupled via the external concentrations of HSL in the local neighbourhood of each cell.

$$\frac{d\zeta_i}{d\tau} = \eta_i - D \left((1 - 2d)\zeta_i + d \sum_{j=\pm 1} (\zeta_i - \zeta_{i+j}) \right), \quad (21)$$

where ζ is the rescaled H_{ex} variable and d is now the fraction of the isotropic single loss that is retained in the single cell-cell interchange.

In this simplified system we can investigate whether it is possible for a single cell $i = 1$ to propagate a signal. To do this we trigger an arbitrary gain in the HSL, sufficient to trigger an excitable response from system $i = 1$, and examine the downstream impact. To do this we increase the levels of HSL by two orders of magnitude from the low steady state (QS off) for the equivalent lifetime of a single cell (corresponding to one dimensionless time unit) after the cells reach steady-state. Three types of solution are presented in Fig. 5,




Fig. 5: Pulse generation in the *las* system for single cells triggers a pulse train when the individual cells are coupled together. x - and y -axes represent time variation of external HSL concentration (ζ) corresponding to the pulse train, which consist of a linear chain of seven cells. Three types of solution are found when the coupling fraction d is varied: (a) $d = 0.03$, (b) $d = 0.1$, (c) $d = 0.3$.

illustrating a range of behaviour from no propagation of the signal for small coupling, propagation of a pulse for intermediate coupling, to propagation from a mutually supported high steady state (QS on) for large coupling.

4 Discussion

We have described a model of the QS system in *P. aeruginosa* by considering non-linear positive and negative feedback loops associated with the production of the synthase LasI and the regulator RsaL (Fagerlind et al 2005; Pearson et al 1997; Rampioni et al 2007b). These non-linear effects create the possibility of novel dynamical behaviours in the model. We have created a dimensionless set of equations to describe these behaviours and explored how the five dimensionless parameters affect the results whilst maintaining biological plausibility. Where possible we have taken parameters from the biological literature and this has led to significant deviation in our parameter choices from existing mathematical biology manuscripts, notably the work of Fagerlind et al (2005); Fagerlind (2008), which needs commenting upon. When the parameters in these articles, which subsequently have been adopted in other later texts, were compared to biological estimates (presented for example in Alon (2006)) they were found to be significantly different. For example the parameters used in (Fagerlind et al 2005) suggest that the typical lifetime of a transcription factor is of the order of seconds, when biological estimates typically describe transcription factors as stable proteins with lifetimes of the same order as the cellular turnover time, i.e. hours for *Pseudomonas aeruginosa*. Furthermore, it suggests mRNA molecules are *more* stable than the proteins they are translated to, which is clearly at odds with biological knowledge about the bursty nature of protein production (Xie et al 2008).

We are confident that our dimensionless parameters thus lie within a biologically plausible region, though we acknowledge that the effects we report here are sensitive to the values of these parameters. An additional complication is the possibility of other functional forms for the rate of transcription of mRNA from the *lasI* and *rsaL* genes as a function of the active forms of both LasR and RsaL. The form we present is one from a family of different choices that result in the same qualitative behaviour: our analysis, presented in the appendix, suggests that competitive binding between RsaL and LasR is a requirement for excitability but symmetrical effects, resulting in the negative action of RsaL on its own production, are not required (as is shown on many diagrams of the *las* system (Dockery and Keener 2001)). In the absence of specific biochemical data of this relationship at the intergenic region our analysis represents a significant step forward in incorporating biochemical knowledge in mathematical models of quorum sensing. There are biological instances where the binding in the intergenic region decouples and the *rsaL* and *lasI* genes are promoted and repressed independently, notably in the genetic isolates obtained from cystic fibrosis patients (Rampioni et al 2007b). In this case the

mathematical modelling of the system becomes trivial as the two systems are decoupled, and there is no possibility of excitation.

A survey of a realistic region of parameter space revealed a range of interesting numerical solution behaviour, such as limit cycles. The full equations, without approximations to reduce the number of equations, were also solved numerically and revealed similar solution behaviour (not shown). Continuation methods were employed for the reduced system in the a_1 - a_3 plane to track bifurcations of the system of co-dimension one and two. The parameters a_1 and a_3 represent the information outside and inside the cell, respectively. We found fold and Hopf bifurcations, both of co-dimension one. Furthermore, there are Bogdanov-Takens and generalized Hopf bifurcations, each of co-dimension two. For example, at a Bogdanov-Takens bifurcation, Hopf and fold curves in the plane intersect. However, most of the complex bifurcation structure is confined to a relatively small region of parameter space; the dominant behaviour is that of the fold bifurcation outside which are regions of excitable solutions.

We have demonstrated the potential for the *las* system to act as a pulse generator. As there is no explicit feedback from *rhl* to *las*, we may consider the *las* system as a black-box controller for the *rhl* system. We have demonstrated that this can lead to the generation of a pulse train when the individual cells are coupled by diffusion of the HSL signal molecule. These observations allow for a novel stigmergy (Gloag et al 2015) in the coupled *las/rhl* system; that of a *quorum memory*. The *las* system is coupled to the *rhl* system in two distinct ways; the active LasR transcription factor promotes the transcription of its counterpart in the *rhl* system, RhlR, and due to competitive binding on the RhlR molecule of the two homoserine lactones the presence of 3-oxo-C12-HSL acts to prevent the activation of RhlR by C4-HSL. This creates an interesting unreported effect in the combined system: what one might call a ‘handbraked acceleration’. Assuming the *rhl* system follows the same Lux-like dynamics of the *las* system the additional production of RhlR increases the likelihood of the system switching to its higher steady state, but this is tempered, or handbraked, by the presence of significant amounts of 3-oxo-C12-HSL preventing the activation of the rhlR system and its downstream consequences, notably rhamnolipid production. However, the effect of the pulses (the ‘revs’ of the ‘acceleration’) is still to increase the amount of RhlR, sustained due to its relative stability compared to the diffusing homoserine lactones, so that when the handbrake of 3-oxo-C12-HSL is removed the system has an increased likelihood of activating the *rhl* system. In effect the cells have been primed by the pulses and thus have a memory of experiencing higher densities of cells (or low diffusion regions (Redfield 2002)). Cells will lose memory only when they no longer experience a local environment rich in the 3-oxo-C12-HSL from the *las* system. Therefore, the competition between the two signal molecules, one produced by the *las* system and the other produced by the *rhl* system, may enable cells to trigger rhamnolipid production only when they are at the edge of an established aggregation. This suggests a previously unreported reason for the coupled nature of the *las* and *rhl* systems and a mechanism for how

they act in tandem to create a sophisticated control system for sociality and virulence in this important pathogenic organism.

Acknowledgements C. A. was financially supported by a PhD scholarship from RISTEK-DIKTI (Ministry of Research, Technology and Higher Education, Indonesia).

Appendix I

RsaL is thought to work as an inhibitor in the *las* system on *P. aeruginosa* and, therefore, the Michaelis-Menten equation can be applied. While there are a considerable number of studies about the QS system for *P. aeruginosa*, to the best knowledge of the authors, none of these studies provide sufficient biochemical evidence to determine whether the binding of RsaL in the intergenic region of the *las* system is competitive, uncompetitive, non-competitive, or mixed-competitive inhibition (using the standard biochemical descriptors). Here, we restrict attention to competitive, uncompetitive and non-competitive inhibition (mixed-competitive inhibition represents a combination of both competitive and uncompetitive inhibition, such that the RsaL inhibitor binds LasR/3O-C12-HSL or LasR/3O-C12-HSL+*lasI* genes with different affinities; mixed-competitive is the same as non-competitive inhibition if genes are bound with equal affinity). Furthermore, there is insufficient biochemical evidence whether the expression rates for *lasI* and *rsaL* genes are symmetric or not.

Competitive inhibition

Competitive binding occurs when both the LasR/3O-C12-HSL enzyme and RsaL inhibitor compete for binding to the active sites of the *lasI* gene (see Fig. 7a). Thus the governing equations result from a blockade of either positive or negative *lasI* gene interaction by either LasR/3O-C12-HSL or RsaL, respectively. Furthermore, this competitive inhibition can be classified into non-symmetric and symmetric binding based on expression rates for LasR/3O-C12-HSL to *lasI* and *rsaL* genes.

Due to the overlapping of binding sites, LasI is completely inactive when the RsaL inhibitor binds *lasI* genes. For the non-symmetric case, the affinity constant between LasR/3O-C12-HSL and *rsaL* is relatively small. Meanwhile, in the symmetric case the affinity constant between LasR/3O-C12-HSL and *rsaL* gene is identical to the affinity constant between LasR/3O-C12-HSL and the *lasI* gene.




Fig. 6: Binding types of inhibition: (a) competitive, (b) uncompetitive, and (c) non-competitive inhibition.




Fig. 7: Intersection between LasI and RsaL concentration nullclines with different types of inhibition: (a) non-symmetric competitive, (b) symmetric competitive, (c) non-symmetric uncompetitive, (d) symmetric uncompetitive, (e) non-symmetric non-competitive, and (f) symmetric non-competitive.

Non-symmetric competitive inhibition.

Here, the governing equations are

$$\frac{d\hat{I}_L}{dt} = \beta_L \frac{R_{LH}^2}{K_L^2(1 + S_L/K_{SL})^2 + R_{LH}^2} - \gamma_{mL}\hat{I}_L + \beta_{L0}, \quad (22)$$

$$\frac{d\hat{S}_L}{dt} = \beta_S \frac{R_{LH}}{K_S + R_{LH}} - \gamma_{mS}\hat{S}_L + \beta_{S0}. \quad (23)$$

Therefore, the non-dimensional model becomes

$$\frac{d\eta}{d\tau} = \frac{(\eta + a_1)^2}{a_2(1 + \xi)^2(\eta + a_1 + 1)^2 + (\eta + a_1)^2} - a_3\eta, \quad (24)$$

$$\frac{d\xi}{d\tau} = \frac{a_4(\eta + a_1)}{a_6(\eta + a_1 + 1) + (\eta + a_1)} - a_5\xi. \quad (25)$$

Symmetric competitive inhibition.

In this case, we find that

$$\frac{d\hat{I}_L}{dt} = \beta_L \frac{R_{LH}^2}{K_L^2(1 + S_L/K_{SL})^2 + R_{LH}^2} - \gamma_{mL}\hat{I}_L + \beta_{L0}, \quad (26)$$

$$\frac{d\hat{S}_L}{dt} = \beta_S \frac{R_{LH}^2}{K_L^2(1 + S_L/K_{SL})^2 + R_{LH}^2} - \gamma_{mS}\hat{S}_L + \beta_{S0}, \quad (27)$$

yielding the non-dimensional model

$$\frac{d\eta}{d\tau} = \frac{(\eta + a_1)^2}{a_2(1 + \xi)^2(\eta + a_1 + 1)^2 + (\eta + a_1)^2} - a_3\eta, \quad (28)$$

$$\frac{d\xi}{d\tau} = \frac{a_4(\eta + a_1)^2}{a_2(1 + \xi)^2(\eta + a_1 + 1)^2 + (\eta + a_1)^2} - a_5\xi. \quad (29)$$

Uncompetitive inhibition

Uncompetitive binding occurs when the RsaL inhibitor binds to a site which only becomes available after the LasR/3O-C12-HSL enzyme has bound to the active site of the *lasI* gene. It can also be classified into non-symmetric and symmetric binding types, as indicated below.

Non-symmetric uncompetitive inhibition.

Here,

$$\frac{d\hat{I}_L}{dt} = \beta_L \frac{R_{LH}^2}{K_L^2 + R_{LH}^2 (1 + S_L/K'_{SL})^2} - \gamma_{mL} \hat{I}_L + \beta_{L0}, \quad (30)$$

$$\frac{d\hat{S}_L}{dt} = \beta_S \frac{R_{LH}}{K_S + R_{LH}} - \gamma_{mS} \hat{S}_L + \beta_{S0}, \quad (31)$$

so that the non-dimensional model becomes

$$\frac{d\eta}{d\tau} = \frac{(\eta + a_1)^2}{a_2 (\eta + a_1 + 1)^2 + (\eta + a_1)^2 (1 + \xi')^2} - a_3 \eta, \quad (32)$$

$$\frac{d\xi}{d\tau} = \frac{a_4 (\eta + a_1)}{a_6 (\eta + a_1 + 1) + (\eta + a_1)} - a_5 \xi, \quad (33)$$

where ξ' is a rescaled ξ .

Symmetric uncompetitive inhibition.

For this case,

$$\frac{d\hat{I}_L}{dt} = \beta_L \frac{R_{LH}^2}{K_L^2 + R_{LH}^2 (1 + S_L/K'_{SL})^2} - \gamma_{mL} \hat{I}_L + \beta_{L0}, \quad (34)$$

$$\frac{d\hat{S}_L}{dt} = \beta_S \frac{R_{LH}^2}{K_L^2 + R_{LH}^2 (1 + S_L/K'_{SL})^2} - \gamma_{mS} \hat{S}_L + \beta_{S0}, \quad (35)$$

such that the non-dimensional model is

$$\frac{d\eta}{d\tau} = \frac{(\eta + a_1)^2}{a_2 (\eta + a_1 + 1)^2 + (\eta + a_1)^2 (1 + \xi')^2} - a_3 \eta, \quad (36)$$

$$\frac{d\xi}{d\tau} = \frac{a_4 (\eta + a_1)^2}{a_2 (\eta + a_1 + 1)^2 + (\eta + a_1)^2 (1 + \xi')^2} - a_5 \xi. \quad (37)$$

Here, K'_{SL} represents the dissociation constant for the RsaL inhibition of binding of *lasI* genes to the LasR/3O-C12-HSL enzyme. It is different to K_{SL} , which represents the dissociation constant of inhibitor RsaL on *lasI* genes.

Non-competitive inhibition

Non-competitive inhibition occurs when the RsaL inhibitor can bind to the free *lasI* gene or the *lasI*-bound LasR/3O-C12-HSL enzyme. As explained above, the non-competitive inhibition form is a special case of mixed-competitive inhibition, where it binds to the target with two equal inhibition constants.

Non-symmetric non-competitive inhibition.

Here,

$$\frac{d\hat{I}_L}{dt} = \beta_L \frac{R_{LH}^2}{K_L^2 + R_{LH}^2} \frac{1}{(1 + S_L/K_{SL})^2} - \gamma_{mL}\hat{I}_L + \beta_{L0}, \quad (38)$$

$$\frac{d\hat{S}_L}{dt} = \beta_S \frac{R_{LH}}{K_S + R_{LH}} - \gamma_{mS}\hat{S}_L + \beta_{S0}, \quad (39)$$

providing the non-dimensional model

$$\frac{d\eta}{d\tau} = \frac{(\eta + a_1)^2}{a_2 (\eta + a_1 + 1)^2 + (\eta + a_1)^2} \frac{1}{(1 + \xi)^2} - a_3\eta, \quad (40)$$

$$\frac{d\xi}{d\tau} = \frac{a_4 (\eta + a_1)}{a_6 (\eta + a_1 + 1) + (\eta + a_1)} - a_5\xi. \quad (41)$$

Symmetric non-competitive inhibition.

Here, the governing equations are

$$\frac{d\hat{I}_L}{dt} = \beta_L \frac{R_{LH}}{K_L + R_{LH}} \frac{1}{(1 + S_L/K_{SL})} - \gamma_{mL}\hat{I}_L + \beta_{L0}, \quad (42)$$

$$\frac{d\hat{S}_L}{dt} = \beta_S \frac{R_{LH}}{K_L + R_{LH}} \frac{1}{(1 + S_L/K_{SL})} - \gamma_{mS}\hat{S}_L + \beta_{S0}, \quad (43)$$

and the non-dimensional model is


$$\frac{d\eta}{d\tau} = \frac{(\eta + a_1)^2}{a_2 (\eta + a_1 + 1)^2 + (\eta + a_1)^2} \frac{1}{(1 + \xi)^2} - a_3\eta, \quad (44)$$

$$\frac{d\xi}{d\tau} = \frac{a_4 (\eta + a_1)^2}{a_2 (\eta + a_1 + 1)^2 + (\eta + a_1)^2} \frac{1}{(1 + \xi)^2} - a_5\xi. \quad (45)$$

Using the simplification steps highlighted in the main text, we determine appropriate nullclines for each binding type (see Fig. 7). The equations resulting from non-symmetric and symmetric competitive behaviour are presented in Fig. 7a and Fig. 7b, revealing the same qualitative behaviour as for the system in the main text. This suggests that competitive binding between RsaL and LasR/3O-C12-HSL with or without symmetry is sufficient for excitable behaviour. Therefore, although (Rampioni et al 2007b) suggest there may be negative feedback from RsaL to its own production, this symmetrical binding is not required for excitable behaviour (c.f. general diagrams of the *las* system in Van Delden and Iglewski 1998; De Kievit et al 2002; Dockery and Keener 2001; Fagerlind et al 2005; Schaadt et al 2013).

Appendix II

Further investigation of the structure in Fig. 4 finds that the (a_1, a_3) parameter plane is divided by bifurcation lines into six regions. The six corresponding phase diagrams of the dynamic solution behaviour for LasI and RsaL are illustrated below. In order to get clear understanding of qualitative behaviour for region \mathcal{R}_1 to \mathcal{R}_4 , we fix parameter values for a_1, a_2, a_4, a_5 (see Table 3) and vary a_3 from a high to a low value. For figure 8a to 8i, $a_3 = 0.6, 0.3, 0.281, 0.28, 0.278, 0.275, 0.27, 0.2$, and 0.1 , respectively.



In the region \mathcal{R}_1 , the system has a single low stable steady state, i.e. with 0 eigenvalues with positive real part. In this region, solutions have two qualitatively different behaviours (see Fig. 8a and 8b). The difference between those two behaviours is that the large perturbation in Fig. 8b has a large excursion around the phase plane before it reaches the stable fixed point. In the region \mathcal{R}_2 after a saddle-node bifurcation, system has 0, 1, and 2 eigenvalues with positive real part in the low, middle and high steady states, respectively. Thus it consists of one stable low fixed point, one saddle and one unstable fixed




Fig. 8: Phase diagram corresponding to the labelled regions in Figure 4 (figure a and b in region \mathcal{R}_1 ; figure c and d in region \mathcal{R}_2 ; figure e, f, g, and h in region \mathcal{R}_3 ; figure i in region \mathcal{R}_4 ; figure j in region \mathcal{R}_5). Stable (unstable) steady states are denoted by solid (open) circles.

point. There is a change from unstable fixed point to unstable spiral (see Fig. 8c and 8d).

In the region \mathcal{R}_3 after a sub-critical Hopf bifurcation, the system has 0, 1, and 0 eigenvalues with positive real part in the low, middle and high states, respectively. In this region, the system has four different qualitative behaviours. Upon decreasing a_3 , the journey of qualitative dynamical behaviour can be seen clearly in Fig 8e, 8f, 8g, and 8h. First, the unstable limit cycle coexists with one saddle and two stable steady states (see Fig. 8e). Then, a homoclinic bifurcation occurs (a collision between the unstable limit cycle and the saddle; see Fig. 8e) resulting in a high state stable spiral with large basin of attraction (see Fig. 8f). Lastly, the saddle node moves closer to the low stable fixed point (see Fig. 8h) eventually colliding via a saddle-node bifurcation to yield region \mathcal{R}_4 . This system has one high state fixed point with 0 eigenvalues with positive real part (see Fig. 8i).

In Fig. 4 we find two other regions, \mathcal{R}_5 and \mathcal{R}_6 . In region \mathcal{R}_5 , a stable limit cycle and one unstable steady state exists (see Fig. 8j, $a_1 = 0.03298$, $a_2 = 0.3$, $a_3 = 0.3226$, $a_4 = 0.4$ and $a_5 = 0.075$). In region \mathcal{R}_6 the system has one unstable and two stable fixed points, but the region is very small as it is close to the cusp where the two saddle-node bifurcations collide.

Appendix III




Fig. 9: (a) Time dependent solutions from the full system of equations before systematic reduction, corresponding to the excitable trajectory of LasI. (b) Excitable dynamics in the LasI-RsaL phase plane

In this paper, the reduced model of the QS system for *P. aeruginosa* explored in the main text reveals the potential for excitable pulse generation in the *las* subsystem. However, numerical solution of the full system of seven differential equations reveals very similar solution behaviour, including excitable

pulse generation. Excitation occurs when there is a sufficiently large perturbation of (*N*-(3-oxododecanoyl)-HSL) leading to pulse production of LasI, pushing the system to the right in the LasI-RsaL projected phase plane. Typically, this is followed by a large excursion around phase space before returning to the stable fixed point (see Fig. 9a and 9b). Other solutions are also possible analogous to those in the reduced system. These results provide confidence that the behaviour of the full system has been adequately captured by the reduced model.

References

- Alon U (2006) An Introduction to System Biology: Design Principles of Biological Circuits. Chapman and Hall or CRC Press, Boca Raton, FL
- Alves F, Dilo R (2005) A simple framework to describe the regulation of gene expression in prokaryotes. *Comptes Rendus - Biologies* 328(5):429–444
- Angeli D, Ferrell Jr JE, Sontag ED (2004) Detection of multistability, bifurcations, and hysteresis in a large class of biological positive-feedback systems. *Proceedings of the National Academy of Sciences of the United States of America* 101(7):1822–1827
- Chen F, Chen CC, Riadi L, Ju LK (2004) Modeling rhl quorum-sensing regulation on rhamnolipid production by *pseudomonas aeruginosa*. *Biotechnol Prog* 20(5):1325–31
- Daniels R, Vanderleyden J, Michiels J (2004) Quorum sensing and swarming migration in bacteria. *FEMS Microbiol Rev* 28(3):261–89
- De Kievit TR, Kakai Y, Register JK, Pesci EC, Iglewski BH (2002) Role of the *pseudomonas aeruginosa* las and rhl quorum-sensing systems in rhlI regulation. *FEMS Microbiology Letters* 212(1):101–106
- Dockery JD, Keener JP (2001) A mathematical model for quorum sensing in *pseudomonas aeruginosa*. *Bull Math Biol* 63(1):95–116, DOI 10.1006/bulm.2000.0205
- Dubern JF, Diggle SP (2008) Quorum sensing by 2-alkyl-4-quinolones in *pseudomonas aeruginosa* and other bacterial species. *Mol Biosyst* 4(9):882–8
- Fagerlind MG (2008) Mathematical modelling of bacterial quorum sensing and biofilm formation
- Fagerlind MG, Rice SA, Nilsson P, Harlen M, James S, Charlton T, Kjelleberg S (2003) The role of regulators in the expression of quorum-sensing signals in *pseudomonas aeruginosa*. *J Mol Microbiol Biotechnol* 6:88–100
- Fagerlind MG, Nilsson P, Harln M, Karlsson S, Rice SA, Kjelleberg S (2005) Modeling the effect of acylated homoserine lactone antagonists in *pseudomonas aeruginosa*. *BioSystems* 80:201–213, DOI 10.1016/j.biosystems.2004.11.008
- Fuqua C, Parsek MR, Greenberg EP (2001) Regulation of gene expression by cell-to-cell communication: acyl-homoserine lactone quorum sensing. *Annu Rev Genet* 35:439–68, DOI 10.1146/annurev.genet.35.102401.090913
- Glick D, Barth S, Macleod KF (2010) Autophagy: Cellular and molecular mechanisms. *Journal of Pathology* 221(1):3–12
- Gloag ES, Turnbull L, Whitchurch CB (2015) Bacterial stigmergy: an organising principle of multicellular collective behaviours of bacteria. *Scientifica* 2015:8
- Hat B, Kochaczyk M, Bogda MN, Lipniacki T (2016) Feedbacks, bifurcations, and cell fate decision-making in the p53 system. *PLoS Computational Biology* 12(2)
- Ishihama A, Kori A, Koshio E, Yamada K, Maeda H, Shimada T, Makinoshima H, Iwata A, Fujitac N (2014) Intracellular concentrations of 65 species of transcription factors with known regulatory functions in *escherichia coli*. *Journal of Bacteriology* 196(15):2718–2727
- James S, Nilsson P, James G, Kjelleberg S, Fagerstrom T (2000) Luminescence control in the marine bacterium *vibrio fischeri*: an analysis of the dynamics of lux regulation. *J Mol Biol* 296:1127–1137, DOI doi:10.1006/jmbi.1999.3484

- Kohler T, Curty LK, Barja F, Van Delden C, Pechere JC (2000) Swarming of *pseudomonas aeruginosa* is dependent on cell-to-cell signaling and requires flagella and pili. *Journal of Bacteriology* 182(21):5990–5996
- Latifi A, Winson MK, Foglino M, Bycroft BW, Stewart GS, Lazdunski A, Williams P (1995) Multiple homologues of luxR and luxI control expression of virulence determinants and secondary metabolites through quorum sensing in *pseudomonas aeruginosa* pao1. *Mol Microbiol* 17(2):333–43
- Latifi A, Foglino M, Tanaka K, Williams P, Lazdunski A (1996) A hierarchical quorum-sensing cascade in *pseudomonas aeruginosa* links the transcriptional activators lasR and rhIR (vsmR) to expression of the stationary-phase sigma factor rpos. *Mol Microbiol* 21(6):1137–46
- Miller MB, Bassler BL (2001) Quorum sensing in bacteria. *Annu Rev Microbiol* 55:165–99, DOI 10.1146/annurev.micro.55.1.165
- Murray JD (1993) *Mathematical Biology*, 2nd ed. Springer-Verlag, New York
- O’Loughlin CT, Miller LC, Siryaporn A, Drescher K, Semmelhack MF, Bassler BL (2013) A quorum-sensing inhibitor blocks *pseudomonas aeruginosa* virulence and biofilm formation. *Proceedings of the National Academy of Sciences of the United States of America* 110(44):17,981–17,986
- Pai A, You L (2009) Optimal tuning of bacterial sensing potential. *Molecular Systems Biology* 5
- Pearson JP, Pesci EC, Iglewski BH (1997) Roles of *pseudomonas aeruginosa* las and rhl quorum-sensing systems in control of elastase and rhamnolipid biosynthesis genes. *J Bacteriol* 179(18):5756–67
- Pesci EC, Pearson JP, Seed PC, Iglewski BH (1997) Regulation of las and rhl quorum sensing in *pseudomonas aeruginosa*. *Journal of Bacteriology* 179(10):3127–3132
- Pfeuty B, Kaneko K (2009) The combination of positive and negative feedback loops confers exquisite flexibility to biochemical switches. *Physical Biology* 6(4)
- Poignard C (2014) Inducing chaos in a gene regulatory network by coupling an oscillating dynamics with a hysteresis-type one. *Journal of Mathematical Biology* 69(2):335–368
- Rampioni G, Polticelli F, Bertani I, Righetti K, Venturi V, Zennaro E, Leoni L (2007a) The *pseudomonas* quorum-sensing regulator rsal belongs to the tetrahelical superclass of h-t-h proteins. *J Bacteriol* 189(5):1922–30
- Rampioni G, Schuster M, Greenberg EP, Bertani I, Grasso M, Venturi V, Zennaro E, Leoni L (2007b) Rsal provides quorum sensing homeostasis and functions as a global regulator of gene expression in *pseudomonas aeruginosa*. *Mol Microbiol* 66(6):1557–65
- Raychaudhuri A, Jerga A, Tipton PA (2005) Chemical mechanism and substrate specificity of rhII, an acylhomoserine lactone synthase from *pseudomonas aeruginosa*. *Biochemistry* 44(8):2974–2981
- Redfield RJ (2002) Is quorum sensing a side effect of diffusion sensing? *Trends Microbiol* 10(8):365–70
- Schaadt NS, Steinbach A, Hartmann RW, Helms V (2013) Rule-based regulatory and metabolic model for quorum sensing in *p. aeruginosa*. *BMC Syst Biol* 7:81
- Schuster M, Greenberg EP (2006) A network of networks: Quorum-sensing gene regulation in *pseudomonas aeruginosa*. *International Journal of Medical Microbiology* 296(2-3):73–81
- Shadel BS, Baldwin TO (1991) The *vibrio fischeri* luxR protein is capable of bidirectional stimulation of transcription and both positive and negative regulation of the luxR gene. *Journal of Bacteriology* 173(2):568–574
- Shih PC, Huang CT (2002) Effects of quorum-sensing deficiency on *pseudomonas aeruginosa* biofilm formation and antibiotic resistance. *Journal of Antimicrobial Chemotherapy* 49(2):309–314
- Shrout JD, Chopp DL, Just CL, Hentzer M, Givskov M, Parsek MR (2006) The impact of quorum sensing and swarming motility on *pseudomonas aeruginosa* biofilm formation is nutritionally conditional. *Molecular Microbiology* 62(5):1264–1277
- Singh D, Manjunatha BS, Meena RP, Taria S (2015) Bacterial biofilm and quorum sensing. *Annals of Biology* 31(1):13–17
- Sitnikov DM, Schineller JB, Baldwin TO (1995) Transcriptional regulation of bioluminescence genes from *vibrio fischeri*. *Molecular Microbiology* 17(5):801–812

- Skindersoe ME, Alhede M, Phipps R, Yang L, Jensen PO, Rasmussen TB, Bjarnsholt T, Tolker-Nielsen T, Hiby N, Givskov M (2008) Effects of antibiotics on quorum sensing in *pseudomonas aeruginosa*. *Antimicrobial Agents and Chemotherapy* 52(10):3648–3663
- Song H, Smolen P, Av-Ron E, Baxter DA, Byrne JH (2007) Dynamics of a minimal model of interlocked positive and negative feedback loops of transcriptional regulation by camp-response element binding proteins. *Biophysical Journal* 92(10):3407–3424
- Van Delden C, Iglewski BH (1998) Cell-to-cell signaling and *pseudomonas aeruginosa* infections. *Emerging Infectious Diseases* 4(4):551–560
- Viretta AU, Fussenegger M (2004) Modeling the quorum sensing regulatory network of human-pathogenic *pseudomonas aeruginosa*. *Biotechnology Progress* 20(3):670–678
- Welch M, Todd DE, Whitehead NA, McGowan SJ, Bycroft BW, Salmond GPC (2000) N-acyl homoserine lactone binding to the carr receptor determines quorum-sensing specificity in *erwinia*. *EMBO Journal* 19(4):631–641
- Xie XS, Choi PJ, Li GW, Nam KL, Lia G (2008) Single-molecule approach to molecular biology in living bacterial cells. In: *Annual Review of Biophysics*, vol 37, pp 417–444