



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/116776/>

Version: Accepted Version

Article:

Giannoudis, PV, Giannoudis, VP and Horwitz, DS (2017) Time to think outside the box: 'Prompt-Individualised-Safe Management' (P.R.I.S.M.) should prevail in patients with multiple injuries. *Injury*, 48 (7). pp. 1279-1282. ISSN: 0020-1383

<https://doi.org/10.1016/j.injury.2017.05.026>

© 2017 Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.


Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Accepted Manuscript

Title: Time to think outside the box:
'Prompt-Individualised-Safe Management' (P.R.I.S.M.)
should prevail in patients with multiple injuries

Authors: P.V. Giannoudis, Vasileios P. Giannoudis



PII: S0020-1383(17)30351-0
DOI: <http://dx.doi.org/doi:10.1016/j.injury.2017.05.026>
Reference: JINJ 7252

To appear in: *Injury, Int. J. Care Injured*

Author: Daniel S. Horwitz

PII: S0020-1383(17)30351-0
DOI: <http://dx.doi.org/doi:10.1016/j.injury.2017.05.026>
Reference: JINJ 7252

To appear in: *Injury, Int. J. Care Injured*

Please cite this article as: Horwitz Daniel S. Time to think outside the box: 'Prompt-Individualised-Safe Management' (P.R.I.S.M.) should prevail in patients with multiple injuries. *Injury* <http://dx.doi.org/10.1016/j.injury.2017.05.026>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Time to think outside the box: 'Prompt-Individualised-Safe Management' (P.R.I.S.M.) should prevail in patients with multiple injuries

Peter V Giannoudis, Vasileios Giannoudis, Daniel S Horwitz

P.V. Giannoudis MD, FACS, FRCS

Professor, Academic Department of Trauma & Orthopaedics, School of Medicine, University of Leeds

Vasileios P. Giannoudis MSc

Academic Department of Trauma & Orthopaedics, School of Medicine, University of Leeds

Daniel S Horwitz MD

Chief Orthopaedic Trauma Geisinger Health System, Danville, Pennsylvania

* No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article. No funds were received in support of this study.

Corresponding Author Address:

Professor Peter V Giannoudis MB, BSc, MD, FACS, FRCS(Eng)
Academic Department Trauma & Orthopaedic Surgery
School of Medicine
University of Leeds
Leeds, UK

Email: pgiannoudi@aol.com

Tel: +44-113-3922750

Fax: +44-13-3923290

Keywords: polytrauma, management, Damage control orthopaedics, Early total care, Early appropriate care, prompt individualised safe management.

The management of polytrauma patients has undergone a revolution over the past 80 years. In its infancy in the 1940s, orthopaedic management of polytrauma patients predominately involved the use of traction, casts and splints.

The levels of morbidity and mortality which ensued following long bone fractures was problematic and there was a common consensus that “The patient was too sick to operate on” and surgeons were fearful of operating on patients.

The 1970s brought the first shift in the treatment paradigm of polytrauma patients with long bone fractures. On-going advances in osteosyntheses yielded positive results with reductions in pulmonary complications and time to mobilisation⁽¹⁾. In 1985 Johnson et al. published a case series of 132 patients investigating the incidence of adult respiratory distress syndrome (ARDS). His results showed that patients with an ISS of greater than 40 who did not receive early operative fracture stabilisation within the first 24 hours had a 75% chance of developing ARDS compared to a 17% chance in those who did ($p < 0.001$).⁽²⁾ Subsequently, several other retrospective studies were published which confirmed Johnson’s findings^(3,4), and Bone’s publication a few years later being the first prospective randomised controlled trial built the foundations of Early Total Care (ETC).⁽⁵⁾ His study showed reduced rates of fat embolism, ARDS and sepsis related mortality in patients who underwent definitive femoral fracture stabilisation within 24 hours. Delay in stabilisation after that time resulted in a five times greater risk of adult respiratory distress syndrome (ARDS).⁽⁵⁾ Not surprisingly therefore, the argument became “patients are too sick not to be operated on”.

The change in treatment paradigm was successful and many patients benefit from this approach. However, in subsequent studies, it was noted that ETC itself may be detrimental to patients who were most severely injured or had associated chest^(6,7) or head injuries^(8,9). These findings were particularly pertinent from the trauma centre in Hannover, Germany. This paved the way to the development of Damage Control Orthopaedics (DCO).⁽¹⁰⁾

DCO is a term originally used in naval and maritime industry implemented in emergency situations to prevent the sinking of ships through the implementation of several measures. The principles of DCO were transferred to medicine and first used in general surgery and were subsequently adapted to orthopaedics for the management of extremity injuries including pelvic fractures.

DCO includes 4 stages of management: ^(11,12)

1. Performance of life saving procedures as deemed necessary.
2. Control of haemorrhage, stabilisation of long bone fracture often through use of external fixators and the management of soft tissue injuries if present.
3. Vital organ monitoring and aggressive resuscitation in the intensive care setting
4. Definitive fracture fixation.

Scalea et al. first described the application of DCO in a case series of 43 patients (median ISS-27) with femoral fractures. ⁽¹³⁾ These patients underwent temporary external fixation for a median of 4 days before staged nailing occurred. Moving on, as our knowledge of the immunological system improved, the 'second hit theory' was proposed, ⁽¹⁴⁾ following operative interventions in patients with polytrauma. It was noted that trauma itself was the 'first hit' and the subsequent operation that was performed the 'second hit' inducing further increase in several molecular mediators (ie. IL 6, TNF α and IL 1 β), markers of immune reactivity, demonstrating a surgical stress reaction which when becoming exaggerated could lead to ARDS/MODS due to excessive systemic inflammation. ⁽¹⁵⁾

The issue then became in identifying which of these polytrauma patients would be the ones who lacked the physiological capacity for further significant insult, such as the complete fixation of all fractures within 24 hours (ETC). The review published in 2005 by Pape et al. attempted to classify patients into 4 categories: Stable (grade I); Borderline (grade II), Unstable (Grade III) and in Extremis (Grade IV) physiological stage. The authors used physiological parameters to assess the four pathophysiologic cascades seen in polytrauma (shock, coagulation abnormalities, hypothermia and soft tissue injuries). ⁽¹⁶⁾

For both the 'unstable' and 'in extremis' groups, consisting of patients with the highest risk of sustaining complications post surgery, DCO was recommended. Conversely, patients who are physiologically stable are eligible for ETC, which allows for the total fixation of fractures in one theatre setting. The 'Borderline' patient group, also known as group 'at risk', continues to cause debate amongst the orthopaedic community. This subset's response to surgery could be unpredictable

as their physiological parameters are not grossly abnormal and they may be susceptible to the 'second hit' phenomenon. Ultimately, this should come down to surgeons' experience and discretion in opting to follow DCO or ETC.⁽¹⁶⁾

In the Adams Cowley shock trauma centre in Maryland the authors recommend DCO in patients with the following characteristics: closed head injury; poor response to resuscitation in the first 12 hours and poor respiratory status at the time of fracture treatment.⁽¹⁷⁾

Overall, the use of DCO has its benefits for any patient not deemed to be physiologically 'stable'. Studies have also shown its successful applications in natural disasters in being able to deal with and turnover high volumes of patients.⁽¹⁸⁾

DCO also poses some negative aspects, predominately the increased risk of infection associated with both the external fixator and the subsequent second visit to theatre.⁽¹⁹⁾ Furthermore, there is still no clear consensus on when the definitive fixation should be performed. Several different time points are suggested according to the long bone affected (femur 5-7 days, pelvis 5-9 days).⁽²⁰⁾ Several studies in Germany suggest a minimal delay up until day 5 where decrease hepatic and pulmonary complications were noted in a set of over 4,000 patients.⁽²¹⁾ Of note, there has been limited Level I RCT literature published with adequate power investigating this.

Most recently, Vallier's research group have proposed the application of Early Appropriate Care (EAC) following resuscitative efforts in the polytrauma patient.⁽²²⁾ This approach can be applied through the evaluation of three physiological parameters: lactate, acid base excess and pH as long as initial acidosis improved to at least one of the following parameters: lactate is <4 , pH is ≥ 7.25 or base excess is ≥ 5.5 . According to the authors, you can proceed to the definitive management of mechanically unstable fractures of the femur, pelvis and acetabulum and thoracolumbar spine in the form of early total care within 36 hours. However, DCO is only recommended for those who do not respond within the first eight hours to resuscitation. The authors believe that this protocol is readily and easily applicable in most trauma centres, it is affordable and can provide continued guidance to the surgeons of a patient's physiological status allowing for decisions regarding operations to be made.⁽²³⁾ Furthermore, in privatised healthcare systems application

of EAC can increase profitability margins and allow better allocation of trauma services.⁽²⁴⁾ In terms of patient outcomes Vallier et al. have demonstrated a reduction in patient complications and a shorter length of hospital stay since the application of this protocol. Their study investigating EAC of axial and femoral fractures in multiply injured patients yielded favourable results in using the EAC algorithm.⁽²⁵⁾ There were no differences noted in the resuscitative needs of the elderly cohort (≥ 60 years old) and the 2 younger groups. Furthermore, the complication rates of the elderly cohort (16.2%) were similar to those of the younger one (15.8 %). This study was underpowered and the authors have acknowledged this so further studies with larger sample sizes are required to validate the use of EAC in patients whose physiological parameters are distorted.

Lately, Pape et al. raised concerns about EAC and its appropriateness for certain subsets of patients who suffer from multiple traumatic injuries.⁽²⁶⁾ The 3 parameters of interest suggested by Vallier et al. (lactate, PH and base excess) can be distorted in patients who are diabetic, in the elderly, and those suffering from renal failure and therefore the true physiological profile of these patients may be masked. Even the type of fluid received during the resuscitative attempts can also influence lactate levels. More over, Pape et al highlighted that the protocol does not state the type of fluids used. Furthermore, when reaching to the conclusion of using the above 3 parameters alterations were made weighting of several parameters recommended in Pape et al. publication of 2005.⁽¹⁶⁾ The differing weighting statistically would increase the likelihood of producing improvements in the findings of their protocol. Finally, Pape et al raised questions concerning the data originally available in Vallier's research group. Several parameters assessing coagulopathy, clotting factors, circulatory and pulmonary functions were missing.

Pape et al went on to suggest the so called 'Safe Definitive Surgery approach'.⁽²⁶⁾ According to this strategy, the three subgroups of polytrauma patients, borderline, unstable, extremis would undergo initial assessment in the Emergency room in terms of their physiological state. Parameters for the initial physiological assessment and resuscitation requirements include 'multiple end points'. If the patient has been

declared as stable, safe definitive treatment can be applied. Alternatively, if the patient continues to be unstable or extremis, DCO surgery is recommended. After 24hrs reassessment is necessary examining coagulation, fluid balance, lung function and vasopressor need. If the patient is deemed to be 'stable' then again, the surgeon can proceed with safe definitive surgery. On the other hand, if the condition of the patient remains borderline repeated daily assessment is necessary until the patient will be declared stable and fit for definitive surgery. ⁽²⁶⁾

Overall, it is clear that a debate is ongoing amongst orthopaedic trauma surgeons as to whether one should treat a multiply injured patient with ETC? DCO? or EAC? Each strategy assumes specific indications and contraindications. However, the time has come to think outside the box and be less opinionated and leave behind us disagreements and conflicts. We should stop arguing or defending each one of these treatment modalities. Our decision making process should be based on the principles of doing no 'further harm to the patient', intervening promptly, and utilizing the concept of individualised/personalised medicine. When we manage a polytrauma patient, we should apply the dogma of 'Prompt-Individualised-Safe Management' (P.R.I.S.M.), (**Figure 1**). Our philosophy is based on the understanding that every patient responds in a different way to the same degree of injury, every patient has a different genetic constitution, and the fact that the trauma service provision and resources including man power are dissimilar from country to country. In trying to categorise polytrauma patients into specific pathways we feel that the importance of patient specific treatment may be overlooked. It doesn't matter which treatment algorithm you decide to follow as long as it is in-keeping with the specific clinical and physiological parameters the patient in front of you presents with. Ultimately we aim to do no further harm as surgeons and the clear sole aim should be to save the patient's life and to apply the differing strategies available in the patient's best interest. Therefore, we should stop discussing ETC, DCO, EAC and we should talk, implement the philosophy of P.R.I.S.M. This should become the new dogma of treating polytrauma patients. Out of this one should direct the patient to what type of treatment fits best, remain flexible to this direction in a continuous manner.

In terms of the specific criteria required to assess the physiological state of the patient, each unit should utilise whatever they have available in their armamentarium and should apply the in house protocol that they have developed. We acknowledge that not every unit can measure markers of immune reactivity such as IL-6, but they can utilise parameters that have been described over the years (lactate, acid base balance, PH, lung function, blood pressure, pulse rate, haemoglobin, etc), consider the presence of injury patterns, comorbidities and the age of the patient. All these should be taken into account to ascertain a complete picture of the state of the patient and then decide what is the safest way to proceed in an individualised/personalised manner. ⁽²⁷⁻³⁰⁾ Such an approach, should be established as an institutional protocol in order to simplify each individual evaluation and to standardize criteria best applied based on each unit's capabilities and the injury patterns involved.

It should not be forgotten that the initial factors which were evaluated can easily and rapidly change, related to blood loss and the quality of ongoing resuscitation. It is therefore crucial that the orthopaedic surgeon revisit the 'in house chosen' physiological assessment parameters frequently particularly during any extended surgery during ETC, coordinate with anesthesia, and be ready to revert to DCO as needed. Of interest, analysis of in house results (data not shown) failed to prove that the protocol of EAC was valid for our polytrauma patient cohort.

Safety of the patient remains our priority and direct comparisons between different health care systems in polytrauma protocols should become constructive not destructive. We therefore suggest that the dogma of 'PRISM' should be considered as a unified concept to allow surgeons to speak the same language, manage the patient promptly (no timeline cut offs ie 24 and/or 36 hours) and consider an individualised patient approach minimising the risk of unpredicted complications and mortality.

Conflict of Interest:

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Acknowledgements:


The authors acknowledge to contribution of Mr Matt Smith for the development of the acronym PR.I.S.M.

References

1. Seibel R, LaDuca J, Hassett JM, Babikian G, Mills B, Border DO, et al. Blunt multiple trauma (ISS 36), femur traction, and the pulmonary failure-septic state. *Ann Surg*. 1985 Sep;202(3):283–95.
2. Johnson KD, Cadambi A, Seibert GB. Incidence of adult respiratory distress syndrome in patients with multiple musculoskeletal injuries: effect of early operative stabilization of fractures. *J Trauma*. 1985 May;25(5):375–84.
3. Behrman SW, Fabian TC, Kudsk KA, Taylor JC. Improved outcome with femur fractures: early vs. delayed fixation. *J Trauma*. 1990 Jul;30(7):792–7; discussion 797–8.
4. Charash WE, Fabian TC, Croce MA. Delayed surgical fixation of femur fractures is a risk factor for pulmonary failure independent of thoracic trauma. *J Trauma*. 1994 Oct;37(4):667–72.
5. Bone LB, Johnson KD, Weigelt J, Scheinberg R. Early versus delayed stabilization of femoral fractures. A prospective randomized study. *J Bone Joint Surg Am*. 1989 Mar;71(3):336–40.
6. Goris RJ, Gimbrère JS, van Niekerk JL, Schoots FJ, Booy LH. Early osteosynthesis and prophylactic mechanical ventilation in the multitrauma patient. *J Trauma*. 1982 Nov;22(11):895–903.
7. Pape HC, Auf'm Kolk M, Paffrath T, Regel G, Sturm JA, Tscherne H. Primary intramedullary femur fixation in multiple trauma patients with associated lung contusion—a cause of posttraumatic ARDS? *J Trauma*. 1993 Apr;34(4):540–7; discussion 547–8.
8. Townsend RN, Lheureau T, Protech J, Riemer B, Simon D. Timing fracture repair in patients with severe brain injury (Glasgow Coma Scale score <9). *J Trauma*. 1998 Jun;44(6):977–82; discussion 982–3.
9. Jaicks RR, Cohn SM, Moller BA. Early fracture fixation may be deleterious after head injury. *J Trauma*. 1997 Jan;42(1):1–5; discussion 5–6.
10. Roberts CS, Pape HC, Jones AL, Malkani AL, Rodriguez JL, Giannoudis PV. Damage control orthopaedics: evolving concepts in the treatment of patients who have sustained orthopaedic trauma. *Instr Course Lect*. 2005;54:447–62.
11. Giannoudis PV. Surgical priorities in damage control in polytrauma. *J Bone Joint Surg Br*. 2003 May;85(4):478–83. Review. PubMed PMID: 12793548.
12. Giannoudis PV, Pape HC. Damage control orthopaedics in unstable pelvic ring injuries. *Injury*. 2004 Jul;35(7):671–7. Review. PubMed PMID: 15203307.

13. Scalea TM, Boswell SA, Scott JD, Mitchell KA, Kramer ME, Pollak AN. External fixation as a bridge to intramedullary nailing for patients with multiple injuries and with femur fractures: damage control orthopedics. *J Trauma*. 2000 Apr;48(4):613–21; discussion 621–3.
14. Giannoudis PV, Smith RM, Bellamy MC, Morrison JF, Dickson RA, Guillou PJ. Stimulation of the inflammatory system by reamed and unreamed nailing of femoral fractures. An analysis of the second hit. *J Bone Joint Surg Br*. 1999 Mar;81(2):356-61.
15. Giannoudis PV, Pape HC. Trauma and immune reactivity: too much, or too little immune response? *Injury*. 2007 Dec;38(12):1333-5.
16. Pape H-C, Giannoudis PV, Krettek C, Trentz O. Timing of fixation of major fractures in blunt polytrauma: role of conventional indicators in clinical decision making. *J Orthop Trauma*. 2005 Sep;19(8):551–62.
17. D’Alleyrand J-CG, O’Toole RV. The evolution of damage control orthopedics: current evidence and practical applications of early appropriate care. *Orthop Clin North Am*. 2013 Oct;44(4):499–507.
18. Awais S, Saeed A, Ch A. Use of external fixators for damage-control orthopaedics in natural disasters like the 2005 Pakistan earthquake. *International Orthopaedics*. 2014 Aug;38(8):1563–8.
19. Nicola R. Early Total Care versus Damage Control: Current Concepts in the Orthopedic Care of Polytrauma Patients. *ISRN Orthopedics*. 2013;2013:1–9.
20. Karadimas EJ, Nicolson T, Kakagia DD, Matthews SJ, Richards PJ, Giannoudis PV. Angiographic embolisation of pelvic ring injuries. Treatment algorithm and review of the literature. *International Orthopaedics*. 2011 Sep;35(9):1381–90.
21. Pape H, Stalp M, v Griensven M, Weinberg A, Dahlweit M, Tscherne H. [Optimal timing for secondary surgery in polytrauma patients: an evaluation of 4,314 serious-injury cases]. *Chirurg*. 1999 Nov;70(11):1287–93.
22. Vallier HA, Wang X, Moore TA, Wilber JH, Como JJ. Timing of orthopaedic surgery in multiple trauma patients: development of a protocol for early appropriate care. *J Orthop Trauma*. 2013 Oct;27(10):543–51.
23. Vallier HA, Moore TA, Como JJ, Wilczewski PA, Steinmetz MP, Wagner KG, et al. Complications are reduced with a protocol to standardize timing of fixation based on response to resuscitation. *Journal of Orthopaedic Surgery and Research [Internet]*. 2015 Dec [cited 2017 May 12];10(1). Available from: <http://www.josr-online.com/content/10/1/155>
24. Vallier HA, Dolenc AJ, Moore TA. Early Appropriate Care: A Protocol to Standardize Resuscitation Assessment and to Expedite Fracture Care Reduces

- Hospital Stay and Enhances Revenue. *Journal of Orthopaedic Trauma*. 2016 Jun;30(6):306–11.
25. Reich MS, Dolenc AJ, Moore TA, Vallier HA. Is Early Appropriate Care of axial and femoral fractures appropriate in multiply-injured elderly trauma patients? *Journal of Orthopaedic Surgery and Research* [Internet]. 2016 Dec [cited 2017 May 12];11(1). Available from: <http://josr-online.biomedcentral.com/articles/10.1186/s13018-016-0441-7>
 26. Pape HC, Andruszkow H, Pfeifer R, Hildebrand F, Barkatali BM. Options and hazards of the early appropriate care protocol for trauma patients with major fractures: Towards safe definitive surgery. *Injury*. 2016 Apr;47(4):787–91.
 27. Tebby J, Lecky F, Edwards A, Jenks T, Bouamra O, Dimitriou R, Giannoudis PV. Outcomes of polytrauma patients with diabetes mellitus. *BMC Med*. 2014 Jul 16;12:111.
 28. Giannoudis V, Panteli M, Giannoudis PV. Management of polytrauma patients in the UK: Is there a 'weekend effect'? *Injury*. 2016 Nov;47(11):2385-2390.
 29. Llompart-Pou JA, Chico-Fernández M, Sánchez-Casado M, Alberdi-Odrizola F, Guerrero-López F, Mayor-García MD, González-Robledo J, Ballesteros-Sanz MÁ, Herrán-Monge R, León-López R, López-Amor L, Bueno-González A; Trauma Neurointensive Care Working Group of the Spanish Society of Intensive Care Medicine (SEMICYUC). Age-related injury patterns in Spanish trauma ICU patients. Results from the RETRAUCI. *Injury*. 2016 Sep;47 Suppl 3:S61-S65.
 30. Sammy I, Lecky F, Sutton A, Leaviss J, O'Cathain A. Factors affecting mortality in older trauma patients-A systematic review and meta-analysis. *Injury*. 2016 Jun;47(6):1170-83.

Legends:**Figure 1:** PRISM Concept of management of polytrauma patients

PRISM= Prompt - Individualised - Safe – Management

ATLS: Advanced Trauma life support

CT: Computed tomography

ETC: Early total care

DCO: Damage control orthopaedics

EAC: Early appropriate care