

This is a repository copy of *Feeding management strategies among the early Neolithic pigs in the NE of the Iberian Peninsula*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/117508/>

Version: Accepted Version

Article:

Navarrete, Vanessa, Colonese, Andre Carlo orcid.org/0000-0002-0279-6634, Tornero, Carlos et al. (6 more authors) (2017) Feeding management strategies among the early Neolithic pigs in the NE of the Iberian Peninsula. *International Journal of Osteoarchaeology*. pp. 1-26. ISSN: 1047-482X

<https://doi.org/10.1002/oa.2598>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Feeding management strategies among the early Neolithic pigs in the NE of the Iberian Peninsula

Journal:	<i>International Journal of Osteoarchaeology</i>
Manuscript ID	OA-17-0026.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Navarrete, Vanessa; Universitat Autònoma de Barcelona, Prehistoria</p> <p>Colonese, Andre; University of York, BioArCh, Department of Archaeology</p> <p>Tornero, Carlos; Institute of Human Paleoecology and Social Evolution , Archaeology</p> <p>Antolín, Ferran; University of Basel , Integrative Prehistoric and Archaeological Science</p> <p>von Tersch, Matthew; University of York, BioArCh, Department of Archaeology</p> <p>Subirà, M. Eulàlia; Universitat Autònoma de Barcelona, Unitat d'Antropologia Biològica, Dpt. BABVE</p> <p>Comes, Pau; Universitat Autònoma de Barcelona, Institute of Environmental Science and Technology</p> <p>Rosell, Antoni; Universitat Autònoma de Barcelona, Institute of Environmental Science and Technology</p> <p>Saña Seguí, Maria; UAB Barcelona, Departament de Prehistòria</p>
Keywords:	Iberian Peninsula, Early Neolithic, husbandry practices, pig diet, stable carbon and nitrogen isotopes

SCHOLARONE™
Manuscripts

Feeding management strategies among the early Neolithic pigs in the NE of the Iberian Peninsula

Vanessa Navarrete^{1*}, André Carlo Colonese², Carlos Tornero³, Ferran Antolín⁴, Matthew Von Tersch², M. Eulàlia Subirà⁵, Pau Comes⁶, Antoni Rosell-Melé^{6,7}, Maria Saña¹

1 Laboratori d'Arqueozoologia, Department of Prehistory, Autonomous University of Barcelona, 08193, Catalonia, Spain.

2 BioArCh, Department of Archaeology, University of York, York, YO10 5DD, United Kingdom.

3 Institute of Human Paleoecology and Social Evolution (IPHES), 43007, Tarragona, Spain.

4 Integrative Prehistoric and Archaeological Science (University of Basel), Spalenring 145, 4055 Basel, Switzerland.

5 GRAPAC, Unitat d'Antropologia Biològica, Departament Animal Biology, Plant Biology and Ecology, Autonomous University of Barcelona, 08193, Catalonia, Spain.

6 Institute of Environmental Science and Technology, 08193, Autonomous University of Barcelona, Catalonia, Spain.

7 Institució Catalana de Recerca i Estudis Avançats (ICREA), 08193, Catalonia, Spain.

***Corresponding author:** Vanessa Navarrete. Laboratori d'Arqueozoologia, Department of Prehistory, Autonomous University of Barcelona, 08193, Catalonia, Spain. vanessa.navarrete@uab.cat

Keywords: Iberian Peninsula, Early Neolithic, husbandry practices, pig diet, stable carbon and nitrogen isotopes

Abstract

The socio-economic relevance of domesticated animals during the Early Neolithic in the Iberian Peninsula is indisputable, yet we essentially know little about the way they were managed. Among domesticated animals, pig (*Sus domesticus*) was a common food source and previous studies have shown the potential of stable isotopes for assessing variability in pig diet in relation to husbandry practices. Nevertheless, this approach has never been applied to the earliest pigs in the Iberian Peninsula. We analyzed the carbon and nitrogen stable isotope composition of pig bone collagen from several Early Neolithic sites in the NE Iberian Peninsula. While pig $\delta^{13}\text{C}$ values were similar across different populations, there were significant differences in $\delta^{15}\text{N}$ values between sites. These are attributed to different pig husbandry systems, which may reflect distinct social and spatial organization and interaction

with environmental conditions during the Early Neolithic in this region.

Introduction

The establishment of farmers and their domesticates marked a turning point in the socio-economic and cultural landscape of the western Mediterranean in the Middle Holocene. Livestock management practices in the Early Neolithic contributed to the emergence of new work processes, integrated new products in the human diet, and catalyzed the transformation of natural environments (Barker, 2005; Blondel, 2006; Zeder, 2008). With some exceptions in the north of the Iberian Peninsula, the hunting of wild game declined in most regions, while livestock became the most important source of animal meat and other secondary products (Saña, 1998; Guerra et al., 2008; Marín & Morales, 2009; Altuna & Mariezkurrena, 2011). The farming practices included complementary exploitation of pig, cattle, sheep and goat (Altuna & Mariezkurrena, 2011; Saña, 2013).

Among domesticated animals, pigs (*Sus domesticus*) played a role as meat supplier during the establishment of farmers in the Iberian Peninsula (Saña, 1998, 2013). Their relative abundance varied considerably between sites dated to early Neolithic periods (from 4 % to 23.9 % of domestic animals) suggesting different scales of husbandry regimes (Saña, 2013; Saña et al., 2015), but the form of these practices still remains highly elusive. Traditional husbandry practices may have involved home-based systems with complete to partial stabling of herds in close proximity to settlements, or extensive management of herds in semi-free to free-range regimes. While these management practices are known from modern traditional communities in northern Mediterranean areas (Albarella et al., 2007; Hadjikoumis, 2012), and have been also postulated for prehistoric groups in Europe (Balasse et al., 2016), their occurrence during the Neolithic in the NE Iberian Peninsula remains unclear. This gap in our knowledge prevents a full understanding of the socio-environmental significance of animal management practices during the establishment of early farming communities in this region.

Husbandry practices usually imply some degree of human control over animal diet in order to ensure their adaptation to and performance in local environmental conditions, promote health and prevent diseases. Due to their omnivorous nature, pigs have access to a broad range of food sources which can be mediated by cultural practices. Under human control, pig diets can be manipulated and supplemented with agricultural and animal products. Individual animal diets can be investigated using stable carbon ($^{13}\text{C}/^{12}\text{C}$) and nitrogen ($^{15}\text{N}/^{14}\text{N}$) isotope compositions of organic tissues, such as collagen in bone (DeNiro & Epstein, 1978; Schwarcz & Schoeninger, 1991; Ambrose, 1993; Sealy, 2001). Previous studies have demonstrated the potential of bulk collagen carbon and nitrogen stable isotopes for determining the degree of omnivory in pig feeding behaviour, and this information has been used to reconstruct pig management strategies in prehistoric and historic times (Matsui et al., 2005; Pechekina et al., 2005; Hamilton et al. 2009; Hamilton & Thomas, 2012; Madgwick et al., 2012; Balasse et al.,

2015, 2016).

In this paper we investigate the stable isotope ecology of pigs from six Early Neolithic sites in the NE Iberian Peninsula dating between ca. 5700 and 4200 cal BC (Fig. 1). Using stable carbon and nitrogen isotope analysis we assess the main dietary components of the pigs **in relation to wild and domestic herbivores, omnivores and carnivores**, and discuss the implications for understanding the variability of husbandry practices in this region.

Stable isotope analysis of bulk collagen

Collagen is the major protein in bone and its carbon and nitrogen isotope composition are indicative of individual diet over a relatively long period of time (e.g. **5-15 years depending on the age**; Hedges et al., 2007; Hedges & Reynard, 2007). Dietary information derived from **collagen** $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values is generally biased toward the protein fraction in the diet (Schwarcz & Schoeninger, 1991; Ambrose & Norr, 1993; Sealy, 2001). In diets with inadequate amounts of proteins for tissues synthesis the carbon isotopes may also originate from other macronutrients such as lipids and carbohydrates (Tieszen et al., 1983; Ambrose & Norr, 1993; Jim et al., 2004; Craig et al., 2013). Stable carbon isotopes differ greatly between terrestrial, marine and estuary ecosystems (Schoeninger & DeNiro, 1984; Bocherens et al., 1991, 1995), and between plants with different photosynthetic pathways (C_3 and C_4 plants) (O’Leary, 1988), thus providing a means of discriminating between macronutrients from distinct isotopic sources in diet.

Nitrogen isotopes are obtained exclusively from dietary protein, with $\delta^{15}\text{N}$ values generally reflecting the trophic position of the food source due to relatively large and predictable isotopic fractionations (ca. +3‰ to +6‰) through the foodweb (Schoeninger & DeNiro, 1984; Ambrose, 1993; O’Connell et al., 2012). Moreover, bulk collagen $\delta^{15}\text{N}$ values of humans and animals from agricultural contexts can potentially record changes in the abundance of ^{15}N in plant-soil systems due to land management practices, such as the addition of animal fertilizers (Bogaard et al., 2007; Szpak, 2014). When placed within the **context of the** local ecosystem, nitrogen stable isotopes can thus provide valuable information about the trophic level of food sources in pig diet, as well as spatio-temporal changes in agricultural strategies (Müldner & Richards, 2005; Hamilton & Thomas, 2012; Madgwick et al., 2012; Balasse et al., 2015, 2016).

Archaeological settings

Faunal remains were sampled from six early dated Neolithic sites located in the NE Iberian Peninsula (Table 1), distributed from the Eastern Pyrenees to the Mediterranean coast. Their elevations range from 960 m asl (cova del Frare) to 7 m asl (Caserna de Sant Pau). The majority of the sites are open settlements in coastal (Reina Amàlia, Caserna de Sant Pau and Serra de Mas Bonet) and inland areas (La Draga), while others are **inland** cave sites (cova del

Frare and Can Sadurní). Faunal and plant remains have been widely recovered in all sites during excavations, with zooarchaeological and archaeobotanical studies being developed over the past several years.

La Draga is a lakeshore settlement ^{14}C dated to 5201-4721 cal BC at Lake Banyoles, 170 m asl (Palomo et al., 2014; Terradas et al., 2015). The site is estimated to encompass approximately 8000 m², while an area of about 800 m² has been excavated. A large fraction of the archaeological record is underwater or in waterlogged environments, which has allowed excellent preservation of organic remains. The subsistence economy involved the exploitation of domestic and wild animals including mammals, fish, birds and molluscs (Saña, 2011, 2013). Agricultural practices are attested to by rich macrobotanical assemblages including *Hordeum distichum*, *Triticum durum/turgidum*, *Triticum aestivum*, *Triticum dicoccum*, *Triticum monococcum*, *Triticum* sp., *Vicia faba* and *Pisum sativum* (Buxó, 2007; Antolín & Buxó, 2012; Antolín et al., 2014). Farming was the main economic activity, followed by hunting and gathering of wild plants (Saña, 2013; Antolín et al., 2014).

Reina Amàlia and the adjacent site of Caserna de Sant Pau are coastal settlements located in Barcelona. At 25 m asl and ^{14}C dated to 4611-4373 cal BC (González et al., 2011), the site at Reina Amàlia has a surface area of at least 200m², consisting of a group of habitation structures (50m²), including burials, where the subsistence economy involved the exploitation of domesticated animals (*Sus domesticus*, *Ovis aries*, *Capra hircus* and *Bos taurus*), complemented with wild fauna (Saña & Navarrete, 2016). Agricultural practices are similarly attested to by the remains of *Hordeum vulgare* var. *nudum* and *Triticum aestivum/durum/turgidum* and *Triticum dicoccum* (Antolín, 2016). The site of Caserna de Sant Pau has been successively occupied during the Neolithic, with the phase analyzed in this study ^{14}C dated to 5372-5076 cal BC (Molist et al., 2008). The excavated area corresponds to approximately 800m². This site also consists of structures (silos), and individual and double burials. Subsistence included the exploitation of domestic animals (*Sus domesticus*, *Ovis aries*, *Capra hircus* and *Bos taurus*) (Colominas et al., 2008) and plants (*Hordeum vulgare*, *Hordeum vulgare* var. *nudum*, *Triticum aestivum/durum*, *Triticum dicoccum*, *Vicia faba* and *Pisum sativum*) (Buxó & Canal, 2008), along with wild resources (Buxó & Canal, 2008; Colominas et al., 2008; Molist et al., 2008).

Cova del Frare is a cave located in the massif of Sant Llorenç de Munt at 960 m asl, and was occupied from the Early Neolithic to the Bronze Age. The Early Neolithic deposits analysed in this work were ^{14}C dated to 5216-4993 cal BC (Martín et al., 2009). The cave has been interpreted as a seasonal dwelling for Neolithic shepherds, with storage bins, household waste and tool production debris having been recovered at the site (Martín et al., 2010). The local economy included four main domestic animals - *Sus domesticus*, *Ovis aries*, *Capra hircus* and *Bos taurus*. No archaeobotanical analyses have been performed on the site as it was excavated in the early 1980s when sampling for such analyses was not common practice in the study

area.

Can Sadurní is a cave located in the massif of the Serra de Garraf at 420 m asl. In this work, we analyzed Early Neolithic deposits from the site ¹⁴C dated to 5291-4710 cal BC (Layer 17) and 4456-4335 cal BC (Layer 10b / 11) (Edo et al., 2011). Plant remains suggest that the cave was used as a stall during the Early Neolithic (Edo et al., 2011), and this seems to be supported by archaeozoological analysis (Saña et al., 2015). The subsistence economy involved the exploitation of domesticated animals (*Sus domesticus*, *Ovis aries*, *Capra hircus* and *Bos taurus*) (Saña et al., 2015), and agricultural practices are well represented by several crops, such as *Hordeum vulgare*, *Hordeum distichon*, *Hordeum vulgare* var. nudum, *Triticum aestivum/durum*, *Triticum dicoccum*, *Triticum monococcum*, *Lens culinaris*, *Pisum sativum*, *Papaver somniferum* and possibly also *Linum usitatissimum* (Antolín, 2016; Antolín et al., 2017). Wild resources were also exploited at the site (as evidenced by *Arbutus unedo*, *Pistacia lentiscus*, *Pinus* sp., *Quercus* sp. and *Vitis vinifera* subsp. *sylvestris*, and *Sus scrofa*, *Capra pyrenaica* and *Capreolus capreolus*) (Blasco et al., 1999; Antolín et al., 2015; Saña et al., 2015).

Finally, Serra de Mas Bonet is a settlement in Vilafant, at 75 m asl and ¹⁴C dated to 4900-4600 cal BC (Rosillo et al., 2010). The analysed material derived from two structures interpreted as silos, however their spatial distribution appears random and the total surface area of the site could not be assessed. Farming was the principal economic activity at the site, with the rearing of domesticated animals (*Sus domesticus*, *Ovis aries*, *Capra hircus* and *Bos taurus*) (Saña, unpublished). No direct evidence of plant remains was found for this settlement phase, probably due to poor sampling or preservation conditions of the organic remains at the site (Antolín, 2016).

Material and Methods

Selection of faunal samples

From the six Early Neolithic sites, 92 pig individuals were selected from layers dated between 5372-5076 to 4456-4335 cal BC, for stable carbon and nitrogen isotope analysis (Tables 1, 2). In order to establish the local $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ isotopic baselines of the pigs, we also selected a range of herbivores (n = 70), including domestic (*Ovis aries*, *Capra hircus*, *Bos taurus*) and wild animals (*Cervus elaphus*, *Capra pyrenaica*, *Capreolus capreolus*, *Oryctolagus cuniculus*), carnivores (n = 15) (*Meles meles*, *Martes martes*, *Felis sylvestris*, *Canis familiaris*, *Vulpes vulpes*) and a few wild boar (n = 5) (*Sus scrofa*). Wild boar, which represents only a minor fraction of suids found at these sites, were previously distinguished from domesticated pigs based on osteometric data (Hain, 1982; Albarella et al., 2005; Altuna & Mariezkurrena, 2011), using a data-driven approach of Payne & Bull (1988) and Albarella et al., (2009) (SI1-2). Samples for stable isotope analyses consisted of adult specimens, and

included the diaphysis of long bones, maxilla and mandibular diastema. Whenever possible, specimens were selected to represent individual animals by sampling the same-sided portion of a specific element. Specimens available from Reina Amàlia and Caserna de Sant Pau were combined to form a single assemblage (Reina Amàlia-Caserna de Sant Pau) as they were located adjacent to one another.

Collagen extraction and stable isotope analysis

Collagen extraction and stable isotope analysis were performed at the Unitat d'Antropologia Biològica (Departament Animal Biology, Plant Biology and Ecology) and Laboratori d'Arqueozoologia (Department of Prehistory) at the Autonomous University of Barcelona (Spain). Some samples were also extracted and analysed at the BioArCh facilities in the Department of Archaeology, University of York (UK). Collagen extraction followed similar protocols in both labs. Bones were cleaned mechanically to remove the surface and the extraction followed a modified Longin method (Brown et al., 1988); details can be found in previous studies (e.g. Craig et al., 2010). In short, shards of bones (ca. 200 to 300 mg) were demineralised using 0.6 M HCl, at 4°C for several days, then rinsed with ultrapure water (milli-Q®) and gelatinised with 0.001 M HCl at 80°C for 48 h. Samples were then ultrafiltered (30 kDa, Amicon® Ultra-4 centrifugal filter units; Millipore, MA, USA), frozen and freeze dried.

Collagen samples (0.3 mg) were analysed in duplicate using a Thermo Flash 1112 elemental analyser (EA) coupled to a Thermo Delta V Advantage isotope ratio mass spectrometer (IRMS) with a ConFlo III interface, at the Institute of Environmental Science and Technology, Autonomous University of Barcelona. The international laboratory standard IAEA 600 (caffeine) was used as a control. The average analytical error was <0.2‰ (1σ) as determined from the duplicate analyses of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. Some samples were analyzed at the University of York (1 mg) in duplicate or triplicate on another EA-IRMS in a GSL analyser coupled to a 20-22 mass spectrometer (Sercon, Crewe, UK). The analytical error for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values, calculated from repeated measurements of each sample and measurements of the bovine control from multiple extracts, was also <0.2‰ (1σ). The standard used for $\delta^{13}\text{C}$ was Vienna PeeDee Belemnite (V-PDB), and the standard for $\delta^{15}\text{N}$ was air N_2 (AIR). Samples ($n = 3$) and in-house collagen standards ($n = 1$ bovine control) were analyzed in both IRMS system to ensure accuracy.

Comparison between $\delta^{13}\text{C}$ values of wild and domestic herbivores, pig, wild boar, and carnivores was performed using one-way ANOVA ($\alpha = 0.05$), after checking for normal distribution with the Shapiro-Wilk test for normality ($\alpha = 0.05$). The null hypothesis that the data were normally distributed was rejected for the $\delta^{15}\text{N}$ values. Thus comparison between the $\delta^{15}\text{N}$ values was performed using the Kruskal-Wallis test ($\alpha = 0.05$). All statistical tests were performed in PAST 3.x (Hammer et al., 2001).

Results

Collagen preservation

The results of the stable isotopes and collagen quality indicators are reported in the figure 2 and table 2. Out of a total of 182 specimens, collagen was successfully extracted from 100 (55%). Collagen yields were extremely variable, ranging from 0.4 to 10.6 mg. The C% and N% ranged from 27% to 47%, and 10% to 17% respectively, with C:N ratios ranging from 3.1 to 3.6 and falling within the values proposed by DeNiro (1985) and Van Klinken (1999) for preserved collagen. We also applied a cut-off of 13% and 4.8% for C% and N%, respectively, as recommended by Ambrose (1990, 1993).

$\delta^{13}C$ values

The average $\delta^{13}C$ values of pigs exhibited little variability between sites (Fig. 2), ranging from $-20.6 \pm 0.6\text{‰}$ (Reina Amàlia-Caserna de Sant Pau; $n = 16$) to $-19.6 \pm 0.7\text{‰}$ (Can Sadurní; $n = 3$). However, pigs from Can Sadurní were significantly enriched in ^{13}C by approximately 1‰ compared to La Draga ($n = 24$) and Reina Amàlia-Caserna de Sant Pau ($p = 0.003$, $f = 4.37$). The average $\delta^{13}C$ values of herbivores ranged from $-20.6 \pm 0.7\text{‰}$ (Reina Amàlia-Caserna de Sant Pau; $n = 6$) to $-20.2 \pm 0.7\text{‰}$ (cova del Frare; $n = 12$), and were statistically indistinguishable between sites ($p = 0.6031$, $f = 0.516$). Moreover, no significant differences were found between the $\delta^{13}C$ values of wild and domestic herbivores at cova del Frare ($n = 12$) and La Draga ($n = 10$; $p > 0.05$), the only sites with sufficient specimens for statistical comparison. Comparison between pig and herbivore $\delta^{13}C$ values was possible for cova del Frare, Reina Amàlia-Caserna de Sant Pau and La Draga (Fig. 3A). At all of these sites, the $\delta^{13}C$ values of pigs and herbivores were statistically indistinguishable ($p > 0.05$). Due to the low number of sampled specimens ($n = 5$), the average $\delta^{13}C$ values of wild boar could not be compared between sites; however, the lowest wild boar $\delta^{13}C$ value was found at Reina Amàlia-Caserna de Sant Pau (-20.3‰), while the highest was found at La Draga (-19.3‰). Finally, the $\delta^{13}C$ values of carnivores varied between sites, but the highest variability was found at La Draga ($n = 3$), where the $\delta^{13}C$ values of European badger (*Meles meles*) ranged from -19‰ to -21.5‰ .

$\delta^{15}N$ values

The average $\delta^{15}N$ values of pigs ranged from $+5.1 \pm 1.6\text{‰}$ (La Draga) to $+7.6 \pm 1.1\text{‰}$ (Reina Amàlia-Caserna de Sant Pau), and these differences were statistically significant ($p < 0.001$). In particular, pigs from Reina Amàlia-Caserna de Sant Pau and from Can Sadurní were enriched in ^{15}N by an average of 2.5‰ compared to specimens from La Draga and cova del Frare ($n = 13$) ($p = 0.030$). The average $\delta^{15}N$ values of herbivores ranged from $+4.9 \pm 1.1\text{‰}$ (La Draga; $n = 10$) to $+6.1 \pm 1\text{‰}$ (Reina Amàlia-Caserna de Sant Pau; $n = 6$). Although the herbivores from Reina Amàlia-Caserna de Sant Pau were on average enriched in ^{15}N by

~1.2‰ compared to those from La Draga and Can Sadurní, these differences were statistically insignificant ($p = 0.070$). The $\delta^{15}\text{N}$ values of wild and domestic herbivores were statistically indistinguishable at cova del Frare ($n = 12$) ($p = 0.852$). However, the domestic herbivores were significantly ^{15}N enriched by 0.6‰ compared to the wild ones at La Draga ($p = 0.028$). The $\delta^{15}\text{N}$ values of pigs were statistically indistinguishable from those of wild and domestic herbivores at cova del Frare and La Draga ($p > 0.05$). Conversely, the $\delta^{15}\text{N}$ values of pigs from Reina Amàlia-Caserna de Sant Pau were significantly higher by an average of 1.6‰ compared to the herbivores ($p = 0.008$) (Fig. 3B). The $\delta^{15}\text{N}$ values of wild boar were highly variable, with the highest values found at Can Sadurní (+7.6‰) and La Draga (+7.4‰), and the lowest at cova del Frare (+4.5‰) and Reina Amàlia-Caserna de Sant Pau (+5.6‰). Interestingly, the $\delta^{15}\text{N}$ values of wild boar from Reina Amàlia-Caserna de Sant Pau fall within the range of herbivores, and are lower than domestic pigs across most sites. The $\delta^{15}\text{N}$ values of carnivores were also variable across all sites, ranging from +6.5‰ (dog, *Canis familiaris*) to +12.6‰ (European badger) at La Draga. The $\delta^{15}\text{N}$ values of badgers justify their inclusion in the carnivore group.

Discussion

Source of carbon and nitrogen in pig diet

The $\delta^{13}\text{C}$ values of the analyzed specimens fall within the expected values for C_3 plant ecosystems (O'Leary, 1988), which dominated the vegetation composition in the NE Iberian Peninsula during the early-middle Holocene (Burjachs, 2000; Jalut et al., 2009; Revelles et al., 2016). Given the insignificant or small difference between the $\delta^{13}\text{C}$ (0.3‰) and $\delta^{15}\text{N}$ (0.6‰) values of domestic and wild herbivores for each site we combined them to obtain a more robust average isotopic baselines for interpreting pig $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values. The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of herbivores show very small to insignificant differences between sites, and present narrow ranges (3.3‰ and 3.7‰ respectively). Using a carbon isotope fractionation of ~5‰ between whole plant and consumer's bone collagen (Ambrose & Norr, 1993), and a correction (~+1.5‰) due to the fossil fuel effect when applied to pre-industrial ecosystems (Friedli et al., 1986; Hellevang & Aagaard, 2015), the highest $\delta^{13}\text{C}$ values in herbivorous animals (e.g. >-19‰) could be explained by the consumption of drought-resistant vegetation, such as shrubs (e.g. -23‰; Filella & Peñuelas, 2003). The $\delta^{13}\text{C}$ range of the pigs from all sites is even narrower compared to the herbivores (2.3‰) and their absolute $\delta^{13}\text{C}$ values generally overlap with the herbivore data (Fig. 3A). The results therefore indicate that pig diets were in general dominated by plant products.

Conversely to carbon, the average $\delta^{15}\text{N}$ values of pigs varied significantly among populations. The lowest $\delta^{15}\text{N}$ values were found in pigs from La Draga, followed by specimens from cova del Frare and Serra de Mas Bonet. In these sites the $\delta^{15}\text{N}$ values were generally comparable with their local wild and domestic herbivores (Fig. 2 and Fig. 3B), therefore likely reflecting a

diet predominantly based on plant products. The highest $\delta^{15}\text{N}$ values were found in pigs from Can Sadurní and Reina Amàlia-Caserna de Sant Pau, where the $\delta^{15}\text{N}$ values were on average higher than the local herbivores by 2.5‰ and 1.4‰ respectively, and in some cases similar to carnivore $\delta^{15}\text{N}$ values (e.g. Can Sadurní).

Implications for pig husbandry practices

There is a general consensus that the variable contribution of animal and plant macronutrients to pig diet can potentially reflect the scale of husbandry practice in the past (Minagawa et al., 2005; Müldner & Richards, 2005; Pechenkina et al., 2005; Fuller et al., 2012; Hamilton & Thomas, 2012; Madgwick et al., 2012; Hammond & O'Connor, 2013; Halley & Rosvold, 2014; Balasse et al., 2015, 2016). This appears to be supported by ethnographic studies on traditional husbandry systems in Southern Europe (Italy, Spain, Greece and Corsica) that also reveal some variability in pig diet in relation to management strategies (Albarella et al., 2007, 2011; Isaakidou, 2011; Hadjikoumis, 2012). Given their omnivory and foraging habits, the nitrogen isotope composition of pig bone collagen can be derived from both animal and plant proteins. Pigs raised in a home-based system with complete or temporary stabling are expected to have a more controlled diet than free-range animals. Enclosed pigs may feed predominantly on plant materials (ground cereals and legumes), but due to a higher degree of human control they may have diets supplemented with domestic left-overs, including animal products. For pigs raised under this regime we could hypothetically expect collagen $\delta^{15}\text{N}$ values higher than local herbivores, although this would also depend upon other factors such as the proportions and quality of meat protein in their diets. By contrast, pigs reared in semi-free or free-range systems will likely obtain most of their nutrients from available plants, although this does not exclude some consumption of wild animals. Within this management strategy we might expect pig collagen $\delta^{15}\text{N}$ values to be compatible or very close to local herbivores. However, as discussed below, other interplaying factors may affect bulk collagen $\delta^{15}\text{N}$ values (Szpak, 2014).

The $\delta^{15}\text{N}$ values of pigs at Reina Amàlia-Caserna de Sant Pau and Can Sadurní suggest a relatively high proportion of animal sources contributed to their dietary proteins, which may be associated with a home-based management system, where pigs are enclosed or relatively free to forage within the settlement. This home-based system occurs in contexts where pigs are relatively abundant, such as at Reina Amàlia-Caserna de Sant Pau where they accounted for the 23.6% and 23.9% of domesticates, as well as in contexts where pigs were a minor component of livestock, such as at Can Sadurní, where pigs accounted for 7.5% of domesticates. In addition, the presence of neonate pigs found at Reina Amàlia-Caserna de Sant Pau (2.7% of pigs remains) supports the hypothesis that pigs were kept within the settlement. However, it has to be said that herbivores from Reina Amàlia-Caserna de Sant Pau, predominantly domestic animals, were on average enriched in ^{15}N by ~1.2‰ compared to those from La Draga and Can Sadurní. They suggest some degree of manuring effect (see

below), which may have been subsequently propagated to pig collagen through the consumption of animal products.

Home-based management at Can Sadurní seems to be corroborated by other lines of evidence. Archaeobotanical and zooarchaeological analyses suggest that the site was used as a stall for sheep and goats (Saña et al., 2015). Moreover, organic residue analysis recently detected dairy products in Early Neolithic ceramic vessels from this site, suggesting that the cave may also have been used in dairy production (Debono Spiteri et al., 2016). Presumably the small number of pigs at Can Sadurní may have been kept in enclosed regime, and could have been foddered on a range of locally available sources, including animal products. Interestingly, the only specimen of wild boar analysed from Can Sadurní had a high $\delta^{15}\text{N}$ value, comparable with those of the pigs. Whether this indicates that wild boars may have been kept enclosed along with domesticated pigs, or if their natural diet also included several animal products, remains a matter of debate. However the number of specimens from Can Sadurní is too small to lead to conclusive interpretations.

Conversely, pigs from La Draga, cova del Frare and Serra de Mas Bonet had $\delta^{15}\text{N}$ values consistent with their respective local herbivores, which might be associated with a free-range system. Our results indicate that pigs were fed predominantly on plant products, which might indicate that they were free to roam in forested environments. However, the relatively elevated frequency of neonates at La Draga, where no bias is expected against adult specimens due to differential preservation processes (Saña et al., 2014), suggests that pigs were kept within the settlement and possibly managed at the household level (Antolín et al., 2014). If this was the case, their feeding management may have predominantly included plant products, perhaps by-products of crop production (e.g. cereals, pulses). Similar interpretations have been proposed for prehistoric pigs possibly raised at the household level but on a dominantly herbivorous diet (Balasse et al., 2016), and for modern pigs raised in traditional farming communities (Hadjikoumis, 2012). Two specimens from La Draga had remarkably high $\delta^{15}\text{N}$ values, perhaps indicating variability in husbandry regimes at the local scale. This could be due to selective feeding practices, for reasons that remain unclear for us, or variability of protein intake between households (e.g. Sykes, 2014). These two specimens form a distinctive isotopic group (Fig. 2), plotting along with one wild boar and one European badger, which may also indicate that some pigs were not local, but rather were raised elsewhere then brought to the site at a later stage. This seems to be supported by the large range of $\delta^{13}\text{C}$ values of local herbivores, which again could be tentatively associated with non-local animals. Moreover, shellfish and lithic raw material sourced from coastal areas were used by Neolithic groups at La Draga (Terradas et al., 2012). These independent lines of evidence suggest that La Draga was integrated within a regional trading network that may have involved circulation of livestock.

Recent studies suggest that pigs from La Draga were integrated within intensive mixed

farming, where they would have had access to the fields for grazing on leftover crops while also manuring the plots (Antolín et al., 2014). Interestingly, domestic herbivores at La Draga were significantly ^{15}N enriched by 0.6‰ compared to the wild ones. This slight enrichment is also observed in the average $\delta^{15}\text{N}$ values of pigs, which is higher by 0.7‰ compared with the average values of the wild herbivores. However, the enrichment in ^{15}N is too low for a manuring effect if we consider the $\delta^{15}\text{N}$ values of manured crops in several Neolithic contexts in Europe (Fraser et al., 2011; Bogaard et al., 2013). This could, however, be due to the isotopic resolution not being adequate to resolve short-term feeding practices. Moreover, the effect of animal dung on plant $\delta^{15}\text{N}$ values is highly variable, and depends on, among other factors, the rate of manuring and the type of fertilizer (Fraser et al., 2011; Szpak, 2014). Furthermore, the increase in ^{15}N is not homogenous throughout the plant, and manured cereal straws may be depleted in ^{15}N relative to the grains (Bogaard et al., 2007, 2013). Legumes were also documented (though scarcely) at La Draga (Berrocal et al., in press), but these are typically ^{15}N depleted compared to cereal grains, and their $^{15}\text{N}/^{14}\text{N}$ only respond to a very high level of manuring (Bogaard et al., 2013). The isotope results of pigs from La Draga provide new insights into feeding management practices, but the scale of husbandry remains unclear. The same can be stated for cova del Frare and Serra de Mas Bonet, where the scarce archaeological information prevents conclusive interpretations to be drawn.

Our results indicate the coexistence of distinct foddering strategies among pig populations during the Early Neolithic in the NE Iberian Peninsula, but the scale of husbandry still remains elusive. The relative importance of pigs does not appear to be homogeneous among sites, but this does not seem to affect pig diet. However some variability was also observed in the size of specimens found at these sites. A large size variability indeed characterized pig populations in the Iberian Peninsula during the Neolithic, resulting from different selective environmental and social pressures (Navarrete & Saña, 2017). Stable isotope analysis on single amino acids from bone collagen may provide in the future additional complementary information on pig foddering strategies. This may be complemented with nitrogen isotope analysis of plant remains that are currently under investigation.

Determining the diet of domestic animals is also essential for developing appropriate isotopic baselines for interpreting human $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in order to derive paleodietary information. For example, relatively high $\delta^{15}\text{N}$ values were recently observed in Neolithic human individuals in this region (Fontanals-Coll et al., 2015, 2017) and were tentatively interpreted as the occasional consumption of freshwater fish. The isotopic data discussed in this work suggests that variability in pig foddering strategies provides an alternative explanation to freshwater fish consumption, assuming that pigs were a relevant source of animal protein to human diet.

Conclusion

In this study we analyzed bulk collagen carbon and nitrogen stable isotopes of pigs from six Early Neolithic sites in the northeastern Iberian Peninsula dated between ca. 5700 and 4200 cal BC. When compared with herbivores, we observed significant differences in pig $\delta^{15}\text{N}$ values between the sites that likely reflect variable foddering strategies. These differences may be indicative of variable management strategies, perhaps resulting from distinct adaptive responses to natural and/or social factors. In particular, a mainly herbivorous diet was detected in pig specimens from cova del Frare, La Draga and Serrat de Mas Bonet, which could reflect free-range or semi-free management systems. On the other hand, a higher consumption of animal protein could be postulated for pig specimens from Can Sadurní and Reina Amàlia-Caserna de Sant Pau, which in turn would reflect household-level regimes or the presence of pigs raised in enclosures. In summary, multiple factors may have contributed to the variability in foddering strategies among the earliest farmers in this area. This study offers new elements for discussing and opening new perspectives into early animal management strategies and the implications for understanding management strategies during the regional development of the Neolithic economy in the Iberian Peninsula.

Acknowledgements

This work was funded by the Ministerio de Economía y Competitividad (Spain), under the project “Producción animal y cerámica en el neolítico peninsular. Estudio biogeoquímico integrado del consumo y las prácticas culinarias” (HAR2014-60081-R). The authors thank the directors of the research in archaeological sites studied in this work for providing access to the samples. The authors thank Krista McGrath for kindly revising the English and providing constructive comments on the early drafts of this manuscript. The authors also wish to thank Gundula Müldner and Oliver Craig for productive discussions, and Matthew Collins for his constructive comments on the manuscript.

Table caption

Table 1. Conventional radiocarbon date along with calibrated radiocarbon ages using OxCal 4.2 software (IntCal13; Reimer et al., 2013).

Table 2. Results from isotopic analysis of faunal samples.

Figure caption

Figure 1. Location of the sites in the northeastern Iberian Peninsula.

Figure 2. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of animals from cova del Frare, Reina Amàlia-Caserna de Sant Pau, Can Sadurní, La Draga and Serra de Mas Bonet.

Figure 3A-B. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of pigs (unfilled circle) corrected for the average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of herbivores ($\Delta\text{‰}$) for cova del Frare (CF), La Draga (DR), Reina Amàlia-Caserna de Sant Pau (RA-CSP) and Can Sadurní (CS). We also corrected the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of herbivores for their average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (filled circles) in order to show their isotopic variability.

Supplementary information

SI 1. Summary statistics for pig measurements from the sites in the northern Iberian Peninsula, Valencina de la Concepcion, Zambujal, Durrington Walls, La Draga, Reina Amàlia, Caserna de Sant Pau, cova del Frare and Can Sadurní. N = number of specimens, MIN = minimum, MAX = maximum, VAR = coefficient of variation. List of the measurements taken, according to von den Driesch, 1976.

SI 2. Pig measurements for specimens selected for isotopic analysis from La Draga, Reina Amàlia, Caserna de Sant Pau, cova del Frare, Can Sadurní and Serra de Mas Bonet. List of the measurements taken according to von den Driesch, 1976.

References

Albarella U, Davis S, Detry C, Rowley-Conwy P. 2005. Pigs of the ‘Far West’: The biometry of *Sus* from archaeological sites in Portugal. *Anthropozoologica* **40**: 27-54.

Albarella U, Dobney K, Rowley-Conwy P. 2009. Size and shape of the Eurasian wild boar (*Sus scrofa*) with a view to the reconstruction of its Holocene history. *Environmental Archaeology* **14**: 103–136.

Albarella U, Manconi F, Trentacoste A. 2011. A week on the plateau: pig husbandry, mobility and resources exploitation in central Sardinia. In *Ethnoarchaeology: the present past of human: animal relationships*, Albarella U (ed.). Oxford: Oxbow; 143-159.

Albarella U, Manconi F, Vigne JD, Rowley-Conwy P. 2007. Ethnoarchaeology of pig husbandry in Sardinia and Corsica. In *Pigs and Humans. 10,000 years of Interactions*, Albarella U, Dobney K, Ervynck A, Rowley-Conwy P (eds.). Oxford University Press: Oxford; 285-307.

Albarella U, Payne S. 2005. Neolithic pigs from Durrington Walls, Wiltshire, England: A biometrical database. *Journal of Archaeological Science* **32**: 589-599.

Altuna J, Mariezkurrena K. 2011. Diferenciación biométrica de *Sus scrofa* y *Sus domesticus* en yacimientos arqueológicos del norte de la Península Ibérica. *Kobie Serie Paleoantropología* **30**: 5-22.

Ambrose SH, Norr L. 1993. Experimental Evidence for the Relationship of the Carbon

Isotope Ratios of Whole Diet and Dietary Protein to Those of Bone Collagen and Carbonate. In *Prehistoric Human Bone-Archaeology at the Molecular Level*, Lambert JB, Grupe G (eds.). Springer Verlag: Berlin; 1-37.

Ambrose SH. 1990. Preparation and characterization of bone and tooth collagen for isotopic analysis. *Journal of Archaeological Science* **17**: 431–451.

Ambrose SH. 1993. *Isotopic analysis of paleodiets: methodological and interpretive considerations*. Food and nutrition in history and anthropology: USA.

Antolín F, Buxó R, Jacomet S, Navarrete V, Saña M. 2014. An integrated perspective on farming in the early Neolithic lakeshore site of La Draga (Banyoles, Spain). *Environmental Archaeology* **19**: 241-255.

Antolin F, Buxó R. 2012. Chasing the traces of diffusion of agriculture during the early neolithic in the western mediterranean coast. *Rubricatum: revista del Museu de Gavà* **0**: 95-102.

Antolín F, Jacomet S, Buxó R. 2015. The hard knock life. Archaeobotanical data on farming practices during the Neolithic (5400-2300 cal BC) in the NE of the Iberian Peninsula. *Journal of Archaeological Science* **61**: 90-104.

Antolín F, Navarrete V, Saña M, Viñerta A, Gassiot E. **Submitted**. Herders in the mountains and farmers in the plains? A comparative evaluation of the archaeobiological record from Neolithic sites in the eastern Iberian Pyrenees and the southern lower lands. *Quaternary International*.

Antolín F. 2016. *Local, intensive and diverse? Early farmers and plant economy in the North-East of the Iberian Peninsula (5500-2300 cal BC)*. Barkhuis: Gröningen.

Balasse M, Balasescu A, Tornero C, Frémondeau D, Hovsepyan R, Gillis R, Popovici D. 2015. Investigating the scale of herding in chalcolithic pastoral communities settled by the Danube river in the 5th millennium BC: a case study at Bordusani-Popina and Hârsova-tell (Romania). *Quaternary International*. DOI: 10.1016/j.quaint.2015.07.030.

Balasse M, Evin A, Tornero C, Radu V, Fiorillo D, Popovici D, Andreescu R, Dobney K, Cucchi T, Bălăşescu A. 2016. Wild, domestic and feral? Investigating the status of suids in the Romanian Gumelnița (5th mil. cal BC) with biogeochemistry and geometric morphometrics. *Journal of Anthropological Archaeology* **42**: 27–36.

Barker G. 2005. Agriculture, pastoralism, and Mediterranean landscapes in prehistory. In *The archaeology of Mediterranean prehistory*, Blake M, Knapp B (eds.). Blackwell Publishing Ltd: United Kingdom; 46-76.

1
2
3 Berrocal A, Antolín F, Buxó R. in press. Actividades agrícolas en La Draga (Banyoles, Pla
4 de l'Estany): resultados del análisis carpológico de nuevos contextos excavados en el
5 sector A. *Actas del 6º Congreso del Neolítico Peninsular*.

6
7 Blasco A, Edo M, Villalba MJ, Buxó R, JuanTresserras J, Saña M. 1999. Del Cardial al
8 Postcardial en la Cueva de Can Sadurní (Begues, Barcelona). Primeros datos sobre su
9 secuencia estratigráfica, paleoeconómica y ambiental. *Saguntum Extra* **2**: 5968.

10
11 Blondel J. 2006. The 'design' of Mediterranean landscapes: a millennial story of humans
12 and ecological systems during the historic period. *Human Ecology* **34**: 713-729.

13
14 Bocherens H, Fizet M, Mariotti A, Lange-Badré B, Vandermeersch B, Borel JP, Bellon G.
15 1991. Isotopic biogeochemistry (^{13}C , ^{15}N) of fossil vertebrate collagen: implications for the
16 study of a fossil food web including Neandertal Man. *Journal Human Evolution* **20**: 481–
17 492.

18
19 Bocherens H, Fogel ML, Tuross N, Zeder M. 1995. Trophic structure and climatic
20 information from isotopic signatures in Pleistocene cave fauna of southern England.
21 *Journal of Archaeological Science* **22**: 327-340.

22
23 Bogaard A, Fraser R, Heaton TH, Wallace M, Vaiglova P, Charles M, Jones G, Evershed
24 R, Stryring A, Andersen N, Arbogast RM, Bartosiewicz L, Gardeisen A, Kanstrup M,
25 Maier U, Marinova E, Ninov L, Schäfer M, Stephan, E. 2013. Crop manuring and
26 intensive land management by Europe's first farmers. *Proceedings of the National*
27 *Academy of Sciences* **110**: 12589-12594.

28
29 Bogaard A, Heaton THE, Poulton P, Merbach I. 2007. The impact of manuring on nitrogen
30 isotope ratios in cereals: archaeological implications for reconstruction of diet and crop
31 management practices. *Journal of archaeological science* **34**: 335-343.

32
33 Brown TA, Nelson EE, Vogel SJ, Southon JR. 1988. Improved Collagen Extraction by
34 Modified Longin Method. *Radiocarbon* **30**: 171-177.

35
36 Burjachs, F. 2000. El paisatge del neolític antic. Les dades palinològiques. In *El poblat*
37 *lacustre neolític de La Draga. Excavacions del 1990 1998*, Bosch A, Chinchilla J, Tarrús J
38 (eds.). Monografies del CASC2: Girona; 46-54.

39
40 Buxó R, Canal D. 2008. L'agricultura i l'alimentació vegetal. *Quarhis* **4**: 54-56.

41
42 Buxó R. 2007. Crop Evolution: New Evidence from the Neolithic of the West
43 Mediterranean Europe. In *The Origins and Spread of Domestic Plants in Southwest Asia*
44 *and Europe*, Colledge S, Conolly J (eds.). UCL Institute of Archaeology Publication:
45 Routledge; 155-71.

Colominas L, Aguillo E, Saña M, Tornero C. 2008. La gestió dels recursos animals durant les ocupacions de l'assentament de la caserna de Sant Pau. *Quarhis: Quaderns d'Arqueologia i Història de la Ciutat de Barcelona* **4**: 57-63.

Craig OE, Biazzo M, Colonese AC, Di Giuseppe Z, Martinez-Labarga C, Lo Vetro D, Lelli R, Martini F, Rickards O. 2010. Stable isotope analysis of Late Upper Palaeolithic human and faunal remains from Grotta del Romito (Cosenza), Italy. *Journal of archaeological science* **37**: 2504-2512.

Craig OE, Bondioli L, Fattore L, Higham T, Hedges R. 2013. Evaluating Marine Diets through Radiocarbon Dating and Stable Isotope Analysis of Victims of the AD79 Eruption of Vesuvius. *American Journal of Physical Anthropology* **152**: 345-52.

Debono Spiteri C, Gillis R, Roffet-Salque M, Castells L, Guilaine J, Manen C, Muntoni I. 2016. Regional Asynchronicity in Dairy Production and Processing in Early Farming Communities of the Northern Mediterranean. *Proceedings of the National Academy of Sciences of the United States of America* **113**: 13594-99.

DeNiro MJ, Epstein S. 1978. Influence of diet on the distribution of carbon isotopes in animals. *Geochimica et Cosmochimica Acta* **42**: 495-506.

DeNiro MJ. 1985. Post-mortem preservation and alteration of in vivo bone collagen isotope ratios in relation to paleodietary reconstructions. *Nature* **317**: 806-809.

Driesch A. 1976. *A guide to the measurement of animal bones from archaeological sites*. Peabody Museum of Archaeology and Ethnology: Cambridge.

Edo M, Blasco A, Villalba MJ. 2011. La cova de Can Sadurní, guió sintètic de la prehistòria recent de Garraf. *La Cova de Can Sadurní i la prehistòria de Garraf* **30**: 13-97.

Filella I, Peñuelas J. 2003. Partitioning of water and nitrogen in co-occurring Mediterranean woody shrub species of different evolutionary history. *Oecologia* **137**: 51-61.

Fontanals-Coll M, Subirà ME, Bonilla MD-Z, Duboscq S, Gibaja JF. 2015. Investigating palaeodietary and social differences between two differentiated sectors of a Neolithic community, La Bòbila Madurell-Can Gambús (north-east Iberian Peninsula). *Journal of Archaeological Science: Reports* **3**: 160-170.

Fontanals-Coll M, Subirà ME, Díaz-Zorita Bonilla M, Gibaja JF. 2017. First insight into the Neolithic subsistence economy in the north-east Iberian Peninsula: paleodietary reconstruction through stable isotopes. *American Journal of Physical Anthropology* **162**: 36-50.

Fraser RA, Bogaard A, Heaton T, Charles M, Jones G, Christensen BT, Halstead P, Merbach I, Poulton P, Sparkes D, Styring AK. 2011. Manuring and stable nitrogen isotope ratios in cereals and pulses: towards a new archaeobotanical approach to the inference of land use and dietary practices. *Journal of archaeological science* **38**: 2790–2804.

Friedli H, Löttscher H, Oeschger H, Siegenthaler U, Stauffer B. 1986. Ice core record of the $^{13}\text{C}/^{12}\text{C}$ ratio of the atmospheric CO_2 in the past two centuries. *Nature* **324**: 237–238.

Fuller BT, De Cupere B, Marinova E, Van Neer W, Waelkens M, Richards MP. 2012. Isotopic reconstruction of human diet and animal husbandry practices during the Classical-Hellenistic, imperial, and Byzantine periods at Sagalassos, Turkey. *American Journal of Physical Anthropology* **149**: 157-171.

González J, Molist M, Spezzia KH. 2011. Un nou assentament del V mil.lenni a la costa de Barcelona. *Quarhis: Quaderns d'Arqueologia i Història de la Ciutat de Barcelona* **7**: 86-100.

Guerra MAR, Pena RG, de Lagrán IGM, Kunst M. 2008. Los recintos del poblado del Neolítico Antiguo de la Revilla del Campo (Ambrona, Soria). In *IV Congreso del Neolítico Peninsular: 27-30 de noviembre de 2006*. Museo Arqueológico de Alicante-MARQ; 252-258.

Hadjikoumis A. 2012. Traditional Pig Herding Practices in Southwest Iberia: Questions of Scale and Zooarchaeological Implications. *Journal of Anthropological Archaeology* **31**: 353-64.

Hain FH. 1982. Kupferzeitliche Tierknochenfunde aus Valencina de la Concepcion/Sevilla. *Studien über frühe Tierknochenfunde von der Iberischen Halbinsel* **8**: 1-178.

Halley DJ, Rosvold J. 2014. Stable isotope analysis and variation in medieval domestic pig husbandry practices in northwest Europe: absence of evidence for a purely herbivorous diet. *Journal of Archaeological Science* **49**: 1-5.

Halstead P, Isaakidou V. 2011. A pig fed by hand is worth two in the bush: ethnoarchaeology of pig husbandry in Greece and its archaeological implications. In *Ethnozooarchaeology: The present and past of human-animal relationships*, Albarella U, Trentacoste A. (eds.). Archaeology and Cultural Mixture: Cambridge; 160-74

Hamilton J, Hedges REM, Robinson M. 2009. Rooting for pigfruit: pig feeding in Neolithic and Iron Age Britain compared. *Antiquity* **83**: 998–1011.

Hamilton J, Thomas R. 2012. Pannage, Pulses and Pigs: Isotopic and Zooarchaeological

Evidence for Changing Pig Management Practices in Later Medieval England. *Medieval archaeology* **56**: 234-259.

Hammer Ø, Harper D, Ryan PD. 2001. PAST: Paleontological Statistics Software Package for education and data analysis. *Palaeontologia Electronica* **4**.

Hammond C, O'Connor T. 2013. Pig diet in medieval York: carbon and nitrogen stable isotopes. *Archaeological and Anthropological Science* **5**: 123-127.

Hedges RE, Clement JG, Thomas CDL, O'connell TC. 2007. Collagen turnover in the adult femoral mid-shaft: Modeled from anthropogenic radiocarbon tracer measurements. *American Journal of Physical Anthropology* **133**: 808-816.

Hedges RE, Reynard LM. 2007. Nitrogen isotopes and the trophic level of humans in archaeology. *Journal of Archaeological Science* **34**: 1240-1251.

Hellevang H, Aagaard P. 2015. Constraints on natural global atmospheric CO₂ fluxes from 1860 to 2010 using a simplified explicit forward model. *Scientific Reports* **5**: 17352. <http://doi.org/10.1038/srep17352>

Isaakidou, V. 2011. Farming regimes in Neolithic Europe: gardening with cows and other models. In *Dynamics of Neolithisation in Europe. Studies in Honour of Andrew Sherratt*. Hadjikoumis A, Robinson E, Viner S. (eds.). Oxford: Oxbow Books; 90-112

Jalut G, Dedoubat JJ, Fontugne M, Otto T. 2009. Holocene circum-Mediterranean vegetation changes: Climate forcing and human impact. *Quaternary international: the journal of the International Union for Quaternary Research* **200**: 4-18.

Jim S, Ambrose SH, Evershed R. 2004. Stable Carbon Isotopic Evidence for Differences in the Dietary Origin of Bone Cholesterol, Collagen and Apatite: Implications for Their Use in Palaeodietary Reconstruction. *Geochimica et Cosmochimica Acta* **68**: 61-72.

Madgwick R, Mulville J, Stevens RE. 2012. Diversity in foddering strategy and herd management in late Bronze Age Britain: An isotopic investigation of pigs and other fauna from two midden sites. *Environmental Archaeology* **17**: 126-140.

Marín AB, Morales MR. 2009. Comportamiento económico de los últimos cazadores-recolectores y primeras evidencias de domesticación en el occidente de Asturias. La Cueva de Mazaculos II. *Trabajos de prehistoria* **66**: 47-74.

Martín A, Edo M, Tarrus M, Clop X. 2010. Le Néolithique ancien de Catalogne (VI-première moitié du Ve millénaire av. J.-C.). Les séquences chronoculturelles. In *Premières sociétés paysannes de Méditerranée occidentale: structure des productions céramiques*, Manen C, Convertini F, Binder D, Sénépart I (eds.). Société Préhistorique Française:

Toulouse; 197-214.

Martín A, Estévez J. 1991. Funció de la Cova del Frare de St. Llorenç de Munt (Matadepera, Barcelona) al Neolític Antic en relació a la ramaderia. Estat de la Investigació sobre el Neolític a Catalunya. 9è Col·loqui Internacional d'Arqueologia de Puigcerdà Institut d'Estudis Ceretans, Puigcerdà: 105-108.

Martín A, Guilaine J, Thommeret Y. 2009. Estratigrafia y dataciones C14 del yacimiento de la «Cova del Frare» de St. Llorenç del Munt (Matadepera, Barcelona). *Zephyrus* **32**: 33.

Matsui A, Ishiguro N, Hongo H, Minagawa M. 2005. Wild pig? Or domesticated boar? An archaeological view on the domestication of *Sus scrofa* in Japan. In *The first steps of animal domestication*, Vigne JD, Peters J, Helmer D (eds.). Oxbow Books; 148-159.

Minagawa M, Matsui A, Ishiguro N. 2005. Patterns of prehistoric boar *Sus scrofa* domestication, and inter-islands pig trading across the East China Sea, as determined by carbon and nitrogen isotope analysis. *Chemical Geology* **218**: 91-102.

Molist M, Campos OV, Farré R. 2008. El jaciment de la Caserna de Sant Pau del Camp: aproximació a la caracterització d'un assentament del Neolític antic. *Quarhis: Quaderns d'Arqueologia i Història de la Ciutat de Barcelona* **4**: 14-24.

Müldner G, Richards MP. 2005. Fast or feast: reconstructing diet in later medieval England by stable isotope analysis. *Journal of archaeological Science* **32**: 39-48.

Navarrete V, Saña M. 2017. Size changes in wild and domestic pig populations between 10,000-800 cal BC in the Iberian Peninsula: evaluation of natural versus social impacts in animal populations during the first domestication stages. *The Holocene*. <http://dx.doi.org/10.1177%2F0959683617693902>

O'Leary MH. 1988. Carbon isotopes in photosynthesis: Fractionation techniques may reveal new aspects of carbon dynamics in plants. *Bioscience* **38**: 328-336.

O'Connell TC, Kneale CJ, Tasevska N, Kuhnle G. 2012. The diet-body offset in human nitrogen isotopic values: A controlled dietary study. *American Journal of Physical Anthropology* **149**: 426-434.

Oms X. 2014. *La neolitització del Nord-Est de la Península Ibèrica a partir de les datacions de 14 (C) i les primeres ceràmiques impreses c. 5600-4900 cal BC*. Phd Thesis. Universitat de Barcelona.

Palomo A, Piqué R, Terradas X, Bosch À, Buxo R, Chinchilla J, Tarrus J. 2014. Prehistoric occupation of Banyoles lakeshore: results of recent excavations at La Draga site, Girona, Spain. *Journal of Wetland Archaeology* **14**: 58-73.

Payne S, Bull G. 1988. Components of variation in measurements of pig bones and teeth, and the use of measurements to distinguish wild from domestic pig remains. *Archaeozoologia* **2**: 27–65.

Pechenkina EA, Ambrose SH, Xiaolin M, Benfer R A. 2005. Reconstructing northern Chinese Neolithic subsistence practices by isotopic analysis. *Journal of Archaeological Science* **32**: 1176–1189.

Reimer PJ, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, Grootes PM, Guilderson TP, Hafliðason H, Hajdas I, Hatt C, Heaton TJ, Hoffmann DL, Hogg AG, Hughen KA, Kaiser KF, Kromer B, Manning SW, Niu M, Reimer RW, Richards DA, Scott EM, Southon JR, Staff RA, Turney CSM, van der Plicht J. 2013. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon* **55**.

Revelles J, Burjachs F, van Geel B. 2016. Pollen and non-pollen palynomorphs from the Early Neolithic settlement of La Draga (Girona, Spain). *Review of Palaeobotany and Palynology* **225**: 1–20.

Rosillo R, Tarrús J, Palomo A, García R. 2010. Les esteles amb banyes de la Serra del Mas Bonet (Vilafant, Alt Empordà) dins de l'art megalític de Catalunya. *Cypsela: revista de prehistòria i protohistòria* **18**: 43–59.

Saña M, Antolín F, Zapata M, Castells L, Craig OE, Benaiges ME, Spiteru C. 2015. Prácticas agropecuarias durante el Neolítico antiguo y medio en la cueva de Can Sadurní (Begues, Barcelona) In *5º Congresso do Neolítico Peninsular*, Gonçalves V, Diniz M, Sousa AC (eds.). Centro do Arqueologia da Universidade de Lisboa: Lisboa; 57–66.

Saña M, Bogdanovic I, Navarrete V. 2014. Taphonomic evaluation of the degree of historical representation of the archaeological bone samples in anaerobic versus aerobic environments: The Neolithic site of La Draga (Banyoles, Spain). *Quaternary International* **330**: 72–87.

Saña M, Navarrete V. 2016. Gestió ramadera al pla de Barcelona durant la prehistòria. In *La prehistòria al pla de Barcelona. Nous documents per a una síntesi*, Molist M, Gómez A (eds.). MUHBA. Ajuntament de Barcelona: Barcelona; 97–104.

Saña M. 1998. Entorn a la dinàmica del procés de domesticació animal. *Cypsela* **12**: 99–110.

Saña M. 2011. La gestió dels recursos animals. In *El poblat lacustre del Neolític antic de la Draga, Excavacions 2000–2005*, Bosch A, Chinchilla J, Tarrús J (eds.). Monografies del CASC 9: Girona; 50–64.

Saña M. 2013. Domestication of animals in the Iberian Peninsula. In *The origins and spread of domestic animals in southwest Asia and Europe*, Colledge S, Conolly J, Dobney K, Manning K, Shennan S (eds.). UCL Institute of Archaeology Publications: London; 195-220.

Saña M. 2014. *Anàlisi Arqueozoològica del jaciment de Serra de Mas Bonet (Vilafant, Girona)*. Laboratori d'Arqueozoològia. Universitat Autònoma de Barcelona.

Schoeninger MJ, DeNiro MJ. 1984. Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochimica et Cosmochimica Acta* **48**: 625-639.

Szwarcz HP, Schoeninger MJ. 1991. Stable isotope analyses in human nutritional ecology. *American Journal of Physical Anthropology* **34**: 283-321.

Sealy J. 2001. Body tissue chemistry and palaeodiet. In *Handbook of archaeological sciences*, Brothwell DR, Pollard AM (eds.). John Wiley & Sons: Michigan; 269-279.

Sykes N. 2014. *Beastly questions: Animal answers to archaeological issues*. Bloomsbury Publishing: London.

Szpak P. 2014. Complexities of nitrogen isotope biogeochemistry in plant-soil systems: implications for the study of ancient agricultural and animal management practices. *Frontiers in plant science* **5**: 288.

Terradas X, Antolín F, Bosch A, Buxó R, Chinchilla J, Clop J, Gibaja JF, Oliva M, Palomo A, Piqué R, Saña M, Tarrús J. 2012. Áreas de aprovisionamiento, territorios de subsistencia y producciones técnicas en el Neolítico Antiguo de La Draga. *Revista del Museu de Gava* **5**: 441-448.


Terradas-Batlle X, Piqué R, Palomo A, Bosch À, Buxó R, Chinchilla J, Rosillo R. 2015. Darreres intervencions arqueològiques al poblat neolític de La Draga al l'estany de Banyoles (Banyoles, Pla de l'Estany). In *Tribuna d'Arqueologia 2012-2013*, Departament de Cultura de la Generalitat de Catalunya (ed.). Entitat Autònoma del Diari Oficial i de Publicacions: Barcelona; 33-47.

Tieszen L, Boutton L, Tesdahl K, Slade N. 1983. Fractionation and Turnover of Stable Carbon Isotopes in Animal Tissues: Implications for $\delta^{13}\text{C}$ Analysis of Diet. *Oecologia* **57**: 32-37.

Van Klinken GJ. 1999. Bone collagen quality indicators for palaeodietary and radiocarbon measurements. *Journal of Archaeological Science* **26**: 687-695.

Zeder MA. 2008. Domestication and early agriculture in the Mediterranean Basin: Origins, diffusion, and impact. *Proceedings of the national Academy of Sciences* **105**: 11597-11604.

For Peer Review



Location of the sites in the northeastern Iberian Peninsula.

64x58mm (300 x 300 DPI)





Figure 2. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of animals from cova del Frare, Reina Amàlia-Caserna de Sant Pau, Can Sadurní, La Draga and Serra de Mas Bonet.
 Figure 3A-B. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of pigs (unfilled circle) corrected for the average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of herbivores ($\Delta\text{‰}$) for cova del Frare (CF), La Draga (DR), Reina Amàlia-Caserna de Sant Pau (RA-CSP) and Can Sadurní (CS). We also corrected the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of herbivores for their average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (filled circles) in order to show their isotopic variability.

155x108mm (300 x 300 DPI)



$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of pigs (unfilled circle) corrected for the average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of herbivores ($\Delta\text{‰}$) for cova del Frare (CF), La Draga (DR), Reina Amàlia-Caserna de Sant Pau (RA-CSP) and Can Sadurní (CS). We also corrected the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of herbivores for their average $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (filled circles) in order to show their isotopic variability.

133x56mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	Location	Level	¹⁴ C yr BP	Material	¹⁴ C yr cal BP (2σ)	¹⁴ C yr cal BC (2σ)	Reference
6	Gerona	-	-	Animal bone	-	4900-4600	Rosillo et al., 2010
7	Gerona	-	6010±70	Animal bone	7150-6670	5201-4721	Terradas et al., 2015
8	Barcelona	C6	6150±40	Animal bone	7165-6942	5216-4993	This study
9	Barcelona	C5	6020±40	Animal bone	6950-6770	5000-4820	Oms, 2014
10	Barcelona	I-II-III	5670±40	Animal bone	6560-6322	4611-4373	González et al., 2011
11	Barcelona	-	6290±50	Animal bone	7321-7025	5372-5076	Molist et al., 2008
12	Barcelona	10b/11	5540±40	Animal bone	6405-6284	4456-4335	Edo et al., 2011
13	Barcelona	17	6050±110	Charcoal	7240-6659	5291-4710	Blasco et al., 1999

Site	Specimens ID	Niv	Species ID	Skeleton part	%yield	%C	%N	$\delta^{13}\text{C}(\text{‰})$	$\delta^{15}\text{N}(\text{‰})$	C:N	LAB
Cova del Frare	48	C5	<i>Sus domesticus</i>	Maxilla	1.4	36	13	-20.1	6.3	3.2	UAB
Cova del Frare	50	C5	<i>Sus domesticus</i>	Maxilla	5.8	43	15	-20.0	6.0	3.3	UAB
Cova del Frare	45	C5	<i>Sus domesticus</i>	Mandible	6.7	43	15	-19.6	5.8	3.3	UAB
Cova del Frare	67	C5	<i>Sus domesticus</i>	Mandible	4.7	41	15	-19.9	6.0	3.2	UAB
Cova del Frare	46	C5	<i>Sus scrofa</i>	Mandible	1.3	31	11	-19.9	4.5	3.3	UAB
Cova del Frare	68	C5	<i>Sus domesticus</i>	Mandible	4.3	39	15	-20.7	6.9	3.0	UAB
Cova del Frare	53	C5	<i>Sus domesticus</i>	Maxilla	4.9	43	15	-20.0	5.0	3.3	UAB
Cova del Frare	49	C6	<i>Sus domesticus</i>	Maxilla	3.0	39	14	-19.8	4.4	3.3	UAB
Cova del Frare	54	C5	<i>Sus domesticus</i>	Maxilla	4.0	42	15	-20.1	6.9	3.3	UAB
Cova del Frare	52	C6	<i>Sus domesticus</i>	Maxilla	2.7	42	15	-20.2	6.3	3.3	UAB
Cova del Frare	47	C5	<i>Sus domesticus</i>	Mandible	1.7	42	14	-19.9	4.4	3.5	UAB
Cova del Frare	51	C6	<i>Sus domesticus</i>	Humerus	5.9	38	13	-20.1	4.2	3.4	UAB
Cova del Frare	66	C5	<i>Sus domesticus</i>	Humerus	4.9	42	15	-19.9	5.7	3.3	UAB
Cova del Frare	36	C6	<i>Sus domesticus</i>	Humerus	3.1	40	14	-20.3	5.5	3.3	UAB
Cova del Frare	60	C6	<i>Bos taurus</i>	Metatarsus	3.7	34	12	-19.4	5.3	3.3	UAB
Cova del Frare	63	C6	<i>Capra hircus</i>	Humerus	2.3	40	15	-20.1	6.9	3.1	UAB
Cova del Frare	57	C5	<i>Capra hircus</i>	Humerus	4.7	41	15	-20.3	3.9	3.2	UAB
Cova del Frare	56	C5	<i>Capra hircus</i>	Humerus	5.2	41	15	-20.6	4.2	3.2	UAB
Cova del Frare	62	C6	<i>Capra hircus</i>	Humerus	2.4	42	15	-20.6	3.5	3.3	UAB
Cova del Frare	58	C5	<i>Capra hircus</i>	Humerus	1.9	33	12	-20.3	6.6	3.2	UAB
Cova del Frare	72	C5	<i>Capra hircus</i>	Metacarpus	6.9	41	15	-20.2	4.1	3.2	UAB
Cova del Frare	65	C6	<i>Capreolus capreolus</i>	Metatarsus	3.2	34	13	-20.2	3.9	3.1	UAB
Cova del Frare	55	C5	<i>Cervus elephus</i>	Metatarsus	4.1	39	14	-18.6	5.4	3.3	UAB
Cova del Frare	69	C5	<i>Cervus elephus</i>	Tibia	4.1	39	15	-21.1	4.8	3.0	UAB
Cova del Frare	61	C6	<i>Ovis aries</i>	Humerus	4.3	40	15	-19.9	6.6	3.1	UAB
Cova del Frare	73	C5	<i>Ovis aries</i>	Tibia	5.7	41	15	-20.9	4.3	3.2	UAB
Caserna de Sant Pau	99	XV	<i>Sus domesticus</i>	Maxilla	1.4	32	11	-20.9	9.0	3.4	UAB

Caserna de Sant Pau	94	XXV	<i>Sus domesticus</i>	Mandible	1.7	40	14	-21.4	7.3	3.3	UAB
Caserna de Sant Pau	82	XXV	<i>Sus domesticus</i>	Humerus	1.1	36	12	-20.7	6.1	3.5	UAB
Caserna de Sant Pau	93	XVIII	<i>Sus domesticus</i>	Radius	1.2	37	14	-20.7	8.9	3.1	UAB
Caserna de Sant Pau	86	XVII	<i>Sus domesticus</i>	Tibia	1.6	43	16	-20.7	6.7	3.1	UAB
Caserna de Sant Pau	92	XXIII	<i>Capra hircus</i>	Humerus	2.5	42	16	-19.8	6.8	3.1	UAB
Caserna de Sant Pau	90	XII	<i>Bos taurus</i>	Metacarpus	2.4	40	14	-21.6	6.6	3.3	UAB
Caserna de Sant Pau	88	XVII	<i>Ovis aries</i>	Metatarsus	2.1	39	14	-21.3	7.2	3.3	UAB
Can Sadurní	44	10	<i>Sus domesticus</i>	Mandible	1.4	33	12	-19.2	7.1	3.2	UAB
Can Sadurní	42	10	<i>Sus domesticus</i>	Mandible	2.7	37	13	-19.8	7.5	3.3	UAB
Can Sadurní	43	11b	<i>Sus domesticus</i>	Mandible	1.9	40	14	-19.9	6.2	3.3	UAB
Can Sadurní	122	17	<i>Sus scrofa</i>	Phalanx	1.0	37	13	-20.0	7.6	3.3	UAB
Can Sadurní	126	17	<i>Capra hircus</i>	Mandible	0.4	31	11	-20.1	4.7	3.3	UAB
Can Sadurní	129	10	<i>Bos taurus</i>	Metatarsus	2.7	35	13	-20.1	4.1	3.1	UAB
Can Sadurní	132	10	<i>Canis familiaris</i>	Metatarsus	2.3	43	16	-19.0	8.0	3.1	UAB
La Draga	2	B	<i>Sus domesticus</i>	Humerus	3.0	43	16	-21.1	4.1	3.1	UAB
La Draga	16	B	<i>Sus domesticus</i>	Mandible	7.2	42	15	-20.6	4.0	3.3	UAB
La Draga	20	B	<i>Sus domesticus</i>	Maxilla	9.4	42	14	-20.5	4.5	3.5	UAB
La Draga	10	B	<i>Sus domesticus</i>	Tibia	4.3	42	15	-20.0	5.1	3.3	UAB
La Draga	18	B	<i>Sus domesticus</i>	Mandible	2.9	42	14	-20.7	4.3	3.5	UAB
La Draga	1	B	<i>Sus domesticus</i>	Humerus	4.1	42	15	-20.7	4.6	3.3	UAB
La Draga	5	B	<i>Sus domesticus</i>	Humerus	1.3	36	12	-21.4	5.6	3.5	UAB
La Draga	7	A	<i>Sus domesticus</i>	Radius	5.6	39	14	-19.1	7.5	3.3	UAB
La Draga	D-26	A	<i>Sus domesticus</i>	Tibia	3.3	41	15	-20.3	5.1	3.2	York
La Draga	21	B	<i>Sus domesticus</i>	Maxilla	4.9	42	14	-20.8	5.2	3.5	UAB
La Draga	17	B	<i>Sus domesticus</i>	Mandible	5.1	43	15	-20.9	4.0	3.3	UAB
La Draga	15	B	<i>Sus domesticus</i>	Tibia	10.6	44	15	-21.0	4.9	3.4	UAB
La Draga	12	B	<i>Sus domesticus</i>	Radius	9.7	42	15	-21.0	6.7	3.3	UAB
La Draga	11	B	<i>Sus domesticus</i>	Radius	5.1	42	15	-20.4	5.0	3.3	UAB

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

La Draga	8	B	<i>Sus domesticus</i>	Radius	3.1	42	15	-20.6	5.3	3.3	UAB
La Draga	9	B	<i>Sus domesticus</i>	Tibia	2.3	42	15	-20.9	4.7	3.3	UAB
La Draga	19	B	<i>Sus domesticus</i>	Maxilla	6.2	43	14	-19.8	5.1	3.6	UAB
La Draga	4	A	<i>Sus domesticus</i>	Humerus	1.2	40	13	-21.0	3.6	3.6	UAB
La Draga	28	D	<i>Sus domesticus</i>	Humerus	1.1	37	12	-20.7	5.6	3.6	UAB
La Draga	26	D	<i>Sus domesticus</i>	Tibia	3.8	42	15	-20.2	4.3	3.3	UAB
La Draga	29	B	<i>Sus domesticus</i>	Radius	6.7	43	15	-20.4	4.8	3.3	UAB
La Draga	27	B	<i>Sus domesticus</i>	Humerus	4.2	42	15	-20.3	5.0	3.3	UAB
La Draga	D-18	B	<i>Sus domesticus</i>	Humerus	6.9	42	15	-20.2	5.2	3.3	York
La Draga	D-34	B	<i>Meles meles</i>	Humerus	1.2	35	12	-21.3	12.6	3.5	York
La Draga	165	B	<i>Bos taurus</i>	Humerus	7.6	36	13	-18.8	6.3	3.2	UAB
La Draga	163	B	<i>Bos taurus</i>	Tibia	2.3	36	13	-19.8	5.4	3.2	UAB
La Draga	D-28	B	<i>Canis familiaris</i>	Tibia	6.6	41	15	-21.5	6.5	3.2	York
La Draga	159	B	<i>Capra hircus</i>	Humerus	2.8	38	13	-20.3	5.9	3.4	UAB
La Draga	10A	B	<i>Capra pyrenaica</i>	Humerus	5.7	42	15	-21.8	4.5	3.3	UAB
La Draga	20A	B	<i>Capra pyrenaica</i>	Humerus	3.2	35	12	-20.1	3.7	3.4	UAB
La Draga	13A	B	<i>Capra pyrenaica</i>	Metacarpus	9.1	36	12	-20.3	5.0	3.5	UAB
La Draga	D-10	B	<i>Capreolus capreolus</i>	Humerus	4.2	29	10	-21.9	4.4	3.4	York
La Draga	D-6	B	<i>Cervus elephus</i>	Phalanx	2.1	45	15	-21.0	4.3	3.5	York
La Draga	D-11	B	<i>Meles meles</i>	Humerus	7.8	47	17	-19.0	7.7	3.2	York
La Draga	162	B	<i>Ovis aries</i>	Tibia	1.0	34	12	-19.0	4.7	3.3	UAB
La Draga	160	B	<i>Ovis aries</i>	Radius	2.2	38	13	-20.5	4.6	3.4	UAB
Reina Amàlia	137	FIII	<i>Sus scrofa</i>	Humerus	0.7	29	10	-19.3	7.4	3.4	UAB
Reina Amàlia	136	FII	<i>Sus domesticus</i>	Humerus	1.7	34	13	-19.5	8.0	3.1	UAB
Reina Amàlia	149	FII	<i>Sus domesticus</i>	Mandible	1.4	34	13	-20.3	8.6	3.1	UAB
Reina Amàlia	140	FI	<i>Sus domesticus</i>	Radius	1.8	42	16	-19.6	7.8	3.1	UAB
Reina Amàlia	143	FI	<i>Sus domesticus</i>	Maxilla	1.2	32	12	-20.1	7.9	3.1	UAB
Reina Amàlia	147	FI	<i>Sus domesticus</i>	Mandible	1.0	35	13	-21.0	7.3	3.1	UAB

Reina Amàlia	139	FI	<i>Sus domesticus</i>	Humerus	1.6	34	13	-20.8	7.9	3.1	UAB
Reina Amàlia	138	FII	<i>Sus domesticus</i>	Humerus	1.3	35	13	-19.4	5.3	3.1	UAB
Reina Amàlia	145	FII	<i>Sus domesticus</i>	Mandible	1.6	36	13	-20.2	8.7	3.2	UAB
Reina Amàlia	146	FI	<i>Sus domesticus</i>	Mandible	0.7	33	11	-21.1	7.6	3.5	UAB
Reina Amàlia	41	FI	<i>Sus domesticus</i>	Mandible	1.3	40	14	-21.0	6.3	3.3	UAB
Reina Amàlia	40	FI	<i>Sus domesticus</i>	Mandible	1.9	42	15	-20.6	8.5	3.3	UAB
Reina Amàlia	144	FI	<i>Sus domesticus</i>	Mandible	1.5	40	14	-21.4	7.3	3.3	UAB
Reina Amàlia	142	FI	<i>Sus scrofa</i>	Metatarsus	1.3	36	13	-20.3	6.6	3.2	UAB
Reina Amàlia	141	FII	<i>Sus scrofa</i>	Metatarsus	1.1	36	14	-20.0	5.6	3.0	UAB
Reina Amàlia	150	FII	<i>Bos taurus</i>	Metatarsus	1.3	40	15	-20.2	4.9	3.1	UAB
Reina Amàlia	151	FIV	<i>Bos taurus</i>	Tibia	1.0	31	11	-20.5	6.1	3.3	UAB
Reina Amàlia	155	FII	<i>Capra pyrenaica</i>	Radius	1.1	38	14	-20.2	4.8	3.2	UAB
Serra de Mas Bonet	181	-	<i>Sus domesticus</i>	Tibia	1.5	32	11	-20.4	5.0	3.4	UAB
Serra de Mas Bonet	180	-	<i>Sus domesticus</i>	Metacarpus	1.6	37	13	-19.7	7.4	3.3	UAB
Serra de Mas Bonet	182	-	<i>Sus domesticus</i>	Phalanx	1.7	37	12	-20.5	6.2	3.6	UAB
Serra de Mas Bonet	171	-	<i>Canis familiaris</i>	Mandible	0.6	36	12	-19.0	11.1	3.5	UAB
Serra de Mas Bonet	174	-	<i>Capra hircus</i>	Humerus	0.6	27	10	-20.8	6.5	3.2	UAB