



This is a repository copy of *Provision of long-term monitoring and late effects services following adult allogeneic haematopoietic stem cell transplant: a survey of UK NHS-based programmes.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/118154/>

Version: Supplemental Material

---

**Article:**

Hamblin, A., Greenfield, D.M., Gilleece, M. et al. (9 more authors) (2017) Provision of long-term monitoring and late effects services following adult allogeneic haematopoietic stem cell transplant: a survey of UK NHS-based programmes. *Bone Marrow Transplantation*, 52. pp. 889-894. ISSN 0268-3369

<https://doi.org/10.1038/bmt.2017.67>

---

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**Supplementary Table 1**

Time Point	Characteristic	Number of Centres			P value
		Percentage of patients being followed-up			
		11-50	51-90	91-100	
<b>Number of adult allogeneic HSCTs performed in 2014 (n=27)</b>					
1-5 years	1-59 <sup>a</sup> (n=16)	0	5	11	0.44
	>60 <sup>b</sup> (n=11)	0	2	9	
5-10 years	1-59	2	7	7	0.86
	>60	1	4	6	
Beyond 10 years	1-59	4	6	6	0.39
	>60	2	7	2	
<b>Centre also undertaking paediatric allogeneic HSCTs<sup>c</sup> (n=27)</b>					
1-5 years	Yes (n=14)	0	5	9	0.23
	No (n=13)	0	2	11	
5-10 years	Yes	2	7	5	0.40
	No	1	4	8	
Beyond 10 years	Yes	4	7	3	0.55
	No	2	6	5	

Comparison of reported follow-up patterns for centres according to allogeneic HSCT activity, with or without co-existent paediatric activity.

Chi-square tests were performed to determine statistical significance.

<sup>a</sup>Range 3-51, median 34 allogeneic HSCTs performed.

<sup>b</sup>Range 60-105, median 80 allogeneic HSCTs performed.

<sup>c</sup>Centres were defined as having co-existent paediatric allogeneic HSCT activity if they were either officially designated as a paediatric centre for allogeneic HSCTs by BSBMT or undertook an allogeneic HSCT in an individual under 18 years of age in 2014.