



This is a repository copy of *The Near East as a cradle of biodiversity: a phylogeography of banded newts (genus Ommatotriton) reveals extensive inter- and intraspecific genetic differentiation.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/118294/>

Version: Published Version

Article:

van Riemsdijk, I., Arntzen, J.W., Bogaerts, S. et al. (4 more authors) (2017) The Near East as a cradle of biodiversity: a phylogeography of banded newts (genus Ommatotriton) reveals extensive inter- and intraspecific genetic differentiation. *Molecular Phylogenetics and Evolution*, 114. pp. 73-81. ISSN 1055-7903

<https://doi.org/10.1016/j.ympev.2017.05.028>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



The Near East as a cradle of biodiversity: A phylogeography of banded newts (genus *Ommatotriton*) reveals extensive inter- and intraspecific genetic differentiation



Isolde van Riemsdijk^{a,*}, Jan W. Arntzen^a, Sergé Bogaerts^b, Michael Franzen^c, Spartak N. Litvinchuk^d, Kurtuluş Olgun^e, Ben Wielstra^{a,f,g}

^a Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands

^b Lupinelaan 25, 5582 CG Waalre, The Netherlands

^c Zoologische Staatssammlung München (ZSM-SNSB), Münchhausenstraße 21, 81247 München, Germany

^d Institute of Cytology, Russian Academy of Sciences, Tikhoretsky pr. 4, 194064 St. Petersburg, Russia

^e Department of Biology, Faculty of Arts and Sciences, Adnan Menderes University, 09010 Aydın, Turkey

^f Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA 90095, USA

^g Department of Animal and Plant Sciences, University of Sheffield, S10 2TN Sheffield, UK

ARTICLE INFO

Article history:

Received 7 September 2016

Revised 4 April 2017

Accepted 31 May 2017

Available online 13 June 2017

Keywords:

Anatolia

Historical biogeography

Levant

Molecular dating

Phylogeny

Species distribution modelling

ABSTRACT

The banded newt (genus *Ommatotriton*) is widely distributed in the Near East (Anatolia, Caucasus and the Levant) – an understudied region from the perspective of phylogeography. The genus is polytypic, but the number of species included and the phylogenetic relationships between them are not settled. We sequenced two mitochondrial and two nuclear DNA markers throughout the range of *Ommatotriton*. For mtDNA we constructed phylogenetic trees, estimated divergence times using fossil calibration, and investigated changes in effective population size with Bayesian skyline plots and mismatch analyses. For nuDNA we constructed phylogenetic trees and haplotype networks. Species trees were constructed for all markers and nuDNA only. Species distribution models were projected on current and Last Glacial Maximum climate layers. We confirm the presence of three *Ommatotriton* species: *O. nesterovi*, *O. ophryticus* and *O. vittatus*. These species are genetically distinct and their most recent common ancestor was dated at ~25 Ma (Oligocene). No evidence of recent gene flow between species was found. The species show deep intraspecific genetic divergence, represented by geographically structured clades, with crown nodes of species dated ~8–13 Ma (Miocene to Early Quaternary); evidence of long-term *in situ* evolution and survival in multiple glacial refugia. While a species tree based on nuDNA suggested a sister species relationship between *O. vittatus* and *O. ophryticus*, when mtDNA was included, phylogenetic relationships were unresolved, and we refrain from accepting a particular phylogenetic hypothesis at this stage. While species distribution models suggest reduced and fragmented ranges during the Last Glacial Maximum, we found no evidence for strong population bottlenecks. We discuss our results in the light of other phylogeographic studies from the Near East. Our study underlines the important role of the Near East in generating and sustaining biodiversity.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Phylogeography reconstructs phylogenetic patterns in space and time to identify the drivers that shape biodiversity within taxa (Avice, 2000; Hickerson et al., 2010). The Mediterranean Basin is considered a near-ideal laboratory for phylogeographic studies (Hewitt, 2011). The collision of the Eurasian, African and Arabian

tectonic plates, from the Late Eocene (37–34 Ma) onwards, caused major geological changes in the Mediterranean Basin, as marine barriers were re-arranged and mountains formed (Meulenkamp and Sissingh, 2003; Popov et al., 2006). This process left its signature in the phylogeography of the species inhabiting the region (Hewitt, 2011). While the Quaternary (2.6 Ma – present) glacial cycles resulted in drastic climate shifts in temperate Eurasia, the climate in the Mediterranean Basin remained relatively stable (Stewart et al., 2009; Hewitt, 2011). As a consequence, many species had a continuous presence within the Mediterranean Basin,

* Corresponding author.

E-mail address: isolde.vanriemsdijk@naturalis.nl (I. van Riemsdijk).

while they only periodically colonized more northern areas during interglacials – and subsequently were wiped out during the next glacial cycle (Hewitt, 2000).

Because of the interaction of geographic instability and climatic stability, many Mediterranean species are characterized by deep intraspecific lineages (Oosterbroek and Arntzen, 1992; Bilgin, 2011). The Mediterranean Basin features a high concentration of biodiversity and is considered a biodiversity hotspot (Myers et al., 2000). Complex phylogeographic patterns are particularly well documented for the Iberian, Italian and Balkan peninsulas (Hewitt, 2011). Despite the similar geologic and climatologic background of the Near East, historical biogeographical studies of this region are underrepresented in the phylogeographic literature (Beheregaray, 2008; Riddle, 2016).

Most amphibian species, especially salamanders, are particularly influenced by geological disruptions and climate oscillations, due to their limited dispersal capabilities and strong dependence on water, often resulting in a pronounced phylogenetic structure (Vences and Wake, 2007). Seven Salamandridae genera (*Lissotriton*, *Lyciasalamandra*, *Mertensiella*, *Neurergus*, *Ommatotriton*, *Salamandra*, and *Triturus*) occur in the Near East. Phylogeographic studies of these genera revealed long term *in situ* evolution and survival during glacial cycles (Steinfartz et al., 2000; Tarkhnishvili et al., 2000; Veith et al., 2008; Wielstra et al., 2010; Hendrix et al., 2014; Pabijan et al., 2015). However, banded newts (genus *Ommatotriton*) remain to be studied in-depth. Because *Ommatotriton* is the most wide-spread salamander genus within the Near East (Sparreboom, 2014), a phylogeographic survey is expected to be insightful regarding the region's historical biogeography.

Ommatotriton is endemic to the Near East (Fig. 1). The genus was considered a single species, *O. vittatus* (Gray, 1835), with a disjunct northern and southern distribution, until Litvinchuk et al. in 2005 proposed that the northern banded newt range segment should be regarded as a separate, distinct species, *O. ophryticus* (Berthold, 1846). This treatment is now generally accepted (e.g.

Sparreboom, 2014; Frost, 2016). A recent proposal by Bülbül and Kutrup (2013) to split the northern banded newt range segment into two species, *O. ophryticus* in the east and *O. nesterovi* (Litvinchuk et al., 2005) in the west, is not universally supported. For convenience we refer to the three taxa as species from here on. The phylogeny of *Ommatotriton* is unclear. Considering three *Ommatotriton* species, three branching orders are possible and each is supported by existing data. Phylogenetic hypothesis I ((*O. ophryticus*, *O. vittatus*), *O. nesterovi*) is supported by mtDNA data (Bülbül and Kutrup, 2013), II ((*O. ophryticus*, *O. nesterovi*), *O. vittatus*) by allozyme data (Arntzen and Olgun, 2000; Litvinchuk et al., 2005), and III ((*O. vittatus*, *O. nesterovi*), *O. ophryticus*) by the number of rib-bearing vertebrae (Arntzen and Olgun, 2000; Arntzen et al., 2015; Litvinchuk et al., 2005).

Previous studies dealing with the inter- and intraspecific phylogenetic patterns of *Ommatotriton* suffered from limited sampling. We here conduct range-wide sampling. We use mitochondrial and nuclear DNA (mtDNA and nuDNA) markers to determine inter- and intraspecific phylogenetic relationships, conduct fossil-based time calibration and test for interspecific hybridization. Additionally, species distribution models are constructed to evaluate the impact of the Quaternary climate oscillations, and compared to historical demographic reconstructions. We identify interspecific and intraspecific phylogeographic patterns of *Ommatotriton* and discuss our findings in the context of a comparative phylogeography of the Near East. Our study highlights the importance of the Near East as a biodiversity hotspot.

2. Material and methods

2.1. Sampling and sequencing

We collected tissue samples of 1–3 individuals (average 2.6, total 106) from 40 localities (Fig. 1, Table A.1). The Thermo Scien-




Fig. 1. Distribution of and sampling scheme for banded newts (genus *Ommatotriton*). Coloured dashed lines roughly delineate the three species' ranges and black dashed lines delineate mtDNA-based phylogeographic groups (A–M). Numbered dots (1–45) represent sampled localities and small dots additional localities used for species distribution modelling. Grey shading reflects elevation. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

tific KingFisher Purification System was used to extract DNA. Fragments of two mitochondrial protein coding markers were sequenced: cytochrome *b* (cyt *b*, 786 bp) and cytochrome *c* oxidase I (COI, 658 bp). We added COI sequences for 13 individuals from five additional localities from Smith et al. (2008). Fragments of two unlinked nuclear protein coding markers (Hendrix et al., 2014; Shen et al., 2012), KIAA1239 (600 bp; hereafter KIAA) and SACS (624 bp), were sequenced for a single individual per locality, but included all three individuals for the three *O. ophryticus* and three *O. nesterovi* locations closest to their parapatric contact zone as defined by the mtDNA data (see Results, total 47, Table A.1).

We adjusted the ‘universal’ cyt *b* primers (MVZ-15-NP, MVZ-16-NP) from Moritz et al. (1992) and COI primers (VF1-d, VR1-d) from Ivanova et al. (2006) to match the full mitogenome of *O. nesterovi* (Zhang et al., 2008; Table A.2). We designed *Ommatotriton*-specific primers based on a KIAA and a SACS (Shen et al., 2012) sequence for *O. nesterovi* presented in Hendrix et al. (2014). In cases where no PCR product could be obtained, presumably due to degraded DNA, internal primers were designed to amplify stretches of 100–300 bp (three primer sets for COI, KIAA and SACS, and four for cyt *b*; Table A.3).

PCR conditions for DNA amplification were: initial denaturation for 180 s at 94 °C, 35 cycles with 30 s denaturation at 94 °C, 30 s annealing at 58 °C and 60 s extension at 72 °C, and a 240 s final extension step at 72 °C. Sanger sequencing was done commercially on an ABI 3730xl DNA analyser at BaseClear B.V., Leiden, the Netherlands. Sequences were assembled in Sequencher 4.10.1 (Gene Codes Corporation, MI USA) and aligned in Mesquite 3.04 (Maddison and Maddison, 2015). Nuclear alleles were phased with the Phase 2.1 algorithm (Stephens and Donnelly, 2003) implemented in DnaSP 5 (Librado and Rozas, 2009). GenBank accession numbers are listed in Table A.1.

2.2. Phylogenetic analyses

The mtDNA alignment was divided into six partitions, by marker (COI and cyt *b*) and by codon position (first, second and third). *Lissotriton vulgaris* and *Neurergus strauchii* were added as outgroups (data from Zhang et al., 2008; Table A.2). Maximum likelihood (ML) analysis was performed with RAxML 8.2.4 (Stamatakis, 2014), with 1000 rapid bootstrap replicates and a search for the best scoring maximum likelihood tree. The nucleotide substitution model GTR+G was used for each partition, as recommended in the RAxML manual. Bayesian phylogenetic reconstruction was performed with MrBayes 3.2.6 (Ronquist et al., 2012). We determined the best fitting models of sequence evolution for each of the six partitions based on the Bayesian information criterion (Guindon and Gascuel, 2003) in jModelTest 2.1.7 (Darriba et al., 2012; Table A.4). A Markov Chain Monte Carlo search was run for a million generations under default parameters (two runs, four chains, with temperature 0.2), a sampling frequency of 0.001 and a burn-in of 25% of generations. State frequencies were unlinked and rates were set to vary across each of six partitions. Tracer 1.6 (Rambaut et al., 2014) was used to confirm stabilisation within and convergence between runs (all ESS values >200). RAxML and MrBayes were run on the Cipres Science Gateway (Miller et al., 2010). Trees were also reconstructed for each mtDNA marker separately, following the same procedure as for the concatenated mtDNA data.

Phylogenetic trees for nuDNA were constructed using the same approach as for mtDNA, with *Neurergus strauchii* as outgroup (data taken from Hendrix et al., 2014). Because jModelTest suggested two out of three substitution models to be identical for codon positions within nuclear markers (Table A.4), each marker was analysed unpartitioned to avoid over-parameterisation of the analysis. Because these trees showed little divergence (Figs. B.7–B.10),

we additionally constructed minimum spanning haplotype networks (Bandelt et al., 1999) for each nuclear marker in PopArt 1.7 (Leigh et al., 2016).

*BEAST, implemented in BEAST 1.8.3 (Heled and Drummond, 2010; Drummond et al., 2012), was used to construct a species tree for the nuDNA markers. Species were set as operational taxonomic units. We selected the best fitting model per marker in jModelTest (Table A.4). We performed a species tree analysis including all mtDNA and nuclear DNA markers (two twenty million generation runs), and a species tree analysis using only the two nuclear DNA markers (one ten million generation run) with a sampling frequency of 0.001. Tracer 1.6 was used to confirm stabilisation and convergence as for MrBayes. A burn-in of 10% was removed when combining log and tree files in LogCombiner (BEAST 1.8.3). TreeAnnotator (BEAST 1.8.3) was used to construct a maximum clade credibility tree with mean node height.

2.3. Molecular dating

COI and cyt *b* sequences for all Salamandridae genera (Table A.2) were taken from Zhang et al. (2008). We included 14 *Ommatotriton* haplotypes, one randomly chosen haplotype from each phylogeographic group (see Results, Fig 2a) and a distinct haplotype found at a single locality only (Nes_08 from locality 4). We excluded the cyt *b* sequence for *Ichthyosaura alpestris*, because it contained a large number of unique non-synonymous substitutions suggestive of a pseudogene (see also Steinfartz et al., 2007). BEAST 1.8.3 was used to generate a calibrated phylogeny, employing seven fossil-based calibration points within Salamandridae, the family to which *Ommatotriton* belongs, following the procedure outlined in Wiens et al. (2011; Table A.5). To ensure that the calibration points fitted within the age constraints, we constructed a starting tree adapted from a RAxML tree with the R package ‘APE’ (Paradis et al., 2004). Four runs of 20 million generations were executed on the Cipres Science Gateway with nucleotide substitution models selected with jModelTest (Table A.4) and a sampling frequency of 0.001. Log and tree files were combined and convergence was checked as above.

2.4. Species distribution modelling

We combined our localities of species occurrence with those in Borkin et al. (2003). This yielded 75 localities for *O. vittatus*, 53 for *O. nesterovi* and 233 for *O. ophryticus*. Localities were assigned to species based on ranges suggested by our genetic analyses. We used bioclimatic variables from the WorldClim database 1.4 (Hijmans et al., 2005) as explanatory variables. Following Guisan and Thuiller (2005) and Peterson (2011), we selected variables with a Pearson’s correlation <0.7 that were considered biologically meaningful: isothermality (bio3), temperature seasonality (bio4), mean temperature of wettest quarter (bio8), mean temperature of driest quarter (bio9), precipitation of wettest quarter (bio16) and precipitation of driest quarter (bio17).

We used Maxent 3.3.3 k (Phillips et al., 2006) to construct species distribution models for each *Ommatotriton* species. Because the area on which species distribution models are calibrated influences model parameters considerably (Stokland et al., 2011), we restricted sampling of pseudo-absence data to a 200 km buffer zone around *Ommatotriton* localities (VanDerWal et al., 2009). Feature type was restricted to hinge features for a smoother model fit, emphasising trends rather than idiosyncrasies of the data (Elith et al., 2010). We confirmed that the models performed better than random by testing them for significance against a null model derived from random localities, following Raes and ter Steege (2007). Random point data (99 replicates per analysis) were created with ENMTools 1.3 (Warren et al., 2010). Models were




Fig. 2. Mitochondrial DNA phylogenetic tree and dated tree for banded newts (genus *Ommatotriton*). Panel a) Bayesian consensus tree with Bayesian posterior probability values above and Maximum Likelihood bootstrap support values below branches. Asterisks indicate nodes supported with a posterior probability >0.95 and a bootstrap support value >80 (values for nodes below the crown nodes of phylogeographic groups are not shown). Tips are haplotypes with codes corresponding to Table A.1 and localities in parentheses. Panel b) dated phylogeny with Bayesian posterior probability value left above branches (asterisks indicate posterior probability >0.95) and node age on the left below with 95% HPD confidence interval in parentheses right of nodes. In a) and b) capital letters correspond to the phylogeographic groups delineated in Fig. 1; the outgroup is not shown. Panel c) males of the three banded newt species in breeding condition. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

projected on climate reconstructions for the Last Glacial Maximum (~21 ka), based on the MIROC (Sueyoshi et al., 2013) and CCSM (Brady et al., 2013) climate simulations, available from WorldClim.

2.5. Population dynamics

Bayesian skyline plots (Ho and Shapiro, 2011) were generated with BEAST for 40 *O. nesterovi*, 40 *O. ophryticus*, and 39 *O. vittatus* mtDNA sequences with six million generations. Parameters were kept at default except for the nucleotide substitution models (selected with jModelTest; Table A.4) and the estimated total substitution rate (4.802×10^{-3} substitutions My^{-1} from the molecular dating analysis). Mismatch analyses were performed per species in DnaSP (Table A.6). We used COI data only because DnaSP is unable to handle missing data, and several *cyt b* sequences were not available (Table A.1). The observed haplotype frequencies were compared to haplotype frequencies expected at constant population size. The statistics R_2 and Fu 's F_s (Ramos-Onsins and Rozas, 2002) were calculated to test for significance ($R_2 < 0.05$ and Fu 's $F_s < 0.02$), using coalescent simulations with 1000 generations.

3. Results

For COI the number of segregating sites (S) is 214, nucleotide diversity (π) is 0.119, and total number of mutations (θ) is 71.26, for *cyt b* 100, 0.114, and 30.55, for KIAA 20, 0.007 and 4.11 and for SACS 17, 0.006 and 3.34. The mtDNA phylogenetic tree revealed three monophyletic groups, corresponding to the three *Ommatotriton* species (summarized in Fig. 2a, detailed MrBayes and RAXML trees in Figs. B.1 and B.2). The most recent common ancestor of *Ommatotriton* was dated at 24.5 Ma [95% HPD confidence interval (CI) 14.6–35.8 Ma, Fig. 2b]. *Ommatotriton nesterovi*, *O. ophryticus* and *O. vittatus* crown nodes were estimated at 13.0 Ma (CI 4.9–22.3 Ma), 8.6 Ma (CI 2.1–17.2 Ma) and 12.5 Ma (CI 5.5–20.8 Ma). The phylogenetic tree based on both mtDNA markers suggested *O. nesterovi* and *O. ophryticus* as sister species, but with low support, independent of the method used (Fig. 2a). Phylogenetic trees for the individual mtDNA markers strongly supported the three monophyletic clades, while relationship among clades were poorly supported (Figs. B.3–B.6). The species tree based on nuDNA only suggested that *O. ophryticus* and *O. vittatus* are sister species, with high support (posterior probability of 0.98; Fig. 3c), while adding




Fig. 3. Haplotype networks and species tree based on nuclear markers for banded newts (genus *Ommatotriton*). In the haplotype networks for KIAA (a) and SACS (b) pie diameter represents the number of individuals per haplotype and bars the number of inferred substitutions. Panel c) BEAST species tree with posterior values for all markers above, and only nuDNA markers below the branches. Colours indicate species as in Fig. 1. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

mtDNA to the *BEAST analysis resulted in a lower support for this clade (0.75).

For nuDNA phylogenetic trees (BI and ML, Figs. B.7–B.10), weak support was found for a monophyletic *O. nesterovi*. *Ommatotriton vittatus* was suggested, albeit with low support, to be either monophyletic or paraphyletic and nested within *O. ophryticus*, reflecting the relationships found in the haplotype network. In the KIAA network haplotypes for each species are clustered, while in the SACS haplotype network *O. nesterovi* was split up in two groups (Fig. 3a, b). We found no evidence for nuDNA haplotype sharing between species as defined by mtDNA (Fig. 2a, b).

Within species, 13 distinct monophyletic mtDNA clades representing geographically coherent groups were found (Figs. 1 and 2a). The sole exception was haplotype Nes_08 from locality 4, which is placed in a sister position to phylogeographic groups A and B and occurs in the geographic range of group A. Intraspecific phylogeographic groups were found to be more closely related to geographically more closely located phylogeographic groups than to groups located further apart, with the exception of the I+M-clade from Israel and the Adana Basin. For nuDNA, intraspecific differentiation was low (Figs. 3 and B.7–B.10). For the SACS gene, individuals from group I (localities 30 and 31) were relatively diverged from other individuals of *O. vittatus* (groups J–M). The KIAA marker suggested divergence of western (group E; localities 15–17) and eastern (groups F–H; localities 19–29) *O. ophryticus*, in line with the mtDNA-based phylogenetic tree. The mtDNA phylogeographic groups, however, were not recovered in the nuDNA analyses.

Projections on Last Glacial Maximum climate reconstructions showed that the potential distribution for the three species was reduced and fragmented compared to the present (Fig. 4 and B.11). For *O. vittatus*, the range reduction was less pronounced than

for *O. nesterovi* and *O. ophryticus*. The potential distributions of *O. nesterovi* and *O. ophryticus* overlap considerably at the present day and much less so at the Last Glacial Maximum. Unsuitable environmental conditions separated *O. vittatus* from the other two species during either period. Bayesian skyline plots based on mtDNA indicated a more pronounced dip in effective population size for *O. nesterovi* at ~0.5 Ma than for *O. ophryticus* and *O. vittatus* at ~0.05 Ma and at ~0.5 Ma (Fig. B12). Mismatch analyses indicated that allele frequencies did not differ from the allele frequency expected for a stable population over time (Fig. B.11, Table A.6).

4. Discussion

4.1. Biogeographical scenario for three genetically distinct species

The mtDNA phylogeny for *Ommatotriton* reveals three genetically distinct clades, which is in line with the proposal to recognize three banded newt species: *O. nesterovi*, *O. ophryticus* and *O. vittatus*. While reciprocal monophyly is not significantly supported for individual nuDNA markers, crucially none of the nuclear haplotypes are shared between species, not even in the region where the ranges of *O. nesterovi* and *O. ophryticus* approach one another, and where we sampled more densely (Fig. 1). On the basis of these results we accept the three species treatment.

Hybridization is not a likely cause of the lack of reciprocal monophyly of *O. vittatus* since it is separated from the remaining *Ommatotriton* range by the Anatolian Diagonal (Fig. 1). This mountain range was formed at 30–25 Ma as a consequence of the Arabia-Eurasia collision (Jolivet and Faccenna, 2000) and acts as a barrier for a wide range of taxa (Kapli et al., 2013). Uplift of the




Fig. 4. Species distribution models for banded newts (genus *Ommatotriton*). Model projections for the three species are arranged from top to bottom, with projections on current (left) and MIROC-derived Last Glacial Maximum (right) climate layers. Predicted suitability is reflected by a colour scale running from deep blue (highly unsuitable) to deep red (highly suitable). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Anatolian Diagonal likely isolated the stock that would become *O. vittatus* at ~25 Ma (Fig. 2b). Last Glacial Maximum conditions appear to have exacerbated this geographical isolation (Fig. 4).

The Arabia-Eurasia collision resulted in the uplift of the Anatolian Plateau (Hearn and Ni, 1994), which likely initiated speciation of what would become *O. nesterovi* and *O. ophryticus* at ~25 Ma (Fig. 2b). The two species were isolated during the Last Glacial Maximum (Fig. 4), implying that the current contact is of secondary, postglacial origin. Bilgin (2011) documented a north-south oriented suture zone, related to postglacial expansion from refugia in western and eastern Anatolia for plants, insects and vertebrates. The two other pairs of parapatric newt species distributed along the southern Black Sea coast show evidence of hybridisation upon secondary postglacial contact with pronounced mtDNA introgression, from the western (*Lissotriton vulgaris* and *Triturus ivanbureschi*) to the eastern species (*L. kosswigi* and *T. anatolicus*; Nadachowska and Babik, 2009; Wielstra et al., 2013, 2017), but there is no such mismatch between mtDNA and nuDNA for *O. nesterovi* and *O. ophryticus*. While we did not find evidence of (recent) introgression between the two species, to be conclusive a denser sampling, filling the 60 km gap between the closest *O. nesterovi* and *O. ophryticus* localities (localities 14 and 15, Fig. 1), is required.

4.2. Ancient and rapid radiation

The fossil-based time calibration suggests the *Ommatotriton* species radiated ~25 Ma, during a short time span. The mtDNA phylogeny suggests the three species form a polytomy. While our nuDNA species tree supported a ((*O. ophryticus*, *O. vittatus*), *O. nesterovi*) phylogeny, support decreased when adding mtDNA. We conclude that our data provide no convincing support of a particu-

lar branching order, and we consider the *Ommatotriton* phylogeny unresolved.

Conflicting results among markers can be expected for ancient and rapid radiations because few informative substitutions accumulate along internal branches (Philippe et al., 2011), little time is available for lineage sorting, resulting in conflicting gene genealogies (Pamilo and Nei, 1988) and uninformative substitutions accumulate along terminal branches, causing long-branch attraction (Felsenstein, 1978). Although comprehensive sampling of phylogenetic diversity can break up long branches (Bergsten, 2005), our sampling is comprehensive. However, we merely employed three unlinked genetic markers. Multispecies coalescence methods, as implemented in e.g. *BEAST, explicitly take incomplete lineage sorting into account (Heled and Drummond, 2010) and benefit from sampling more unlinked markers (Maddison and Knowles, 2006). Hence, we suggest a phylogenomic approach is required to resolve the phylogeny of banded newts.

4.3. Deep intraspecific phylogeographic structure

A deep genetic differentiation within *Ommatotriton* species is reflected by four to five intraspecific mtDNA phylogeographic groups per species. Splits between *O. nesterovi* groups located southeast of the Sea of Marmara (A, B) and along the central southern Black Sea shore (C, D), and between *O. vittatus* groups from the Adana Basin and Israel (I, M) and Turkey east of the Amanus Mountains and Syria (J, K, L; Fig. 1, Fig. 2b), coincide with the Middle Miocene disruption, a sudden period of climate cooling between 14.0 and 13.5 Ma (Flower and Kennett, 1994). This period is linked to vicariance and genetic divergence in the Near East for skinks (*Ablepharus*; Skourtanioti et al., 2016), toads (*Bufo*; Garcia-Porta

et al., 2012) and scorpions (*Mesobuthus*; Shi et al., 2013). The basal split between *O. ophryticus* groups located in central Turkey (E) and north-eastern Turkey, Georgia and on the western side of the Greater Caucasus (F, G, H) coincides with the uplift of the Armenian plateau during the Serravallian, caused by the Arabia-Eurasia collision (13–11 Ma; Gelati, 1975). This event was also inferred to have caused the split between the crested newts *Triturus karelinii* versus *T. ivanbureschi* and *T. anatolicus* (Wielstra et al., 2010).

We observe further subdivision into *O. nesterovi* groups (A and B, C and D), during the Late Miocene and Pliocene. This time period was associated with large-scale orogeny throughout the region (Oosterbroek and Arntzen, 1992; Bilgin, 2011). The different *O. nesterovi* groups are currently separated by protrusions of the Pontic Mountains, but the presence of a distinct haplotype from inside the range of group A, basal to the clade of groups A and B, shows the pattern is more complex. Divergence of the three groups distributed in the eastern part of the *O. ophryticus* range (F, G, H) suggests the Quaternary climate oscillations (2.6 Ma – present) initiated increased population fragmentation.

For *O. vittatus* we observe two allopatrically distributed groups that, even though they are sister clades, occur in the Adana Basin (I) and Israel (M). The Adana Basin is isolated from the remainder of the *Ommatotriton* range by the Amanus Mountains lifted during the lower-middle Miocene (~23–11 Ma; Aksu et al., 2005). Desiccation of the Mediterranean Sea during the Messinian salinity crisis (~5.96–5.33 Ma) would have established a direct connection allowing colonization of the Adana Basin (Duggen et al., 2003). Subsequent expansion of the stock that became groups J, K and L would explain the disjunct distribution pattern observed today. Interpreting the phylogeographic pattern of groups J, K, and L is difficult based on the current sampling. Wider sampling in this presently hostile area would be required to further elucidate the historical biogeography of *O. vittatus*. Deep phylogenetic splits and complex phylogeographic patterns caused by the uplift of the Amanus Mountains and dispersal routes facilitated by the Messinian salinity crisis in southern Anatolia and the Levant were previously suggested for amphibians (Akin et al., 2010; Skourtanioti et al., 2016) and reptiles (Guicking et al., 2009; Kapli et al., 2013).

4.4. Multiple intraspecific glacial refugia

The species distribution models for *O. vittatus* suggest relatively little range contraction during the Last Glacial Maximum (Fig. 4), supported by the Bayesian skyline plot suggesting a relatively small population decrease during the Quaternary (Fig. B.12). The deep phylogeographic structure suggests multiple glacial refugia (i.e. “refugia-within-refugia”) were present within the current range of *O. vittatus*. Multiple refugia within the current range of *O. vittatus* have been found for reptiles (Kornilios et al., 2011; Stümpel et al., 2016), insects (Simonato et al., 2007) and plants (Médail and Diadema, 2009).

Compared to *O. vittatus*, a more pronounced habitat reduction at the Last Glacial Maximum was suggested by the species distribution models for *O. ophryticus* and, especially, *O. nesterovi* (Fig. 4). This is supported by the reduced population size indicated by the Bayesian skyline plots (Fig. B.12). The deep genetic structure within the two northern species also survived the Last Glacial Maximum in multiple glacial refugia, located south of the Marmara Sea, and various locations along the southern Black Sea shore and in the Colchis (Fig. 1). The Marmara region acted as a glacial refugium for newts (Wielstra et al., 2010; Pabijan et al., 2015), multiple glacial refugia along the southern Black Sea coast were found for amphibians (Wielstra et al., 2010; Dufresnes et al., 2016), reptiles (Guicking et al., 2009) and mammals (Dubey et al., 2006, 2007; Bilgin et al., 2008) and multiple glacial refugia in the Colchis were found for plants, insects and snails (Tarkhnishvili et al., 2012),

amphibians (Tarkhnishvili et al., 2000; Veith et al., 2003), lizards (Freitas et al., 2016), and mammals (Seddon et al., 2002; Tarkhnishvili et al., 2012).

5. Conclusions

The genus *Ommatotriton* represents a rapid radiation, initiated by orogeny associated with the Arabia-Eurasia collision during late Oligocene. Lack of gene flow supports the recognition of three species of banded newt, but relationships among these species remain unresolved. Each species is composed of distinct, geographically coherent mtDNA clades, with splits dated from the onset of the Quaternary to as far back in time as the Middle Miocene. Multiple glacial refugia per species are positioned along the coasts of the Black and Mediterranean Seas, suggesting a refugia-within-refugia scenario for the Near East, as has been previously inferred for the Iberian (Gómez and Lunt, 2007), Italian (Canestrelli et al., 2012) and Balkan peninsulas (Poulakakis et al., 2015). *Ommatotriton* is exemplary for co-distributed taxa and the overarching pattern emerging from cumulative phylogeographic studies underlines the Near East as an important biodiversity hotspot, on par with hotspots revealed, through comparative phylogeography, for the better studied Mediterranean peninsulas (Riddle, 2016).

Acknowledgements

IvR is supported by the ‘Nederlandse organisatie voor Wetenschappelijk Onderzoek’ (NWO Open Programme 824.14.014). This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 655487. Permits to collect were made available to KO by Türkiye Bilimsel ve Teknolojik Araştırma Kurumu. We thank Michael Fahrback and the late Max Sparreboom for providing *O. nesterovi* and *O. ophryticus* pictures and Aroub Almasri, Frank Pasmans and the late Max Sparreboom for help during fieldwork. We thank two anonymous reviewers who provided helpful comments.

Appendix A. Supplementary data

GenBank accession numbers are available in Table A.1. The raw and edited sequence data, in- and output for phylogenetic analyses and species distribution model analyses are available at the Dryad database (<http://dx.doi.org/10.5061/dryad.4q08q>). Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ympbev.2017.05.028>.

References

- Akin, C., Bilgin, C.C., Beerli, P., Westaway, R., Ohst, T., Litvinchuk, S.N., Uzzell, T., Bilgin, M., Hotz, H., Guex, G.D., Plötner, J., 2010. Phylogeographic patterns of genetic diversity in eastern Mediterranean water frogs were determined by geological processes and climate change in the Late Cenozoic. *J. Biogeogr.* 37, 2111–2124.
- Aksu, A.E., Calon, T.J., Hall, J., Mansfield, S., Yaşar, D., 2005. The Cilicia-Adana basin complex, Eastern Mediterranean: Neogene evolution of an active fore-arc basin in an obliquely convergent margin. *Mar. Geol.* 221, 121–159.
- Arntzen, J.W., Olgun, K., 2000. Taxonomy of the banded newt, *Triturus vittatus*: morphological and allozyme data. *Amphib. Reptil.* 21, 155–168.
- Arntzen, J.W., Beukema, W., Galis, F., Ivanović, A., 2015. Vertebral number is highly evolvable in salamanders and newts (family Salamandridae) and variably associated with climatic parameters. *Contrib. Zool.* 84, 85–113.
- Avise, J.C., 2000. *Phylogeography: The History and Formation of Species*. Harvard University Press, Cambridge, Massachusetts.
- Bandelt, H.J., Forster, P., Rohlf, A., 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* 16, 37–48.
- Beheregaray, L.B., 2008. Twenty years of phylogeography: the state of the field and the challenges for the Southern Hemisphere. *Mol. Ecol.* 17, 3754–3774.
- Bergsten, J., 2005. A review of long-branch attraction. *Cladistics* 21, 163–193.

- Berthold, A.A., 1846. Über das Vorkommen von Tritonen am Kaukasus. Nachrichten von der Georg-Augusts-Universität und der Königlichen Gesellschaft der Wissenschaften, Göttingen.
- Bilgin, R., 2011. Back to the suture: the distribution of intraspecific genetic diversity in and around Anatolia. *Int. J. Mol. Sci.* 12, 4080–4103.
- Bilgin, R., Karatas, C., Coraman, E., Disotell, T., Morales, J.C., 2008. Regionally and climatically restricted patterns of distribution of genetic diversity in a migratory bat species, *Miniopterus schreibersii* (Chiroptera: Vespertilionidae). *BMC Evol. Biol.* 8, 209–223.
- Borkin, L.J., Litvinchuk, S.N., Zuiderwijk, A., 2003. *Triturus vittatus* (Gray, 1835) – Bandmolch. In: Grossenbacher, K., Thiesmeier, B. (Eds.), *Handbuch Der Reptilien Und Amphibien Europas*. Aula-Verlag, Wiebelsheim, pp. 555–605.
- Brady, E.C., Otto-Bliessner, B.L., Kay, J.E., Rosenbloom, N., 2013. Sensitivity to glacial forcing in the CCSM4. *J. Clim.* 26, 1901–1925.
- Bülbül, U., Kuttrup, B., 2013. Morphological and genetic variations of *Ommatotriton* in Turkey. *Anim. Biol.* 63, 297–312.
- Canestrelli, D., Salvi, D., Maura, M., Bologna, M.A., Nascetti, G., 2012. One species, three Pleistocene evolutionary histories: phylogeography of the Italian crested newt, *Triturus carnifex*. *PLoS One* 7, e41754.
- Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2012. JModelTest 2: more models, new heuristics and parallel computing. *Nat. Methods* 9, 772.
- Drummond, A.J., Suchard, M.A., Xie, D., Rambaut, A., 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* 29, 1969–1973.
- Dubey, S., Zaitsev, M., Cosson, J.-F., Abdoukadir, A., Vogel, P., 2006. Pliocene and Pleistocene diversification and multiple refugia in a Eurasian shrew (*Crocidura suaveolens* group). *Mol. Phylogenet. Evol.* 38, 635–647.
- Dubey, S., Cosson, J.-F., Vohralik, V., Krystufek, B., Diker, E., Vogel, P., 2007. Molecular evidence of Pleistocene bidirectional faunal exchange between Europe and the Near East: the case of the bicoloured shrew (*Crocidura leucodon*, Soricidae). *J. Evol. Biol.* 20, 1799–1808.
- Dufresnes, C., Litvinchuk, S.N., Leuenberger, J., Ghali, K., Zinenko, O., Stöck, M., Perrin, N., 2016. Evolutionary melting pots: a biodiversity hotspot shaped by ring diversifications around the Black Sea in the Eastern tree frog (*Hyla orientalis*). *Mol. Ecol.* 25 (17), 4285–4300. <http://dx.doi.org/10.1111/mec.13706>.
- Duggen, S., Hoernle, K., van den Bogaard, P., Rüpke, L., Phipps, Morgan J., 2003. Deep roots of the Messinian salinity crisis. *Nature* 422, 602–606.
- Elith, J., Kearney, M., Phillips, S., 2010. The art of modelling range-shifting species. *Methods Ecol. Evol.* 1, 330–342.
- Felsenstein, J., 1978. Cases in which parsimony or compatibility methods will be positively misleading. *Syst. Zool.* 27, 401–410.
- Flower, B.P., Kennett, J.P., 1994. The middle Miocene climatic transition: East Antarctic ice sheet development, deep ocean circulation and global carbon cycling. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 108, 537–555.
- Freitas, S., Vavakou, A., Arakelyan, M., Drovetski, S.V., Crnobrnja-Isailović, J., Kidov, A.A., Cogălniceanu, D., Corti, C., Lymberakis, P., Harris, D.J., Carretero, M.A., 2016. Cryptic diversity and unexpected evolutionary patterns in the meadow lizard, *Darevskia praticola* (Eversmann, 1834). *Syst. Biodivers.* 14, 184–197.
- Frost, D., 2016. Amphibian species of the world 6.0, Online reference. Available at: <http://research.amnh.org/vz/herpetology/amphibia/Amphibia/Caudata/Salamandridae/Pleurodelinae/Ommatotriton> (Last visited: 12 August 2016).
- García-Porta, J., Litvinchuk, S.N., Crochet, P.A., Romano, A., Geniez, P.H., Lo-Valvo, M., Lymberakis, P., Carranza, S., 2012. Molecular phylogenetics and historical biogeography of the west-palaearctic common toads (*Bufo bufo* species complex). *Mol. Phylogenet. Evol.* 63, 113–130.
- Gelati, R., 1975. Miocene marine sequence from lake Van eastern Turkey. *Riv. Ital. Paleontol. Stratigr.* 81, 477–480.
- Gómez, A., Lunt, D.H., 2007. Refugia within refugia: patterns of phylogeographic concordance in the Iberian Peninsula. In: Weiss, S., Ferrand, N. (Eds.), *Phylogeography of Southern European Refugia*. Springer, Netherlands, Dordrecht, pp. 155–188.
- Gray, 1835. *A Manual of British Vertebrate Animals*. Pitt Press, Cambridge.
- Guicking, D., Joger, U., Wink, M., 2009. Cryptic diversity in a Eurasian water snake (*Natrix tessellata*, Serpentes: Colubridae): evidence from mitochondrial sequence data and nuclear ISSR-PCR fingerprinting. *Org. Divers. Evol.* 9, 201–214.
- Guindon, S., Gascuel, O., 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* 52, 696–704.
- Guisan, A., Thuiller, W., 2005. Predicting species distribution: offering more than simple habitat models. *Ecol. Lett.* 8, 993–1009.
- Hearn, T.M., Ni, J.F., 1994. Pn velocities beneath continental collision zones: the Turkish-Iranian Plateau. *Geol. J. Int.* 117, 273–283.
- Heled, J., Drummond, A.J., 2010. Bayesian inference of species trees from multilocus data. *Mol. Biol. Evol.* 27, 570–580.
- Hendrix, R., Fleck, J., Schneider, W., Schneider, C., Geller, D., Avci, A., Olgun, K., Steinfartz, S., 2014. First comprehensive insights into nuclear and mitochondrial DNA based population structure of Near East mountain brook newts (Salamandridae: genus *Neurergus*) suggest the resurrection of *Neurergus derjugini*. *Amphib. Reptil.* 35, 172–187.
- Hewitt, G., 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405, 907–913.
- Hewitt, G.M., 2011. Mediterranean peninsulas: the evolution of hotspots. In: Zachos, F.E., Habel, J.C. (Eds.), *Biodiversity Hotspots: Distribution and Protection of Conservation Priority Areas*. Springer, Berlin, pp. 123–147.
- Hickerson, M.J., Carstens, B.C., Cavender-Bares, J., Crandall, K.A., Graham, C.H., Johnson, J.B., Rissler, L., Victoriano, P.F., Yoder, A.D., 2010. Phylogeography's past, present, and future: 10 years after Avise, 2000. *Mol. Phylogenet. Evol.* 54, 291–301.
- Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G., Jarvis, A., 2005. Very high resolution interpolated climate surfaces for global land areas. *Int. J. Climatol.* 25, 1965–1978.
- Ho, S.Y.W., Shapiro, B., 2011. Skyline-plot methods for estimating demographic history from nucleotide sequences. *Mol. Ecol. Resour.* 11, 423–434.
- Ivanova, N.V., Dewaard, J.R., Hebert, P.D.N., 2006. An inexpensive, automation-friendly protocol for recovering high-quality DNA. *Mol. Ecol. Notes* 6, 998–1002.
- Jolivet, L., Faccenna, C., 2000. Mediterranean extension and the Africa-Eurasia collision. *Tectonics* 19, 1095–1106.
- Kapli, P., Botoni, D., Ilgaz, C., Kumlutas, Y., Avci, A., Rastegar-Pouyani, N., Fathinia, B., Lymberakis, P., Ahmadzadeh, F., Poulakakis, N., 2013. Molecular phylogeny and historical biogeography of the Anatolian lizard *Apathya* (Squamata, Lacertidae). *Mol. Phylogenet. Evol.* 66, 992–1001.
- Kornilios, P., Ilgaz, C., Kumlutas, Y., Giokas, S., Fragedakis-Tsolis, S., Chondropoulos, B., 2011. The role of Anatolian refugia in herpetofaunal diversity: an mtDNA analysis of *Typhlops vermicularis* Merrem, 1820 (Squamata, Typhlopidae). *Amphib. Reptil.* 32, 351–363.
- Leigh, J., Bryant, D., Steel, M., 2016. PopArt. Version 1, 7. Available at: <http://popart.otago.ac.nz>.
- Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics (Oxford, England)* 25, 1451–1452.
- Litvinchuk, S.N., Zuiderwijk, A., Borkin, L.J., Rosanov, J.M., 2005. Taxonomic status of *Triturus vittatus* (Amphibia: Salamandridae) in western Turkey: trunk vertebrae count, genome size and allozyme data. *Amphib. Reptil.* 26, 305–323.
- Maddison, W.P., Knowles, L.L., 2006. Inferring phylogeny despite incomplete lineage sorting. *Syst. Biol.* 55, 21–30.
- Maddison, W.P., Maddison, D.R., 2015. Mesquite: a modular system for evolutionary analysis. Version 3, 10. Available at: <http://mesquiteproject.org>.
- Médail, F., Diadema, K., 2009. Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *J. Biogeogr.* 36, 1333–1345.
- Meulenckamp, J.E., Sissingh, W., 2003. Tertiary palaeogeography and tectonostratigraphic evolution of the Northern and Southern Peri-Tethys platforms and the intermediate domains of the African-Eurasian convergent plate boundary zone. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 196, 209–228.
- Miller, M.A., Pfeiffer, W., Schwartz, T., 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing Environments Workshop*. 14 Nov., New Orleans, LA, pp. 1–8.
- Moritz, C., Schneider, C.J., Wake, D.B., 1992. Evolutionary relationships within the *Ensatina eschscholtzii* complex confirm the ring species interpretation. *Syst. Biol.* 41, 273–291.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., Kent, J., 2000. Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858.
- Nadachowska, K., Babik, W., 2009. Divergence in the face of gene flow: the case of two newts (Amphibia: Salamandridae). *Mol. Biol. Evol.* 26, 829–841.
- Oosterbroek, P., Arntzen, J.W., 1992. Area-cladograms of circum-Mediterranean taxa in relation to Mediterranean palaeogeography. *J. Biogeogr.* 19, 3–20.
- Pabijan, M., Zieliński, P., Dudek, K., Chloupek, M., Sotiropoulos, K., Liana, M., Babik, W., 2015. The dissection of a Pleistocene refugium: phylogeography of the smooth newt, *Lissotriton vulgaris*, in the Balkans. *J. Biogeogr.* 42, 671–683.
- Pamilo, P., Nei, M., 1988. Relationships between gene trees and species trees. *Mol. Biol. Evol.* 5, 568–583.
- Paradis, E., Claude, J., Strimmer, K., 2004. APE: Analyses of Phylogenetics and Evolution in R language. *Bioinformatics* 20, 289–290.
- Peterson, A.T., 2011. Ecological niche conservatism: a time-structured review of evidence. *J. Biogeogr.* 38, 817–827.
- Philippe, H., Brinkmann, H., Lavrov, D.V., Littlewood, D.T.J., Manuel, M., Wörheide, G., Baurain, D., 2011. Resolving difficult phylogenetic questions: why more sequences are not enough. *PLoS Biol.* 9, e1000602.
- Phillips, S.J., Anderson, R.P., Schapire, R.E., 2006. Maximum entropy modeling of species geographic distributions. *Ecol. Model.* 190, 231–259.
- Popov, S.V., Shcherba, I.G., Ilyina, L.B., Nevesskaya, L.A., Paramonova, N.P., Khondkarian, S.O., Magyar, I., 2006. Late Miocene to Pliocene palaeogeography of the Paratethys and its relation to the Mediterranean. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 238, 91–106.
- Poulakakis, N., Kapli, P., Lymberakis, P., Trichas, A., Vardinoyiannis, K., Sfenthourakis, S., Mylonas, M., 2015. A review of phylogeographic analyses of animal taxa from the Aegean and surrounding regions. *J. Zool. Syst. Evol. Res.* 53, 18–32.
- Raes, N., ter Steege, H., 2007. A null-model for significance testing of presence-only species distribution models. *Ecography* 30, 727–736.
- Rambaut, A., Suchard, M.A., Drummond, A.J., 2014. Tracer. Version 1.6. Available at: www.tree.bio.ed.ac.uk/software/tracer/.
- Ramos-Onsins, S.E., Rozas, J., 2002. Statistical properties of new neutrality tests against population growth. *Mol. Biol. Evol.* 19, 2092–2100.
- Riddle, B.R., 2016. Comparative phylogeography clarifies the complexity and problems of continental distribution that drove A.R. Wallace to favor islands. *PNAS* 113, 7970–7977.
- Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 539–542.
- Seddon, J.M., Santucci, F., Reeve, N., Hewitt, G.M., 2002. Caucasus Mountains divide postulated postglacial colonization routes in the white-breasted hedgehog, *Erinaceus concolor*. *J. Evol. Biol.* 15, 463–467.

- Shen, X.-X., Liang, D., Zhang, P., 2012. The development of three long universal nuclear protein-coding locus markers and their application to osteichthyan phylogenetics with nested PCR. *PLoS One* 7, e39256.
- Shi, C.M., Ji, Y.J., Liu, L., Wang, L., Zhang, D.X., 2013. Impact of climate changes from Middle Miocene onwards on evolutionary diversification in Eurasia: insights from the mesobuthid scorpions. *Mol. Ecol.* 22, 1700–1716.
- Simonato, M., Mendel, Z., Kerdelhué, C., Rousset, J., Magnoux, E., Salvato, P., Roques, A., Battisti, A., Zane, L., 2007. Phylogeography of the pine processionary moth *Thaumetopoea wilkinsoni* in the Near East. *Mol. Ecol.* 16, 2273–2283.
- Skourtanioti, E., Kapli, P., Ilgaz, Ç., Kuşlutaş, Y., Avcı, A., Ahmadzadeh, F., Crnobrnja-Isailović, J., Gherghel, I., Lymberakis, P., Poulakakis, N., 2016. A reinvestigation of phylogeny and divergence times of the *Ablepharus kitaibelii* species complex (Sauria, Scincidae) based on mtDNA and nuDNA genes. *Mol. Phylogenet. Evol.* 103, 199–214.
- Smith, M.A., Poyarkov, N.A., Hebert, D.N., 2008. CO1 DNA barcoding amphibians: take the chance, meet the challenge. *Mol. Ecol. Resour.* 8, 235–246.
- Sparreboom, M., 2014. Salamanders of the Old World. KNNV Publishing, Zeist, The Netherlands.
- Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30, 1312–1313.
- Steinfartz, S., Veith, M., Tautz, D., 2000. Mitochondrial sequence analysis of *Salamandra* taxa suggests old splits of major lineages and postglacial recolonizations of Central Europe from distinct source populations of *Salamandra salamandra*. *Mol. Ecol.* 9, 397–410.
- Steinfartz, S., Vicario, S., Arntzen, J.W., Caccone, A., 2007. A Bayesian approach on molecules and behavior: reconsidering phylogenetic and evolutionary patterns of the Salamandridae with emphasis on *Triturus* newts. *J. Experim. Zool. (Mol. Develop. Evol.)* 306B, 1–24.
- Stephens, M., Donnelly, P., 2003. A comparison of Bayesian methods for haplotype reconstruction from population genotype data. *Am. J. Hum. Genet.* 73, 1162–1169.
- Stewart, J.R., Lister, A.M., Barnes, I., Dalén, L., 2009. Refugia revisited: individualistic responses of species in space and time. *Proc. R. Soc. Biol. Sci.* 277, 661–671.
- Stokland, J.N., Halvorsen, R., Støa, B., 2011. Species distribution modelling – effect of design and sample size of pseudo-absence observations. *Ecol. Model.* 222, 1800–1809.
- Stümpel, N., Rajabizadeh, M., Avcı, A., Wüster, W., Joger, U., 2016. Phylogeny and diversification of mountain vipers (Montivipera, Nilson et al., 2001) triggered by multiple Plio-Pleistocene refugia and high-mountain topography in the Near and Middle East. *Mol. Phylogenet. Evol.* 101, 336–351.
- Sueyoshi, T., Ohgaito, R., Yamamoto, A., Chikamoto, M.O., Hajima, T., Okajima, H., Yoshimori, M., Abe, M., Oishi, R., Saito, F., Watanabe, S., Kawamiya, M., Abe-Ouchi, A., 2013. Set-up of the PMIP3 paleoclimate experiments conducted using an Earth system model, MIROC-ESM. *Geosci. Model Develop.* 6, 819–836.
- Tarkhnishvili, D.N., Thorpe, R.S., Arntzen, J.W., 2000. Pre-pleistocene refugia and differentiation between populations of the Caucasian salamander (*Mertensiella caucasica*). *Mol. Phylogenet. Evol.* 14, 414–422.
- Tarkhnishvili, D., Gavashelishvili, A., Mumladze, L., 2012. Palaeoclimatic models help to understand current distribution of Caucasian forest species. *Biol. J. Linn. Soc.* 105, 231–248.
- VanDerWal, J., Shoo, L.P., Graham, C., Williams, S.E., 2009. Selecting pseudo-absence data for presence-only distribution modeling: how far should you stray from what you know? *Ecol. Model.* 220, 589–594.
- Veith, M., Schmidler, J.F., Kosuch, J., Baran, I., Seitz, A., 2003. Palaeoclimatic changes explain Anatolian mountain frog evolution: a test for alternating vicariance and dispersal events. *Mol. Ecol.* 12, 185–199.
- Veith, M., Lipscher, E., Oz, M., Kiefer, A., Baran, I., Polymeni, R.M., Steinfartz, S., 2008. Cracking the nut: geographical adjacency of sister taxa supports vicariance in a polytomic salamander clade in the absence of node support. *Mol. Phylogenet. Evol.* 47, 916–931.
- Vences, M., Wake, D.B., 2007. Speciation, species boundaries and phylogeography of amphibians. In: Heatwole, H., Tyler, M. (Eds.), *Amphibian Biology*, vol. 6, Systematics. Surrey Beatty & Sons, Chipping Norton, Australia, pp. 2613–2660.
- Warren, D.L., Glor, R.E., Turelli, M., 2010. ENMTools: a toolbox for comparative studies of environmental niche models. *Ecography* 33, 607–611.
- Wielstra, B., Espregueira, Themudo G., Güçlü, Ö., Olgun, K., Poyarkov, N.A., Arntzen, J.W., 2010. Cryptic crested newt diversity at the Eurasian transition: the mitochondrial DNA phylogeography of Near Eastern *Triturus* newts. *Mol. Phylogenet. Evol.* 56, 888–896.
- Wielstra, B., Baird, A.B., Arntzen, J.W., 2013. A multimarker phylogeography of crested newts (*Triturus cristatus* superspecies) reveals cryptic species. *Mol. Phylogenet. Evol.* 67, 167–175.
- Wielstra, B., Burke, T., Butlin, R.K., Avcı, A., Üzümlü, N., Bozkurt, E., Olgun, K., Arntzen, J.W., 2017. A genomic footprint of hybrid zone movement in crested newts. *Evol. Lett.* <http://dx.doi.org/10.1002/evl3.9>.
- Wiens, J.J., Sparreboom, M., Arntzen, J.W., 2011. Crest evolution in newts: implications for reconstruction methods, sexual selection, phenotypic plasticity and the origin of novelties. *J. Evol. Biol.* 24, 2073–2086.
- Zhang, P., Papenfuss, T.J., Wake, M.H., Qu, L., Wake, D.B., 2008. Phylogeny and biogeography of the family Salamandridae (Amphibia: Caudata) inferred from complete mitochondrial genomes. *Mol. Phylogenet. Evol.* 49, 586–597.