



UNIVERSITY OF LEEDS

This is a repository copy of *Pilot study to determine whether microbial contamination levels in hospital washrooms are associated with hand-drying method.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/119067/>

Version: Accepted Version

Article:

Wilcox, MH orcid.org/0000-0002-4565-2868, Best, EL and Parnell, P (2017) Pilot study to determine whether microbial contamination levels in hospital washrooms are associated with hand-drying method. *Journal of Hospital Infection*, 97 (2). pp. 201-203. ISSN 0195-6701

<https://doi.org/10.1016/j.jhin.2017.07.007>

Crown Copyright © 2017 Published by Elsevier Ltd on behalf of The Healthcare Infection Society. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.




eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Accepted Manuscript

Pilot study to determine if microbial contamination levels in hospital washrooms are associated with hand drying method

Mark H. Wilcox, Emma L. Best, Peter Parnell



PII: S0195-6701(17)30389-4

DOI: [10.1016/j.jhin.2017.07.007](https://doi.org/10.1016/j.jhin.2017.07.007)

Reference: YJHIN 5162

To appear in: *Journal of Hospital Infection*

Received Date: 22 June 2017

Revised Date: 0195-6701 0195-6701

Accepted Date: 7 July 2017

Please cite this article as: Wilcox MH, Best EL, Parnell P, Pilot study to determine if microbial contamination levels in hospital washrooms are associated with hand drying method, *Journal of Hospital Infection* (2017), doi: 10.1016/j.jhin.2017.07.007.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Pilot study to determine if microbial contamination levels in hospital washrooms are associated with hand drying method

Mark H. Wilcox, Emma L. Best, Peter Parnell

Microbiology, Leeds Teaching Hospitals NHS Trust & University of Leeds, Leeds, UK.

Correspondence:

Professor Mark H. Wilcox

Consultant / Head of Microbiology Research & Development
Leeds Teaching Hospitals NHS Trust

Professor of Medical Microbiology

Sir Edward Brotherton Chair of Bacteriology
University of Leeds

Microbiology,
Old Medical School,
Leeds General Infirmary,
Leeds LS1 3EX,
W. Yorks., UK.

mark.wilcox@nhs.net

Tel +44 113 392 6818

Fax +44 113 392 2696

Sir,

We and others have demonstrated that some hand drying methods are associated with a greater risk of dissemination of residual microbes from hands after (particularly sub-optimal) hand washing.¹⁻⁴ For example, air bacterial counts in close proximity to hand drying by a jet air dryer were 27-fold higher than measured next to use of paper towels ($P < 0.001$).¹ Such results suggest that air dryers may be unsuitable for use in healthcare settings, where they may facilitate microbial cross-contamination via airborne droplet dispersal. Such risks could have very differing implications depending on multiple factors, including the magnitude of dispersal, the particular microorganisms involved and, of course, the setting. In hospitals, general infection prevention and control principles aim to limit the spread of microorganisms given the increased susceptibility of patients to infection, and the greater prevalence of potential and/or antimicrobial-resistant pathogens.

We have carried out a pilot study to demonstrate the feasibility of testing strategies to examine prospectively the levels of environmental bacterial contamination in hospital washrooms associated with two hand drying methods: paper hand towels (Tork H3 classic dispenser with Tork Advanced Towels, MRT213), and jet air dryer (Dyson Airblade, Dyson, UK). We sampled, on 26 occasions over 3 months, two washrooms (for males) within one hospital, both accessed via a large entrance foyer/thoroughfare. Each washroom was utilised by hospital staff, patients and visitors, and had similar footfalls (mean 26-32 people per daytime hour). There were no windows in either washroom and the air in both was maintained by standard ventilation without air conditioning. Both washrooms were routinely cleaned on three occasions daily (am, midday and pm); sampling took place immediately before scheduled cleaning. Each washroom sampling session involved taking two 5-minute air samples as described elsewhere,¹ up to five surface swabs (Polywipe sponges; Microbial wire, Corsham, UK) and collecting a sample of dust using a high-efficiency vacuum cleaner with a flexible extension hose (Dyson, UK). Environmental samples were cultured on both selective and non-selective agars. Bacteria were identified using the Bruker MALDI-Biotyper (Coventry, UK) and antibiotic susceptibilities were tested using a Vitek 2 (Biomérieux, UK). All statistical comparisons were made using the Mann-Whitney U test.

Trends towards and some significant differences were seen, generally showing lower levels of bacterial contamination in the paper towel washroom (washroom 2) than in the jet air dryer washroom (washroom 1). A greater range of bacteria were recovered from washroom 1 compared with washroom 2; in general, the floor, dryer unit and (vacuum cleaner) dust were more heavily contaminated in washroom 1. Over the 7-day testing period, the mean number of micro-organisms recovered from air was 1.6 fold higher in washroom 1 versus 2 ($P=0.14$). Bacterial burdens on floors were significantly higher in washroom 1 versus 2 (2.0×10^4 CFU/ml vs. 3.3×10^3 CFU/ml) ($P=0.002$). Also, the jet air dryer casing had significantly higher counts than the paper towel dispenser (1.2×10^5 CFU/ml versus 2.4×10^4 CFU/ml, $P=0.01$). Notably, higher counts of *Enterococcus faecalis*, which could be related to toileting followed by sub-optimal hand washing, were recovered from washroom 1 versus 2 (Figure). Significantly more *E. faecalis* was recovered from the dryer unit (3.4×10^3 CFU/ml) in washroom 1 compared with the paper towel dispenser in washroom 2 (71.4 CFU/ml) ($P=0.04$).

Measuring within-day levels of environmental contamination (three testing sessions per each washroom during the same day) was not particularly helpful, and may be more prone to confounding by behaviours of people visiting washrooms. Recovery of potential pathogens that are antibiotic resistant bacteria was uncommon in this pilot study. Study of washrooms that are closer to clinical areas may be expected to increase the yield of environmental antibiotic resistant bacteria.

Interestingly, current National Health Service (NHS) building guidance states: 'Hot-air hand dryers reduce paper waste and may be considered for use in public areas of healthcare facilities, but should not be installed in clinical areas as they are noisy and could disturb patients.'⁵ Such advice may need to be strengthened to take into account emerging data on the potential for microbe dissemination associated with electric hand dryers. Notably, current World Health Organisation guidance for healthcare settings advocates to 'dry hands thoroughly with a single use towel.'⁶ The key issue relating to increased levels of washroom environment microbial contamination is whether this could have adverse infection consequences for washroom users or, in clinical setting, patients. This point remains unstudied. We conclude from this pilot study that it is feasible to carry out longitudinal testing to examine the levels of environmental contamination that is associated with

different hand drying methods. Our pilot data suggest that bacterial burdens may be higher in hospital washrooms employing jet air versus paper towel hand drying, consistent with in situ testing data. We encourage further studies to determine the risks associated with hand drying method associated environmental microbial contamination.

Conflict of interest statement

M.H.W. has received honoraria from ETS for microbiological advice and travel expenses to attend ETS meetings.

Funding sources

The project was designed by the authors but funded by the European Tissue Symposium (ETS).

References

1. Best EL, Parnell P, Wilcox MH. Microbiological comparison of hand-drying methods: the potential for contamination of the environment, user, and bystander. *J Hosp Infect* 2014;88:199-206.
2. Kimmitt PT, Redway KF. Evaluation of the potential for virus dispersal during hand drying: a comparison of three methods. *J Appl Microbiol* 2016;120:478-86.
3. Margas E, Maguire E, Berland CR, Welander F, Holah JT. Assessment of the environmental microbiological cross contamination following hand drying with paper hand towels or an air blade dryer. *J Appl Microbiol* 2013;115:572-82.
4. Huang C, Ma W, Stack S. The hygienic efficacy of different hand-drying methods: a review of the evidence. *Mayo Clin Proc* 2012;87:791-8.
5. Department of Health. HBN 00-09 – Infection control in the built environment. Gateway Reference 18521. 2013.

6. World Health Organisation. Hand Hygiene: Why, How & When? Available at [http://who.int/gpsc/5may/Hand Hygiene Why How and When Brochure.pdf](http://who.int/gpsc/5may/Hand_Hygiene_Why_How_and_When_Brochure.pdf) Last accessed 19 June 2017.

ACCEPTED MANUSCRIPT

Figure

Mean results for recovery of *E. faecalis* from environmental sites in two washrooms over 7 days of sampling

