

This is a repository copy of *Novel substrates as sources of ancient DNA: : prospects and hurdles*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/119087/>

Version: Published Version

---

**Article:**

Green, Eleanor Joan and Speller, Camilla Filomena orcid.org/0000-0001-7128-9903  
(2017) Novel substrates as sources of ancient DNA: : prospects and hurdles. *Genes*. pp. 1-26. ISSN 2073-4425

<https://doi.org/10.3390/genes8070180>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:


<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

Review

# Novel Substrates as Sources of Ancient DNA: Prospects and Hurdles

Eleanor Joan Green and Camilla F. Speller \* 

BioArCh, Department of Archaeology, University of York, Wentworth Way, York YO10 5DD, UK; eg715@york.ac.uk

\* Correspondence: camilla.speller@york.ac.uk; Tel.: +44-1904-328868

Academic Editor: Michael Hofreiter

Received: 30 May 2017; Accepted: 10 July 2017; Published: 13 July 2017

**Abstract:** Following the discovery in the late 1980s that hard tissues such as bones and teeth preserve genetic information, the field of ancient DNA analysis has typically concentrated upon these substrates. The onset of high-throughput sequencing, combined with optimized DNA recovery methods, has enabled the analysis of a myriad of ancient species and specimens worldwide, dating back to the Middle Pleistocene. Despite the growing sophistication of analytical techniques, the genetic analysis of substrates other than bone and dentine remain comparatively “novel”. Here, we review analyses of other biological substrates which offer great potential for elucidating phylogenetic relationships, paleoenvironments, and microbial ecosystems including (1) archaeological artifacts and ecofacts; (2) calcified and/or mineralized biological deposits; and (3) biological and cultural archives. We conclude that there is a pressing need for more refined models of DNA preservation and bespoke tools for DNA extraction and analysis to authenticate and maximize the utility of the data obtained. With such tools in place the potential for neglected or underexploited substrates to provide a unique insight into phylogenetics, microbial evolution and evolutionary processes will be realized.


**Keywords:** ancient DNA; methodological advances; PCR; NGS

## 1. Introduction

The first successful recovery of ancient DNA (aDNA) in 1984 was from a piece of dried muscle, still connected to the salt-preserved skin of a quagga (*Equus quagga*), which had died ~140 years previously [1]. The demonstration that mitochondrial DNA (mtDNA) could be isolated and cloned from museum specimens influenced the choice of ancient samples for the next decade. Early aDNA research focused on soft tissue [2–8], however, the first successful extraction of aDNA from archaeological bones by three independent laboratories in 1989–1990 revolutionized the field [9–11]. The ability to amplify DNA from (sub)fossils gave-way to a new branch of aDNA studies; paleo-geneticists no longer focused their attention solely on relatively rare soft tissue specimens, but instead began exploiting the component of a vertebrate’s body most likely to survive through time: bone.

For the last 30 years, bone and teeth have been the most frequently studied substrates in paleogenetic research. Most paleontological hard tissues, however, contain very low proportions of endogenous DNA [12], and thus the search for methods to maximize the utility of endogenous DNA data from bony specimens has been extensive, including rapid column-based extractions [13–15], and the use of alternative skeletal elements (e.g., the petrous bone, tooth cementum [16–19]). However, it has been the development of massive parallel DNA sequencing (so called “next generation sequencing” or NGS), often coupled with enriched capture-based methods [12,20] that have resolved two fundamental limiting factors of aDNA research: cost and time. With improved accessibility to

highly decayed and fragmentary genetic information, facilitated by the NGS revolution, aDNA is increasingly being extracted from what are now considered novel or alternative substrates (Figure 1).



**Figure 1.** The relative proportion of studies discussed in this review (126 papers from 1988 to May 2017) targeting various alternative substrates for the recovery of ancient DNA. NHC: natural history collections.

## 2. Novel Substrates

Although all biological materials have the potential to preserve DNA, relatively few will resist decay over time, unless protected within an inorganic matrix or deliberately preserved through human intervention. Here, we review the potential for DNA preservation within three broad classes of materials: (1) archaeological artifacts and ecofacts; (2) calcified or mineralized substrates; and (3) biological and cultural archives.

### 2.1. Archaeological Artifacts and Ecofacts

Archaeological excavations worldwide recover millions of artifacts drawn from all time periods in all stages of decay—yet the only materials routinely analyzed genetically are hard tissues. Bones and teeth are mostly targeted due to scientific focus on human (and, to a lesser extent, animal) evolution, while the extraction of DNA from other hard tissues like preserved antler, or antler artifacts has facilitated research into cervid population genetics and object manufacture [21,22]. Researchers are now extracting DNA from an array of artifacts (i.e., objects manufactured by human beings) and ecofacts (i.e., organic materials recovered from archaeological sites that carry anthropological significance) including archaeobotanical remains (seeds, wood, worked plant remains), keratinous

and collagenous tissues (feathers, hair, nail and leather), and lithics and ceramics. Although DNA preservation is variable, these items provide unique access to evolutionary pathways, taxonomy and phylogeny that may be simply unobtainable from the analysis of hard tissue.

#### 2.1.1. Archaeobotanical Remains

The genetic analysis of archaeobotanical specimens is relatively well established, and thus no longer “novel”; however, compared to vertebrates, archaeobotanical studies remain vastly underrepresented. A range of early polymerase chain reaction (PCR) based approaches (e.g., [23–29]) and more recent whole genome studies (e.g., [30–35]) have demonstrated the potential to access genetic information held within preserved plant remains. Seeds tend to preserve in archaeological sites only when charred, desiccated, frozen, or deposited in anoxic conditions. Charred seeds, making up the vast majority of recovered archaeobotanical materials, have variable degrees of DNA preservation, depending, in part, on the extent of charring, as well as their age and depositional environment [36–38]. For example, in Bunning et al.’s [30] analysis of 3000 year-old charred, mixed cereals from Assiros Toumba (Greece), less than 0.1% of the over 21 M recovered sequences could be assigned to the kingdom Viridiplantae. Likewise, Nistelberger et al. [34] obtained 0–0.12% endogenous DNA from 4450 to 550 year-old charred barley, grape seeds, maize cobs, and rice remains from a range of contexts worldwide, even using a targeted enrichment approach. Variation in DNA preservation is also observed within desiccated remains. For example, in their analysis of five 6000 year-old barley grains from a single site in Israel, Mascher et al. [33] reported endogenous DNA ranging from 0.4 to 96.4%.

Successful DNA recovery has also been achieved using other archaeological plant tissues, including maize cobs [32,39,40], fruit stones [41], pollen [42–44], grains and seeds [45–48], rind [37], and chaff [49,50]. Plant-based artifacts can also be analyzed for their human DNA content—for example, LeBlanc et al. [51] extracted DNA from quids (chewed plant material) and fiber aprons (sanitary wear) to identify human mtDNA haplotypes from Southwest America. Despite PCR methods suggesting there were few prospects for retrieving aDNA from wood [52,53] due to PCR-inhibiting metabolites [54,55], recently the application of NGS to ~9000 year-old archaeological wood has recovered DNA fragments of 87–596 base pairs (bp) [56]. Using more advanced DNA recovery and bioinformatic techniques such as whole genome sequencing, single nucleotide polymorphism (SNP) capture, exome analyses, RNA analysis, and methylation patterns, these archaeobotanical projects are demonstrating the vast potential of ancient plant remains to address significant debates around the origins, movement and adaptation of domestic crops [32,37], reconstruction of paleo-environments [57] and models of DNA decay within plant remains [34,36].

#### 2.1.2. Keratinous and Collagenous Tissues

Leather, hair, baleen, claws and feathers are all composed of layers of collagen or keratin, which frequently decompose when deposited underground. In some extraordinary contexts, (e.g., burial in permafrost), such items may be preserved in the archaeological record and retain host DNA. Although reports on DNA extraction from archaeological leathers are few, mtDNA within this substrate seems to be particularly resilient, with previous studies recovering 70–177 bp fragments from medieval [58] and Neolithic [59] leather; although the tanning process may be particularly damaging for nuclear DNA [58,60].

The keratin structure of hair, claws and baleen is thought to protect endogenous DNA from contamination [61,62], and several studies have successfully retrieved endogenous DNA from human and animal hair [62–64] and whale baleen [65,66] preserved in Danish and Greenlandic archaeological contexts. Preservation of claws in archaeological contexts is rare; however, research on natural history collections (NHC) indicates that DNA can be sufficiently well preserved for analysis of this substrate [67,68], with potentially the same success rate as ancient bone [69]. The base of feathers (the calamus) has been frequently exploited as a source of high-quality DNA in bird phylogenetics and conservation biology research, and the potential for feathers in NHC to preserve DNA for several

hundred years has been known for more than two decades (e.g., [70–72]). In archaeological and subfossil environments, the upper shaft and the feather vane are the only part of the feathers that typically survive and were thought to be unsuitable for aDNA analysis. Using Moa feathers preserved in rock shelter sites in New Zealand, Rawlence et al. (2009) demonstrated that both the rachis and barbs, as well as the calamus, can preserve DNA for potentially up to 3000 years. Subsequently, Speller et al. [73] were able to recover mtDNA fragments from as few as two feather barbs, suggesting that feathers may represent a robust source of aDNA when preserved in favorable contexts.

The potential for mtDNA to survive in these keratinous substrates has been well documented, however, the potential for nuclear DNA survival is less well explored [66,74]. Future metagenomic analyses are required to assess both the relative proportion and survival of mitochondrial versus nuclear DNA, and to test susceptibility of different keratinous substrates to exogenous contamination from the burial environment.

### 2.1.3. Lithics and Ceramics

Lithics represent some of the most abundant and durable artifacts within the archaeological record, particularly in prehistoric contexts. As such, studies endeavoring to understand the day-to-day use of various stone artifacts have a long history, employing methods such as experimental archaeology [75]; usewear analysis [76]; microwear polish examination [77]; and microscopic and chemical residue analysis [78,79]. Unsurprisingly, researchers attempted to extract DNA from lithics as early as the 1990s. Loy [80] reported the identification of bovine satellite DNA from a 2200 year-old stone tool from British Columbia using nested PCR, while Hardy et al. [81] attempted even older samples, reporting the amplification of short fragments of *Sus scrofa* cytochrome b (cytb) mtDNA, from a single stone tool dating to 35,000–65,000 years ago. In 2001, Shanks et al. [82] reported observation of fluorescently labeled blood protein and DNA within the microcracks of stone tools, and later [83] the purported recovery of 116 bp fragments of cytb mtDNA from these microcracks, the majority corresponding to the canid family. The findings of these early studies have been called into question, first by the lack of authentication measures and contamination controls required to validate aDNA results [84], as well as by observations of domestic animal DNA within laboratory reagents [85]. Coupled with extensive controversy regarding the ability to extract authentic proteins from ancient lithic materials (e.g., [86–91]), the potential for stone tools to entrap and preserve DNA had, for many years, seemingly been laid to rest. The application of NGS approaches however, may resurrect this old idea in a new form, again through the analysis of another novel substrate: historic building materials. Two recent studies have investigated the potential for metagenomic analysis of ancient brick and stone work to elucidate building histories and investigate microbial factors influencing biodeterioration of built heritage [92,93], opening up potentially more fruitful research directions for ancient lithics.

Like lithic materials, ancient ceramics make up a large percentage of archaeological finds; however, attempts to mine this substrate for preserved DNA has lagged significantly compared to the analysis of other surface or adsorbed organic molecules, including lipids, amino acids, alkaloids, waxes, etc. [94–97]. Attempts to extract DNA from ceramic objects have involved scraping or drilling the interior of the vessel, as well as non-destructive swabbing of the interior face [98,99]. Recently, aDNA identifications have been reported from pottery vessels recovered from environments assumed to be hostile to DNA. Foley et al. [100] reported the amplification of 69–188 bp fragments of plant chloroplast DNA from 5<sup>th</sup> to 3<sup>rd</sup> Century B.C. empty amphora recovered from Mediterranean Sea floor, identifying a range of plants known to have been exploited by the Greeks. Based on their results, Foley et al. claimed that sufficient DNA could be recovered merely by swabbing the inner surfaces of ceramic fragments that had been lying beneath the ocean floor for over two millennia. Robinson et al. [101] also reported the recovery of aDNA from terracotta libation figurines from pre-colonial Ghana using a swabbing approach. Here, DNA from three plant types—plantain, pine and grasses—were identified using generic plant primers, recovering fragments up to 257 bp, significantly beyond the average length of DNA fragments they predicted by thermal age modeling. The latter

study applied extensive contamination controls and numerous measures to validate their sequences, but, in so doing, highlighted a major challenge when utilizing atypical substrates: due to the infancy of the research, the proper authentication of results is imperative; paradoxically, there is currently little understanding of if and how DNA is expected to preserve within such substrates—an issue we discuss in greater detail below (Section 3.1).

## 2.2. Calcified and Mineralized Substrates

The predominant use of skeletal hard tissues in aDNA analysis is related to their ubiquity in the archaeological and paleontological record, but also because the inorganic fraction of hydroxyapatite in bone and teeth is thought to stabilize and preserve DNA through adsorption [102,103]. More recently, the potential for other mineralized substrates to preserve ancient biomolecules in deep time is also being realized [104], including substrates such as archaeological dental calculus, coprolites, calcified soft tissues, invertebrate shells and ancient eggshells. Unlike bones and teeth, these substrates may be exploited not only as a source of host genetic information, but also to provide insight into ancient microbiomes and paleoenvironments.

### 2.2.1. Calculus

Dental calculus (tartar) is a bacterial biofilm composed from dental plaque, saliva and gingival crevicular fluid, mineralized within a matrix of multiple calcium phosphates, forming a cement-like substrate on the surface of teeth [105–107]. Although found virtually ubiquitously on archaeological human skeletons without modern dental hygiene interventions, the potential for this substrate to preserve abundant and varied ancient biomolecules has only recently been discovered. The preservation of ancient oral bacterial DNA was first recognized through gold-labeled antibody transmission electron microscopy [108] and subsequently through targeted PCR and 16S ribosomal RNA (rRNA) amplicon metagenomics [108–110]. Shotgun metagenomic approaches have demonstrated that although calculus is dominated by bacterial (and to a lesser extent fungal, archeal and viral) DNA derived from the oral microbiome, minute quantities of host DNA as well as inhaled or ingested eukaryotic DNA also survive [111–113]. Research to date has demonstrated that calculus preserves an exceptional abundance of entrapped DNA and proteins [109,112–114], and even ancient metabolites [115]. These combined studies have provided insight into the oral ecology of ancient humans and hominids, and even allowed for microbial genome reconstruction dating back to nearly 50 k years before present (BP) [112,113]. In some studies, calculus has been noted to be relatively resistant to exogenous bacterial colonization from the burial environment [113]. In combination, this recent work highlights the promise of calculus as a “novel” substrate for reconstructing ancient microbial genomes, tracking evolutionary changes in oral ecology, detecting dietary and environmental information, and analyzing host population demographics through mitochondrial genome capture [105,116,117].

### 2.2.2. Paleofeces and Coprolites

Unlike calculus, paleofeces and coprolites have been long been recognized for their value as a source of ancient biomolecules and paleodietary information. Paleofeces can be preserved via various mechanisms such as rapid desiccation or waterlogging, while coprolites form through the precipitation of calcium and phosphate from ingested bone (especially in carnivores and scavengers) or through the acquisition of preservational constituents via groundwater [118]. Due to the specific environmental conditions required for their preservation, paleofeces and coprolites are also relatively rare archaeological finds compared with the ubiquity of dental calculus. At the turn of the millennium, Poinar et al. [119–121] and Hofreiter et al. [122] first demonstrated the molecular potential of coprolites by amplifying and cloning DNA from extinct ground sloth and human coprolites to confirm the depositing species, and to investigate the diversity of ingested plant and animal material. These studies laid the foundation for other human and animal paleofeces and coprolite analyses using targeted



PCR to identify the presence and genetic diversity of humans in the Americas [123–125], elucidate the taxonomy and diet of extinct [126–129] and extant birds [130,131] as well as to taxonomically identify parasite eggs within coprolites [132–134].

More recently, high-throughput approaches have been applied to recover more representative observations of dietary components within paleofeces and coprolites. Last year, Wood et al. [135] applied metabarcoding of eukaryotic 18S rDNA and plant chloroplast *trnL* to investigate the diet of Polynesian domestic dogs, identifying the main dietary components as marine bony fishes (*Euteleosteiomorpha*) and plants (*Cucurbitaceae*). Analysis of retroviral DNA has also been attempted to access paleodietary information. In their metagenomic analysis of ancient human coprolites from the Caribbean, Rivera-Perez et al. [136] reported a range of eukaryotic retroviruses through a BLASTX comparison of the non-redundant NCBI database. Although the identified retroviruses corresponded with expected dietary patterns within the region, the dominance of well-characterized taxa (e.g., fowlpox virus) and model organisms (e.g., *Xenopus*) within the identified sequences, coupled with the detection of viruses infecting European plants (e.g., *Morus notabilis*) in pre-Columbian contexts raises questions about the extent to which the quantity and quality of reference sequences within public databases may bias the identifications (see “Section 3.2”).

Paleofeces and coprolites, however, provide genetic information beyond dietary inclusions. Tito et al. [137] were the first to recognize that coprolites and desiccated feces can facilitate the study of ancient gut microbiomes by reconstructing the microbial ecology of two 1300 year-old coprolites from Mexico through shotgun metagenomics. In a follow up study using a 16S rRNA metabarcoding approach on an expanded sample set, they revealed that not all fecal deposits may preserve the signature of endogenous ecology at the time of deposition [138]. While rapidly desiccated feces from cave environments were found to preserve the integrity of the gut ecology, others were found to have undergone self-digestion and decomposition as well as bacterial infiltration from the burial environment. The portion of coprolite sampled for analysis can also bias the identified microbial communities. Cano et al. [139] noted in their study of human coprolites from Central America that larger proportions of soil microbes were observed within the cortices compared to the inner core, likely reflecting environmental contamination on the coprolites surface; subsequently Wood and Wilmshurst [140] have recommended subsampling protocols to target the coprolite core. Paleofeces and coprolites can also reveal the presence of ancient viruses. Multiple-displacement amplification of viral genomes within a 14<sup>th</sup> Century human coprolite revealed an ancient virome comprised predominantly of double-stranded viral DNA (85.21%), with the majority of identifiable sequences corresponding to bacteriophages, especially *Siphoviridae* [141]. Although the virome was similar to that found in modern fecal matter and soil, further characterization of virome taxonomy and function was limited by the relatively high proportion of sequences of unknown origin.

### 2.2.3. Calcified Nodules

The human body may produce a variety of calcified tissues or “biological stones” with highly diverse structures and chemical compositions [142,143]. Although found principally within the digestive or urinary tracts, calcified tissues and stony neoplastic tumors or exudates may form within the lungs, vascular system, tear ducts, tendons, or skin, and are occasionally recovered during archaeological excavations. Two recent studies have exemplified the wealth of genetic information that may be preserved within these serendipitous finds. Kay et al. [144] applied metagenomic analysis to a calcified nodule recovered from the pelvic girdle of a 14<sup>th</sup> Century male skeleton from Geridu (Sardinia). Although tuberculosis was initially suspected as the cause of the pathology, the study instead identified sequences matching *Brucella melitensis* (0.48%), ultimately obtaining sufficient sequences to allow the reconstruction of a complete genome with 6.5× coverage. The proportion of human DNA within the nodule was relatively high for archaeological material (23%), demonstrating the potential of these substrates to provide information both on host genetics and bacterial pathogens. In a subsequent study, scanning electron microscopy and metagenomic analysis were used to investigate calcified abscesses

recovered from a woman's skeletal remains at a Byzantine cemetery in Troy [145]. Microscopic analysis revealed images of "ghost cells" preserved in a mineralized layer resulting from dystrophic calcification. Like the calcified nodule from Sardinia, the tumor contained an amalgam of human and microbial DNA. The human sex chromosome component of the DNA revealed a dominance of X chromosome sequences, with minute quantities of conserved Y chromosome sequences, interpreted to result from the presence of a male fetus, and suggesting that the growths developed in placental tissue [145]. The bacterial DNA recovered from shotgun sequencing identified *Staphylococcus saprophyticus* and *Gardnerella vaginalis* in high abundance, allowing for genome reconstruction and strain level analysis. Although calcified nodules may represent a rather unusual archaeological find, these case studies display the potential of these substrates to provide insight both into host genetics and ancient health and disease.

#### 2.2.4. Invertebrate Shell

There is a long history of research around organic molecules preserved within the calcareous exoskeleton of marine invertebrates. Previous research has focused almost exclusively on the preservation of proteins, particularly those responsible for biomineralization [146,147], and the use of these proteins for amino acid geochronology [148–151] and more recently, taxonomic identification [152]. The potential for the calcium carbonate of marine invertebrate shell to preserve genetic material has only recently been explored. Pawłowska et al. [153,154] demonstrated that well-preserved DNA could be recovered from micropaleontological taxa, like foraminifera, to reconstruct past micro-eukaryotic diversity and its relationship to climate change. Likewise, Villanea et al. [155] successfully recovered fragments of gastropod DNA from *Naesiotus* shells. Very recently, Der Sarkissian et al. [156] applied high-throughput shotgun DNA sequencing to mollusk shells up to 7000 years old from various countries (France, Korea, Japan, Norway, Italy, Denmark and the UK). They recovered average fragment lengths of 43–50 bp, enabling positive taxonomic identification of various mollusk species, marine organism pathogens as well as minute quantities of mtDNA from the marine environment [156]. Considering the wide array of marine species which produce carbonate exoskeletons (e.g., chitons, gastropods, cephalopods, bivalves, scaphopods) and the ubiquity of such shells in archaeological sites dating back to the Middle Pleistocene [157], there is a huge potential for using these substrates to track long term natural and human-induced changes in marine invertebrate paleodemography and biodiversity.

#### 2.2.5. Ancient Eggshell

Like marine invertebrate shell, eggshell represents another calcified deposit where early biomolecular analysis has focused principally on the protein component to reconstruct both geochronology [158,159] and ancient diets [160], with the genetic potential of ancient eggshell recognized only several years later. In modern genetic studies, as well as investigations on museum specimens, DNA extraction concentrates on the membranous layers on the inner face (e.g., [161,162]), prompting researchers to suggest that aDNA is more likely to survive on this inner shell where the membrane desiccates [163]. Using confocal microscopy to visualize DNA hotspots, Oskam et al. [163] demonstrated that DNA was more heavily concentrated around the boundaries of individual mammillary cones (a layer of the matrix close to the inner eggshell), likely resulting from the incorporation of oviduct epithelial cells into the calcite [164]. Oskam et al. [163] went on to sequence DNA extracted from ancient eggshells successfully conducting quantitative PCR (qPCR) on 16 specimens up to 19,000 years old from New Zealand, Madagascar and Australia, amplifying both mtDNA and a conserved region of the nuclear *c-mos* gene.

As with all ancient substrates, depositional environments and overall biomolecular preservation seem to influence ancient eggshell DNA quality and quantity. For example, in the sub-tropical climate of Madagascar, targeted enrichment coupled with NGS on elephant bird (*Aepyornis* sp.) eggshell resulted in just 3% endogenous DNA [165]. In spite of the relatively low recovery rate, eggshell DNA



preservation may prove to be comparable or even superior to bone. For example, previous DNA analysis from elephant bird bone produced only a partial mitogenome [166]—by comparison eggshell produced a complete mitochondrial genome with an additional 2271 bp, approximately twice the average coverage (33.5×) with fewer than 100 ambiguous sites [165]. Furthermore, Grealy et al. [165] report the recovery of nuclear DNA from eggshell, represented by 12,500 bp of exonic sequences. In their qPCR study of eggshells from various sites, Oskam et al. [163] detected proportionally fewer microbial reads in eggshell (1:10.6) than in bone (1:1333) [163], making eggshell a potentially attractive substrate for investigating avian genetics using high-throughput methods.

### 2.3. Biological and Cultural Archives

Natural history museums house millions of specimens from all over the world, and are the cornerstone of biological and biodiversity research. These collections have been exploited for their rich stores of historic and prehistoric genetic information for decades [167,168], with studies targeting DNA preserved in skeletal tissues, hair [169], claws [67], nails [68], skin [69], and invertebrate tissues [170,171] among others. More recently, other repositories such as herbaria and cultural archives (e.g., anthropological museums, documentary archives) have demonstrated their long-term genetic potential through successful recovery of DNA from historic plant collections [172–176] and parchments [177–179]. A comprehensive review of ancient DNA research on all biological and cultural archival material would be outside the scope of this paper; here, we focus on more unusual applications of DNA analysis to NHC, focusing specifically on fixed specimens, which hold particular, yet relatively untapped, promise as long-term reservoirs of ancient genetic material.

#### Fluid-Preserved Specimens

NHC hold vast numbers of fixed specimens collected over the last ~200 years, preserved typically through immersion in formaldehyde, formalin, methanol or ethanol. Visually, fluid-preserved specimens can appear near-perfect; this aesthetic preservation encouraged attempts at DNA extraction from the late 20<sup>th</sup> Century, mostly focused on ethanol-preserved specimens [180–184]. Extracting DNA from fixed materials, however, is problematic due to the molecules being particularly prone to oxidative and hydrolytic damage, fragmentation, base modifications and cross-linkage during the fixation process [185–187]. Compared with ethanol, formaldehyde and formalin treatments appear to be the most damaging to nucleic acids [188], and extensive research on formaldehyde-fixed clinical tissue specimens demonstrated that even relatively short exposure times (e.g., a few hours) induce DNA degradation [189], reduced DNA solubility [186], decreased PCR success rates due to crosslinking of proteins and DNA [190], and resulted in a higher-frequency of sequence alterations [191]. Ten years ago, the routine recovery of DNA from formalin- or formaldehyde-treated museum specimens seemed unlikely (see review in [192]), though more recent studies have expanded the potential of formalin fixed tissues by elucidating molecular preservation patterns [188], optimizing DNA extraction methods [193,194] and maximizing sequence recovery using NGS approaches [195,196].

Fixed specimens offer the potential not only to extract preserved host DNA, but also associated pathogens or even microbiomes. This potential was recognized early by Barnes et al. [181], who attempted to recover both host and pathogen DNA from historic diseased human tissues preserved in ethanol. Although modern biopsies indicated that *Helicobacter pylori* should be present in a ratio of 1:103 compared to human mtDNA, only sporadic evidence of the latter could be recovered from the fixed samples. More recently, Hühns et al. [197] reported success in recovering short fragments of *Mycobacterium tuberculosis* and human papillomavirus DNA from 78 to 12 year-old formalin fixed specimens. With further optimization, sufficient host and pathogenic DNA may be recovered to reconstruct host-pathogen interactions and pathogen evolution over the last several hundred years. Non-invasive approaches may also be an exciting new possibility for these fixed samples. For example, Shokralla et al. [198] have demonstrated the possibility of extracting a specimen's DNA directly from the preservative ethanol. As extraction and analytical methods continue to

evolve, we anticipate exponential growth in the genetic analysis of NHC and the field of “museomics” more broadly [199–201].

### 3. Moving Forward

High-throughput sequencing and omic technologies have revolutionized molecular research on ancient samples. With decreasing costs and increasing capacity, we have witnessed a surge of studies applying these techniques to a range of atypical archaeological materials and historical collections. Although the analysis of unusual biological substrates has the potential to open exciting new research directions in paleontology, paleoecology and paleogenetics, there are a number of key ethical and methodological considerations that need to be addressed if their analysis is to move beyond “novelty” to make lasting scientific contributions. Many of these new technologies have been applied to rare archaeological or paleontological finds, and there are significant ethical issues associated with the destructive analysis of finite archaeological or historical resources. First and foremost, researchers have an obligation to apply destructive techniques only if there is deemed a high likelihood of analytical success. However, our knowledge is limited on how genetic information may be preserved within these different materials. Furthermore, these atypical substrates may require specialized extraction techniques or analytical approaches to maximize the information that can be obtained from these precious resources. Below, we discuss issues of DNA preservation, authentication and analytical approaches in more detail, and make some recommendations for areas of future research to maximize the paleogenetic data that can be obtained from these varied resources.

#### 3.1. Predicting DNA Survival

To date, much of our understanding of DNA preservation in ancient substrates has been limited to hard tissues, such as tooth and bone [202–205]. Although it is expected that DNA within all ancient substrates will be highly fragmented [204], enriched in GC content [206], and feature mis-incorporations due to cytosine deamination and single-stranded fragment overhangs [207], we have little understanding of the extent to which degradation processes will limit analytical success in various biological substrates, especially considering the wide range of environments in which they have been preserved. In addition to confounding variables of time and temperature, archaeological artifacts and NHC may have been subjected to processing (e.g., tanning or dyeing) and/or conservation treatments that may also influence DNA degradation [58,188,208–210].

The majority of models predicting DNA degradation and fragmentation have been based on ancient bone. For example, decision-making models such as *thermal-age.eu* [211] have been developed to predict DNA preservation based on temperature-dependent rates of depurination in ancient bone [203]. Subsequent modeling of DNA preservation in previously published hard tissues genomic datasets suggested that while cytosine deamination is affected by sample age and the temperature of the depositional environment, DNA fragmentation rates do not appear to be influenced by age [202]. Instead of a constant rate of degradation, multiple stages of decay occur. Beginning with an intense initial stage of microbial attack and cellular activity that results in the destruction of significant quantities of endogenous DNA, a threshold is eventually reached, whereupon the DNA appears to be relatively stable, provided that the depositional environment is conducive to preservation [202]. This dual model of fragmentation has also been observed in anthropogenically preserved NHC specimens [188] and herbarium collections [209], suggesting that even tissues collected within the last century will display significant DNA degradation that may influence analytical success. Together, these findings suggest that DNA decay does not operate on a simple linear scale, and predicting the ultimate decay pattern in individual samples involves the detailed consideration of multiple complex processes resulting from primary depositional environments and/or conservation and subsequent storage.

The extent to which DNA preservation mirrors that of other biomolecules is also a key concern. Proteins survive longer than DNA in the archaeological record, but the extent to which proteins can be used to predict DNA survival is still under debate [212–215]. Alternative substrates such

as archaeological dental calculus [113,114], and food residues [216] have yielded some of the most abundant ancient proteomes, while Demarchi et al. [104] demonstrated the persistence of eggshell proteins as far back as 3.8 million years. The remarkable survival of these molecules is thought to result from protein binding strongly to a calcite surface, preventing hydrolytic breakdown. The extent to which similar factors may influence DNA survival in eggshells, and other calcified deposits, warrants further systematic testing to maximize the potential of these substrates. Understanding the methods by which DNA may bind to non-organic substrates like pottery or lithics is of particular importance, as we currently have little experimental data upon which to build reliable degradation models. Experimental methods, involving artificial DNA degradation [36,217–221] and different specimen treatments [58,64,88,210,222], are essential for developing more nuanced and informed models of DNA preservation on various substrates. These systematic studies are essential not only for making more informed choices when selecting rare samples for analysis, but for providing the necessary data to authenticate ancient genetic sequences recovered from atypical samples.

### 3.2. Authenticating Results

Authenticating endogenous DNA is a longstanding challenge faced by those studying aDNA, with multiple authentication criteria put in place and elaborated over time [84,223–228]. As the field advanced in terms of our understanding of DNA degradation, so too has our ability to recognize contaminating sequence by their deviation from “expected” molecular behavior. While the pre-2008 era of targeted PCR and Sanger sequencing relied principally on the criteria of authenticity proposed by Poinar and Cooper [84], more recent NGS studies have relied primarily on demonstrating “typical” signatures of post-mortem damage (PMD), principally cytosine deamination at 5'-overhangs [207,229] and sequence length distribution plots demonstrating DNA fragmented by hydrolysis. Detection of these expected patterns is now facilitated by the application of software such as “mapDamage” [230,231] and “bamdamage” [232], which quantify C-T transitions at the 5' end and G-A transitions at the 3' end of sequences. Over the last few years, researchers have come to rely heavily on damage plotting software as the primary means of authenticating data, even when studying unusual substrates where our knowledge of DNA degradation patterns is lacking. For example, Kay et al. [144] applied a damage pattern assessment in their metagenomic analysis of a 14<sup>th</sup> Century calcified nodule identifying the expected PMD pattern, but also discarded any sequences >150 bp (regardless of PMD). Although their comprehensive controls and conservative approach to data authentication is to be commended, the lack of knowledge of “appropriate molecular behavior” in novel substrates means that we may be potentially discarding authentic genetic data by assuming degradation patterns identical to those of bone or teeth.

A second challenge posed by unconventional substrates is that, unlike skeletal tissue, the taxonomic identity of the source(s) of the DNA (i.e., the species) is often unknown, for example in the metagenomic analysis of coprolites and calculus, or the analysis of artifacts, and thus more difficult to validate. Many of the aforementioned studies have used the BLAST application as a means of taxonomic identification for either metagenomic [34,136] or targeted amplicon sequences [101,135]. Although BLAST searches against the NCBI database is a widely used method for detecting sequence homology [233], taxonomic misidentification is a distinct possibility when working with short, damaged ancient DNA fragments. When studying ancient samples this problem is heightened since for many extinct species there are no well-matched modern genomes that could be used for comparisons [234]. NCBI entries also contain many low quality sequences that have not undergone full annotation and robust quality checks [235], and genomic sequences may include adaptor sequences [236], as well as contaminant human [237], domestic animal [238] and bacterial reads [239], which may result in query sequences being falsely aligned to non-authentic species. This is a particular challenge for microbial analysis, where closely related environmental contaminants and false positive matches against pathogens can easily confound taxonomic identifications [240]. Although using validated, curated databases of reference sequences (e.g., NCBI RefSeq) may be a more secure approach

to taxonomic identification, species identifications resulting from low-coverage data should always be approached with caution.

We agree with Carl Sagan that “extraordinary claims require extraordinary evidence”; until the analysis of atypical substrates becomes more routine, the results require an additional burden of proof for authentication. Many of the early authentication criteria, such as physically isolated laboratories, quantitation, detection of hydrolytic and oxidative damage and independent replication remain applicable to novel substrates [228], although these are never a substitute for critical thought [227]. As with all types of scientific inquiry, multiple lines of evidence are more powerful than one, and the incorporation of multiple biomolecular approaches serves not only to validate the results, but maximize the data that can be obtained from these, often extraordinary, materials.

### 3.3. Maximizing Information

Researchers have an obligation to maximize the data obtained during destructive analysis, but particularly when working with rare or novel specimens. Factors such as sample size, extraction method, sequencing approach and bioinformatic analysis are all important considerations for increasing the amount of retrieved data, while reducing the destructive impact of sampling.

#### 3.3.1. Sampling and DNA Extraction Methods


As most of the ancient genetic work of the last 30 years has been conducted on bone and teeth, sampling protocols and extraction methods have been tailored accordingly to maximize the retrieval of endogenous DNA [14–16,241,242]. With our limited understanding of where and how DNA preserves in non-bone material, there is pressing need for the development and systematic optimization of DNA extraction protocols from a variety of other tissues. The importance of bespoke protocols has recently been demonstrated for herbaria; for example, Weiß et al. [174] estimated that herbaria specimens have a per-nucleotide fragmentation rate of  $1.66 \times 10^{-4}$  meaning that DNA from even relatively recent herbaria specimens is expected to be highly fragmented. Gutaker et al. [175] subsequently modified their herbaria extraction procedure by using a *N*-phenacylthiazolium bromide buffer and altering the binding conditions of DNA to obtain a greater proportion of ultrashort (<50 bp) DNA fragments. Although over the last few years, some studies have begun to investigate the efficacy of different sampling and extraction protocols for degraded plants [243–245], archaeological dental calculus [113], coprolites [125,140], parchment [179] and formalin-fixed tissues [193,246], these have typically been conducted only when sufficient materials are available for systematic analysis. For more rare materials, such as archaeological artifacts, systematic testing may rely more on modern experimental or artificially degraded materials.

The difficulties in authenticating aDNA from atypical substrates, and the desire to maximize information from these unique genetic resources provides a compelling case for “unified protocols” or multi-proxy approaches [140,247] that are capable of extracting multiple biomolecules or organic residues from the same piece of starting material. The aforementioned studies have demonstrated a range of analytical approaches that can both complement and validate ancient genetic information, including microscopy [113,135], lipid analysis [248], proteomics [113,249], metabolomics [93,115], radiocarbon dating [122,128], collagen peptide mass fingerprinting [22], etc. In the majority of these studies, these complementary approaches have required the destruction of additional material; we advocate that future studies attempt to co-extract and analyze multiple biomolecules.

Non-destructive and non-invasive sampling methodologies are increasingly being demanded by museums and archives in order to preserve the integrity of unique accessions and maintain sufficient material for future analyses [250–252]. A variety of studies have developed novel biomolecular sampling techniques for bones and teeth [253,254], insects [170,255,256], shells [152], books and parchment [249,257], and fluid-preserved specimens [198] which do not require invasive sampling. As both our extraction techniques and bioinformatic approaches improve, potentially greater amounts of information may be achieved from minute samples.

### 3.3.2. Sequencing Technologies and Analytical Approaches

While early studies obtained ancient genetic sequences through PCR coupled with Sanger sequencing, since 2010 there has been a greater push towards metabarcoding and whole genome approaches coupled with high-throughput sequencing (Figure 2). Here again, bespoke analytical methods may vastly improve our access to ancient genetic information particularly when studying unusual substrates that are suspected to preserve only highly fragmented DNA. For example, Gansauge et al. [258] and Stiller et al. [259] both recently noted significant improvements in library yields from formalin-fixed tissues using a single-stranded vs. a double stranded library preparation method. Similarly, Gómez-Zeledón et al. [56] achieved success in recovering informative SNPs from highly fragmentary and inhibited archaeological wood samples using optimized TaqMan qPCR assays.



**Figure 2.** Area chart displaying the proportion of studies utilizing various methods in the analysis of neglected substrates, demonstrating the rise of whole genome and metagenomic approaches following the emergence of NGS technologies (includes studies reviewed in this paper, from 1996 to May 2017). WGS: whole genome sequencing; PCR: polymerase chain reaction.

The endogenous DNA content of ancient materials is variable to say the least, and may represent 0% of the total DNA recovered in whole genome approaches. As a response, targeted enrichment or “capture” based methods, where DNA of interest is immobilized on custom-designed baits in solution or on a physical array, is emerging as a more frequent methodological approach for atypical substrates, and has, for example, been applied to archaeological dental calculus [111], ancient eggshells [165], charred archaeobotanical remains [34] and fluid-preserved specimens [195] in order to recover mtDNA genomes, enrich full genome coverage, or capture ultraconserved elements, respectively (Figure 2).

Novel substrates, however, are increasingly being recognized for the genetic information they provide, not only about the host, but also associated microbes and microbiomes. While the sequencing of endogenous DNA facilitates the study of host genetics including ancestry, provenancing, phylogenetic and evolutionary questions (e.g., [260,261]), it is only one aspect of the biological history of an organism. As host microbes outnumber endogenous host cells at a ratio of at least 2:1 [262], and considering that humans (and animals) are critically dependent on these >1000 bacterial species, focusing entirely on the host genome means that an entire ancient ecosystem is disregarded [263,264]. Shotgun sequencing approaches to coprolites, calculus, calcified nodules, and shells have already demonstrated the ancient microbial information that may be obtained from non-skeletal tissues, in some



cases allowing for reconstructions of complete ancient microbial genomes [112,113], demonstrating the vast potential of these substrates to reconstruct the evolution of host-microbe interactions. For example, shotgun sequencing of ancient eggshell may reveal evolutionary changes in eggshell microbiomes, albumen immune defenses and avian pathogens that influence egg mortality [265], while metagenomic sequencing of fluid-preserved specimens may allow for the reconstruction of myriad vertebrate microbiomes in various states of health and disease. Metagenomic approaches may also provide information on the conservation status of museum collections and artifacts [92], for example, by revealing microbes implicated in their deterioration and decay [266].

The move towards whole genome approaches and NGS technologies yields significantly more data than have traditional sequencing methods, with much of this data made up of increasingly short DNA fragments. As bioinformatic methods improve, we will recover ever more genomic and metagenomic information from unconventional substrates, including increasingly high-resolution information on genetic traits underlying local adaptations [29], and phenotypes [267]. A promising advance is the recovery of epigenomic information from tissues other than bones and teeth. The potential of ancient epigenetics has already been demonstrated through the observation of methylation patterns in ancient bison bone [268], human bone [269], and Neanderthal and Denisovan skeletal material [270]. Recently, epigenetics have offered insight into other tissues, such as plant remains and hair [271]. For example, Smith et al. [272] detected genome-wide hypermethylation in ancient barley grains following viral infection, revealing the epigenetic impacts of exposure to biotic stress. The further development of epigenomic recovery and analysis could refine insight into age-at-death for biological organisms [271], provide evidence of how methylation status influenced evolutionary adaptation [270] and elucidate regulatory changes underlying species divergences [273].

#### 4. Conclusions

The past three decades of aDNA analysis have witnessed the recovery of genetic material from a wide range of ancient tissues and artifacts, including inorganic matrices once thought to be barren of surviving DNA (e.g., eggshells, ceramics). Further optimization of DNA extraction and analytical techniques, coupled with high-throughput and single-molecule sequencing will catalyze research in this area, allowing greater access to the often low-template, highly damaged DNA retained within these substrates. Nevertheless, these new technology-driven opportunities also present some challenges that need to be addressed, including:

1. Greater understanding of DNA preservation and degradation within various materials and their implications for selecting the most appropriate material for destructive analysis;
2. Developing appropriate multi-proxy approaches for authenticating ancient genetic data;
3. Raising awareness of the common pitfalls of aDNA analysis on atypical substrates that may be unfamiliar to many “modern” genetic laboratories entering the field.

Evidence for progress in these areas is already apparent—over 45% of the studies reviewed above had either method development or modeling DNA degradation as their primary research goal. A richer understanding of the potentials and pitfalls associated with the genetic analysis of these unconventional substrates will ultimately lead to more robust scientific results and more ethically informed decisions on sampling these irreplaceable historical resources.

**Acknowledgments:** This work was supported by the Arts and Humanities Research Council [grant number AH/N005015/1].

**Author Contributions:** E.J.G. and C.F.S. conceived of and wrote the paper.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

1. Higuchi, R.; Bowman, B.; Freiberger, M.; Ryder, O.A.; Wilson, A.C. DNA sequences from the quagga, an extinct member of the horse family. *Nature* **1984**, *312*, 282–284. [[CrossRef](#)] [[PubMed](#)]
2. Pääbo, S. Preservation of DNA in ancient Egyptian mummies. *J. Archaeol. Sci.* **1985**, *12*, 411–417. [[CrossRef](#)]
3. Pääbo, S. Molecular cloning of Ancient Egyptian mummy DNA. *Nature* **1985**, *314*, 644–645. [[CrossRef](#)] [[PubMed](#)]
4. Doran, G.H.; Dickel, D.N.; Ballinger, W.E., Jr.; Agee, O.F.; Laipis, P.J.; Hauswirth, W.W. Anatomical, cellular and molecular analysis of 8000-year-old human brain tissue from the Windover archaeological site. *Nature* **1986**, *323*, 803–806. [[CrossRef](#)] [[PubMed](#)]
5. Pääbo, S.; Gifford, J.A.; Wilson, A.C. Mitochondrial DNA sequences from a 7000-year old brain. *Nucleic Acids Res.* **1988**, *16*, 9775–9787. [[CrossRef](#)] [[PubMed](#)]
6. Pääbo, S. Ancient DNA: Extraction, characterization, molecular cloning, and enzymatic amplification. *Proc. Natl. Acad. Sci. USA* **1989**, *86*, 1939–1943. [[CrossRef](#)] [[PubMed](#)]
7. Thuesen, I.; Engberg, J. Recovery and analysis of human genetic material from mummified tissue and bone. *J. Archaeol. Sci.* **1990**, *17*, 679–689. [[CrossRef](#)]
8. Rogan, P.K.; Salvo, J.J. Study of nucleic acids isolated from ancient remains. *Am. J. Phys. Anthropol.* **1990**, *33*, 195–214. [[CrossRef](#)]
9. Hagelberg, E.; Sykes, B.; Hedges, R. Ancient bone DNA amplified. *Nature* **1989**, *342*, 485. [[CrossRef](#)] [[PubMed](#)]
10. Hänni, C.; Laudet, V.; Sakka, M.; Bègue, A.; Stéhelin, D. Amplification de fragments d'ADN mitochondrial à partir de dents et d'os humains anciens. *Comptes Rendus de l'Académie des Sciences* **1990**, *310*, 365–370. [[PubMed](#)]
11. Horai, S.; Hayasaka, K.; Murayama, K.; Wate, N.; Koike, H.; Nakai, N. DNA Amplification from Ancient Human Skeletal Remains and Their Sequence Analysis. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.* **1989**, *65*, 229–233. [[CrossRef](#)]
12. Carpenter, M.L.; Buenrostro, J.D.; Valdiosera, C.; Schroeder, H.; Allentoft, M.E.; Sikora, M.; Rasmussen, M.; Gravel, S.; Guillén, S.; Nekhrizov, G.; et al. Pulling out the 1%: Whole-genome capture for the targeted enrichment of ancient DNA sequencing libraries. *Am. J. Hum. Genet.* **2013**, *93*, 852–864. [[CrossRef](#)] [[PubMed](#)]
13. Rohland, N.; Siedel, H.; Hofreiter, M. A rapid column-based ancient DNA extraction method for increased sample throughput. *Mol. Ecol. Resour.* **2010**, *10*, 677–683. [[CrossRef](#)] [[PubMed](#)]
14. Dabney, J.; Knapp, M.; Glocke, I.; Gansauge, M.-T.; Weihmann, A.; Nickel, B.; Valdiosera, C.; García, N.; Pääbo, S.; Arsuaga, J.-L.; et al. Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 15758–15763. [[CrossRef](#)] [[PubMed](#)]
15. Yang, D.Y.; Eng, B.; Wayne, J.S.; Dudar, J.C.; Saunders, S.R. Technical note: Improved DNA extraction from ancient bones using silica-based spin columns. *Am. J. Phys. Anthropol.* **1998**, *105*, 539–543. [[CrossRef](#)]
16. Pinhasi, R.; Fernandes, D.; Sirak, K.; Novak, M.; Connell, S.; Alpaslan-Roodenberg, S.; Gerritsen, F.; Moiseyev, V.; Gromov, A.; Raczky, P.; et al. Optimal Ancient DNA Yields from the Inner Ear Part of the Human Petrous Bone. *PLoS ONE* **2015**, *10*, e0129102. [[CrossRef](#)] [[PubMed](#)]
17. Gamba, C.; Jones, E.R.; Teasdale, M.D.; McLaughlin, R.L.; Gonzalez-Fortes, G.; Mattiangeli, V.; Domboróczki, L.; Kővári, I.; Pap, I.; Anders, A.; et al. Genome flux and stasis in a five millennium transect of European prehistory. *Nat. Commun.* **2014**, *5*, 5257. [[CrossRef](#)] [[PubMed](#)]
18. Hansen, H.B.; Damgaard, P.B.; Margaryan, A.; Stenderup, J.; Lynnerup, N.; Willerslev, E.; Allentoft, M.E. Comparing Ancient DNA Preservation in Petrous Bone and Tooth Cementum. *PLoS ONE* **2017**, *12*, e0170940. [[CrossRef](#)] [[PubMed](#)]
19. Adler, C.J.; Haak, W.; Donlon, D.; Cooper, A. Survival and recovery of DNA from ancient teeth and bones. *J. Archaeol. Sci.* **2011**, *38*, 956–964. [[CrossRef](#)]
20. Burbano, H.A.; Hodges, E.; Green, R.E.; Briggs, A.W.; Krause, J.; Meyer, M.; Good, J.M.; Maricic, T.; Johnson, P.L.F.; Xuan, Z.; et al. Targeted investigation of the Neandertal genome by array-based sequence capture. *Science* **2010**, *328*, 723–725. [[CrossRef](#)] [[PubMed](#)]

21. Røed, K.H.; Bjørnstad, G.; Flagstad, Ø.; Haanes, H.; Huffhammer, A.K.; Jordhøy, P.; Rosvold, J. Ancient DNA reveals prehistoric habitat fragmentation and recent domestic introgression into native wild reindeer. *Conserv. Genet.* **2014**, *15*, 1137–1149. [[CrossRef](#)]
22. Von Holstein, I.C.C.; Ashby, S.P.; Van Doorn, N.L.; Sachs, S.M.; Buckley, M.; Meiri, M.; Barnes, I.; Brundle, A.; Collins, M.J. Searching for Scandinavians in Pre-Viking Scotland: Molecular Fingerprinting of Early Medieval Combs. *J. Archaeol. Sci.* **2014**, *41*, 1–6. [[CrossRef](#)]
23. Rollo, F.; Amici, A.; Salvi, R.; Garbuglia, A. Short but faithful pieces of ancient DNA. *Nature* **1988**, *335*, 774. [[CrossRef](#)] [[PubMed](#)]
24. Brown, T.A.; Allaby, R.G.; Brown, K.A.; O'Donoghue, K.; Sallares, R. DNA in wheat seeds from European archaeological sites. *Experientia* **1994**, *50*, 571–575. [[CrossRef](#)] [[PubMed](#)]
25. Allaby, R.G.; O'Donoghue, K.; Sallares, R.; Jones, M.K.; Brown, T.A. Evidence for the survival of ancient DNA in charred wheat seeds from European archaeological sites. *Anc. Biomol.* **1997**, *1*, 119–129.
26. Allaby, R.G.; Banerjee, M.; Brown, T.A. Evolution of the high molecular weight glutenin loci of the A, B, D, and G genomes of wheat. *Genome* **1999**, *42*, 296–307. [[CrossRef](#)] [[PubMed](#)]
27. Freitas, F.O.; Bendel, G.; Allaby, R.G.; Brown, T.A. DNA from primitive maize landraces and archaeological remains: Implications for the domestication of maize and its expansion into South America. *J. Archaeol. Sci.* **2003**, *30*, 901–908. [[CrossRef](#)]
28. Jaenicke-Després, V.; Buckler, E.S.; Smith, B.D.; Gilbert, M.T.P.; Cooper, A.; Doebley, J.; Pääbo, S. Early allelic selection in maize as revealed by ancient DNA. *Science* **2003**, *302*, 1206–1208. [[CrossRef](#)] [[PubMed](#)]
29. Palmer, S.A.; Moore, J.D.; Clapham, A.J.; Rose, P.; Allaby, R.G. Archaeogenetic evidence of ancient nubian barley evolution from six to two-row indicates local adaptation. *PLoS ONE* **2009**, *4*, e6301. [[CrossRef](#)] [[PubMed](#)]
30. Bunning, S.L.; Jones, G.; Brown, T.A. Next generation sequencing of DNA in 3300-year-old charred cereal grains. *J. Archaeol. Sci.* **2012**, *39*, 2780–2784. [[CrossRef](#)]
31. Allaby, R.G.; Gutaker, R.; Clarke, A.C.; Pearson, N.; Ware, R.; Palmer, S.A.; Kitchen, J.L.; Smith, O. Using archaeogenomic and computational approaches to unravel the history of local adaptation in crops. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **2014**, *370*, 20130377. [[CrossRef](#)] [[PubMed](#)]
32. Ramos-Madrigal, J.; Smith, B.D.; Moreno-Mayar, J.V.; Gopalakrishnan, S.; Ross-Ibarra, J.; Gilbert, M.T.P.; Wales, N. Genome Sequence of a 5310-Year-Old Maize Cob Provides Insights into the Early Stages of Maize Domestication. *Curr. Biol.* **2016**, *26*, 3195–3201. [[CrossRef](#)] [[PubMed](#)]
33. Mascher, M.; Schuenemann, V.J.; Davidovich, U.; Marom, N.; Himmelbach, A.; Hübner, S.; Korol, A.; David, M.; Reiter, E.; Riehl, S.; et al. Genomic analysis of 6000-year-old cultivated grain illuminates the domestication history of barley. *Nat. Genet.* **2016**, *48*, 1089–1093. [[CrossRef](#)] [[PubMed](#)]
34. Nistelberger, H.M.; Smith, O.; Wales, N.; Star, B.; Boessenkool, S. The efficacy of high-throughput sequencing and target enrichment on charred archaeobotanical remains. *Sci. Rep.* **2016**, *6*, 37347. [[CrossRef](#)] [[PubMed](#)]
35. Hagenblad, J.; Morales, J.; Leino, M.W.; Rodríguez-Rodríguez, A.C. Farmer fidelity in the Canary Islands revealed by ancient DNA from prehistoric seeds. *J. Archaeol. Sci.* **2017**, *78*, 78–87. [[CrossRef](#)]
36. Threadgold, J.; Brown, T.A. Degradation of DNA in artificially charred wheat seeds. *J. Archaeol. Sci.* **2003**, *30*, 1067–1076. [[CrossRef](#)]
37. Erickson, D.L.; Smith, B.D.; Clarke, A.C.; Sandweiss, D.H.; Tuross, N. An Asian origin for a 10,000-year-old domesticated plant in the Americas. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 18315–18320. [[CrossRef](#)] [[PubMed](#)]
38. Palmer, S.A.; Smith, O.; Allaby, R.G. The blossoming of plant archaeogenetics. *Ann. Anat.* **2012**, *194*, 146–156. [[CrossRef](#)] [[PubMed](#)]
39. Goloubinoff, P.; Pääbo, S.; Wilson, A.C. Evolution of maize inferred from sequence diversity of an Adh2 gene segment from archaeological specimens. *Proc. Natl. Acad. Sci. USA* **1993**, *90*, 1997–2001. [[CrossRef](#)] [[PubMed](#)]
40. Lia, V.V.; Confalonieri, V.A.; Ratto, N.; Hernández, J.A.C.; Alzogaray, A.M.M.; Poggio, L.; Brown, T.A. Microsatellite typing of ancient maize: Insights into the history of agriculture in southern South America. *Proc. Biol. Sci.* **2007**, *274*, 545–554. [[CrossRef](#)] [[PubMed](#)]
41. Pollmann, B.; Jacomet, S.; Schlumbaum, A. Morphological and genetic studies of waterlogged Prunus species from the Roman vicus Tasgetium (Eschenz, Switzerland). *J. Archaeol. Sci.* **2005**, *32*, 1471–1480. [[CrossRef](#)]
42. Suyama, Y.; Kawamuro, K.; Kinoshita, I.; Yoshimura, K.; Tsumura, Y.; Takahara, H. DNA sequence from a fossil pollen of *Abies* spp. from Pleistocene peat. *Genes Genet. Syst.* **1996**, *71*, 145–149. [[CrossRef](#)] [[PubMed](#)]

43. Parducci, L.; Suyama, Y.; Lascoux, M.; Bennett, K.D. Ancient DNA from pollen: A genetic record of population history in Scots pine. *Mol. Ecol.* **2005**, *14*, 2873–2882. [[CrossRef](#)] [[PubMed](#)]
44. Paffetti, D.; Vettori, C.; Caramelli, D.; Vernesi, C.; Lari, M.; Paganelli, A.; Paule, L.; Giannini, R. Unexpected presence of *Fagus orientalis* complex in Italy as inferred from 45,000-year-old DNA pollen samples from Venice lagoon. *BMC Evol. Biol.* **2007**, *7* (Suppl. 2), S6. [[CrossRef](#)] [[PubMed](#)]
45. Lágler, R.; Gyulai, G.; Humphreys, M.; Szabó, Z.; Horváth, L.; Bittsánszky, A.; Kiss, J.; Holly, L.; Heszky, L. Morphological and molecular analysis of common millet (*P. miliaceum*) cultivars compared to an aDNA sample from the 15th century (Hungary). *Euphytica* **2005**, *146*, 77–85. [[CrossRef](#)]
46. Lágler, R.; Gyulai, G.; Szabó, Z.; Tóth, Z.; Bittsánszky, A.; Horváth, L.; Kiss, J.; Gyulai, F.; Heszky, L. Molecular diversity of common millet (*P. miliaceum*) compared to archaeological samples excavated from the 4th and 15th centuries. *Hung. Agric. Res.* **2006**, *1*, 14–19.
47. Gyulai, G.; Humphreys, M.; Lagler, R.; Szabo, Z.; Toth, Z.; Bittsánszky, A.; Gyulai, F.; Heszky, L. Seed remains of common millet from the 4th (Mongolia) and 15th (Hungary) centuries: AFLP, SSR and mtDNA sequence recoveries. *Seed Sci. Res.* **2006**, *16*, 179–191. [[CrossRef](#)]
48. Kistler, L.; Shapiro, B. Ancient DNA confirms a local origin of domesticated chenopod in eastern North America. *J. Archaeol. Sci.* **2011**, *38*, 3549–3554. [[CrossRef](#)]
49. Blatter, R.H.E.; Jacomet, S.; Schlumbaum, A. Spelt-specific alleles in HMW glutenin genes from modern and historical European spelt (*Triticum spelta* L.). *Theor. Appl. Genet.* **2002**, *104*, 329–337. [[CrossRef](#)] [[PubMed](#)]
50. Blatter, R.H.E.; Jacomet, S.; Schlumbaum, A. Little Evidence for the Preservation of a Single-Copy Gene in Charred Archaeological Wheat. *Anc. Biomol.* **2002**, *4*, 65–77. [[CrossRef](#)]
51. LeBlanc, S.A.; Kreisman, L.S.C.; Kemp, B.M.; Smiley, F.E.; Carlyle, S.W.; Dhody, A.N.; Benjamin, T. Quids and aprons: Ancient DNA from artifacts from the American Southwest. *J. Field Archaeol.* **2007**, *32*, 161–175. [[CrossRef](#)]
52. Deguilloux, M.F.; Bertel, L.; Celant, A.; Pemonge, M.H.; Sadori, L.; Magri, D.; Petit, R.J. Genetic analysis of archaeological wood remains: First results and prospects. *J. Archaeol. Sci.* **2006**, *33*, 1216–1227. [[CrossRef](#)]
53. Liepelt, S.; Sperisen, C.; Deguilloux, M.-F.; Petit, R.J.; Kissling, R.; Spencer, M.; de Beaulieu, J.-L.; Taberlet, P.; Gielly, L.; Ziegenhagen, B. Authenticated DNA from ancient wood remains. *Ann. Bot.* **2006**, *98*, 1107–1111. [[CrossRef](#)] [[PubMed](#)]
54. Lee, A.B.; Cooper, T.A. Improved direct PCR screen for bacterial colonies: Wooden toothpicks inhibit PCR amplification. *Biotechniques* **1995**, *18*, 225–226. [[PubMed](#)]
55. Matheson, C.D.; Gurney, C.; Esau, N.; Lehto, R. Assessing PCR inhibition from humic substances. *Open Enzym. Inhib. J.* **2010**, *3*, 38–45. [[CrossRef](#)]
56. Gómez-Zeledón, J.; Grasse, W.; Runge, F.; Land, A.; Spring, O. TaqMan qPCR pushes boundaries for the analysis of millennial wood. *J. Archaeol. Sci.* **2017**, *79*, 53–61. [[CrossRef](#)]
57. Sønstebo, J.H.; Gielly, L.; Brysting, A.K.; Elven, R.; Edwards, M.; Haile, J.; Willerslev, E.; Coissac, E.; Rioux, D.; Sannier, J.; et al. Using next-generation sequencing for molecular reconstruction of past Arctic vegetation and climate. *Mol. Ecol. Resour.* **2010**, *10*, 1009–1018. [[CrossRef](#)] [[PubMed](#)]
58. Vuissoz, A.; Worobey, M.; Odegaard, N.; Bunce, M.; Machado, C.A.; Lynnerup, N.; Peacock, E.E.; Gilbert, M.T.P. The survival of PCR-amplifiable DNA in cow leather. *J. Archaeol. Sci.* **2007**, *34*, 823–829. [[CrossRef](#)]
59. Schlumbaum, A.; Campos, P.F.; Volken, S.; Volken, M.; Hafner, A.; Schibler, J. Ancient DNA, a Neolithic legging from the Swiss Alps and the early history of goat. *J. Archaeol. Sci.* **2010**, *37*, 1247–1251. [[CrossRef](#)]
60. Burger, J.; Pfeiffer, I.; Hummel, S.; Fuchs, R.; Brenig, B.; Herrmann, B. Mitochondrial and nuclear DNA from (pre) historic hide-derived material. *Anc. Biomol.* **2001**, *3*, 227–238.
61. Gilbert, M.T.P.; Tomsho, L.P.; Rendulic, S.; Packard, M.; Drautz, D.I.; Sher, A.; Tikhonov, A.; Dalén, L.; Kuznetsova, T.; Kosintsev, P.; et al. Whole-genome shotgun sequencing of mitochondria from ancient hair shafts. *Science* **2007**, *317*, 1927–1930. [[CrossRef](#)] [[PubMed](#)]
62. Rasmussen, M.; Li, Y.; Lindgreen, S.; Pedersen, J.S.; Albrechtsen, A.; Moltke, I.; Metspalu, M.; Metspalu, E.; Kivisild, T.; Gupta, R.; et al. Ancient human genome sequence of an extinct Palaeo-Eskimo. *Nature* **2010**, *463*, 757–762. [[CrossRef](#)] [[PubMed](#)]
63. Sinding, M.-H.S.; Arneborg, J.; Nyegaard, G.; Gilbert, M.T.P. Ancient DNA unravels the truth behind the controversial GUS Greenlandic Norse fur samples: The bison was a horse, and the muskox and bears were goats. *J. Archaeol. Sci.* **2015**, *53*, 297–303. [[CrossRef](#)]



64. Brandt, L.Ø.; Tranekjer, L.D.; Mannering, U.; Ringgaard, M.; Frei, K.M.; Willerslev, E.; Gleba, M.; Gilbert, M.T.P. Characterising the potential of sheep wool for ancient DNA analyses. *Archaeol. Anthropol. Sci.* **2011**, *3*, 209–221. [[CrossRef](#)]
65. Sinding, M.-H.S.; Gilbert, M.T.P.; Grønnow, B.; Gulløv, H.C.; Toft, P.A.; Foote, A.D. Minimally destructive DNA extraction from archaeological artefacts made from whale baleen. *J. Archaeol. Sci.* **2012**, *39*, 3750–3753. [[CrossRef](#)]
66. Sinding, M.-H.S.; Tervo, O.M.; Grønnow, B.; Gulløv, H.C.; Toft, P.A.; Bachmann, L.; Fietz, K.; Rekdal, S.L.; Christoffersen, M.F.; Heide-Jørgensen, M.P.; et al. Sex determination of baleen whale artefacts: Implications for ancient DNA use in zooarchaeology. *J. Archaeol. Sci. Rep.* **2016**, *10*, 345–349. [[CrossRef](#)]
67. Hedmark, E.; Ellegren, H. Microsatellite genotyping of DNA isolated from claws left on tanned carnivore hides. *Int. J. Legal Med.* **2005**, *119*, 370–373. [[CrossRef](#)] [[PubMed](#)]
68. Willerslev, E.; Gilbert, M.T.P.; Binladen, J.; Ho, S.Y.W.; Campos, P.F.; Ratan, A.; Tomsho, L.P.; da Fonseca, R.R.; Sher, A.; Kuznetsova, T.V.; et al. Analysis of complete mitochondrial genomes from extinct and extant rhinoceroses reveals lack of phylogenetic resolution. *BMC Evol. Biol.* **2009**, *9*, 95. [[CrossRef](#)] [[PubMed](#)]
69. Casas-Marce, M.; Revilla, E.; Godoy, J.A. Searching for DNA in museum specimens: A comparison of sources in a mammal species. *Mol. Ecol. Resour.* **2009**, *10*, 502–507. [[CrossRef](#)] [[PubMed](#)]
70. Ellegren, H. Genomic DNA from Museum Bird Feathers. In *Ancient DNA*; Herrmann, B., Hummel, S., Eds.; Springer: New York, NY, USA, 1994; pp. 211–217.
71. Sefc, K.M.; Payne, R.B.; Sorenson, M.D.; Fleischer, R.C. Microsatellite amplification from museum feather samples: Effects of fragment size and template concentration on genotyping errors. *Auk* **2003**, *120*, 982–989. [[CrossRef](#)]
72. Horváth, M.B.; Martínez-Cruz, B.; Negro, J.J.; Kalmár, L.; Godoy, J.A. An overlooked DNA source for non-invasive genetic analysis in birds. *J. Avian Biol.* **2004**, *36*, 84–88. [[CrossRef](#)]
73. Speller, C.F.; Nicholas, G.P.; Yang, D.Y. Feather barbs as a good source of mtDNA for bird species identification in forensic wildlife investigations. *Investig. Genet.* **2011**, *2*, 16. [[CrossRef](#)] [[PubMed](#)]
74. Hartnup, K.; Huynen, L.; Te Kanawa, R.; Shepherd, L.D.; Millar, C.D.; Lambert, D.M. Ancient DNA recovers the origins of Māori feather cloaks. *Mol. Biol. Evol.* **2011**, *28*, 2741–2750. [[CrossRef](#)] [[PubMed](#)]
75. Crabtree, D.E. *An Introduction to Flintworking*; Idaho State University Museum Occasional Paper No. 28; Idaho State University Museum: Pocatello, IA, USA, 1972.
76. Odell, G.H.; Odell-Vereecken, F. Verifying the reliability of lithic use-wear assessments by “blind tests”: The low-power approach. *J. Field Archaeol.* **1980**, *7*, 87–120. [[CrossRef](#)]
77. Keeley, L.H. *Experimental Determination of Stone Tool Uses: A Microwear Analysis*; University of Chicago Press: Chicago, IL, USA, 1980.
78. Loy, T.H. Prehistoric blood residues: Detection on tool surfaces and identification of species of origin. *Science* **1983**, *220*, 1269–1271. [[CrossRef](#)] [[PubMed](#)]
79. Newman, M.; Julig, P. The Identification of Protein Residues on Lithic Artifacts from a Stratified Boreal Forest Site. *Can. J. Archaeol.* **1989**, *13*, 119–132.
80. Loy, T.H. The artifact as site: An example of the biomolecular analysis of organic residues on prehistoric tools. *World Archaeol.* **1993**, *25*, 44–63. [[CrossRef](#)] [[PubMed](#)]
81. Hardy, B.L.; Raff, R.A.; Raman, V. Recovery of Mammalian DNA from Middle Paleolithic Stone Tools. *J. Archaeol. Sci.* **1997**, *24*, 601–611. [[CrossRef](#)]
82. Shanks, O.C.; Bonnichsen, R.; Vella, A.T.; Ream, W. Recovery of Protein and DNA Trapped in Stone Tool Microcracks. *J. Archaeol. Sci.* **2001**, *28*, 965–972. [[CrossRef](#)]
83. Shanks, O.C.; Hodges, L.; Tilley, L.; Kornfeld, M.; Lou Larson, M.; Ream, W. DNA from ancient stone tools and bones excavated at Bugas-Holding, Wyoming. *J. Archaeol. Sci.* **2005**, *32*, 27–38. [[CrossRef](#)]
84. Poinar, H.N.; Cooper, A. Ancient DNA: Do it right or not at all. *Science* **2000**, *5482*, 416. [[CrossRef](#)]
85. Leonard, J.A.; Shanks, O.; Hofreiter, M.; Kreuz, E.; Hodges, L.; Ream, W.; Wayne, R.K.; Fleischer, R.C. Animal DNA in PCR reagents plagues ancient DNA research. *J. Archaeol. Sci.* **2007**, *34*, 1361–1366. [[CrossRef](#)]
86. Kooyman, B.; Newman, M.E.; Ceri, H. Verifying the reliability of blood residue analysis on archaeological tools. *J. Archaeol. Sci.* **1992**, *19*, 265–269. [[CrossRef](#)]
87. Craig, O.E.; Collins, M.J. The Removal of Protein from Mineral Surfaces: Implications for Residue Analysis of Archaeological Materials. *J. Archaeol. Sci.* **2002**, *29*, 1077–1082. [[CrossRef](#)]



88. Tuross, N.; Barnes, I.; Potts, R. Protein Identification of Blood Residues on Experimental Stone Tools. *J. Archaeol. Sci.* **1996**, *23*, 289–296. [[CrossRef](#)]
89. Loy, T.H.; Hardy, B.L. Blood residue analysis of 90,000-year-old stone tools from Tabun Cave, Israel. *Antiquity* **1992**, *66*, 24–35. [[CrossRef](#)]
90. Eisele, J.A.; Fowler, D.D.; Haynes, G.; Lewis, R.A. Survival and detection of blood residues on stone tools. *Antiquity* **1995**, *69*, 36–46. [[CrossRef](#)]
91. Fiedel, S. Blood from Stones? Some Methodological and Interpretive Problems in Blood Residue Analysis. *J. Archaeol. Sci.* **1996**, *23*, 139–147. [[CrossRef](#)]
92. Adamiak, J.; Otlewska, A.; Tafer, H.; Lopandic, K.; Gutarowska, B.; Sterflinger, K.; Piñar, G. First evaluation of the microbiome of built cultural heritage by using the Ion Torrent next generation sequencing platform. *Int. Biodeterior. Biodegrad.* **2017**. [[CrossRef](#)]
93. Gutarowska, B.; Celikkol-Aydin, S.; Bonifay, V.; Otlewska, A.; Aydin, E.; Oldham, A.L.; Brauer, J.I.; Duncan, K.E.; Adamiak, J.; Sunner, J.A.; et al. Metabolomic and high-throughput sequencing analysis—modern approach for the assessment of biodeterioration of materials from historic buildings. *Front. Microbiol.* **2015**, *6*, 979. [[CrossRef](#)] [[PubMed](#)]
94. Evershed, R.P.; Heron, C.; Charters, S.; Goad, L.J. The survival of food residues: New methods of analysis, interpretation and application. *Proc. Br. Acad.* **1992**, *77*, 187–208.
95. Charters, S.; Evershed, R.P.; Goad, L.J.; Leyden, A.; Blinkhorn, P.W.; Denham, V. Quantification and distribution of lipid in archaeological ceramics: Implications for sampling potsherds for organic residue analysis and the classification of vessel use. *Archaeometry* **1993**, *35*, 211–223. [[CrossRef](#)]
96. Regert, M.; Bland, H.A.; Dudd, S.N.; Bergen, P.F.V.; Evershed, R.P. Free and bound fatty acid oxidation products in archaeological ceramic vessels. *Proc. R. Soc. B* **1998**, *265*, 2027–2032. [[CrossRef](#)]
97. Reber, E.A.; Dudd, S.N.; Van der Merwe, N.J.; Evershed, R.P. Direct detection of maize in pottery residues via compound specific stable carbon isotope analysis. *Antiquity* **2004**, *78*, 682–691. [[CrossRef](#)]
98. Cavalieri, D.; McGovern, P.E.; Hartl, D.L.; Mortimer, R.; Polsinelli, M. Evidence for *S. cerevisiae* fermentation in ancient wine. *J. Mol. Evol.* **2003**, *57* (Suppl. 1), S226–S232. [[CrossRef](#)] [[PubMed](#)]
99. Hansson, M.C.; Foley, B.P. Ancient DNA fragments inside Classical Greek amphoras reveal cargo of 2400-year-old shipwreck. *J. Archaeol. Sci.* **2008**, *35*, 1169–1176. [[CrossRef](#)]
100. Foley, B.P.; Hansson, M.C.; Kourkoumelis, D.P.; Theodoulou, T.A. Aspects of ancient Greek trade re-evaluated with amphora DNA evidence. *J. Archaeol. Sci.* **2012**, *39*, 389–398. [[CrossRef](#)]
101. Robinson, H.A.; Insoll, T.; Kankpeyeng, B.W.; Brown, K.A.; Brown, T.A. Ritual complexity in a past community revealed by ancient DNA analysis of pre-colonial terracotta items from Northern Ghana. *J. Archaeol. Sci.* **2017**, *79*, 10–18. [[CrossRef](#)]
102. Lindahl, T. Instability and Decay of the Primary Structure of DNA. *Nature* **1993**, *362*, 709–715. [[CrossRef](#)] [[PubMed](#)]
103. Campos, P.F.; Craig, O.E.; Turner-Walker, G.; Peacock, E.; Willerslev, E.; Gilbert, M.T.P. DNA in ancient bone—Where is it located and how should we extract it? *Ann. Anat.* **2012**, *194*, 7–16. [[CrossRef](#)] [[PubMed](#)]
104. Demarchi, B.; Hall, S.; Roncal-Herrero, T.; Freeman, C.L.; Woolley, J.; Crisp, M.K.; Wilson, J.; Fotakis, A.; Fischer, R.; Kessler, B.M.; et al. Protein sequences bound to mineral surfaces persist into deep time. *eLife* **2016**, *5*, e17092. [[CrossRef](#)] [[PubMed](#)]
105. Warinner, C.; Speller, C.; Collins, M.J. A new era in palaeomicrobiology: Prospects for ancient dental calculus as a long-term record of the human oral microbiome. *Philos. Trans. R. Soc. B* **2015**, *370*, 20130376. [[CrossRef](#)] [[PubMed](#)]
106. Schroeder, H.E. Formation and inhibition of dental calculus. *J. Periodontol.* **1969**, *40*, 643–646. [[CrossRef](#)] [[PubMed](#)]
107. Jin, Y.; Yip, H.-K. Supragingival calculus: Formation and control. *Crit. Rev. Oral Biol. Med.* **2002**, *13*, 426–441. [[CrossRef](#)] [[PubMed](#)]
108. Preus, H.R.; Marvik, O.J.; Selvig, K.A.; Bennike, P. Ancient bacterial DNA (aDNA) in dental calculus from archaeological human remains. *J. Archaeol. Sci.* **2011**, *38*, 1827–1831. [[CrossRef](#)]
109. Adler, C.J.; Dobney, K.; Weyrich, L.S.; Kaidonis, J.; Walker, A.W.; Haak, W.; Bradshaw, C.J.A.; Townsend, G.; Sołtysiak, A.; Alt, K.W.; et al. Sequencing ancient calcified dental plaque shows changes in oral microbiota with dietary shifts of the Neolithic and Industrial revolutions. *Nat. Genet.* **2013**, *45*, 450–455. [[CrossRef](#)] [[PubMed](#)]

110. Ziesemer, K.A.; Mann, A.E.; Sankaranarayanan, K.; Schroeder, H.; Ozga, A.T.; Brandt, B.W.; Zaura, E.; Waters-Rist, A.; Hoogland, M.; Salazar-García, D.C.; et al. Intrinsic challenges in ancient microbiome reconstruction using 16S rRNA gene amplification. *Sci. Rep.* **2015**, *5*, 16498. [[CrossRef](#)] [[PubMed](#)]
111. Ozga, A.T.; Nieves-Colón, M.A.; Honap, T.P.; Sankaranarayanan, K.; Hofman, C.A.; Milner, G.R.; Lewis, C.M., Jr.; Stone, A.C.; Warinner, C. Successful enrichment and recovery of whole mitochondrial genomes from ancient human dental calculus. *Am. J. Phys. Anthropol.* **2016**, *160*, 220–228. [[CrossRef](#)] [[PubMed](#)]
112. Weyrich, L.S.; Duchene, S.; Soubrier, J.; Arriola, L.; Llamas, B.; Breen, J.; Morris, A.G.; Alt, K.W.; Caramelli, D.; Dresely, V.; et al. Neanderthal behaviour, diet, and disease inferred from ancient DNA in dental calculus. *Nature* **2017**, *544*, 357–361. [[CrossRef](#)] [[PubMed](#)]
113. Warinner, C.; Rodrigues, J.F.M.; Vyas, R.; Trachsel, C.; Shved, N.; Grossmann, J.; Radini, A.; Hancock, Y.; Tito, R.Y.; Fiddyment, S.; et al. Pathogens and host immunity in the ancient human oral cavity. *Nat. Genet.* **2014**, *46*, 336–344. [[CrossRef](#)] [[PubMed](#)]
114. Warinner, C.; Hendy, J.; Speller, C.; Cappellini, E.; Fischer, R.; Trachsel, C.; Arneborg, J.; Lynnerup, N.; Craig, O.E.; Swallow, D.M.; et al. Direct evidence of milk consumption from ancient human dental calculus. *Sci. Rep.* **2014**, *4*, 7104. [[CrossRef](#)] [[PubMed](#)]
115. Velsko, I.M.; Overmyer, K.A.; Speller, C.; Collins, M.; Loe, L.; Frantz, L.A.F.; Martinez, J.B.R.; Chavez, E.; Klaus, L.; Sankaranarayanan, K.; et al. The dental calculus metabolome in modern and historic samples. *bioRxiv* **2017**, 136176. [[CrossRef](#)]
116. Warinner, C.; Speller, C.; Collins, M.J.; Lewis, C.M., Jr. Ancient human microbiomes. *J. Hum. Evol.* **2015**, *79*, 125–136. [[CrossRef](#)] [[PubMed](#)]
117. Weyrich, L.S.; Dobney, K.; Cooper, A. Ancient DNA analysis of dental calculus. *J. Hum. Evol.* **2014**, *6*, 5–12. [[CrossRef](#)] [[PubMed](#)]
118. Hollocher, K.; Hollocher, T.C. Early processes in the fossilization of terrestrial feces to coprolites, and microstructure preservation. In *Vertebrate Coprolites: Bulletin 57*; Hunt, A.P., Milàn, J., Lucas, S.G., Spielmann, J.A., Eds.; New Mexico Museum of Natural History and Science: Albuquerque, NM, USA, 2012; pp. 93–98.
119. Poinar, H.N.; Hofreiter, M.; Spaulding, W.G.; Martin, P.S.; Stankiewicz, B.A.; Bland, H.; Evershed, R.P.; Possnert, G.; Pääbo, S. Molecular coproscopy: Dung and diet of the extinct ground sloth *Nothotheriops shastensis*. *Science* **1998**, *281*, 402–406. [[CrossRef](#)] [[PubMed](#)]
120. Poinar, H.N.; Kuch, M.; Sobolik, K.D.; Barnes, I.; Stankiewicz, A.B.; Kuder, T.; Spaulding, W.G.; Bryant, V.M.; Cooper, A.; Pääbo, S. A molecular analysis of dietary diversity for three archaic Native Americans. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 4317–4322. [[CrossRef](#)] [[PubMed](#)]
121. Poinar, H.; Kuch, M.; McDonald, G.; Martin, P.; Pääbo, S. Nuclear gene sequences from a late pleistocene sloth coprolite. *Curr. Biol.* **2003**, *13*, 1150–1152. [[CrossRef](#)]
122. Hofreiter, M.; Poinar, H.N.; Spaulding, W.G.; Bauer, K.; Martin, P.S.; Possnert, G.; Pääbo, S. A molecular analysis of ground sloth diet through the last glaciation. *Mol. Ecol.* **2000**, *9*, 1975–1984. [[CrossRef](#)] [[PubMed](#)]
123. Gilbert, M.T.P.; Jenkins, D.L.; Götherstrom, A.; Naveran, N.; Sanchez, J.J.; Hofreiter, M.; Thomsen, P.F.; Binladen, J.; Higham, T.F.G.; Yohe, R.M.; et al. DNA from Pre-Clovis Human Coprolites in Oregon, North America. *Science* **2008**, *320*, 786–789. [[CrossRef](#)] [[PubMed](#)]
124. Jenkins, D.L.; Davis, L.G.; Stafford, T.W.; Campos, P.F.; Hockett, B.; Jones, G.T.; Cummings, L.S.; Yost, C.; Connolly, T.J.; Yohe, R.M.; et al. Clovis Age Western Stemmed Projectile Points and Human Coprolites at the Paisley Caves. *Science* **2012**, *337*, 223–228. [[CrossRef](#)] [[PubMed](#)]
125. Kemp, B.M.; Monroe, C.; Smith, D.G. Repeat silica extraction: A simple technique for the removal of PCR inhibitors from DNA extracts. *J. Archaeol. Sci.* **2006**, *33*, 1680–1689. [[CrossRef](#)]
126. Wood, J.R.; Rawlence, N.J.; Rogers, G.M.; Austin, J.J.; Worthy, T.H.; Cooper, A. Coprolite deposits reveal the diet and ecology of the extinct New Zealand megaherbivore moa (*Aves, Dinornithiformes*). *Quat. Sci. Rev.* **2008**, *27*, 2593–2602. [[CrossRef](#)]
127. Wood, J.R.; Wilmshurst, J.M.; Worthy, T.H.; Cooper, A. First coprolite evidence for the diet of *Anomalopteryx didiformis*, an extinct forest ratite from New Zealand. *N. Z. J. Ecol.* **2012**, *36*, 164–170.
128. Wood, J.R.; Wilmshurst, J.M.; Wagstaff, S.J.; Worthy, T.H.; Rawlence, N.J.; Cooper, A. High-Resolution Coproecology: Using Coprolites to Reconstruct the Habits and Habitats of New Zealand's Extinct Upland Moa (*Megalapteryx didinus*). *PLoS ONE* **2012**, *7*, e40025. [[CrossRef](#)] [[PubMed](#)]

129. Wood, J.R.; Wilmshurst, J.M.; Richardson, S.J.; Rawlence, N.J.; Wagstaff, S.J.; Worthy, T.H.; Cooper, A. Resolving lost herbivore community structure using coprolites of four sympatric moa species (*Aves: Dinornithiformes*). *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 16910–16915. [[CrossRef](#)] [[PubMed](#)]
130. Speller, C.F.; Kemp, B.M.; Wyatt, S.D.; Monroe, C.; Lipe, W.D.; Arndt, U.M.; Yang, D.Y. Ancient mitochondrial DNA analysis reveals complexity of indigenous North American turkey domestication. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 2807–2812. [[CrossRef](#)] [[PubMed](#)]
131. Wood, J.R.; Wilmshurst, J.M.; Worthy, T.H.; Holzapfel, A.S.; Cooper, A. A lost link between a flightless parrot and a parasitic plant and the potential role of coprolites in conservation paleobiology. *Conserv. Biol.* **2012**, *26*, 1091–1099. [[CrossRef](#)] [[PubMed](#)]
132. Iñiguez, A.M.; Reinhard, K.J.; Araújo, A.; Ferreira, L.F.; Vicente, A.C.P. Enterobius vermicularis: Ancient DNA from North and South American human coprolites. *Mem. Inst. Oswaldo Cruz* **2003**, *98* (Suppl. 1), 67–69. [[CrossRef](#)] [[PubMed](#)]
133. Iñiguez, A.M.; Reinhard, K.; Carvalho Gonçalves, M.L.; Ferreira, L.F.; Araújo, A.; Paulo Vicente, A.C. SL1 RNA gene recovery from Enterobius vermicularis ancient DNA in pre-Columbian human coprolites. *Int. J. Parasitol.* **2006**, *36*, 1419–1425. [[CrossRef](#)] [[PubMed](#)]
134. Leles, D.; Araújo, A.; Ferreira, L.F.; Vicente, A.C.P.; Iñiguez, A.M. Molecular paleoparasitological diagnosis of *Ascaris* sp. from coprolites: New scenery of ascariasis in pre-Colombian South America times. *Mem. Inst. Oswaldo Cruz* **2008**, *103*, 106–108. [[CrossRef](#)] [[PubMed](#)]
135. Wood, J.R.; Crown, A.; Cole, T.L.; Wilmshurst, J.M. Microscopic and ancient DNA profiling of Polynesian dog (kurī) coprolites from northern New Zealand. *J. Archaeol. Sci. Rep.* **2016**, *6*, 496–505. [[CrossRef](#)]
136. Rivera-Perez, J.I.; Cano, R.J.; Narganes-Storde, Y.; Chanlatte-Baik, L.; Toranzos, G.A. Retroviral DNA Sequences as a Means for Determining Ancient Diets. *PLoS ONE* **2015**, *10*, e0144951. [[CrossRef](#)] [[PubMed](#)]
137. Tito, R.Y.; Macmil, S.; Wiley, G.; Najar, F.; Cleeland, L.; Qu, C.; Wang, P.; Romagne, F.; Leonard, S.; Ruiz, A.J.; et al. Phylotyping and functional analysis of two ancient human microbiomes. *PLoS ONE* **2008**, *3*, e3703. [[CrossRef](#)] [[PubMed](#)]
138. Tito, R.Y.; Knights, D.; Metcalf, J.; Obregon-Tito, A.J.; Cleeland, L.; Najar, F.; Roe, B.; Reinhard, K.; Sobolik, K.; Belknap, S.; et al. Insights from characterizing extinct human gut microbiomes. *PLoS ONE* **2012**, *7*, e51146. [[CrossRef](#)] [[PubMed](#)]
139. Cano, R.J.; Rivera-Perez, J.; Toranzos, G.A.; Santiago-Rodriguez, T.M.; Narganes-Storde, Y.M.; Chanlatte-Baik, L.; García-Roldán, E.; Bunkley-Williams, L.; Massey, S.E. Paleomicrobiology: Revealing Fecal Microbiomes of Ancient Indigenous Cultures. *PLoS ONE* **2014**, *9*, e106833. [[CrossRef](#)] [[PubMed](#)]
140. Wood, J.R.; Wilmshurst, J.M. A protocol for subsampling Late Quaternary coprolites for multi-proxy analysis. *Quat. Sci. Rev.* **2016**, *138*, 1–5. [[CrossRef](#)]
141. Appelt, S.; Fancello, L.; Le Bailly, M.; Raoult, D.; Drancourt, M.; Desnues, C. Viruses in a 14th-century coprolite. *Appl. Environ. Microbiol.* **2014**, *80*, 2648–2655. [[CrossRef](#)] [[PubMed](#)]
142. Lonsdale, K. Human stones. *Science* **1968**, *159*, 1199–1207. [[CrossRef](#)] [[PubMed](#)]
143. Gładkowska-Rzeczycka, J.J.; Nowakowski, D. A biological stone from a medieval cemetery in Poland. *PLoS ONE* **2014**, *9*, e109096. [[CrossRef](#)] [[PubMed](#)]
144. Kay, G.L.; Sergeant, M.J.; Giuffra, V.; Bandiera, P.; Milanese, M.; Bramanti, B.; Bianucci, R.; Pallen, M.J. Recovery of a medieval *Brucella melitensis* genome using shotgun metagenomics. *mBio* **2014**, *5*, e01337-14. [[CrossRef](#)] [[PubMed](#)]
145. Devault, A.M.; Mortimer, T.D.; Kitchen, A.; Kiesewetter, H.; Enk, J.M.; Golding, G.B.; Southon, J.; Kuch, M.; Duggan, A.T.; Aylward, W.; et al. A molecular portrait of maternal sepsis from Byzantine Troy. *eLife* **2017**, *6*, e20983. [[CrossRef](#)] [[PubMed](#)]
146. Weiner, S.; Traub, W.; Parker, S.B. Macromolecules in Mollusc Shells and Their Functions in Biomineralization. *Philos. Trans. R. Soc. B* **1984**, *304*, 425–434. [[CrossRef](#)]
147. Marin, F.; Marie, B.; Hamada, S.B.; Ramos-Silva, P.; Le Roy, N.; Guichard, N.; Wolf, S.E.; Montagnani, C.; Joubert, C.; Piquemal, D.; et al. Shellome': Proteins involved in mollusk shell biomineralization-diversity, functions. *Recent Adv. Pearl Res.* **2013**, 149–166.
148. Miller, G.H.; Brigham-Grette, J. Amino acid geochronology: Resolution and precision in carbonate fossils. *Quat. Int.* **1989**, *1*, 111–128. [[CrossRef](#)]

149. Miller, G.H.; Hare, P.E. Amino acid geochronology: Integrity of the carbonate matrix and potential of molluscan fossils. In *Biogeochemistry of Amino Acids*; Hare, P.E., King, K., Hoering, I.C., Eds.; John Wiley & Sons: New York, NY, USA, 1980; pp. 415–442.
150. Penkman, K.E.H.; Preece, R.C.; Keen, D.H.; Maddy, D.; Schreve, D.C.; Collins, M.J. Testing the aminostratigraphy of fluvial archives: The evidence from intra-crystalline proteins within freshwater shells. *Quat. Sci. Rev.* **2007**, *26*, 2958–2969. [[CrossRef](#)] [[PubMed](#)]
151. Demarchi, B.; Williams, M.G.; Milner, N.; Russell, N.; Bailey, G.; Penkman, K. Amino acid racemization dating of marine shells: A mound of possibilities. *Quat. Int.* **2011**, *239*, 114–124. [[CrossRef](#)] [[PubMed](#)]
152. Demarchi, B.; O'Connor, S.; de Lima Ponzoni, A.; de Almeida Rocha Ponzoni, R.; Sheridan, A.; Penkman, K.; Hancock, Y.; Wilson, J. An integrated approach to the taxonomic identification of prehistoric shell ornaments. *PLoS ONE* **2014**, *9*, e99839. [[CrossRef](#)] [[PubMed](#)]
153. Pawłowska, J.; Lejzerowicz, F.; Esling, P.; Szczuciński, W.; Zajączkowski, M.; Pawłowski, J. Ancient DNA sheds new light on the Svalbard foraminiferal fossil record of the last millennium. *Geobiology* **2014**, *12*, 277–288. [[CrossRef](#)] [[PubMed](#)]
154. Pawłowska, J.; Zajączkowski, M.; Łacka, M.; Lejzerowicz, F.; Esling, P.; Pawłowski, J. Palaeoceanographic changes in Hornsund Fjord (Spitsbergen, Svalbard) over the last millennium: New insights from ancient DNA. *Clim. Past Discuss.* **2015**, *11*, 3665–3698. [[CrossRef](#)]
155. Villanea, F.A.; Parent, C.E.; Kemp, B.M. Reviving Galápagos snails: Ancient DNA extraction and amplification from shells of probably extinct endemic land snails. *J. Molluscan* **2016**, *82*, 449–456. [[CrossRef](#)]
156. Der Sarkissian, C.; Pichereau, V.; Dupont, C.; Ilse, P.C.; Perrigault, M.; Butler, P.; Chauvaud, L.; Eiriksson, J.; Scourse, J.; Paillard, C.; et al. Ancient DNA analysis identifies marine mollusc shells as new metagenomic archives of the past. *Mol. Ecol. Resour.* **2017**. [[CrossRef](#)] [[PubMed](#)]
157. Marean, C.W.; Bar-Matthews, M.; Bernatchez, J.; Fisher, E.; Goldberg, P.; Herries, A.I.R.; Jacobs, Z.; Jerardino, A.; Karkanas, P.; Minichillo, T.; et al. Early human use of marine resources and pigment in South Africa during the Middle Pleistocene. *Nature* **2007**, *449*, 905–908. [[CrossRef](#)] [[PubMed](#)]
158. Miller, G.H.; Beaumont, P.B.; Jull, A.J.; Johnson, B. Pleistocene geochronology and palaeothermometry from protein diagenesis in ostrich eggshells: Implications for the evolution of modern humans. *Philos. Trans. R. Soc. B* **1992**, *337*, 149–157. [[CrossRef](#)] [[PubMed](#)]
159. Brooks, A.S.; Hare, P.E.; Kokis, J.E.; Miller, G.H.; Ernst, R.D.; Wendorf, F. Dating Pleistocene archaeological sites by protein diagenesis in ostrich eggshell. *Science* **1990**, *248*, 60–64. [[CrossRef](#)] [[PubMed](#)]
160. Hobson, K.A. Reconstructing Avian Diets Using Stable-Carbon and Nitrogen Isotope Analysis of Egg Components: Patterns of Isotopic Fractionation and Turnover. *Condor* **1995**, *97*, 752–762. [[CrossRef](#)]
161. Bush, K.L.; Vinsky, M.D.; Aldridge, C.L.; Paszkowski, C.A. A comparison of sample types varying in invasiveness for use in DNA sex determination in an endangered population of greater Sage-Grouse (*Centrocercus urophasianus*). *Conserv. Genet.* **2005**, *6*, 867–870. [[CrossRef](#)]
162. Lee, P.L.M.; Prys-Jones, R.P. Extracting DNA from museum bird eggs, and whole genome amplification of archive DNA. *Mol. Ecol. Resour.* **2008**, *8*, 551–560. [[CrossRef](#)] [[PubMed](#)]
163. Oskam, C.L.; Haile, J.; McLay, E.; Rigby, P.; Allentoft, M.E.; Olsen, M.E.; Bengtsson, C.; Miller, G.H.; Schwenninger, J.-L.; Jacomb, C.; et al. Fossil avian eggshell preserves ancient DNA. *Proc. Biol. Sci.* **2010**, *277*, 1991–2000. [[CrossRef](#)] [[PubMed](#)]
164. Egloff, C.; Labrosse, A.; Hebert, C.; Crump, D. A nondestructive method for obtaining maternal DNA from avian eggshells and its application to embryonic viability determination in herring gulls (*Larus argentatus*). *Mol. Ecol. Resour.* **2009**, *9*, 19–27. [[CrossRef](#)] [[PubMed](#)]
165. Grealy, A.; Phillips, M.; Miller, G.; Gilbert, M.T.P.; Rouillard, J.-M.; Lambert, D.; Bunce, M.; Haile, J. Eggshell palaeogenomics: Palaeognath evolutionary history revealed through ancient nuclear and mitochondrial DNA from Madagascan elephant bird (*Aepyornis* sp.) eggshell. *Mol. Phylogenet. Evol.* **2017**, *109*, 151–163. [[CrossRef](#)] [[PubMed](#)]
166. Mitchell, K.J.; Llamas, B.; Soubrier, J.; Rawlence, N.J.; Worthy, T.H.; Wood, J.; Lee, M.S.Y.; Cooper, A. Ancient DNA reveals elephant birds and kiwi are sister taxa and clarifies ratite bird evolution. *Science* **2014**, *344*, 898–900. [[CrossRef](#)] [[PubMed](#)]
167. Thomas, R.H. Analysis of DNA from natural history museum collections. In *Molecular Ecology and Evolution: Approaches and Applications. Experientia Supplementum, vol 69*; Schierwater, B., Streit, B., Wagner, G.P., DeSalle, R., Eds.; Birkhäuser: Basel, Switzerland, 1994; pp. 311–321.



168. McLean, B.S.; Bell, K.C.; Dunnum, J.L.; Abrahamson, B.; Colella, J.P.; Deardorff, E.R.; Weber, J.A.; Jones, A.K.; Salazar-Miralles, F.; Cook, J.A. Natural history collections-based research: Progress, promise, and best practices. *J. Mammal.* **2016**, *97*, 287–297. [[CrossRef](#)] [[PubMed](#)]
169. Gilbert, M.T.P.; Wilson, A.S.; Bunce, M.; Hansen, A.J.; Willerslev, E.; Shapiro, B.; Higham, T.F.G.; Richards, M.P.; O'Connell, T.C.; Tobin, D.J.; et al. Ancient mitochondrial DNA from hair. *Curr. Biol.* **2004**, *14*, R463–R464. [[CrossRef](#)] [[PubMed](#)]
170. Thomsen, P.F.; Elias, S.; Gilbert, M.T.P.; Haile, J.; Munch, K.; Kuzmina, S.; Froese, D.G.; Sher, A.; Holdaway, R.N.; Willerslev, E. Non-destructive sampling of ancient insect DNA. *PLoS ONE* **2009**, *4*, e5048. [[CrossRef](#)] [[PubMed](#)]
171. Tagliavia, M.; Massa, B.; Albanese, I.; La Farina, M. DNA Extraction from Orthoptera Museum Specimens. *Anal. Lett.* **2011**, *44*, 1058–1062. [[CrossRef](#)]
172. Yoshida, K.; Schuenemann, V.J.; Cano, L.M.; Pais, M.; Mishra, B.; Sharma, R.; Lanz, C.; Martin, F.N.; Kamoun, S.; Krause, J.; et al. The rise and fall of the *Phytophthora infestans* lineage that triggered the Irish potato famine. *eLife* **2013**, *2*, e00731. [[PubMed](#)]
173. Martin, M.D.; Zimmer, E.A.; Olsen, M.T.; Foote, A.D.; Gilbert, M.T.P.; Brush, G.S. Herbarium specimens reveal a historical shift in phylogeographic structure of common ragweed during native range disturbance. *Mol. Ecol.* **2014**, *23*, 1701–1716. [[CrossRef](#)] [[PubMed](#)]
174. Weiß, C.L.; Schuenemann, V.J.; Devos, J.; Shirsekar, G.; Reiter, E.; Gould, B.A.; Stinchcombe, J.R.; Krause, J.; Burbano, H.A. Temporal patterns of damage and decay kinetics of DNA retrieved from plant herbarium specimens. *R. Soc. Open Sci.* **2016**, *3*, 160239. [[CrossRef](#)] [[PubMed](#)]
175. Gutaker, R.M.; Reiter, E.; Furtwängler, A.; Schuenemann, V.J.; Burbano, H.A. Extraction of ultrashort DNA molecules from herbarium specimens. *Biotechniques* **2017**, *62*, 76–79. [[CrossRef](#)] [[PubMed](#)]
176. Bakker, F.T. Herbarium genomics: Skimming and plastomics from archival specimens. *Webbia* **2017**, *72*, 35–45. [[CrossRef](#)]
177. Campana, M.G.; Bower, M.A.; Bailey, M.J.; Stock, F.; O'Connell, T.C.; Edwards, C.J.; Checkley-Scott, C.; Knight, B.; Spencer, M.; Howe, C.J. A flock of sheep, goats and cattle: Ancient DNA analysis reveals complexities of historical parchment manufacture. *J. Archaeol. Sci.* **2010**, *37*, 1317–1325. [[CrossRef](#)]
178. Teasdale, M.D.; van Doorn, N.L.; Fiddymont, S.; Webb, C.C.; O'Connor, T.; Hofreiter, M.; Collins, M.J.; Bradley, D.G. Paging through history: Parchment as a reservoir of ancient DNA for next generation sequencing. *Philos. Trans. R. Soc. B* **2015**, *370*, 20130379. [[CrossRef](#)] [[PubMed](#)]
179. Lech, T. Ancient DNA in historical parchments—Identifying a procedure for extraction and amplification of genetic material. *Genet. Mol. Res.* **2016**, *15*. [[CrossRef](#)] [[PubMed](#)]
180. Beebe, T.J.; Rowe, G.; Burke, T. Archive contributions to a molecular phylogeography of the toad *Bufo calamita* in Britain. *Biochem. Genet.* **1998**, *36*, 219–228. [[CrossRef](#)] [[PubMed](#)]
181. Barnes, I.; Holton, J.; Vaira, D.; Spigelman, M.; Thomas, M.G. An assessment of the long-term preservation of the DNA of a bacterial pathogen in ethanol-preserved archival material. *J. Pathol.* **2000**, *192*, 554–559. [[CrossRef](#)]
182. Spigelman, M.; Barnes, I.; Holton, J.; Vaira, D.; Thomas, M.G. Long-term DNA survival in ethanol-preserved archival material. *Ann. R. Coll. Surg. Engl.* **2001**, *83*, 283–284. [[PubMed](#)]
183. Stuart, B.L.; Dugan, K.A.; Allard, M.W.; Kearney, M. Extraction of nuclear DNA from bone of skeletonized and fluid-preserved museum specimens. *Syst. Biodivers.* **2006**, *4*, 133–136. [[CrossRef](#)]
184. Miller, J.A.; Beentjes, K.K.; van Helsdingen, P.; Ijland, S. Which specimens from a museum collection will yield DNA barcodes? A time series study of spiders in alcohol. *Zookeys* **2013**, *365*, 245–261. [[CrossRef](#)] [[PubMed](#)]
185. Huang, H.; Hopkins, P.B. DNA interstrand cross-linking by formaldehyde: Nucleotide sequence preference and covalent structure of the predominant cross-link formed in synthetic oligonucleotides. *J. Am. Chem. Soc.* **1993**, *115*, 9402–9408. [[CrossRef](#)]
186. Douglas, M.P.; Rogers, S.O. DNA damage caused by common cytological fixatives. *Mutat. Res. Fundam. Mol. Mech. Mutagen.* **1998**, *401*, 77–88. [[CrossRef](#)]
187. Vink, C.J.; Thomas, S.M.; Paquin, P.; Hayashi, C.Y.; Hedin, M. The effects of preservatives and temperatures on arachnid DNA. *Invertebr. Syst.* **2005**, *19*, 99–104. [[CrossRef](#)]



188. Zimmermann, J.; Hajibabaei, M.; Blackburn, D.C.; Hanken, J.; Cantin, E.; Posfai, J.; Evans, T.C., Jr. DNA damage in preserved specimens and tissue samples: A molecular assessment. *Front. Zool.* **2008**, *5*, 18. [[CrossRef](#)] [[PubMed](#)]
189. Srinivasan, M.; Sedmak, D.; Jewell, S. Effect of fixatives and tissue processing on the content and integrity of nucleic acids. *Am. J. Pathol.* **2002**, *161*, 1961–1971. [[CrossRef](#)]
190. Ben-Ezra, J.; Johnson, D.A.; Rossi, J.; Cook, N.; Wu, A. Effect of fixation on the amplification of nucleic acids from paraffin-embedded material by the polymerase chain reaction. *J. Histochem. Cytochem.* **1991**, *39*, 351–354. [[CrossRef](#)] [[PubMed](#)]
191. Williams, C.; Pontén, F.; Moberg, C.; Söderkvist, P.; Uhlén, M.; Pontén, J.; Sitbon, G.; Lundeberg, J. A high frequency of sequence alterations is due to formalin fixation of archival specimens. *Am. J. Pathol.* **1999**, *155*, 1467–1471. [[CrossRef](#)]
192. Tang, E. *Path to Effective Recovering of DNA from Formalin-Fixed Biological Samples in Natural History Collections: Workshop Summary*; National Academies Press: Washington, DC, USA, 2006. [[CrossRef](#)]
193. Paireder, S.; Werner, B.; Bailer, J.; Werther, W.; Schmid, E.; Patzak, B.; Cichna-Markl, M. Comparison of protocols for DNA extraction from long-term preserved formalin fixed tissues. *Anal. Biochem.* **2013**, *439*, 152–160. [[CrossRef](#)] [[PubMed](#)]
194. Lin, J.; Kennedy, S.H.; Svarovsky, T.; Rogers, J.; Kemnitz, J.W.; Xu, A.; Zondervan, K.T. High-quality genomic DNA extraction from formalin-fixed and paraffin-embedded samples deparaffinized using mineral oil. *Anal. Biochem.* **2009**, *395*, 265–267. [[CrossRef](#)] [[PubMed](#)]
195. Ruane, S.; Austin, C.C. Phylogenomics using formalin-fixed and 100+ year-old intractable natural history specimens. *Mol. Ecol. Res.* **2017**. [[CrossRef](#)] [[PubMed](#)]
196. Hykin, S.M.; Bi, K.; McGuire, J.A. Fixing Formalin: A Method to Recover Genomic-Scale DNA Sequence Data from Formalin-Fixed Museum Specimens Using High-Throughput Sequencing. *PLoS ONE* **2015**, *10*, e0141579. [[CrossRef](#)] [[PubMed](#)]
197. Hühns, M.; Erbersdobler, A.; Obliers, A.; Röpenack, P. Identification of HPV Types and Mycobacterium Tuberculosis Complex in Historical Long-Term Preserved Formalin Fixed Tissues in Different Human Organs. *PLoS ONE* **2017**, *12*, e0170353. [[CrossRef](#)] [[PubMed](#)]
198. Shokralla, S.; Singer, G.A.C.; Hajibabaei, M. Direct PCR amplification and sequencing of specimens' DNA from preservative ethanol. *Biotechniques* **2010**, *48*, 233–234. [[CrossRef](#)] [[PubMed](#)]
199. Guschanski, K.; Krause, J.; Sawyer, S.; Valente, L.M.; Bailey, S.; Finstermeier, K.; Sabin, R.; Gilissen, E.; Sonet, G.; Nagy, Z.T.; et al. Next-generation museomics disentangles one of the largest primate radiations. *Syst. Biol.* **2013**, *62*, 539–554. [[CrossRef](#)] [[PubMed](#)]
200. Fabre, P.-H.; Vilstrup, J.T.; Raghavan, M.; Sarkissian, C.D.; Willerslev, E.; Douzery, E.J.P.; Orlando, L. Rodents of the Caribbean: Origin and diversification of hutias unravelled by next-generation museomics. *Biol. Lett.* **2014**, *10*, 20140266. [[CrossRef](#)] [[PubMed](#)]
201. Silva, C.; Besnard, G.; Piot, A.; Razanatsoa, J.; Oliveira, R.P.; Vorontsova, M.S. Museomics resolve the systematics of an endangered grass lineage endemic to north-western Madagascar. *Ann. Bot.* **2017**, *119*, 339–351. [[CrossRef](#)] [[PubMed](#)]
202. Kistler, L.; Ware, R.; Smith, O.; Collins, M.; Allaby, R.G. A new model for ancient DNA decay based on paleogenomic meta-analysis. *Nucleic Acids Res* **2017**, *45*, 6310–6320. [[CrossRef](#)] [[PubMed](#)]
203. Smith, C.I.; Chamberlain, A.T.; Riley, M.S.; Stringer, C.; Collins, M.J. The thermal history of human fossils and the likelihood of successful DNA amplification. *J. Hum. Evol.* **2003**, *45*, 203–217. [[CrossRef](#)]
204. Allentoft, M.E.; Collins, M.; Harker, D.; Haile, J.; Oskam, C.L.; Hale, M.L.; Campos, P.F.; Samaniego, J.A.; Gilbert, M.T.P.; Willerslev, E.; et al. The half-life of DNA in bone: Measuring decay kinetics in 158 dated fossils. *Proc. R. Soc. Lond. B Biol. Sci.* **2012**, *279*, 4724–4733. [[CrossRef](#)] [[PubMed](#)]
205. Collins, M.J.; Nielsen-Marsh, C.M.; Hiller, J.; Smith, C.I.; Roberts, J.P.; Prigodich, R.V.; Wess, T.J.; Csapò, J.; Millard, A.R.; Turner-Walker, G. The survival of organic matter in bone: A review. *Archaeometry* **2002**, *44*, 383–394. [[CrossRef](#)]
206. Wales, N.; Carøe, C.; Sandoval-Velasco, M.; Gamba, C.; Barnett, R.; Samaniego, J.A.; Madrigal, J.R.; Orlando, L.; Gilbert, M.T.P. New insights on single-stranded versus double-stranded DNA library preparation for ancient DNA. *Biotechniques* **2015**, *59*, 368–371. [[CrossRef](#)] [[PubMed](#)]

207. Briggs, A.W.; Stenzel, U.; Johnson, P.L.F.; Green, R.E.; Kelso, J.; Prüfer, K.; Meyer, M.; Krause, J.; Ronan, M.T.; Lachmann, M.; et al. Patterns of damage in genomic DNA sequences from a Neandertal. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 14616–14621. [[CrossRef](#)] [[PubMed](#)]
208. Wang, X.; Li, L.; Zhao, J.; Li, F.; Guo, W.; Chen, X. Effects of different preservation methods on inter simple sequence repeat (ISSR) and random amplified polymorphic DNA (RAPD) molecular markers in botanic samples. *C. R. Biol.* **2017**, *340*, 204–213. [[CrossRef](#)] [[PubMed](#)]
209. Staats, M.; Cuenca, A.; Richardson, J.E.; Ginkel, R.V.; Petersen, G.; Seberg, O.; Bakker, F.T. DNA Damage in Plant Herbarium Tissue. *PLoS ONE* **2011**, *6*, e28448. [[CrossRef](#)] [[PubMed](#)]
210. Atanesyan, L.; Steenkamer, M.J.; Horstman, A.; Moelans, C.B.; Schouten, J.P.; Savola, S.P. Optimal Fixation Conditions and DNA Extraction Methods for MLPA Analysis on FFPE Tissue-Derived DNA. *Am. J. Clin. Pathol.* **2017**, *147*, 60–68. [[CrossRef](#)] [[PubMed](#)]
211. Thermal Age Web Tool. Available online: <http://thermal-age.eu/> (accessed on 11 July 2017).
212. Wadsworth, C.; Procopio, N.; Anderung, C.; Carretero, J.-M.; Iriarte, E.; Valdiosera, C.; Elburg, R.; Penkman, K.; Buckley, M. Comparing ancient DNA survival and proteome content in 69 archaeological cattle tooth and bone samples from multiple European sites. *J. Proteom.* **2017**, *158*, 1–8. [[CrossRef](#)] [[PubMed](#)]
213. Cappellini, E.; Collins, M.J.; Gilbert, M.T.P. Biochemistry. Unlocking ancient protein palimpsests. *Science* **2014**, *343*, 1320–1322. [[CrossRef](#)] [[PubMed](#)]
214. Charlton, S.; Alexander, M.; Collins, M.; Milner, N.; Mellars, P.; O’Connell, T.C.; Stevens, R.E.; Craig, O.E. Finding Britain’s last hunter-gatherers: A new biomolecular approach to “unidentifiable” bone fragments utilising bone collagen. *J. Archaeol. Sci.* **2016**, *73*, 55–61. [[CrossRef](#)]
215. Collins, M.J.; Penkman, K.E.H.; Rohland, N.; Shapiro, B.; Dobberstein, R.C.; Ritz-Timme, S.; Hofreiter, M. Is amino acid racemization a useful tool for screening for ancient DNA in bone? *Proc. Biol. Sci.* **2009**, *276*, 2971–2977. [[CrossRef](#)] [[PubMed](#)]
216. Yang, Y.; Shevchenko, A.; Knaust, A.; Abuduresule, I.; Li, W.; Hu, X.; Wang, C.; Shevchenko, A. Proteomics evidence for kefir dairy in Early Bronze Age China. *J. Archaeol. Sci.* **2014**, *45*, 178–186. [[CrossRef](#)]
217. Waite, E.R.; Child, A.M.; Craig, O.E.; Collins, M.J.; Gelsthorpe, K.; Brown, T.A. A preliminary investigation of DNA stability in bone during artificial diagenesis. *Bull. Soc. Géol. Fr.* **1997**, *168*, 547–554.
218. Dobberstein, R.C.; Huppertz, J.; von Wurmb-Schwark, N.; Ritz-Timme, S. Degradation of biomolecules in artificially and naturally aged teeth: Implications for age estimation based on aspartic acid racemization and DNA analysis. *Forensic Sci. Int.* **2008**, *179*, 181–191. [[CrossRef](#)] [[PubMed](#)]
219. Banerjee, M.; Brown, T.A. Non-random DNA damage resulting from heat treatment: Implications for sequence analysis of ancient DNA. *J. Archaeol. Sci.* **2004**, *31*, 59–63. [[CrossRef](#)]
220. Von Wurmb-Schwark, N.; Harbeck, M.; Wiesbrock, U.; Schroeder, I.; Ritz-Timme, S.; Oehmichen, M. Extraction and amplification of nuclear and mitochondrial DNA from ancient and artificially aged bones. *Leg. Med.* **2003**, *5* (Suppl. 1), S169–S172. [[CrossRef](#)]
221. Chalfoun, D.J.; Tuross, N. Botanical remains: Utility in protein and DNA research. *Anc. Biomol.* **1999**, *3*, 67–79.
222. Ottoni, C.; Koon, H.E.C.; Collins, M.J.; Penkman, K.E.H.; Rickards, O.; Craig, O.E. Preservation of ancient DNA in thermally damaged archaeological bone. *Naturwissenschaften* **2009**, *96*, 267–278. [[CrossRef](#)] [[PubMed](#)]
223. Lindahl, T. Recovery of antediluvian DNA. *Nature* **1993**, *365*, 700. [[CrossRef](#)] [[PubMed](#)]
224. Handt, O.; Höss, M.; Krings, M.; Pääbo, S. Ancient DNA: Methodological challenges. *Experientia* **1994**, *50*, 524–529. [[CrossRef](#)] [[PubMed](#)]
225. Hofreiter, M.; Serre, D.; Poinar, H.N.; Kuch, M.; Pääbo, S. Ancient DNA. *Nat. Rev. Genet.* **2001**, *2*, 353–359. [[CrossRef](#)] [[PubMed](#)]
226. Knapp, M.; Lalueza-Fox, C.; Hofreiter, M. Re-inventing ancient human DNA. *Investig. Genet.* **2015**, *6*, 4. [[CrossRef](#)] [[PubMed](#)]
227. Gilbert, M.T.P.; Bandelt, H.-J.; Hofreiter, M.; Barnes, I. Assessing ancient DNA studies. *Trends Ecol. Evol.* **2005**, *20*, 541–544. [[CrossRef](#)] [[PubMed](#)]
228. Poinar, H.N. The top 10 list: Criteria of authenticity for DNA from ancient and forensic samples. *Int. Congr. Ser.* **2003**, *1239*, 575–579. [[CrossRef](#)]
229. Brotherton, P.; Endicott, P.; Sanchez, J.J.; Beaumont, M.; Barnett, R.; Austin, J.; Cooper, A. Novel high-resolution characterization of ancient DNA reveals C > U-type base modification events as the sole cause of post mortem miscoding lesions. *Nucleic Acids Res.* **2007**, *35*, 5717–5728. [[CrossRef](#)] [[PubMed](#)]

230. Ginolhac, A.; Rasmussen, M.; Gilbert, M.T.P.; Willerslev, E.; Orlando, L. Mapdamage: Testing for damage patterns in ancient DNA sequences. *Bioinformatics* **2011**, *27*, 2153–2155. [[CrossRef](#)] [[PubMed](#)]
231. Jónsson, H.; Ginolhac, A.; Schubert, M.; Johnson, P.L.F.; Orlando, L. Mapdamage2.0: Fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* **2013**, *29*, 1682–1684. [[CrossRef](#)] [[PubMed](#)]
232. Malaspinas, A.-S.; Tange, O.; Moreno-Mayar, J.V.; Rasmussen, M.; DeGiorgio, M.; Wang, Y.; Valdiosera, C.E.; Politis, G.; Willerslev, E.; Nielsen, R. Bammds: A tool for assessing the ancestry of low-depth whole-genome data using multidimensional scaling (MDS). *Bioinformatics* **2014**, *30*, 2962–2964. [[CrossRef](#)] [[PubMed](#)]
233. Altschul, S.F.; Gish, W.; Miller, W.; Myers, E.W.; Lipman, D.J. Basic local alignment search tool. *J. Mol. Biol.* **1990**, *215*, 403–410. [[CrossRef](#)]
234. Shapiro, B.; Hofreiter, M. Analysis of ancient human genomes. *Bioessays* **2010**, *32*, 388–391. [[CrossRef](#)] [[PubMed](#)]
235. Pop, M.; Salzberg, S.L.; Shumway, M. Genome sequence assembly: Algorithms and issues. *Computer* **2002**, *35*, 47–54. [[CrossRef](#)]
236. Schmieder, R.; Edwards, R. Fast identification and removal of sequence contamination from genomic and metagenomic datasets. *PLoS ONE* **2011**, *6*, e17288. [[CrossRef](#)] [[PubMed](#)]
237. Longo, M.S.; O'Neill, M.J.; O'Neill, R.J. Abundant human DNA contamination identified in non-primate genome databases. *PLoS ONE* **2011**, *6*, e16410. [[CrossRef](#)] [[PubMed](#)]
238. Merchant, S.; Wood, D.E.; Salzberg, S.L. Unexpected cross-species contamination in genome sequencing projects. *PeerJ* **2014**, *2*, e675. [[CrossRef](#)] [[PubMed](#)]
239. Laurence, M.; Hatzis, C.; Brash, D.E. Common contaminants in next-generation sequencing that hinder discovery of low-abundance microbes. *PLoS ONE* **2014**, *9*, e97876. [[CrossRef](#)] [[PubMed](#)]
240. Warinner, C.; Herbig, A.; Mann, A.; Fellows Yates, J.A.; Weiß, C.L.; Burbano, H.A.; Orlando, L.; Krause, J. A Robust Framework for Microbial Archaeology. *Annu. Rev. Genom. Hum. Genet.* **2017**. [[CrossRef](#)] [[PubMed](#)]
241. Meyer, M.; Kircher, M. Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harb. Protoc.* **2010**, *6*. [[CrossRef](#)] [[PubMed](#)]
242. Gamba, C.; Hanghøj, K.; Gaunitz, C.; Alfarhan, A.H.; Alquraishi, S.A.; Al-Rasheid, K.A.S.; Bradley, D.G.; Orlando, L. Comparing the performance of three ancient DNA extraction methods for high-throughput sequencing. *Mol. Ecol. Resour.* **2016**, *16*, 459–469. [[CrossRef](#)] [[PubMed](#)]
243. Giles, R.J.; Brown, T.A. Improved methodology for extraction and amplification of DNA from single grains of charred wheat. *J. Archaeol. Sci.* **2008**, *35*, 2585–2588. [[CrossRef](#)]
244. Kistler, L. Ancient DNA extraction from plants. *Methods Mol. Biol.* **2012**, *840*, 71–79. [[PubMed](#)]
245. Telle, S.; Thines, M. Amplification of *cox2* (approximately 620 bp) from 2 mg of up to 129 years old herbarium specimens, comparing 19 extraction methods and 15 polymerases. *PLoS ONE* **2008**, *3*, e3584. [[CrossRef](#)] [[PubMed](#)]
246. Coombs, N.J.; Gough, A.C.; Primrose, J.N. Optimisation of DNA and RNA extraction from archival formalin-fixed tissue. *Nucleic Acids Res.* **1999**, *27*, e12. [[CrossRef](#)] [[PubMed](#)]
247. Mackie, M.E.; Radini, A.; Speller, C. The Sustainability of Dental Calculus for Archaeological Research. In *Shallow Pasts, Endless Horizons: Sustainability & Archaeology. Proceedings of the 48th Annual Chacmool Conference*; Favreau, J., Patalano, R., Eds.; The Chacmool Archaeological Association of the University of Calgary: Calgary, AB, Canada, 2017; pp. 74–81.
248. O'Donoghue, K.; Clapham, A.; Evershed, R.P.; Brown, T.A. Remarkable preservation of biomolecules in ancient radish seeds. *Proc. Biol. Sci.* **1996**, *263*, 541–547. [[CrossRef](#)] [[PubMed](#)]
249. Kraková, L.; Šoltys, K.; Otlewska, A.; Pietrzak, K.; Purkrťová, S.; Savická, D.; Puškárová, A.; Bučková, M.; Szemes, T.; Budiš, J.; et al. Comparison of methods for identification of microbial communities in book collections: Culture-dependent (sequencing and MALDI-TOF MS) and culture-independent (Illumina MiSeq). *Int. Biodeterior. Biodegrad.* **2017**. [[CrossRef](#)]
250. Wandeler, P.; Hoeck, P.E. A.; Keller, L.F. Back to the future: Museum specimens in population genetics. *Trends Ecol. Evol.* **2007**, *22*, 634–642. [[CrossRef](#)] [[PubMed](#)]
251. Bi, K.; Linderoth, T.; Vanderpool, D.; Good, J.M.; Nielsen, R.; Moritz, C. Unlocking the vault: Next-generation museum population genomics. *Mol. Ecol.* **2013**, *22*, 6018–6032. [[CrossRef](#)] [[PubMed](#)]
252. Kemp, C. Museums: The endangered dead. *Nature* **2015**, *518*, 292–294. [[CrossRef](#)] [[PubMed](#)]

253. Van Doorn, N.L.; Hollund, H.; Collins, M.J. A Novel and Non-Destructive Approach for Zooms Analysis: Ammonium Bicarbonate Buffer Extraction. *Archaeol. Anthropol. Sci.* **2011**, *3*, 281–289. [[CrossRef](#)]
254. Gomes, C.; Palomo-Díez, S.; Roig, J.; López-Parra, A.M.; Baeza-Richer, C.; Esparza-Arroyo, A.; Gibaja, J.; Arroyo-Pardo, E. Nondestructive extraction DNA method from bones or teeth, true or false? *Forensic Sci. Int. Genet.* **2015**, *5*, e279–e282. [[CrossRef](#)]
255. Andersen, J.C.; Mills, N.J. DNA extraction from museum specimens of parasitic Hymenoptera. *PLoS ONE* **2012**, *7*, e45549. [[CrossRef](#)] [[PubMed](#)]
256. Gilbert, M.T.P.; Moore, W.; Melchior, L.; Worobey, M. DNA extraction from dry museum beetles without conferring external morphological damage. *PLoS ONE* **2007**, *2*, e272. [[CrossRef](#)] [[PubMed](#)]
257. Fiddyment, S.; Holsinger, B.; Ruzzier, C.; Devine, A.; Binois, A.; Albarella, U.; Fischer, R.; Nichols, E.; Curtis, A.; Cheese, E.; et al. Animal origin of 13th-century uterine vellum revealed using noninvasive peptide fingerprinting. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 15066–15071. [[CrossRef](#)] [[PubMed](#)]
258. Gansauge, M.-T.; Gerber, T.; Glocke, I.; Korlević, P.; Lippik, L.; Nagel, S.; Riehl, L.M.; Schmidt, A.; Meyer, M. Single-stranded DNA library preparation from highly degraded DNA using T4 DNA ligase. *Nucleic Acids Res.* **2017**, *45*, e79. [[CrossRef](#)] [[PubMed](#)]
259. Stiller, M.; Sucker, A.; Griewank, K.; Aust, D.; Baretton, G.B.; Schadendorf, D.; Horn, S. Single-strand DNA library preparation improves sequencing of formalin-fixed and paraffin-embedded (FFPE) cancer DNA. *Oncotarget* **2016**, *7*, 59115–59128. [[CrossRef](#)] [[PubMed](#)]
260. Meyer, M.; Fu, Q.; Aximu-Petri, A.; Glocke, I.; Nickel, B.; Arsuaga, J.-L.; Martínez, I.; Gracia, A.; de Castro, J.M.B.; Carbonell, E.; et al. A mitochondrial genome sequence of a hominin from Sima de los Huesos. *Nature* **2014**, *505*, 403–406. [[CrossRef](#)] [[PubMed](#)]
261. Berger, L.R.; Hawks, J.; de Ruiter, D.J.; Churchill, S.E.; Schmid, P.; Deleuzene, L.K.; Kivell, T.L.; Garvin, H.M.; Williams, S.A.; DeSilva, J.M.; et al. Homo naledi, a new species of the genus Homo from the Dinaledi Chamber, South Africa. *eLife* **2015**, *4*, e09560. [[CrossRef](#)] [[PubMed](#)]
262. Sender, R.; Fuchs, S.; Milo, R. Revised Estimates for the Number of Human and Bacteria Cells in the Body. *PLoS Biol.* **2016**, *14*, e1002533. [[CrossRef](#)] [[PubMed](#)]
263. Davies, J. In a map for human life, count the microbes, too. *Science* **2001**, *291*, 2316. [[CrossRef](#)] [[PubMed](#)]
264. Schnorr, S.L.; Sankaranarayanan, K.; Lewis, C.M., Jr.; Warinner, C. Insights into human evolution from ancient and contemporary microbiome studies. *Curr. Opin. Genet. Dev.* **2016**, *41*, 14–26. [[CrossRef](#)] [[PubMed](#)]
265. Cook, M.I.; Beissinger, S.R.; Toranzos, G.A. Incubation reduces microbial growth on eggshells and the opportunity for trans-shell infection. *Ecology* **2005**, *8*, 532–537. [[CrossRef](#)] [[PubMed](#)]
266. Piñar, G.; Sterflinger, K.; Pinzari, F. Unmasking the measles-like parchment discoloration: Molecular and microanalytical approach. *Environ. Microbiol.* **2015**, *17*, 427–443. [[CrossRef](#)] [[PubMed](#)]
267. Fortes, G.G.; Speller, C.F.; Hofreiter, M.; King, T.E. Phenotypes from ancient DNA: Approaches, insights and prospects. *Bioessays* **2013**, *35*, 690–695. [[CrossRef](#)] [[PubMed](#)]
268. Llamas, B.; Holland, M.L.; Chen, K.; Cropley, J.E.; Cooper, A.; Suter, C.M. High-resolution analysis of cytosine methylation in ancient DNA. *PLoS ONE* **2012**, *7*, e30226. [[CrossRef](#)] [[PubMed](#)]
269. Smith, R.W.A.; Monroe, C.; Bolnick, D.A. Detection of Cytosine methylation in ancient DNA from five native american populations using bisulfite sequencing. *PLoS ONE* **2015**, *10*, e0125344. [[CrossRef](#)] [[PubMed](#)]
270. Gokhman, D.; Lavi, E.; Prüfer, K.; Fraga, M.F.; Riancho, J.A.; Kelso, J.; Pääbo, S.; Meshorer, E.; Carmel, L. Reconstructing the DNA methylation maps of the Neandertal and the Denisovan. *Science* **2014**, *344*, 523–527. [[CrossRef](#)] [[PubMed](#)]
271. Pedersen, J.S.; Valen, E.; Velazquez, A.M.V.; Parker, B.J.; Rasmussen, M.; Lindgreen, S.; Lilje, B.; Tobin, D.J.; Kelly, T.K.; Vang, S.; et al. Genome-wide nucleosome map and cytosine methylation levels of an ancient human genome. *Genome Res.* **2014**, *24*, 454–466. [[CrossRef](#)] [[PubMed](#)]
272. Smith, O.; Clapham, A.J.; Rose, P.; Liu, Y.; Wang, J.; Allaby, R.G. Genomic methylation patterns in archaeological barley show de-methylation as a time-dependent diagenetic process. *Sci. Rep.* **2014**, *4*, 5559. [[CrossRef](#)] [[PubMed](#)]
273. Orlando, L.; Willerslev, E. An epigenetic window into the past? *Science* **2014**, *345*, 511–512. [[CrossRef](#)] [[PubMed](#)]

