



UNIVERSITY OF LEEDS

This is a repository copy of *Cost-effectiveness of microscopy of urethral smears for asymptomatic Mycoplasma genitalium urethritis in men in England*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/119201/>

Version: Accepted Version

---

**Article:**

Sutton, AJ [orcid.org/0000-0001-7008-4770](http://orcid.org/0000-0001-7008-4770), Roberts, TE, Jackson, L et al. (4 more authors) (2018) Cost-effectiveness of microscopy of urethral smears for asymptomatic Mycoplasma genitalium urethritis in men in England. *International Journal of STD & AIDS*, 29 (1). pp. 72-79. ISSN 0956-4624

<https://doi.org/10.1177/0956462417717651>

---

(c) 2017, Author(s). This is an author produced version of a paper published in *International Journal of STD & AIDS*. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## Introduction

Over the past decade, there has been a change in the clinical investigation and management of men attending sexual health services in the UK. Previously, all men, regardless of symptoms, underwent urethral smears, a process by which a sample is taken from inside the urethra and Gram stained for examination by light microscopy (1). This allowed for the immediate diagnosis of two conditions: presumptive gonorrhoea and non-gonococcal urethritis (inflammation of the urethra in the absence of gonorrhoea). Men with either of these conditions, and their sexual partners, were then offered immediate treatment with appropriate antibiotics whilst waiting several days for more definitive results.

With the widespread use of sensitive and specific, non-invasive urine testing for chlamydia and gonorrhoea, and growing financial pressures on services, guidelines now recommend only performing urethral microscopy in symptomatic men (1). A consequence of this change in practice is that asymptomatic men with urethritis, caused by neither chlamydia nor gonorrhoea (known as non-chlamydial, non-gonococcal urethritis or NCNGU), no longer receive empirical antimicrobial therapy. Their sexual partners are also left untreated. However, at the time of the most recent national audit (1), a small number of clinics continued to provide routine urethral microscopy to asymptomatic men, contrary to the guidelines.

The potential impact of this change in practice on costs and patients outcomes is not clear and has not yet been explored in any depth. Asymptomatic urethritis has many causes, both infectious and non-infectious (1). Notably, *Mycoplasma genitalium* is present in 8-10% of men with asymptomatic urethritis (1) and is associated with both cervicitis and pelvic inflammatory disease in women (2). There is limited access to testing for *M. genitalium* in the UK and few men are tested for this organism. Therefore, whereas previously, men with asymptomatic urethritis secondary to *M. genitalium* and their partners may have received successful treatment as part of empirical therapy for urethritis, this is no longer the case.

The focus of this study is on the potential cost implications of this change in clinical practice assuming that some men with asymptomatic NCNGU have *M. genitalium*, which can have adverse and costly reproductive health outcomes in their female sexual partners. Specifically, the objective of this economic evaluation is to determine whether the screening landscape at the time of the last national audit, in which a small number of clinics continued to perform routine microscopy in asymptomatic men is a cost-effective approach to diagnosing and

treating asymptomatic NCNGU compared to the national guideline recommending not performing microscopy for this patient group.

## Methods

In order to estimate the impact of testing and treatment on the future transmission of possible significant pathogens responsible for asymptomatic NCNGU it is necessary to use an appropriate modelling approach for infectious diseases which can describe the transmission of M genitalium between individuals, namely a transmission dynamic model (TDM) (3, 4). In this study a TDM describing the transmission of M genitalium in the population of 16-30 year olds in England was constructed in order to examine changes in the use of urethral microscopy in asymptomatic men in GUM clinics. This economic evaluation uses outputs from this model, along with secondary data describing resource use and takes the form of a cost-effectiveness analysis carried out from a health care provider perspective, with costs valued at 2014/2015 UK prices.

### Model structure

The output used in this economic analysis is taken from a TDM which has been described in full elsewhere (5). In brief, this is a compartmental transmission model of the natural history of M genitalium, its diagnosis, and treatment levels, and thus only M genitalium was considered in this cost-effectiveness analysis. Heterogeneous sexual behaviour is described in the model which was parameterised by behaviour data from a number of key UK surveys, national surveillance data, and with the natural history of NCNGU being informed from data in the literature. The model describes the incidence and prevalence of symptomatic and asymptomatic infection, PID, care-seeking behaviour due to symptoms, partner notification, and the possibility of treatment failure. The uncertainty of the parameters in the model was also factored into the model parameterisation.


The time horizon for the economic analysis is 20 years, although this is subject to sensitivity analysis. It was felt that a time horizon longer than this would not be appropriate due to the inevitable changes to testing technology and approaches to offering STI screening to the population in the future. A discount rate of 3.5% was applied to costs and outcomes in accordance with NICE guidelines (6).

All settings where sexual health services are provided were initially considered for inclusion in this analysis. However, guidelines detailing the specific pathways and resources used at different sexual health service settings were sparse with the most reliable clinical data and cost data found in the literature being related to GP and GUM settings, with GP consultations being

considered due to the possibility of referral onwards to GUM services for further management. In this study the methodological focus is narrowed to the diagnosis and treatment of NCNGU in general practice and GUM clinics.

#### Testing Pathways for Economic analysis

Three different pathways are compared in terms of their resource use and costs, each representing alternative approaches to the testing and treatment of patients with asymptomatic NCNGU. These pathways represent: 1) the current recommended practice of not offering microscopy to asymptomatic men in GUM settings; 2) offering a small proportion (5%) of asymptomatic men microscopy (i.e. men attending a small number of GUM services) ; and 3) offering microscopy to all asymptomatic men attending all GUM services. These three pathways are referred to in this study as 'Current Recommended Practice', '5% Microscopy', and '100% Microscopy'.



**Figure 1:** Flow diagram showing the test and treatment pathway

Note: NAAT-nucleic acid amplification test for Chlamydia trachomatis & Neisseria gonorrhoeae

Initially, a patient can be either infected or non-infected with *M. genitalium* and either symptomatic or asymptomatic. The patient may attend either a general practice (GP) or a GUM clinic for testing. For those patients that attend a GP setting, all patients (asymptomatic and symptomatic) are tested for chlamydia and gonorrhoea using nucleic acid amplification test (NAAT) but none are offered microscopy. From the GP setting a proportion of the patients are then referred to a GUM clinic for further investigation and management, for example those who are symptomatic or have more complex sexual health needs.

In contrast, in the GUM setting the diagnostic pathway varies depending on which strategy is being considered and whether the patient presents with symptoms or not. For the 'Current Recommended Practice' strategy, microscopy is not offered to asymptomatic patients and these patients receive a NAAT test for chlamydia and gonorrhoea only. All asymptomatic patients in all locations in the 'Current Recommended Practice' scenario receive a NAAT chlamydia and gonorrhoea only. In the 5% Microscopy and '100% Microscopy' scenarios, 5% and 100% of male patients respectively at GUM clinics receive urethral smear microscopy. During the course of the consultation all symptomatic patients in a GUM setting receive partner notification and condoms with the aim of identifying individuals for whom testing and treatment may be appropriate. The testing pathways considered in this study are shown in Figure 1.

In this analysis, treatment can be deemed either a success or failure. Successful treatment indicates that a patient is no longer infected with *M. genitalium* and cannot transmit infection to their sexual partners. Treatment failure indicates that there has been a failure of the drug treatment to clear *M. genitalium*. Female patients who fail treatment or who are not treated can develop PID, a proportion of which cases are treated. Untreated PID cases may go on to experience symptomatic PID, infertility, or experience an ectopic pregnancy.

#### Model assumptions and parameterisation

This cost-effectiveness analysis was parameterised through secondary sources which are described below. It was necessary to make some pragmatic clarifying assumptions in order to carry out the analysis, these are described in the Appendix.

The model parameters used in this analysis are shown in Table 1.

Parameter	Value (range)	Reference
Proportion of times HIV test delivered alongside a NAAT test in a GUM setting	83% (range=0.71-0.97)	(7)
Proportion of times syphilis tests delivered alongside a NAAT test in a GUM setting	84% (range=0.72-0.97)	(7)
Proportion PID cases that give rise to ectopic pregnancy	(99/1309) 7.6% (6.4-8.8%)	(8, 9) Based on number trying to conceive after laparoscopy diagnosed PID case. Range calculated from a beta distribution taking values at 5% and 95% parameterised using method of moments (10)
Proportion PID cases that give rise to infertility	18% (15-21%)	(8, 11) range calculated from a beta distribution taking values at 5% and 95% assuming standard error = mean/10 (10)
Proportion of PID cases that are symptomatic	56% (30%-89%)	Value here from Posterior-mean of infectious disease model
Delay from PID to infertility / ectopic pregnancy manifest	5 years (1-15years)	Expert opinion – study team

**Table 1:** Model parameters used in economic evaluation

### Resource use and costs

The cost of partner notification was adjusted to 2014/15 prices using the pay and price index for Hospital & Community Health Services. Unit staff costs were obtained from Unit Costs of Health & Social Care (2015) (12). The unit costs of each resource used in this economic evaluation are described in the Appendix.



## Outcomes

The main outcome measure for this evaluation is the additional cost incurred per case of PID averted. The second outcome measure is the additional cost incurred per major outcome averted (MOA), where a major outcome is defined as a case of symptomatic PID, case of ectopic pregnancy, or a case of infertility. All major outcomes are reported for completeness. The results presented here use the incremental cost-effectiveness ratio (ICER), which is the difference in costs between two options divided by the difference in their effects (which are the outcome measures described above).

## Analysis

The base case scenario uses the mean results of 215 parameter sets from the dynamic transmission model and applies resource costs to obtain the baseline deterministic results for each of the three testing scenarios. These deterministic results from the TDM are shown in the Appendix along with details of the sensitivity analysis.

## Results

All results presented here are shown for a time horizon of 20 years with discounting unless otherwise stated. In all cases the costs are presented to the nearest thousand, and the outcomes to the nearest hundred. ICER values were calculated using the unrounded cost and outcome values with these then being rounded to the nearest 100.

	Cost	Cases of PID	Major outcomes*	Symptomatic PID	Infertility	Ectopic Pregnancy
No Microscopy	£1,244,736,000	111,800	37,600	23,300	10,000	4,200
5% Microscopy	£1,249,986,000	111,500	37,500	23,200	10,000	4,200
100% Microscopy	£1,350,369,000	105,300	35,600	21,800	9,700	4,100

**Table 2:** Baseline results for the three strategies for cases of PID and all the major outcomes considered in this study

\*where major outcomes are symptomatic PID, infertility or ectopic pregnancy

## Outcomes

As shown in Table 2, providing microscopy to 5% of asymptomatic men in a GUM setting has a positive impact on cases of PID. That is, the number of PID cases is lower for 5% Microscopy compared to No Microscopy. Likewise 5% Microscopy coverage also has a positive impact on reducing the number of major outcomes. In the case of the 100% Microscopy scenario, this has

a greater impact on reducing cases of PID and major outcomes compared to either 5% or No Microscopy.

#### Costs

When only considering costs, it can be seen that the cost of 5% Microscopy coverage is greater than No Microscopy, while 100% Microscopy coverage is the most costly approach. This indicates that any savings that might have been made as a result of a reduction in major outcomes are insufficient to make 5% Microscopy or 100% Microscopy cost saving.

## Incremental Results

	<b>Discounted cost</b>	<b>Cases of PID</b>	<b>ICER (/PID averted)</b>	<b>Major Outcomes</b>	<b>ICER (/MOA)</b>
No Microscopy	£1,244,736,000	111,800		37,600	
5% Microscopy	£1,249,986,000	111,500	£15,700	37,500	£49,900
100% Microscopy	£1,350,369,000	105,300	£16,300	35,600	£51,900

**Table 3:** Incremental cost per case of PID averted and cost per major outcome averted

For the outcome of a case of PID averted the ICER values are shown in Table 3. It can be seen that 5% Microscopy is more effective than no microscopy and has an ICER of £15,700, meaning that an investment of £15,700 is required to avert one case of PID. For the outcome of MOA it can again be seen (Table 3) that 5% Microscopy is more effective than no microscopy, but in this case an investment of £49,900 is required to avert one major outcome. In the case of 100% Microscopy, an investment of £16,300 is required to avert one case of PID, and £51,900 to avert one major outcome compared to 5% Microscopy.

## Sensitivity Analysis

The results of the sensitivity analysis are described in the Appendix.

## **Discussion**

### **Principal Findings**

This economic evaluation utilized the output from a transmission dynamic model (TDM) to estimate whether providing limited microscopy coverage to asymptomatic men to test for NCNGU at a limited number of GUM services (as was the case at the time of the last national audit of practice (1)) is a cost-effective option compared to the recommended current practice of its complete withdrawal.

This economic analysis was based on a principal outcome of cases of PID averted, and a secondary outcome of major outcome averted (MOA) (symptomatic PID, infertility, or ectopic pregnancy). The results at baseline indicate that performing urethral smear microscopy for approximately 5% of asymptomatic men attending GUM has an incremental cost of £15,700 per case of PID averted compared to no microscopy, meaning that this strategy invests approximately £15,700 to avoid one additional case of PID compared to a strategy of no routine microscopy screening where only symptomatic men are tested. Similarly 5% Microscopy coverage requires approximately £49,900 to avert one major outcome compared to a strategy of no routine microscopy screening where only symptomatic men are tested. Hypothetically, if recommended current practice were expanded to performing urethral smear microscopy for 100% of asymptomatic men attending GUM then this would have an additional cost of £16,300 per additional case of PID averted, and an additional £51,900 to avert an additional case of MOA compared to 5% Microscopy. These results also help to show that while conducting microscopy for 5% of asymptomatic men at GUM locations will avert PID and other major outcomes, at a population level it costs more to undertake the microscopy and associated patient management than it does to manage the adverse effects of not preventing the sequelae in a limited number of patients.

Across all the sensitivity analysis undertaken, 5% microscopy coverage was never found to be cost saving but was always found to have a positive impact on reducing cases of PID and major adverse outcomes. Varying the outputs from the TDM provided a range of values for the outcomes in this study. For case of PID averted the ICER values ranged from £9,600-£39,100, while for case of MOA the ICER values ranged from £30,500-£124,400. By varying the time horizon of the analysis it was found that shorter time horizons made the intervention less cost-effective.

## **Strengths & Weaknesses of Study**

This study has utilised the output from a well parameterised dynamic model that describes the transmission of *M. genitalium* in the population of males and females in England aged 16-30 years old. Uncertainty in this model has been considered through the use of multiple parameter sets, while the results from this economic evaluation have been subject to extensive sensitivity analysis. Inevitably this has led to the range of plausible values being obtained from the economic model being quite wide, although this does help to give confidence to the validity of the conclusions that might be drawn from this model.

In this analysis only NCNGU due to *M. genitalium* has been considered in the analysis, and its scope has not been extended to other causes. There are some causes which are innocuous conditions that are not tested for, such as adenovirus and these do not cause reproductive sequelae in women. Consequently had these non-serious causes been taken into account, then it is very likely that the testing strategies would have been even less cost-effective than has been shown here.

A weakness of this study is the inability to conduct joint probabilistic sensitivity analysis (PSA) for both the economic parameters and the parameters utilized in the transmission dynamic model. Although it was possible to conduct PSA for just the economic parameters while maintaining that output from the TDM at constant values, the results describing the probability of a strategy being below a specific acceptable threshold would be meaningless.

## **Comparisons with existing studies**

To our knowledge this is the first economic analysis related to NCNGU in any setting, and thus comparisons with the results from similar economic studies are impossible.

## **Meaning of study**

It is suggested that UK decision makers are unlikely to fund an intervention if it has an ICER value of £30,000 / quality adjusted life year (QALY) or more (13). However, as this study analysed outcomes in terms of cases of PID and major outcomes averted, there are no accepted threshold values which can be used to assess whether providing limited microscopy coverage to asymptomatic men is acceptable or not. It is therefore necessary to link the results here to

the acceptance threshold values for the QALY in order to draw conclusions from this economic analysis.

Taking mean values from the transmission dynamic model, the ICER for a case of PID averted and MOA were £15,700 and £49,900 respectively for 5% microscopy compared to no microscopy. Using the outcome of case of PID averted as an example, and taking into consideration the maximum acceptance threshold of £30,000 / QALY used by the National Institute of Healthcare and Care Excellence (NICE) in England, for this ICER to be deemed cost-effective based on current accepted thresholds, a case of PID averted would have to result in a gain of 0.53 of a QALY. Alternatively, the implication is that having PID would have to be equivalent to losing more than 6 months of perfect health. Likewise for MOA this would have to be more than 1.67 QALYs, meaning that having a major outcome would have to be equivalent to losing more than 18 months of perfect health.

Current evidence suggests that these outcomes are not valued so extremely. Smith (14) in a primary study based on a time trade off approach, asked respondents with a previous history of PID to value alternative conditions. The mean valuations for long term health states associated with PID were: ectopic pregnancy 0.79 (SD=0.34); pelvic pain 0.69 (SD=0.37); Infertility 0.76 (SD=0.34). These values suggest that the mean QALY gain to avert a case of pelvic pain (the state with the reported greatest negative impact on QoL) that lasted one year would be 0.31 QALYs. However as noted above for the results described here, for 5% microscopy coverage to be cost effective, a MOA must lead to a gain of more than 1 QALY, suggesting that current practice is far from being cost-effective.

Given the comparisons described above, it can therefore be concluded that recommended practice of reserving urethral microscopy for symptomatic men is a cost-effective strategy and reintroducing ad-hoc testing for asymptomatic men in GUM locations is unlikely to be cost-effective. Considering the results at baseline in this study, if ad-hoc microscopy testing for asymptomatic men were reintroduced into GUM locations then this would lead to over £5,000,000 (discounted) in costs over a 20 year period, which could then be better spent expanding testing and treatment regimens for different diseases which are more cost-effective.

### **Unanswered questions and future research**

One of the major issues related to any testing and diagnosis strategy is the impact of the testing pathway on patients. Patients may suffer from anxiety while waiting for the result of a test, or

may incur societal costs as a result of having to take time off work to attend for testing. There are also issues specific to the context of sexually transmitted infections where patients may be worried about the stigma of attending for testing and the difficulties surrounding partner notification for NCNGU. In the testing and diagnosis context, future work should focus on these issues, in order to better quantify their impact on patients with the goal of including the impact of these issues in economic studies such as this in order to better describe the true impact of the complete testing pathway on patients.

### **Key Messages**

- Current clinical recommendations for the UK are that urethral microscopy should not be offered to asymptomatic men attending genitourinary medicine clinics for diagnosis of NCNGU
- Offering Microscopy at very low level of coverage where a small number of GUM services in England routinely offered asymptomatic men urethral microscopy for NCNGU is not cost-effective and wastes resources which could be put to better use elsewhere
- Complete withdrawal of microscopy testing for asymptomatic men in a GUM setting could save over £5,000,000 (discounted) over a 20 year period

## **Funding**

This paper presents independent research commissioned by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research scheme (RP-PG-0707-10208). In addition, PJW thanks the UK Medical Research Council for Centre funding (grant MR/K010174/1) and also thanks the UK National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Modelling Methodology at Imperial College London in partnership with Public Health England (PHE) for funding (grant HPRU-2012-10080). AJS was supported by the National Institute for Health Research Diagnostic Evidence Co-operative Leeds.

The views expressed in this paper are those of the authors and not necessarily those of the NHS, the NIHR, the Department of Health, or Public Health England. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.



## References

1. Horner P, Blee K, O'Mahony C, Muir P, Evans C, Radcliffe K, et al. 2015 UK National Guideline on the management of non-gonococcal urethritis. *International journal of STD & AIDS*. 2016;27(2):85-96.
2. Lis R, Rowhani-Rahbar A, Manhart LE. Mycoplasma genitalium infection and female reproductive tract disease: a meta-analysis. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2015;61(3):418-26.
3. Anderson RM, May R. *Infectious Diseases of Humans: Dynamics and Control*. Oxford: Oxford University Press; 1992 1992.
4. Roberts TE, Robinson S, Barton P, Bryan S, Low N. Screening for Chlamydia trachomatis: a systematic review of the economic evaluations and modelling. *Sex Transm Infect*. 2006;82(3):193-200.
5. Estcourt C, Sutcliffe L, Mercer CH, Copas A, Saunders JM, Fuller S, et al. The Ballseye Programme: Targeting men for better sexual health. 2014 2014. Report No.
6. NICE. Guide to the methods of technology appraisal. 2013 2013. Report No.
7. McClean H, Carne CA, Sullivan AK, Menon-Johansson A, Gokhale R, Sethi G, et al. National audit of asymptomatic screening in UK genitourinary medicine clinics: case-notes audit. *Int J STD AIDS*. 2010;21(7):506-11.
8. Westrom LV. Sexually transmitted diseases and infertility. *Sex Transm Dis*. 1994;21(2 Suppl):S32-S7.
9. Adams EJ, Turner KM, Edmunds WJ. The cost effectiveness of opportunistic chlamydia screening in England. *Sex Transm Infect*. 2007;83(4):267-74.
10. Briggs A, Claxton K, Sculpher M. *Decision Modelling for Health Economic Evaluation*. Oxford: Oxford University Press; 2008 2008.
11. Haggerty CL, Gottlieb SL, Taylor BD, Low N, Xu F, Ness RB. Risk of sequelae after Chlamydia trachomatis genital infection in women. *J Infect Dis*. 2010;201 Suppl 2:S134-S55.
12. Curtis L. Unit Costs of Health & Social Care. University of Kent: 2015 2014. Report No.
13. Guide to the methods of technology appraisal. 2008 6/2008. Report No.
14. Smith KJ, Tsevat J, Ness RB, Wiesenfeld HC, Roberts MS. Quality of life utilities for pelvic inflammatory disease health states. *Sex Transm Dis*. 2008;35(3):307-11.