



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/119360/>

Version: Accepted Version

---

**Article:**

Mackie, A (2017) Food: More than the sum of its parts. *Current Opinion in Food Science*, 16. pp. 120-124. ISSN: 2214-7993

<https://doi.org/10.1016/j.cofs.2017.07.004>

---

© 2017 Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## Accepted Manuscript

Title: Food: More than the sum of its parts

Author: Alan Mackie

PII: S2214-7993(17)30045-0

DOI: <http://dx.doi.org/doi:10.1016/j.cofs.2017.07.004>

Reference: COFS 252



To appear in:

Received date: 4-4-2017

Revised date: 7-7-2017

Accepted date: 12-7-2017

Please cite this article as: Mackie, A., Food: More than the sum of its parts, *COFS* (2017), <http://dx.doi.org/10.1016/j.cofs.2017.07.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Food: More than the sum of its parts

Alan Mackie

School of Food Science and Nutrition, University of Leeds, Leeds, LS2 9JT, UK

*Abstract*

The food we eat and enjoy is not just a collection of macro and micro nutrients, it has structure and texture that affect our response to it. This short review of recent research touches on many of the issues linking different types of food structure to the digestion and absorption of macronutrients. The behavior of both protein and lipid in the gastric compartment can be key to what gets emptied when and thus the kinetics of nutrient absorption, which can in turn influence risk factors for disease. Because of this there is increasing emphasis on understanding the role of gastric processing in digestion kinetics. The use of MRI has significantly improved our understanding of gastric behavior and offers new possibilities for controlling digestion kinetics. In addition, the role of fiber in the upper GI tract is also starting to be better understood. This will also lead to further opportunities to improve the health of the Western diet.

*Keywords*

Food structure; digestion; macronutrients, dietary fiber, health

### Introduction

In the study of nutrition, food is generally considered at the nutrient level, meaning that the impact of food structure is often ignored. This reductionist approach linking one nutrient to one health effect, may partly explain some of the discrepancies between a predicted health effect of a food based on its nutrient content and its actual health effect when consumed as a whole food or indeed as part of a diet [1]. Things are now starting to change and so it is worth looking at recent work highlighting the role of food structure in nutrient release and food related diseases. The fundamental dichotomy is that while human studies provide the only way to definitively show causal links between food and health, it can be very difficult to develop a detailed mechanistic understanding of the role of food structure from such studies. Because of this, the work described in this review will cover not just the results of human studies but the data coming from increasingly widely used simulations of digestion, such as that developed in the Infogest project [2] or more sophisticated models [3]. Increasingly, digestion of the same food systems is being investigated both *in vivo* [4] and *in vitro* [5]. It will become evident from what follows that an accurate simulation of the gastric phase of digestion is key to understanding digestion kinetics and the link to food structure as illustrated by Figure 1.




Figure 1 A schematic showing the aspects of digestion where food structure has a key role to play.

Food has complex structures both physically and nutritionally which will affect digestion and absorption and may lead to interactions within the food matrix both pre and post consumption that alter its bioactive properties. Such interactions can alter the extent and kinetics of nutrient absorption in ways that are not always predictable, at least not from the information on nutrition labels [6]. Evidence from recent studies has shown that the food matrix can modify the nutritional properties of a food [7]. For example, plant-based foods contain cellular structures that need to be degraded before the encapsulated nutrients and bioactive compounds can be released and absorbed. This may be achieved by processing

(industrial as well as cooking at home) and by oral processing such as mastication. Despite this, release of nutrients may be limited in some foods such as almonds [8] or apple [9]. Hence, for some nutrient dense foods, such as almonds, the food matrix attenuates postprandial lipaemia after consumption [8]. People spend the majority of the day in a postprandial state and postprandial lipaemia is acknowledged as an independent risk factor for cardiovascular disease (CVD) [10]. A slower release of nutrients from naturally encapsulated systems has also been shown to increase satiety after consumption [11]. This paradigm also applies to other foods and other ways in which food structure alters digestion and is the main point of discussion in the following review.

### *Macronutrients*

Although macronutrients are not eaten in isolation, most studies tend to concentrate on the outcomes associated with a particular nutrient. Thus, the approach taken here will be based on individual macronutrients.

### *Protein*

Protein is a key constituent of the diet and has been much studied, particularly in relation to muscle building. Muscle growth (hypertrophy) after resistance exercise is a fundamental adaptation to an increased resistive workload. However, muscle growth can only occur if there is net anabolism within the muscle. That is, muscle protein net balance (synthesis minus breakdown) is positive during the period in which hypertrophy occurs [12]. This is true not just for athletes but also for the elderly, where a study on men with a mean age of 72 [13] showed that fast-digesting soluble milk proteins improved postprandial muscle protein synthesis and shows that the kinetics of bioavailability of amino acids and especially essential amino acids is important for health. The high bioavailability of peptides and proteins can also be detrimental to health in the case of allergy [14] and much work has been undertaken on the ability of specific protein families to resist processing and subsequent digestion [15, 16]. This link will not be discussed further here as there are other reviews that cover this topic [14, 17].

It is clear that many different processes can have an effect on protein structure, altering properties including allergenicity [18]. A limiting factor for hydrolysis is the accessibility of the enzyme in question to the substrate. Proteolysis initially occurs in the gastric compartment where recent studies have shown that the digestion of the protein gels is influenced by the microstructure of food matrices, caused by immobilization of the substrate in the network, and the steric hindrance of pepsin diffusing into and peptides diffusing out of the gel [19]. Of course heating in the presence of polysaccharides that are not digested in the upper GI tract can also have a marked effect. In a study on the heat induced gelation of beta lactoglobulin-xanthan mixtures at 70-90 °C for 20 minutes, a considerable decrease in the rate of proteolysis was seen. The formation of a dense protein network created a fine pore structure which restricted pepsin access into the gel thereby slowing proteolysis. This is in contrast to thermally induced protein unfolding that may increase hydrolysis [20]. At the supermolecular level, proteins can be used to make a very wide range of different structures and dairy proteins in particular have been used to investigate this. By tuning the conditions under which dairy products are formed, the underlying structure can be altered. This has been shown to have a marked effect on simulated digestion [21, 22] with homogenization increasing protein hydrolysis in milk but solid matrices such as cheeses being more resistant to protein digestion than liquid or semi-solid matrices. Although in the latter study by Islam et

al., no relationship could be seen between disintegration kinetics and rheological properties of the cheese. These results suggest a significant influence of the meal microstructure (resulting from heat treatment) and macrostructure (resulting from gelation processes) on the different phases of protein digestion.

The effect of processing and structure on many other sources of protein has been also investigated, including meat [23] and plants [24]. For example, the cooking temperature of meat has been shown to inversely effect the digestibility, with well cooked meat less digestible than slightly cooked or raw [23, 25]. The extent of digestion was not significantly affected in either study confirming that meat is a consistently good source of amino acids. The results show that the structure of the meat is responsible for only limited changes digestion kinetics. This is in contrast to data on plant proteins such as those from beans that can be very slow to digest depending on the processing used [26].

### *Lipid*

The digestion and absorption of lipid is complicated by the poor solubility of all the components involved except the enzymes and the low density of triglycerides relative to the aqueous environment of the gut [27]. There has been a significant amount of research in this area and a number of recent reviews addressing the effect of structure [28, 29].

Because of the solubility issue with lipid and many bioactives, there has been a significant amount of research into how to optimize delivery of poorly soluble compounds using different structures [30-32]. The size of the lipid droplets and the available surface area is important for the rate of lipolysis, although overall fat absorption in healthy subjects is not affected by differences in initial droplet size because of efficient fat digestion by pancreatic lipase in the small intestine [33]. Despite this, it has been shown that the initial emulsion droplet size can have a significant effect of satiety [34]. This was most likely due to phase separation in the stomach. The behavior of emulsions under different physiological conditions is also important however [22] and numerous efforts have been made to control rates of lipid hydrolysis by changing gastric behavior [35] or tailoring the composition of the interface [36-38]. In particular coating the interface with polysaccharides such as pectin [39] or alginate [40] has been used, largely unsuccessfully to decrease rates of lipolysis. A far more effective way of altering the timing of the fat digestion is by inducing phase separation in the stomach, as demonstrated in studies using MRI [41]. Other factors that may also effect lipid digestion such as coalescence and droplet structure (e.g. emulsion containing solid fat) highlight that the principal cause of altered lipolysis is the destabilization of the emulsion within the stomach [42].

In addition to emulsified lipid, fat may be delivered in an encapsulated form in plant based foods [7, 43]. In such systems, oral processing is important to break open cellular structures [8] because the cell walls that are completely intact limit lipid digestibility, due to an encapsulation mechanism that hinders the diffusion of lipase into the intracellular environment and lipolysis products out of the cells.

### *Complex carbohydrates vs simple sugars*

Carbohydrates in the diet range from simple sugars such as glucose to highly complex polysaccharides but only monosaccharides are absorbable. The slow release and absorption

of sugars through inclusion of cereal fiber has been shown to be beneficial to health [44] by lowering risk factors for metabolic disease. As a result there is significant interest in altering rates of carbohydrate digestion and glycemic response in a predictable way [45]. Starch provides most of the carbohydrate in the human diet and its structure is a key driver of digestion kinetics [46, 47]. Raw granular starch is very resistant to amylolysis. In contrast, freshly gelatinized starches are highly susceptible to hydrolysis. Starch granule characteristics considered to influence susceptibility to attack by  $\alpha$ -amylase include: crystallinity, granule size, amylose to amylopectin ratio, etc. [48]. The retrogradation of the amylose and amylopectin in starch after cooling varies and in amylose is rapid, resulting in high resistance to amylolysis. In contrast, amylopectin is slow to retrograde and may still be degraded by amylase [49, 50]. In addition, the surrounding matrix can also have a significant effect. For example, the gluten proteins in pasta can interact with amylase and slow down starch hydrolysis [51].

By definition, dietary fiber does not provide a source of nutrients directly to the body. However, it is becoming increasingly apparent that it is important for health for multiple reasons. The ability of dietary fibers to alter digestion and metabolism depends upon their physical properties and there are clearly large differences between soluble and insoluble fiber. The benefits of fiber in the upper GI tract are thought to be a consequence of the increase in viscosity that can be induced by soluble fiber [52]. The viscosity that can be imparted by soluble fiber depends on a number of factors including molecular weight and the extent of hydration. In general, higher viscosity meals tend to cause increased gastric retention in comparison to the equivalent lower viscosity meal [53]. As confirmation of this, a recent *in vitro* study, the impact on digestion of the addition of oat bran to biscuits was assessed. In particular, the viscosity of the chyme was measured throughout digestion [54], showing that the viscosity was maintained throughout the gut up to the ileal compartment and that the highest viscosity was obtained with biscuits containing soluble fiber. The effect of viscosity on digestion was also determined and protein hydrolysis decreased with fiber enrichment, i.e. increased viscosity [55]. This increase in viscosity has a number of benefits from a health perspective. Firstly it decreases luminal mixing, which can decrease mass transfer in the gut [56]. This involves the decreased transport of hydrolysis products to the site of absorption and the changes in interactions between enzymes and substrate. It can also effect the composition of chyme that is emptied from the ileum into the cecum. For example decreasing bile recycling by taking more bile into the colon and thus lowering blood cholesterol as the liver uses some of the available cholesterol to replenish bile reserves. In addition to viscosity effects in the intestinal lumen and the consequent impact on gastrointestinal motility, mixing and rates of hydrolysis, the soluble fiber may interact with the intestinal mucus and alter its permeability [57] or secretion rate. Indeed there is some evidence that it can also have an effect of intestinal morphology, at least in the small intestine of pigs [58].

### Conclusions

It is clear that the health benefits of many foods are associated with their structure and the way that they deliver nutrients to the body. However, with a few exceptions, this association has not led to a clear mechanistic understanding of the role of the food structure. In the case of plant based foods, the different roles of cellular structure in combination with soluble and insoluble fiber fractions throughout the GI tract has yet to be elucidated. If we are to increase fiber intake and improve the health of the western diet then we need a better understanding

of the effect of structure on digestion and the evidence above strongly suggests that the gastric phase of digestion is the place to focus.

### Conflicts of interest

I declare that there are no conflicts of interest in the production of the article

### References

1. Fardet, A. and E. Rock, *The Healthy Core Metabolism: A New Paradigm for Primary Preventive Nutrition*. J Nutr Health Aging, 2016. **20**(3): p. 239-47.
- 2.\*\* Egger, L., et al., *The harmonized INFOGEST in vitro digestion method: From knowledge to action*. Food Research International, 2016. **88, Part B**: p. 217-225.  
This article demonstrates the reproducibility and robustness of the Infogest simulation of upper GI tract digestion for following the digestion of protein. This should facilitate comparison between different studies.
3. Bellmann, S., et al., *Development of an advanced in vitro model of the stomach and its evaluation versus human gastric physiology*. Food Research International, 2016. **88**: p. 191-198.
4. Mackie, A.R., et al., *Specific food structures suppress appetite through reduced gastric emptying rate*. American Journal of Physiology - Gut and Liver Physiology, 2013. **304**(11): p. G1038-G1043.
- 5.\*\* Mulet-Cabero, A.I., et al., *Dairy food structures influence the rates of nutrient digestion through different in vitro gastric behaviour*. Food Hydrocolloids, 2017. **67**: p. 63-73.  
This article shows that a relatively simple system can be used to simulate complex gastrointestinal behaviour. It also shows that changes in nutrient bioavailability can be strongly influenced by gastric behaviour.
- 6.\* Dhillion, J., et al., *Effects of food form on appetite and energy balance*. Food Quality and Preference, 2016. **48**: p. 368-375.  
Nice review of the issues around food structure and energy balance
7. Grundy, M.M., et al., *Re-evaluation of the mechanisms of dietary fibre and implications for macronutrient bioaccessibility, digestion and postprandial metabolism*. British Journal of Nutrition, 2016. **116**(5): p. 816-833.
- 8.\* Grundy, M.M.L., et al., *Effect of mastication on lipid bioaccessibility of almonds in a randomized human study and its implications for digestion kinetics, metabolizable energy, and postprandial lipemia*. American Journal of Clinical Nutrition, 2015. **101**(1): p. 25-33.  
This article shows very clearly the role of oral processing but also the fact that it can be insufficient to fully release nutrients from cellular structures
9. Dalmau, M.E., et al., *Effects of freezing, freeze drying and convective drying on in vitro gastric digestion of apples*. Food Chemistry, 2017. **215**: p. 7-16.
10. Jackson, K.G., S.D. Poppitt, and A.M. Minihane, *Postprandial lipemia and cardiovascular disease risk: Interrelationships between dietary, physiological and genetic determinants*. Atherosclerosis, 2012. **220**(1): p. 22-33.
11. Hull, S., et al., *A mid-morning snack of almonds generates satiety and appropriate adjustment of subsequent food intake in healthy women*. Eur J Nutr, 2015. **54**(5): p. 803-10.
12. Phillips, S.M., et al., *Mixed muscle protein synthesis and breakdown after resistance exercise in humans*. American Journal of Physiology-Endocrinology and Metabolism, 1997. **273**(1): p. E99-E107.

- 13.\* Walrand, S., et al., *Fast-digestive protein supplement for ten days overcomes muscle anabolic resistance in healthy elderly men*. *Clinical Nutrition*, 2016. **35**(3): p. 660-668.

In this study of 31 elderly men, fast-digesting soluble milk proteins improved postprandial muscle protein synthesis, highlighting the importance of digestion kinetics

14. Verhoeckx, K.C.M., et al., *Food processing and allergenicity*. *Food and Chemical Toxicology*, 2015. **80**: p. 223-240.
15. van Broekhoven, S., et al., *Influence of processing and in vitro digestion on the allergic cross-reactivity of three mealworm species*. *Food Chemistry*, 2016. **196**: p. 1075-1083.
16. Li, Y. and S. Damodaran, *In vitro digestibility and IgE reactivity of enzymatically cross-linked heterologous protein polymers*. *Food Chemistry*, 2017. **221**: p. 1151-1157.
17. Bogh, K.L. and C.B. Madsen, *Food Allergens: Is There A Correlation between Stability to Digestion and Allergenicity?* *Critical Reviews in Food Science and Nutrition*, 2016. **56**(9): p. 1545-1567.
18. Benede, S., et al., *Egg proteins as allergens and the effects of the food matrix and processing*. *Food & Function*, 2015. **6**(3): p. 694-713.
19. Luo, Q., R.M. Boom, and A.E.M. Janssen, *Digestion of protein and protein gels in simulated gastric environment*. *Lwt-Food Science and Technology*, 2015. **63**(1): p. 161-168.
20. Leeb, E., et al., *Influence of denaturation and aggregation of beta-lactoglobulin on its tryptic hydrolysis and the release of functional peptides*. *Food Chemistry*, 2015. **187**: p. 545-554.
21. Lamothe, S., et al., *Influence of dairy matrices on nutrient release in a simulated gastrointestinal environment*. *Food Research International*, 2017. **92**: p. 138-146.
- 22.\* Islam, M.A., et al., *Ex vivo digestion of raw, pasteurised and homogenised milk - Effects on lipolysis and proteolysis*. *International Dairy Journal*, 2017. **65**: p. 14-19.

This article highlights the impact of homogenisation on milk digestion using human aspirates. Homogenisation increased both gastric proteolysis and duodenal lipid hydrolysis.

23. Wen, S.Y., et al., *Effect of Cooking on in Vitro Digestion of Pork Proteins: A Peptidomic Perspective*. *Journal of Agricultural and Food Chemistry*, 2015. **63**(1): p. 250-261.
24. Chen, F.P., B.S. Li, and C.H. Tang, *Nanocomplexation between Curcumin and Soy Protein Isolate: Influence on Curcumin Stability/Bioaccessibility and in Vitro Protein Digestibility*. *Journal of Agricultural and Food Chemistry*, 2015. **63**(13): p. 3559-3569.
25. Sayd, T., C. Chambon, and V. Sante-Lhoutellier, *Quantification of peptides released during in vitro digestion of cooked meat*. *Food Chemistry*, 2016. **197**: p. 1311-1323.
26. Romano, A., *Impact of dehulling on the physico-chemical properties and in vitro protein digestion of common beans (*Phaseolus vulgaris* L.)*. *Food & Function*, 2015, **6**: p. 1345-1351.
27. Mao, L.K. and S. Miao, *Structuring Food Emulsions to Improve Nutrient Delivery During Digestion*. *Food Engineering Reviews*, 2015. **7**(4): p. 439-451.
28. Norton, J.E., et al., *Functional food microstructures for macronutrient release and delivery*. *Food & Function*, 2015. **6**(3): p. 663-678.
29. Singh, H., A.Q. Ye, and M.J. Ferrua, *Aspects of food structures in the digestive tract*. *Current Opinion in Food Science*, 2015. **3**: p. 85-93.
30. Mun, S., et al., *Control of lipid digestion and nutraceutical bioaccessibility using starch-based filled hydrogels: Influence of starch and surfactant type*. *Food Hydrocolloids*, 2015. **44**: p. 380-389.
31. Zou, L.Q., et al., *Utilizing Food Matrix Effects To Enhance Nutraceutical Bioavailability: Increase of Curcumin Bioaccessibility Using Excipient Emulsions*. *Journal of Agricultural and Food Chemistry*, 2015. **63**(7): p. 2052-2062.
32. Zou, L.Q., et al., *Designing excipient emulsions to increase nutraceutical bioavailability: emulsifier type influences curcumin stability and bioaccessibility by altering gastrointestinal fate*. *Food & Function*, 2015. **6**(8): p. 2475-2486.
33. Armand, M., et al., *Digestion and absorption of 2 fat emulsions with different droplet sizes in the human digestive tract*. *American Journal of Clinical Nutrition*, 1999. **70**(6): p. 1096-1106.

- 34.\*\* Lett, A.M., J.E. Norton, and M.R. Yeomans, *Emulsion oil droplet size significantly affects satiety: A pre-ingestive approach*. *Appetite*, 2016. **96**: p. 18-24.

This study of 34 men was able to show that very large 50 micron emulsion droplets in a preload meal had less effect on subsequent intake than small 2 micron droplets. This is likely to have been due to differences in gastric creaming and emptying.

35. Wooster, T.J., et al., *Impact of different biopolymer networks on the digestion of gastric structured emulsions*. *Food Hydrocolloids*, 2014. **36**: p. 102-114.
- 36.\* Sarkar, A., et al., *In vitro digestion of Pickering emulsions stabilized by soft whey protein microgel particles: influence of thermal treatment*. *Soft Matter*, 2016. **12**(15): p. 3558-3569.

This article highlights the difficulty of delaying lipolysis using protein particles. In the absence of protease a clear decrease in lipolysis was seen but the addition of pepsin removed the effect.

37. Shao, Y. and C.H. Tang, *Gel-like pea protein Pickering emulsions at pH 3.0 as a potential intestine-targeted and sustained-release delivery system for beta-carotene*. *Food Research International*, 2016. **79**: p. 64-72.
38. Pinheiro, A.C., M.A. Coimbra, and A.A. Vicente, *In vitro behaviour of curcumin nanoemulsions stabilized by biopolymer emulsifiers - Effect of interfacial composition*. *Food Hydrocolloids*, 2016. **52**: p. 460-467.
39. Zhao, J.J., et al., *Influence of soybean soluble polysaccharides and beet pectin on the physicochemical properties of lactoferrin-coated orange oil emulsion*. *Food Hydrocolloids*, 2015. **44**: p. 443-452.
40. Fioramonti, S.A., et al., *Multilayer emulsions as a strategy for linseed oil microencapsulation: Effect of pH and alginate concentration*. *Food Hydrocolloids*, 2015. **43**: p. 8-17.
- 41.\*\* Steingoetter, A., et al., *Imaging gastric structuring of lipid emulsions and its effect on gastrointestinal function: a randomized trial in healthy subjects*. *American Journal of Clinical Nutrition*, 2015. **101**(4): p. 714-724.

This excellent study in which participants were fed 4 different emulsions and then gastric behaviour was followed by MRI. The lipid phase behaviour in the stomach is very clearly seen and the outcomes confirm the link to plasma lipid concentrations and bile secretion (gall bladder volume)

42. Day, L., et al., *Tailoring the digestion of structured emulsions using mixed monoglyceride-caseinate interfaces*. *Food Hydrocolloids*, 2014. **36**: p. 151-161.
42. Grundy, M.M.L., et al., *The role of plant cell wall encapsulation and porosity in regulating lipolysis during the digestion of almond seeds*. *Food & Function*, 2016. **7**(1): p. 69-78.
44. Yamini, S. and P.R. Trumbo, *Qualified health claim for whole-grain intake and risk of type 2 diabetes: an evidence-based review by the US Food and Drug Administration*. *Nutrition Reviews*, 2016. **74**(10): p. 601-611.
45. Zeevi, D., et al., *Personalized Nutrition by Prediction of Glycemic Responses*. *Cell*, 2015. **163**(5): p. 1079-1094.
46. Qiao, D.L., et al., *A further understanding of the multi-scale supramolecular structure and digestion rate of waxy starch*. *Food Hydrocolloids*, 2017. **65**: p. 24-34.
47. Xu, J.C., et al., *Insights into molecular structure and digestion rate of oat starch*. *Food Chemistry*, 2017. **220**: p. 25-30.
48. Copeland, L., et al., *Form and functionality of starch*. *Food Hydrocolloids*, 2009. **23**(6): p. 1527-1534.
49. Ashwar, B.A., et al., *Production of resistant starch from rice by dual autoclaving-retrogradation treatment: Invitro digestibility, thermal and structural characterization*. *Food Hydrocolloids*, 2016. **56**: p. 108-117.
50. Hsu, R.J.C., et al., *Effects of added water and retrogradation on starch digestibility of cooked rice flours with different amylose content*. *Journal of Cereal Science*, 2015. **61**: p. 1-7.
51. Zou, W., et al., *Compact structure and proteins of pasta retard in vitro digestive evolution of branched starch molecular structure*. *Carbohydrate Polymers*, 2016. **152**: p. 441-449.

- 52.\*\* Zhang, D.G., et al., *Soluble arabinoxylan alters digesta flow and protein digestion of red meat-containing diets in pigs*. Nutrition, 2015. **31**(9): p. 1141-1147.

Although barbecued meat is not the normal diet of pigs, this study highlights the beneficial inclusion of soluble fibre in the diet. The potential health benefits are through decreased protein digestibility, increased digesta mass, and reduced mean retention time, even for diets with a moderately higher protein content.

53. Juvonen, K.R., et al., *Viscosity of Oat Bran-Enriched Beverages Influences Gastrointestinal Hormonal Responses in Healthy Humans*. Journal of Nutrition, 2009. **139**(3): p. 461-466.
54. Villemejeane, C., et al., *In vitro digestion of short-dough biscuits enriched in proteins and/or fibres, using a multi-compartmental and dynamic system (1): Viscosity measurement and prediction*. Food Chemistry, 2015. **182**: p. 55-63.
55. Villemejeane, C., et al., *In vitro digestion of short-dough biscuits enriched in proteins and/or fibres using a multi-compartmental and dynamic system (2): Protein and starch hydrolyses*. Food chemistry, 2016. **190**: p. 164-72.
56. Gidley, M.J., *Hydrocolloids in the digestive tract and related health implications*. Current Opinion in Colloid & Interface Science, 2013. **18**(4): p. 371-378.
- 57.\* Mackie, A.R., et al., *Sodium alginate decreases the permeability of intestinal mucus*. Food Hydrocolloids, 2015. **52**: p. 749-755.

This article introduces the idea that soluble fibre in the diet may decrease intestinal mucus permeability at least to lipid digestion products. This has clear potential health benefits.

- 58\*. Gunness, P., et al. *Reduction in circulating bile acid and restricted diffusion across the intestinal epithelium are associated with a decrease in blood cholesterol in the presence of oat beta-glucan*. FASEB J, 2016, **30**: p. 4227-4238.

Although this article does not really achieve the goal of elucidating how soluble dietary fibre lowers circulating cholesterol, it does show an interesting range of effects in pigs including changes in intestinal morphology and bile composition.

## Highlights

Food structure can affect the rate and extent of digestion

Digestion kinetics can affect disease risk factors

Methods such as MRI are helping to improve our understanding of gastric behaviour

The role of dietary fibre in the upper GI tract is starting to be better understood