



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/120822/>

Version: Accepted Version

---

**Article:**

Hobbs, C., Reid, J.D. and Shepherd, M. (2017) The coproporphyrin ferrochelatase of *Staphylococcus aureus* : mechanistic insights into a regulatory iron binding site.

Biochemical Journal , 474 (20). pp. 3513-3522. ISSN: 0264-6021

<https://doi.org/10.1042/BCJ20170362>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## The coproporphyrin ferrochelatase of *Staphylococcus aureus*: mechanistic insights into a regulatory iron binding site

Charlie Hobbs\*, James D. Reid<sup>†</sup> and Mark Shepherd\*<sup>1</sup>

\*School of Biosciences, University of Kent, Canterbury, CT2 7NJ, United Kingdom.

<sup>†</sup>Department of Chemistry, University of Sheffield, Sheffield, S3 7HF, United Kingdom.

<sup>1</sup>To whom correspondence should be addressed:

Mark Shepherd

[M.Shepherd@kent.ac.uk](mailto:M.Shepherd@kent.ac.uk)

School of Biosciences,

University of Kent,

Canterbury,

Kent CT2 7NJ,

United Kingdom.

Tel: +44 (0) 1227823988

**Page Heading Title:** Kinetic analysis of a staphylococcal ferrochelatase

### ABSTRACT

The majority of characterised ferrochelatase enzymes catalyse the final step of classical haem synthesis, inserting ferrous iron into protoporphyrin IX. However, for the recently-discovered coproporphyrin-dependent pathway, ferrochelatase catalyses the penultimate reaction where ferrous iron is inserted into coproporphyrin III. Ferrochelatase enzymes from the bacterial phyla Firmicutes and Actinobacteria have previously been shown to insert iron into coproporphyrin, and those from *Bacillus subtilis* and *Staphylococcus aureus* are known to be inhibited by elevated iron concentrations. The work herein reports a  $K_m$  (coproporphyrin III) for *S. aureus* ferrochelatase of 1.5  $\mu\text{M}$  and it is shown that elevating the iron concentration increases the  $K_m$  for coproporphyrin III, providing a potential explanation for the observed iron-mediated substrate inhibition. Together, structural modelling, site-directed mutagenesis, and kinetic analyses confirm residue Glu271 as being essential for the binding of iron to the inhibitory regulatory site on *S. aureus* ferrochelatase, providing a molecular explanation for the observed substrate inhibition patterns. This work therefore has implications for how haem biosynthesis in *S. aureus* is regulated by iron availability.

**KEYWORDS:** Coproporphyrinogen, ferrochelatase, haem biosynthesis

**SUMMARY STATEMENT:** This study provides new kinetic data for the recently-identified haem biosynthetic enzyme, coproporphyrin ferrochelatase, from *Staphylococcus aureus*. In addition, the location of an inhibitory metal binding site is identified for the first time, which has important implications for how haem synthesis in bacterial pathogens is regulated by iron.

### INTRODUCTION

Ferrochelatase enzymes catalyse the insertion of metal ions into porphyrin rings. The insertion of ferrous iron ( $\text{Fe}^{2+}$ ) into protoporphyrin IX is the final step in the well-characterised classical haem synthesis pathway. However, bacterial species in the Firmicutes and Actinobacteria Phyla (*Bacillus subtilis*, *Mycobacterium tuberculosis*, *Propionibacterium acnes*, *S. aureus*) have been shown to utilise a ‘coproporphyrin-dependent’ pathway [1, 2], where  $\text{Fe}^{2+}$  insertion into coproporphyrin III is the penultimate step (Fig. 1). Branching and regulation of these bacterial haem biosynthetic pathways are discussed in a recent review that suggests a series of new abbreviations based on enzyme activities [3]. These new abbreviations provide a useful way of discriminating between protoporphyrin ferrochelatase

(PpfC) and coproporphyrin ferrochelatase (CpfC), and this nomenclature is adopted herein in place of the traditional HemH abbreviation.

Ferrochelatases are membrane-bound proteins in eukaryotes and many prokaryotes, but Gram-positive bacteria have been shown to encode a soluble form of the protein [4-6]. All of these forms of ferrochelatase have been shown to insert a variety of divalent metals into the centre of porphyrin molecules, with zinc being the most common metal used in *in vitro* assays due to being more stable in an aerobic environment than the natural  $\text{Fe}^{2+}$  substrate [7] and allows for fluorescence based assays to be utilised. Catalytic rates for ferrochelatases vary with  $k_{\text{cat}}$  values typically ranging from  $0.11 \text{ min}^{-1}$  to  $15.3 \text{ min}^{-1}$ , although *S. aureus* CpfC was recently reported to have a  $k_{\text{cat}}$  of  $165 \text{ min}^{-1}$  using coproporphyrin III and  $\text{Fe}^{2+}$  as substrates [1]. Parallel experiments performed for the *B. subtilis* CpfC gave a  $k_{\text{cat}}$  of  $78 \text{ min}^{-1}$ , which compares to another report of  $0.11 \text{ min}^{-1}$  [2]. This recent work also identified  $0.8 \mu\text{M}$  as the threshold concentration of  $\text{Fe}^{2+}$  above which inhibition occurs for the *B. subtilis* and *S. aureus* enzymes, which represents an interesting feature unique to coproporphyrin-dependent ferrochelatases.

While bacterial and mammalian ferrochelatase enzymes generally exhibit poor sequence identity (e.g. <10% between Human and *B. subtilis*), the secondary and tertiary structures are similar in the core region [8]. However, human and yeast PpfC enzymes have an extended C-terminal region compared to *B. subtilis* CpfC. The Human enzyme ligates a [2Fe-2S] iron-sulphur cluster in this region, which is not present in the *S. cerevisiae* enzyme. This cluster is ligated by one residue in the core of the protein (Cys 196) and three in the extended C-terminal region (Cys 403, 406, 411), and at the time was thought to be responsible for anchoring the extended C-terminal to the rest of the monomer, and was also suggested to be indirectly involved in stabilising the dimer form of the protein [8]. *S. cerevisiae* does not contain the necessary residues in the same location to form a cluster and the dimer stability is reportedly not affected [9]. Some ferrochelatases from Gram-positive bacteria have been shown to ligate [2Fe-2S] clusters via an extra insertion, although loss of this cluster did not result in a total loss of activity [10] in contrast to the [2Fe-2S] cluster of the human enzyme [11]. The *in vivo* role for these cofactors remains poorly understood.

Ferrochelatase structures exhibit a degree of variability with respect to the residues present in the substrate binding cleft. For eukaryotic PpfC enzymes a series of leucine residues form part of the hydrophobic lip that covers the active site pocket, which is not present in the *B. subtilis* enzyme. The Tyr13 residue in *B. subtilis* CpfC, which has been determined to aid the conserved His183 residue in stabilisation of the B ring of the porphyrin macrocycle, is not present in the membrane-associated ferrochelatases [12]. The residue in this position in the other enzymes is a methionine whose role is not entirely understood [8, 9]. The tyrosine residue in *B. subtilis* has been implicated in a metal-binding function in the catalytic metal binding site along with the conserved histidine and glutamic acid residues [12, 13] and it is possible that the methionine residue in this position aids in the insertion of metals into the porphyrin ring in a similar manner to the tyrosine residue found in the soluble form of ferrochelatase. The amino acid at this position has also been linked to the specificity of metal chelation, with a Y13M mutation of the *B. subtilis* enzyme resulting in a more rapid insertion of  $\text{Co}^{2+}$  and a complete loss of  $\text{Cu}^{2+}$  utilisation [14].

Soluble ferrochelatases have been crystallised with a variety of metals present in the metal binding cleft, and have been shown to have two separate sites for metal binding: one close to the porphyrin binding site stabilised by the conserved His183 and one closer to the surface of the protein approximately  $7\text{\AA}$  from the H183 residue [12]. In *B. subtilis* CpfC, this outer site has been shown to bind  $\text{Mg}^{2+}$  ions, which are co-ordinated by the residues Arg46, Glu268 and Glu272 [12]. This particular metal site has been reported to provide a  $\text{Mg}^{2+}$ -mediated stimulation of zinc insertion into deuteroporphyrin IX, with mutation of the Glu272 residue resulting in abolition of the stimulatory effects of  $\text{Mg}^{2+}$  [13]. The mechanism for this stimulation was proposed to be due to repulsion between metal ions present at both sites pushing the inserted metal at the inner (catalytic) binding site closer to the porphyrin ring [13]. Such a regulatory metal site has not been investigated in the human enzyme, and it seems unlikely that *S. cerevisiae* PpfC could bind metals at this site due to the presence of a

glycine residue instead of a glutamate residue at position 322 [9]. However, it has been suggested that *S. cerevisiae* PpfC can bind the inhibitor  $\text{Cd}^{2+}$  at two sites, with one site slightly further away from the active site than the outer metal binding site for *B. subtilis* CpfC. Human PpfC is susceptible to substrate inhibition when zinc is the metal ion substrate. This zinc inhibition is more pronounced at higher concentrations of the porphyrin substrate, suggesting that an inhibitory metal ion binds to the enzyme-product complex [15]. Residues that are predicted to be predominantly involved in metal binding in *B. subtilis*, where  $\text{Zn}^{2+}$  and  $\text{Cd}^{2+}$  have been observed, are the conserved residues H183 and E264 and the unique Y13 residue that resides on the other side of the porphyrin binding site [13].  $\text{Cd}^{2+}$  has been shown to also bind to the nearby H262, which is located underneath the porphyrin substrate and is proposed to inhibit CpfC by hampering the binding of porphyrin substrate [13].

The current work investigates the kinetic properties of CpfC from the human pathogen *S. aureus* using the native substrates  $\text{Fe}^{2+}$  and coproporphyrin III. Since iron has dramatic inhibitory properties for this class of coproporphyrin ferrochelatases [1], this study aims to test the hypothesis that iron inhibition is mediated by the regulatory metal binding site previously shown to bind  $\text{Mg}^{2+}$  in the *B. subtilis* enzyme [12, 13]. This work has implications for how staphylococci respond to nutritional immunity (e.g. metal sequestration) encountered during infection.

## EXPERIMENTAL PROCEDURES

### Cloning, overexpression, and purification of *S. aureus* ferrochelatase

The *cpfC* (*hemH*) gene was amplified from *S. aureus* strain USA300 [16] via colony PCR using Q5 polymerase (NEB) and cloned into the *NheI/HindIII* sites of plasmid pTrcHis (ThermoFischer), and the correct sequence of the resultant pTrcHis-*Sa-cpfC* vector was confirmed by Sanger sequencing. *E. coli* JM109 cells (SigmaAldrich) were transformed with pTrcHis-*Sa-cpfC* and a 10 mL LB overnight culture of this expression strain was used to inoculate 1L of CircleGrow medium in a 2L baffled flask ( $125 \mu\text{g mL}^{-1}$  ampicillin was included throughout for plasmid selection). Cultures were grown for 24 h at  $37^\circ\text{C}$  and 160 rpm and cells were harvested (4 krpm,  $4^\circ\text{C}$ , 20 min) and stored at  $-20^\circ\text{C}$ . Cell pellets from 1L cultures were re-suspended in protein storage buffer (30 mL) containing 50mM Tris/MOPS pH 8.0, 100mM KCl, 1% sodium cholate. Cell suspensions were sonicated (6 x 30 s on ice at  $10 \mu$ , with 30 s intervals between bursts), and cell debris was removed via centrifugation (18 krpm,  $4^\circ\text{C}$ , 20 min). The supernatant was applied to a column containing Talon metal affinity resin (2.5 mL bed volume) equilibrated with 5 column volumes of protein storage buffer. The column was washed with 20 column volumes of protein storage buffer followed by 5 column volumes of wash buffer (protein storage buffer including 15 mM imidazole), and purified protein was eluted with 5 column volumes of elution buffer (protein storage buffer including 300 mM imidazole). Purified protein was concentrated to  $<1$  mL using a 10 kDa spin concentrator (Millipore) and imidazole was removed using a PD10 desalting column (GE Healthcare) equilibrated with protein storage buffer.

### Assay of *S. aureus* ferrochelatase activity

Purified CpfC was assayed using coproporphyrin III (Frontier Scientific) as the porphyrin substrate and ferrous iron as the inserted metal ion. Coproporphyrin III powder was solubilised using a few drops of ammonium hydroxide and was then diluted in assay buffer (0.1 M Tris-HCl pH 8.1, 0.5 % sodium cholate, 0.5 % Tween 20, 1 mM  $\beta$ -mercaptoethanol). Coproporphyrin III stock solutions were quantified in 0.1 M HCl spectrophotometrically using an extinction coefficient of  $\epsilon_{399.5} = 489 \text{ mM}^{-1}\text{cm}^{-1}$  [17]. Ferrous iron solutions were quantified spectrophotometrically using ferrozine solution and an extinction coefficient of  $\epsilon_{562} = 27.9 \text{ mM}^{-1}\text{cm}^{-1}$  [18]. Iron chelation was measured using a continuous assay essentially as previously described [10]. Activity of 50 nM CpfC was monitored using a Cary 60 spectrophotometer at  $30^\circ\text{C}$ . The reactions were initiated by the addition of enzyme, which was preincubated at  $30^\circ\text{C}$ , and the depletion of coproporphyrin III was monitored at 392 nm for 3 min ( $\epsilon_{392} = 115 \text{ mM}^{-1}\text{cm}^{-1}$ , [1]). All assays contained 0.1 M Tris-HCl pH 8.1, 0.5%

sodium cholate, 0.5 % Tween 20, 1mM  $\beta$ -mercaptoethanol. The steady state rates were estimated using linear regression of the timecourse at the start of the reaction. Michaelis plots ( $v$  vs.  $[S]$ ) were fitted to the Michaelis Menten equation using nonlinear regression.  $K_{i,app}$  values were calculated by fitting  $v$  vs.  $[inhibitor]$  data to a single binding site model described by equation 1 [19], where no mechanistic assumptions are made to perform a quantitative assessment of the key features of inhibition:

$$v = v_{min} + \frac{v_{control} - v_{min}}{1 + \left( \frac{[I]}{K_{i,app}} \right)^n} \quad (1)$$

where  $v$  is the enzymatic rate,  $v_{control}$  is the rate in the absence of inhibitor,  $v_{min}$  is the rate in the presence of saturating inhibitor,  $[I]$  is the inhibitor concentration,  $K_{i,app}$  is the apparent inhibition constant defining the inhibitor concentration that elicits 50 % of the total inhibition, and  $n$  is the Hill coefficient.

### Structural modelling of *S. aureus* ferrochelatase

A structural model for *S. aureus* CpfC was generated using the RaptorX server [20, 21] with *B. subtilis* CpfC as a template (PDB id = 1AK1,  $\alpha$ -carbon RMSD = 0.23 Å). The *S. aureus* CpfC model was superposed ( $\alpha$ -carbon RMSD = 0.66 Å) onto a  $Mg^{2+}$ -bound and porphyrin-bound structure of *B. subtilis* CpfC (PDBid = 1C1H), and the native substrate coproporphyrin III was modelled into the active site in place of N-methyl mesoporphyrin via superposition using CCP4MG software.

## RESULTS

### Measurement of kinetic constants for *S. aureus* ferrochelatase

Since much of the previous work on coproporphyrin ferrochelatases has been performed with non-native substrates (i.e.  $Zn^{2+}$ , protoporphyrin IX, deuteroporphyrin IX), it was of interest to define the kinetic constants for the *in vivo* substrates  $Fe^{2+}$  and coproporphyrin III. Kinetic assays where  $[Fe^{2+}]$  was varied (Fig. 2A) demonstrate substrate inhibition above 0.8  $\mu M$ , and the data below this threshold value were fitted to the Michaelis equation to provide an estimate of  $0.27 \pm 0.04 \mu M$  for an apparent  $K_m$  for iron binding at the inner catalytic metal site. Varying [coproporphyrin III] at a fixed concentration of 0.7  $\mu M$   $Fe^{2+}$  (Fig. 2B) provides the first estimate of an apparent  $K_m$  for coproporphyrin III as  $1.5 \pm 0.2 \mu M$ , and  $k_{cat}$  was fitted as  $10.5 \pm 0.4 \text{ min}^{-1}$  at this sub-inhibitory  $Fe^{2+}$  concentration.

### Elevated iron increases the $K_m$ for coproporphyrin III

To gain an insight into the influence of iron upon coproporphyrin III binding, kinetic plots were produced for each substrate at fixed concentrations of the other substrate (Fig. 3). Firstly, varying the [coproporphyrin III] in the presence of fixed (sub-inhibitory) concentrations of  $Fe^{2+}$  revealed that increasing  $[Fe^{2+}]$  increases the apparent  $K_m$  for coproporphyrin (Fig. 3A). Performing the converse experiment with variation of  $[Fe^{2+}]$  in the presence of fixed concentrations of coproporphyrin III revealed a more conventional pattern of increasing the concentration of coproporphyrin III also decreases the apparent  $K_m$  for  $Fe^{2+}$  (Fig. 3B).

### Structural modelling of *S. aureus* ferrochelatase reveals a regulatory metal binding site

Based on the data in Fig. 3, it was hypothesized that *S. aureus* CpfC has a regulatory metal binding site, and the most simple explanation would be that this metal site is at the same location as the  $Mg^{2+}$ -binding site in *B. subtilis* CpfC. The only difference is that that this site would be inhibitory when occupied by iron, as opposed to stimulatory when occupied by  $Mg^{2+}$  [13]. To investigate the local environment of this putative regulatory metal site, a structural model for *S. aureus* CpfC was generated using the *B. subtilis* CpfC crystal structure

as a template (Fig. 4). The model and template were superposed and the  $Mg^{2+}$  ion from the *B. subtilis* structure was shown in the same spatial location in the *S. aureus* model (Fig. 4A). The metal-binding residues were fully conserved between species, with residues Glu271, Glu267, and Arg45 in the *S. aureus* model all mapping adjacent to the outer regulatory metal site (Fig. 4B).

### **Magnesium potentiates $Fe^{2+}$ -mediated inhibition of *S. aureus* ferrochelatase**

Since  $Mg^{2+}$  has previously been used to investigate the regulatory metal site in *B. subtilis* CpfC and was shown to stimulate chelatase activity [13], it was of interest to determine if  $Mg^{2+}$  could also be used as a molecular probe to study *S. aureus* CpfC. To investigate the effect of  $Mg^{2+}$  upon the  $Fe^{2+}$  kinetics, iron was varied in the presence and absence of  $Mg^{2+}$  (Fig. 5). Surprisingly, addition of  $Mg^{2+}$  potentiates chelatase inhibition, lowering the threshold concentration of  $Fe^{2+}$  required for substrate inhibition.

### **Loss of the regulatory binding site abolishes inhibition by $Mg^{2+}$ or $Fe^{2+}$**

To test the hypothesis that the metal-binding site shown in Fig. 4 is involved in the  $Mg^{2+}$ -mediated inhibition kinetics shown in Fig. 5, Glu271 was mutated to serine and the resultant E271S mutant protein was purified and assayed in the presence and absence of  $Mg^{2+}$  (Fig. 6). Fig. 6A shows that  $Mg^{2+}$ -potentiated inhibition was dramatically diminished in the E271S mutant, and that inhibition by  $Fe^{2+}$  in the absence of  $Mg^{2+}$  was also dramatically diminished. Fig. 6B displays  $K_{i,app}(Mg^{2+})$  values for wild type and E271S CpfC as 66  $\mu M$  and 11 mM, respectively, highlighting a dramatic loss of sensitivity to  $Mg^{2+}$  for the mutant enzyme. Together, these data strongly indicate that in wild type *S. aureus* CpfC, the binding of  $Mg^{2+}$  or  $Fe^{2+}$  at this metal site results in enzyme inhibition. In addition, the data in Figures 4-6 suggest that loss of the inhibitory metal site should reverse the inhibitory effects of increasing iron concentrations seen in Fig. 3A, so a similar kinetic study was carried out with the E271S variant of *S. aureus* CpfC including iron levels well above the inhibitory threshold for the wild type enzyme (Fig. 7). These data confirm that loss of the E271 metal site abolishes the iron-mediated inhibition with respect to coproporphyrin III (Fig. 7A), and the effect of varying [coproporphyrin III] upon the kinetic constants for iron (Fig. 7B) remains similar to the wild type data (Fig. 3B).

## **DISCUSSION**

Preliminary kinetic analysis of *S. aureus* CpfC revealed a  $k_{cat}$  of  $10.5 \pm 0.4 \text{ min}^{-1}$ . This rate is comparable to previous measurements for other bacterial protoporphyrin ferrochelatases [10], and slightly higher than previous measurements of  $1.8 \text{ min}^{-1}$  and  $0.11 \text{ min}^{-1}$  for coproporphyrin ferrochelatases from *M. tuberculosis* and *B. subtilis*, respectively [2]. While a  $k_{cat}$  for *S. aureus* CpfC was recently calculated to be  $165 \text{ min}^{-1}$  via extrapolation of a substrate inhibition curve [1], the maximum rates measured ( $\sim 28 \text{ min}^{-1}$ ) were of a similar magnitude to those presented herein. We report the first measurement of  $K_{app}$  (coproporphyrin III) for *S. aureus* CpfC as  $1.5 \pm 0.2 \mu M$ , which is slightly lower than previous measurements of 10.5  $\mu M$  and 7.8  $\mu M$  for CpfC enzymes from *M. tuberculosis* and *B. subtilis*, respectively [2]. Our estimate of  $0.27 \pm 0.04 \mu M$  for  $K_{app}(Fe^{2+})$  is comparable with a previous estimate of  $0.6 \pm 0.1 \mu M$  for this enzyme [1], although both of these measurements were hampered by substrate inhibition by iron.

Primary and secondary kinetic plots clearly demonstrate that elevating the iron concentration increases the  $K_m$  for coproporphyrin III (Fig. 3A), which is consistent with  $Fe^{2+}$  causing inhibition at higher concentrations. It was therefore hypothesized that  $Fe^{2+}$  is inserted into the porphyrin substrate following binding to a well-known catalytic site adjacent to the tetrapyrrole substrate, and higher concentrations of  $Fe^{2+}$  inhibit chelation by binding at a second regulatory site. Structural modelling (Fig. 4) provides convincing evidence that *S. aureus* possesses a second metal binding site similar to the  $Mg^{2+}$ -binding site in *B. subtilis* CpfC [12], although  $Mg^{2+}$  was found to potentiate  $Fe^{2+}$ -mediated inhibition in the current study (Fig. 5) compared to having a stimulatory role in previous reports [13]. This could be

due to differences in substrates used, as the current study uses the native substrates whereas previous work has used  $Zn^{2+}$  and deuteroporphyrin IX as substrates [13], which could influence the behavior of metals in the active site. Indeed, previous measurements of *S. aureus* CpfC kinetics demonstrate a lack of  $Fe^{2+}$ -mediated inhibition when protoporphyrin IX is used as a substrate [1].

To confirm the involvement of the regulatory metal site (Fig. 4) in allosteric modulation of *S. aureus* CpfC, the E271S mutant enzyme was purified and kinetic analyses were performed in the presence and absence of  $Mg^{2+}$ . A high concentration of 4 mM  $Mg^{2+}$  was used initially to saturate metal binding (i.e.  $Fe^{2+}$  and  $Mg^{2+}$ ) at the inhibitory site. Substitution of this glutamate residue with serine resulted in two dramatic outcomes: substrate inhibition by  $Fe^{2+}$  was lost and  $Mg^{2+}$  was no longer a potent inhibitor of metal chelation (Fig. 6A). This strongly suggests that the binding of  $Fe^{2+}$  or  $Mg^{2+}$  is stabilized by Glu271. To add weight to this conclusion,  $Mg^{2+}$  dose response data confirms that  $Mg^{2+}$  is a potent inhibitor of the wild type enzyme but not the E271S mutant (Fig. 6B). Steady state kinetics for the E271S mutant revealed that loss of metal binding at this residue reverses the  $Fe^{2+}$ -mediated increase in  $K_m$ (coproporphyrin III) (Fig. 7A) that is seen for the wild type enzyme in Fig. 3A. The effect of varying [coproporphyrin III] upon the kinetic constants for iron elicits the same pattern for the wild type (Fig. 3B) and E271S mutant (Fig. 7B), which strongly suggests that metal binding at the E271S site largely inhibits coproporphyrin binding and/or chelation rather than iron binding at the catalytic site. Taken together, all data presented herein support the model for wild type *S. aureus* CpfC in Fig. 8 where iron binding to a lower affinity regulatory site at E271 (blue) diminishes the rate of iron chelation at an inner catalytic site (green) via increasing the  $K_m$  for the coproporphyrin III substrate.

It is well-known that there is fierce competition for iron both during infection resulting from nutritional immunity [22], and during growth in mixed microbial communities mediated by microbial siderophores. Therefore, where iron is scarce it is important for *S. aureus* to tightly regulate how this nutrient is used within the cell. The current work is consistent with a model whereby at very low  $Fe^{2+}$  concentrations ferrochelatase is able to insert iron into coproporphyrin III, whereas at higher iron concentrations flux through this step of the pathway is down-regulated. There may be a number of reasons for this substrate inhibition, including avoiding accumulation of Fe-coproporphyrin III, or perhaps once the haem requirements of the cell are satisfied  $Fe^{2+}$  is made available for other processes (e.g. Fe-S cluster assembly). Future studies on bacteria that perform coproporphyrin-dependent haem biosynthesis are required to reveal the hierarchical channelling of iron for different uses in the cell, which will be important during infection (*S. aureus*, *M. tuberculosis*, *P. acnes*) and growth in microbial communities (*B. subtilis*).

#### **ACKNOWLEDGEMENTS**

This work was funded by a Research Grant from the Royal Society (RG110528 to M.S.). We thank the laboratory of Prof. Harry Dailey (University of Georgia) for supplying expression plasmids for ferrochelatases.

#### **AUTHOR CONTRIBUTION**

CH performed the experiments and MS drove the experimental design. CH and MS analysed the data. MS, JDR, and CH wrote the paper.

#### **FUNDING**

This work was funded by a Research Grant from the Royal Society (RG110528 to MS).

## REFERENCES

1. Lobo, S.A., Scott A., Videira M.A., Winpenny D., Gardner M., Palmer M.J., Schroeder S., Lawrence A.D., Parkinson T., Warren M.J. and Saraiva L.M. (2015) *Staphylococcus aureus* haem biosynthesis: characterisation of the enzymes involved in final steps of the pathway. *Mol. Microbiol.* **97**, 472–487.
2. Dailey, H.A., Gerdes S., Dailey T.A., Burch J.S. and Phillips J.D. (2015) Noncanonical coproporphyrin-dependent bacterial heme biosynthesis pathway that does not use protoporphyrin. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 2210-2215.
3. Dailey, H.A., Dailey T.A., Gerdes S., Jahn D., Jahn M., O'Brian M.R. and Warren M.J. (2017) Prokaryotic heme biosynthesis: multiple pathways to a common essential product. *Microbiol. Mol. Biol. Rev.* **81**, e00048-16.
4. Hansson, M. and Hederstedt L. (1992) Cloning and characterization of the *Bacillus subtilis hemEHY* gene cluster, which encodes protoheme-IX biosynthetic-enzymes. *J. Bacteriol.* **174**, 8081-8093.
5. Hansson, M. and Hederstedt L. (1994) Purification and characterisation of a water-soluble ferrochelatase from *Bacillus subtilis*. *Eur. J. Biochem.* **220**, 201-208.
6. Dailey, T.A. and Dailey H.A. (2002) Identification of 2Fe-2S clusters in microbial ferrochelatases. *J. Bacteriol.* **184**, 2460-2464.
7. Camadro, J.M., Ibrahim N.G. and Levere R.D. (1984) Kinetic studies of human liver ferrochelatase - role of endogenous metals. *J. Biol. Chem.* **259**, 5678-5682.
8. Wu, C.K., Dailey H.A., Rose J.P., Burden A., Sellers V.M. and Wang B.C. (2001) The 2.0 angstrom structure of human ferrochelatase, the terminal enzyme of heme biosynthesis. *Nat. Struct. Biol.* **8**, 156-160.
9. Karlberg, T., Lecerof D., Gora M., Silvegren G., Labbe-Bois R., Hansson M. and Al-Karadaghi S. (2002) Metal Binding to *Saccharomyces cerevisiae* ferrochelatase. *Biochemistry* **41**, 13499-13506.
10. Shepherd, M., Dailey T.A. and Dailey H.A. (2006) A new class of 2Fe-2S -cluster-containing protoporphyrin (IX) ferrochelatases. *Biochem. J.* **397**, 47-52.
11. Dailey, H.A., Sellers V.M. and Dailey T.A. (1994) Mammalian ferrochelatase - expression and characterization of normal and 2 human protoporphyrin ferrochelatases. *J. Biol. Chem.* **269**, 390-395.
12. Lecerof, D., Fodje M., Hansson A., Hansson M. and Al-Karadaghi S. (2000) Structural and mechanistic basis of porphyrin metallation by ferrochelatase. *J. Mol. Biol.* **297**, 221-232.
13. Lecerof, D., Fodje M.N., Alvarez Leon R., Olsson U., Hansson A., Sigfridsson E., Ryde U., Hansson M. and Al-Karadaghi S. (2003) Metal binding to *Bacillus subtilis* ferrochelatase and interaction between metal sites. *J. Biol. Inorg. Chem.* **8**, 452-458.
14. Hansson, M.D., Karlberg T., Soderberg C.A.G., Rajan S., Warren M.J., Al-Karadaghi S., Rigby S.E.J. and Hansson M. (2011) Bacterial ferrochelatase turns human: Tyr13 determines the apparent metal specificity of *Bacillus subtilis* ferrochelatase. *J. Biol. Inorg. Chem.* **16**, 235-242.
15. Davidson, R.E., Chesters C.J. and Reid J.D. (2009) Metal ion selectivity and substrate inhibition in the metal ion chelation catalyzed by human ferrochelatase. *J. Biol. Chem.* **284**, 33795-33799.
16. Kazakova, S.V., Hageman J.C., Matava M., Srinivasan A., Phelan L., Garfinkel B., Boo T., McAllister S., Anderson J., Jensen B., Dodson D., Lonsway D., McDougal L.K., Arduino M., Fraser V.J., Killgore G., Tenover F.C., Cody S. and Jernigan D.B. (2005) A clone of methicillin-resistant *Staphylococcus aureus* among professional football players. *N. Engl. J. Med.* **352**, 468-475.
17. Smith, K.M. (1975) Porphyrins and metalloporphyrins, Elsevier, New York
18. Berlett, B.S., Levine R.L., Chock P.B., Chevion M. and Stadtman E.R. (2001) Antioxidant activity of ferrozine-iron-amino acid complexes. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 451-456.
19. Stein, R.L. (2011) in *Kinetics of Enzyme Action*, pp. 73-87, John Wiley & Sons, Inc.

20. Kaellberg, M., Wang H., Wang S., Peng J., Wang Z., Lu H. and Xu J. (2012) Template-based protein structure modeling using the RaptorX web server. *Nat. Protoc.* **7**, 1511-1522.
21. Peng, J. and Xu J. (2011) RaptorX: Exploiting structure information for protein alignment by statistical inference. *Proteins* **79**, 161-171.
22. Hammer, N.D. and Skaar E.P. (2012) The impact of metal sequestration on *Staphylococcus aureus* metabolism. *Curr. Op. Microbiol.* **15**, 10-14.

## FIGURE LEGENDS

**Figure 1. The coproporphyrin-dependent pathway of haem synthesis.** This figure depicts a recently discovered route for haem synthesis in Firmicutes and Actinobacteria catalysed by coproporphyrinogen oxidase (CgoX), coproporphyrin ferrochelatase (CpfC), and coprohaem decarboxylase (ChdC). These new abbreviations are introduced in a recent review [3].

**Figure 2. Kinetic analysis of *S. aureus* ferrochelatase.** A) 50 nM CpfC was assayed in the presence of varying concentrations of iron and a fixed concentration of 1  $\mu\text{M}$  of coproporphyrin. Data collected at 0 - 0.8  $\mu\text{M}$  iron were fitted to a single rectangular hyperbola to provide an estimate of the apparent  $K_m$  for iron ( $K_{app} = 0.27 \pm 0.04$ ). B) 50 nM CpfC was assayed in the presence of varying concentrations of coproporphyrin III and a fixed iron concentration of 0.7  $\mu\text{M}$  (below the 0.8  $\mu\text{M}$  inhibitory iron concentration). Data were fitted to a single rectangular hyperbola via nonlinear regression:  $k_{cat} = 10.5 \pm 0.4 \text{ min}^{-1}$ ,  $K_{app} = 1.5 \pm 0.2 \mu\text{M}$ .

**Figure 3. Opposing kinetic patterns for the two ferrochelatase substrates.** Primary Michaelis Menten plots were generated with fixed concentrations of iron (panel A) and coproporphyrin III (panel B), and data were fitted to single rectangular hyperbolae via nonlinear regression. Lineweaver-Burk plots (insets) are shown to highlight differences in kinetic patterns between panels A and B. To avoid well-known problems with these secondary plots, trend lines were generated using the kinetic constants obtained from nonlinear regression of the untransformed data.

**Figure 4. Structural modelling of bound coproporphyrin and a regulatory metal binding site for *S. aureus* ferrochelatase.** A) Surface electrostatics view of a *S. aureus* CpfC structural model generated using the RaptorX server with *B. subtilis* CpfC as a template (PDBid = 1AK1,  $\alpha$ -carbon RMSD = 0.23  $\text{\AA}$ ). The *S. aureus* CpfC model was superposed ( $\alpha$ -carbon RMSD = 0.66  $\text{\AA}$ ) onto a  $\text{Mg}^{2+}$ -bound (cyan) and porphyrin-bound structure of *B. subtilis* CpfC (PDBid = 1C1H), and the native substrate coproporphyrin III (yellow) was modelled into the active site in place of N-methyl mesoporphyrin. B) Cartoon representation of the active site of *S. aureus* CpfC showing conserved residues implicated in metal binding at an outer regulatory site (E271, E267, R45) and an inner catalytic site (E263, Y12, H181).

**Figure 5. Impact of magnesium upon ferrochelatase kinetics.**  $v$  vs.  $[\text{Fe}^{2+}]$  plots were generated for *S. aureus* CpfC assayed in the presence ( $\circ$ ) and absence ( $\bullet$ ) of 4 mM  $\text{Mg}^{2+}$ . Fixed concentrations of 50 nM enzyme and 2  $\mu\text{M}$  coproporphyrin III were used, with the  $\text{Fe}^{2+}$  concentration being varied.

**Figure 6. Loss of the metal-binding residue Glu271 abrogates the inhibition of *S. aureus* ferrochelatase by iron and magnesium.** A)  $v$  vs.  $[\text{Fe}^{2+}]$  plots were generated for wild type (red) and an E271S variant of *S. aureus* CpfC (blue) assayed in the presence (squares) and absence (circles) of 4 mM  $\text{Mg}^{2+}$ . Fixed concentrations of 50 nM enzyme and 2  $\mu\text{M}$  coproporphyrin III were used. The fitted blue line provides estimates of apparent kinetic constants when the inhibition by  $\text{Fe}^{2+}$  is removed:  $k_{cat} = 24.1 \pm 2.0 \text{ min}^{-1}$ ,  $K_{app}(\text{Fe}^{2+}) = 1.2 \pm 0.2 \mu\text{M}$ . B) Inhibition dose responses for wild type (red) and an E271S variant of *S. aureus* CpfC

(blue) assayed in the presence of increasing concentrations of  $\text{Mg}^{2+}$ . Fixed concentrations of 50 nM enzyme, 2  $\mu\text{M}$  coproporphyrin III, and 0.7  $\mu\text{M}$   $\text{Fe}^{2+}$  were used. To estimate  $K_{i,\text{app}}$  values the data were fitted to a four parameter Hill equation (Equation 1, [19]) with the minimum activity (i.e.  $v_{\text{min}}$ ) constrained to a positive value.

**Figure 7. Loss of inhibitory metal site restores conventional Michaelis kinetics.** Primary Michaelis Menten plots for an E271S variant of *S. aureus* CpfC were generated with fixed concentrations of iron (panel A) and coproporphyrin III (panel B), and data were fitted to single rectangular hyperbolae via nonlinear regression. Lineweaver-Burk plots (insets) are shown to highlight differences in kinetic patterns between panels A and B. To avoid well-known problems with these secondary plots, trend lines were generated using the kinetic constants obtained from nonlinear regression of the untransformed data.

**Figure 8. Kinetic model for iron-dependent inhibition of *S. aureus* ferrochelatase.** Processes shown in blue and green depict conditions of high and low iron concentrations where the inhibitory regulatory site is occupied or empty, respectively. Since inhibition is only observed at higher iron concentrations, it is unlikely that the regulatory site would be occupied when the catalytic site is empty, so these intermediates are not shown.

**FIGURE 1**




FIGURE 2





FIGURE 3



**FIGURE 4**




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8

