



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/120941/>

Version: Accepted Version

---

**Article:**

Kozłowski, Karol, Sklyanin, Evgeny and Torrielli, Alessandro (2017) Quantization of the Kadomtsev–Petviashvili equation. *Theoretical and Mathematical Physics*. 1162–1183.  
ISSN: 1573-9333

<https://doi.org/10.1134/S0040577917080074>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# QUANTISATION OF KADOMTSEV-PETVIASHVILI EQUATION

K K KOZLOWSKI<sup>1</sup>, E SKLYANIN<sup>2</sup>, AND A TORRIELLI<sup>3</sup>

ABSTRACT. A quantisation of the KP equation on a cylinder is proposed that is equivalent to an infinite system of non-relativistic one-dimensional bosons carrying masses  $m = 1, 2, \dots$ . The Hamiltonian is Galilei-invariant and includes the split  $\Psi_{m_1}^\dagger \Psi_{m_2}^\dagger \Psi_{m_1+m_2}$  and merge  $\Psi_{m_1+m_2}^\dagger \Psi_{m_1} \Psi_{m_2}$  terms for all combinations of particles with masses  $m_1, m_2$  and  $m_1 + m_2$ , with a special choice of coupling constants. The Bethe eigenfunctions for the model are constructed. The consistency of the coordinate Bethe Ansatz, and therefore, the quantum integrability of the model is verified up to the mass  $M = 8$  sector.

## 1. INTRODUCTION

The Kadomtsev-Petviashvili (KP) equation [10]

$$\varphi_{t\sigma} - \varphi_{xx} - 2\beta(\varphi\varphi_\sigma)_\sigma + \gamma\varphi_{\sigma\sigma\sigma} = 0, \quad (1.1)$$

is one of the most studied nonlinear integrable equations in 2+1 variables  $(\sigma, x; t)$ . The aim of the present paper is to construct a quantised version of KP while preserving its integrability.

*A warning:* For reasons explained below, we have deliberately deviated from the standard notation of [10] by changing the conventional variable  $x$  to  $\sigma$  and  $y$  to  $x$ .

Traditionally, KP is considered as an equation in (2+1)-dimensional space-time, both variables  $\sigma$  and  $x$  playing the role of spatial variables. For our purposes, however, we take a different stance, viewing only  $x$  as a genuine spatial variable and downgrading  $\sigma$  to a mere label indexing the continuum of fields  $\varphi$  in (1+1)-dimensional space-time. The notation  $(\sigma, x)$  stresses the changed roles of the two variables.

We also choose  $x$  to run from  $-\infty$  to  $\infty$ , whereas imposing the periodicity condition  $\sigma \equiv \sigma + 2\pi$  on  $\sigma$ . We assume that  $\varphi \rightarrow 0$  sufficiently fast as  $x \rightarrow \pm\infty$ .

We have also introduced two real coupling constants  $\beta$  and  $\gamma$  into the equation. Though, in the classical case, they can be removed by a rescaling of the variables  $x, \sigma, \varphi$ , they are useful for the quantisation and for discussing the limiting cases. Note that the constants  $\beta$  and  $\gamma$  may have arbitrary sign, the case  $\gamma > 0$  corresponding to the so-called KP-I, respectively  $\gamma < 0$  to KP-II, and  $\gamma = 0$  to the so-called dispersionless KP [17]. Since the substitution  $\varphi := -\varphi$  results in changing the sign of  $\beta$ , one may assume that  $\beta \geq 0$ .

The paper is organised as follows. In Section 2, we describe the Poisson structure and the Hamiltonian of the classical model. In Section 3, we quantise the model using the simplest normal ordering prescription for the Hamiltonian. Passing from the field  $\varphi(\sigma, x)$  to its Fourier components in the variable  $\sigma$  we obtain the description of the system in

<sup>1</sup>UNIV LYON, ENS DE LYON, UNIV CLAUDE BERNARD, CNRS, LABORATOIRE DE PHYSIQUE, F-69342 LYON, FRANCE.

<sup>2</sup>DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK YO10 5DD, UK

<sup>3</sup>DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SURREY, GUILDFORD, GU2 7XH, UK

Date: July 27, 2016.

terms of the discrete infinite set of canonical fields  $\Psi_m^\dagger(x)$ ,  $\Psi_m(x)$  labelled by the index  $m = 1, 2, 3, \dots$  and describing scalar nonrelativistic bosons of mass  $m$ . The Hamiltonian is Galilei-invariant and includes the split  $\Psi_{m_1}^\dagger \Psi_{m_2}^\dagger \Psi_{m_1+m_2}$  and merge  $\Psi_{m_1+m_2}^\dagger \Psi_{m_1} \Psi_{m_2}$  terms for all combinations of particles with masses  $m_1$ ,  $m_2$  and  $m_1 + m_2$ , with a special choice of coupling constants.

In Section 4, we describe the Fock space  $\mathcal{F}$  of the system and introduce a convenient notation to handle the infinite number of fields. We realise as well the action of the Hamiltonian on the  $N$ -particle state as a differential operator with singular delta-function coefficients.

Due to the conservation of the total mass  $M$ , the quantum-field-theoretic problem is reduced to a sequence of quantum-mechanical problems in sectors of fixed mass  $M$ . The structure of mass- $M$  sector  $\mathcal{F}_M$  is analysed in Section 5. Since the number of particles is not preserved, the sector  $\mathcal{F}_M$  splits into the orthogonal sum of subspaces  $\mathcal{F}_M^{\mathbf{m}}$  labelled by compositions  $\mathbf{m}$  of number  $M$ . The corresponding wave functions are defined on Weyl alcoves  $x_1 < x_2 < \dots < x_N$ , where  $N$  is the length of  $\mathbf{m} = (m_1, m_2, \dots, m_N)$ .

In Section 6, we interpret the delta-function terms in the Hamiltonian as jump conditions for the derivatives of components of the wave function, and formulate the complete set of differential equations and boundary conditions for the wave functions.

In Section 7, we solve the eigenvalue problem in the sector  $M = 2$ , and compute the two-particle  $S$ -matrix, as a rational function having 3 poles and 3 zeroes. The two possible arrangements of the poles are labelled as quantum KP-I and KP-II cases.

In Section 8, we formulate the Bethe Ansatz in the subsector  $\mathcal{F}_M^{(1 \dots 1)}$  containing only particles of mass-1. The Bethe eigenfunction is written as a linear combination of plain waves with the coefficients that reproduce the correct 2-particle  $S$ -matrices. In Section 9, we extend the Bethe Ansatz to the generic sector of particles with different masses, and formulate the factorisation conjecture that allows one to reduce the verification of the consistency equations to those for the subsector  $\mathcal{F}_M^{(M)}$  containing a single particle of mass  $M$ . In Section 10, we analyse those equations and describe a solution that is verified by means of computer algebra up to  $M \leq 8$ . A more technical discussion of the involved combinatorial issues is left for the appendices.

In the concluding Section 11 we sum up the results and discuss the unsolved questions and perspectives.

## 2. CLASSICAL KP

In this paper, we use the following notation:  $\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$  stands for the set of integers,  $\mathbb{N} = \{1, 2, 3, \dots\}$  the set of natural numbers,  $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$  the set of non-negative integers,  $\mathbb{R}$  the set of real numbers,  $\mathbb{S}^1 = \mathbb{R}/2\pi\mathbb{Z}$  a circle.

The classical Kadomtsev-Petviashvili (KP) integrable hierarchy [10] is formulated in terms of a real-valued scalar field  $\varphi(\sigma, x)$  on the cylinder  $\mathbb{S}^1 \times \mathbb{R}$ . The field  $\varphi$  vanishes sufficiently fast as  $x \rightarrow \pm\infty$  and has Poisson brackets

$$\{\varphi(\sigma, x), \varphi(\tau, y)\} = 2\pi\delta'(\sigma - \tau)\delta(x - y), \quad \sigma, \tau \in \mathbb{S}^1, \quad x, y \in \mathbb{R}. \quad (2.1)$$

Due to the periodicity of  $\varphi(\sigma, x)$  in  $\sigma$ , the average of  $\varphi$  over  $\mathbb{S}^1$  belongs to the center of the bracket (2.1). In what follows we always set it to 0, assuming that

$$\int_0^{2\pi} d\sigma \varphi(\sigma, x) = 0 \quad \forall x \in \mathbb{R}. \quad (2.2)$$

Due to (2.2), the antiderivative  $\partial_\sigma^{-1}$  on the space of functions with zero average over  $\mathbb{S}^1$  is defined correctly (one can always choose the integration constant in a unique way).

There exists an infinite series of commuting Hamiltonians  $H_p$ ,  $p = 0, 1, 2, \dots$

$$\{H_p, H_q\} = 0 \quad (2.3)$$

expressed as integrals of local (w.r.t.  $x$ ) densities

$$H_p = \int_0^{2\pi} \frac{d\sigma}{2\pi} \int_{-\infty}^{\infty} dx h_p(\sigma, x). \quad (2.4)$$

$$h_p(\sigma, x) = \frac{1}{2} (\partial_\sigma^{-p} \varphi) (\partial_x^p \varphi) + O(\beta) + O(\gamma), \quad \beta, \gamma \rightarrow 0 \quad (2.5)$$

such that

$$h_0(\sigma, x) = \frac{1}{2} \varphi^2(\sigma, x), \quad (2.6a)$$

$$h_1(\sigma, x) = \frac{1}{2} (\partial_\sigma^{-1} \varphi) (\partial_x \varphi), \quad (2.6b)$$

$$h_2(\sigma, x) = \frac{1}{2} (\partial_\sigma^{-2} \varphi) (\partial_x^2 \varphi) + \frac{\beta}{3} \varphi^3 + \frac{\gamma}{2} (\partial_\sigma \varphi)^2. \quad (2.6c)$$

The corresponding equations of motion  $\partial_{t_p} = \{\cdot, H_p\}$  are

$$\varphi_{t_0} = \varphi_\sigma, \quad (2.7a)$$

$$\varphi_{t_1} = -\varphi_x, \quad (2.7b)$$

$$\varphi_{t_2} = \partial_\sigma^{-1} \varphi_{xx} + 2\beta \varphi \varphi_\sigma - \gamma \varphi_{\sigma\sigma\sigma}. \quad (2.7c)$$

Note that  $H_0$  and  $H_1$  are generators of translations in  $\sigma$  and  $x$  respectively.

Differentiating (2.7c) in respect to  $\sigma$  we obtain the KP equation in the form (1.1).

Note that the equations of motion (1.1) are invariant under the Galilei transform  $x := x + 2vt$ ,  $\sigma := \sigma + vx + v^2t$ . The infinitesimal Galilei boost

$$B = \int_0^{2\pi} \frac{d\sigma}{2\pi} \int_{-\infty}^{\infty} dx x h_0(\sigma, x), \quad \{\varphi, B\} = x \varphi_\sigma \quad (2.8)$$

commutes with the Hamiltonians as follows:

$$\{H_p, B\} = -p H_{p-1}. \quad (2.9)$$

### 3. QUANTISATION

Using the correspondence principle  $[\cdot, \cdot] \simeq i\hbar\{\cdot, \cdot\}$  and setting  $\hbar = 1$  we obtain from (2.1) the commutation relations for the field  $\varphi(\sigma, x)$ :

$$[\varphi(\sigma, x), \varphi(\tau, y)] = 2\pi i \delta'(\sigma - \tau) \delta(x - y), \quad \varphi^\dagger = \varphi. \quad (3.1)$$

The corresponding Fourier components

$$\varphi(\sigma, x) = \sum_{n \in \mathbb{Z}} a_n(x) e^{-in\sigma}, \quad a_n(x) = \int_0^{2\pi} \frac{d\sigma}{2\pi} \varphi(\sigma, x) e^{in\sigma}, \quad a_0(x) = 0, \quad (3.2)$$

form the Heisenberg (oscillator) Lie algebra [9]

$$[a_m(x), a_n(y)] = m \delta_{m+n, 0} \delta(x - y), \quad a_n^\dagger(x) = a_{-n}(x). \quad (3.3)$$

Consider the highest-weight (h.w.) module generated by the h.w. vector (vacuum)  $|0\rangle$  such that

$$a_n(x)|0\rangle = 0, \quad n > 0. \quad (3.4)$$

Equivalently, the h.w. module is isomorphic to the bosonic Fock space  $\mathcal{F}$  generated by the canonical creation/annihilation operators  $\Psi_n^\dagger(x)$  and  $\Psi_n(x)$

$$\Psi_n(x) = n^{-1/2}a_n(x), \quad \Psi_n^\dagger(x) = n^{-1/2}a_{-n}(x), \quad n \in \mathbb{N}, \quad x \in \mathbb{R}, \quad (3.5)$$

$$[\Psi_m(x), \Psi_n^\dagger(y)] = \delta_{mn}\delta(x-y), \quad \Psi_m(x)|0\rangle = 0, \quad m, n \in \mathbb{N}, \quad x, y \in \mathbb{R}. \quad (3.6)$$

Our quantisation prescription for the Hamiltonians  $H_0$ ,  $H_1$  and  $H_2$  is to take the classical expressions (2.6), replace  $\varphi$  with the quantum operators and apply the Wick normal ordering:  $\Psi^\dagger$  to the left,  $\Psi$  to the right. The result is

$$\mathbf{H}_0 = \int_0^{2\pi} \frac{d\sigma}{2\pi} \int_{-\infty}^{\infty} dx : \frac{1}{2} \varphi^2(\sigma, x) : = \sum_{m \in \mathbb{N}} m \int_{-\infty}^{\infty} dx \Psi_m^\dagger(x) \Psi_m(x), \quad (3.7)$$

$$\mathbf{H}_1 = \int_0^{2\pi} \frac{d\sigma}{2\pi} \int_{-\infty}^{\infty} dx : \frac{1}{2} (\partial_\sigma^{-1} \varphi) (\partial_x \varphi) : = -i \sum_{m \in \mathbb{N}} \int_{-\infty}^{\infty} dx \Psi_m^\dagger(x) \partial_x \Psi_m(x), \quad (3.8)$$

$$\begin{aligned} \mathbf{H}_2 &= \int_0^{2\pi} \frac{d\sigma}{2\pi} \int_{-\infty}^{\infty} dx : \frac{1}{2} (\partial_\sigma^{-2} \varphi) (\partial_x^2 \varphi) + \frac{\beta}{3} \varphi^3 + \frac{\gamma}{2} (\partial_\sigma \varphi)^2 : \\ &= - \sum_{m \in \mathbb{N}} \frac{1}{m} \int_{-\infty}^{\infty} dx \Psi_m^\dagger(x) \partial_x^2 \Psi_m(x) \\ &+ \sum_{m_1, m_2 \in \mathbb{N}} \beta_{m_1 m_2} \int_{-\infty}^{\infty} dx [\Psi_{m_1+m_2}^\dagger(x) \Psi_{m_1}(x) \Psi_{m_2}(x) + \Psi_{m_1}^\dagger(x) \Psi_{m_2}^\dagger(x) \Psi_{m_1+m_2}(x)] \\ &+ \sum_{m \in \mathbb{N}} \gamma_m \int_{-\infty}^{\infty} dx \Psi_m^\dagger(x) \Psi_m(x), \end{aligned} \quad (3.9)$$

where

$$\beta_{m_1 m_2} = \beta_{m_2 m_1} = \beta \sqrt{(m_1 + m_2) m_1 m_2}, \quad \gamma_m = \gamma m^3. \quad (3.10)$$

As in the classical case,  $\mathbf{H}_0$  and  $\mathbf{H}_1$  being, respectively, generators of  $\sigma$ - and  $x$ -translations, commute between themselves and with  $\mathbf{H}_2$ . The quantum Galilei boost

$$\mathbf{B} = \int_{-\infty}^{\infty} dx x : \frac{1}{2} \varphi^2(\sigma, x) : = \sum_{m \in \mathbb{N}} m \int_{-\infty}^{\infty} dx x \Psi_m^\dagger(x) \Psi_m(x) \quad (3.11)$$

commutes with the Hamiltonians as follows:

$$[\mathbf{H}_0, \mathbf{B}] = 0, \quad [\mathbf{H}_1, \mathbf{B}] = -i\mathbf{H}_0, \quad [\mathbf{H}_2, \mathbf{B}] = -2i\mathbf{H}_1. \quad (3.12)$$

Physically, the Hamiltonian  $\mathbf{H}_2$  describes a non-relativistic, Galilei-invariant system of one-dimensional Bose-particles labelled by the integer index  $m$  that can be interpreted as particle's mass. The interaction is local. The cubic  $\beta$ -terms describe processes where 2 particles of masses  $m_1$  and  $m_2$  merge into one of mass  $m_1 + m_2$  and the respective splitting. The unitary transformation  $\Psi_m \mapsto -\Psi_m$ ,  $\Psi_m^\dagger \mapsto -\Psi_m^\dagger$  simply changes the sign of  $\beta$ , so one may assume  $\beta \geq 0$ . For  $\beta=0$  the fields decouple, and one gets the theory of free particles with masses  $m$  and the rest energy  $\gamma m^3$ .

A model with such kind of interaction was first proposed in [14], and its variants and generalisations under the general name 'Lee model' were popular in 1950-60s as toy models

in nuclear physics. Our variant of the Lee model is distinguished on several counts: first, by being 1D, second, by using infinitely many fields, and third, by the specific choice of coupling constants (3.10) that, as we are expecting, makes the theory integrable. Other examples of integrable 1D models of Lee type that have been studied previously include the  $N$ -waves model [12] and continuous magnet [19].

The crucial question is thus whether the integrability of the theory is preserved in the quantum case. One way of checking the integrability would be to construct higher commuting quantum Hamiltonians  $H_n$ ,  $n \geq 3$  for which the normal ordering prescription can not be expected to work. Moreover, the problem of higher local quantum Hamiltonians is notoriously difficult even in a much simpler case of the quantum nonlinear Schrödinger equation [18]: the higher Hamiltonians are known to be extremely singular and do not have well-defined normal symbols [5, 6, 8].

As our test of integrability, we choose instead to construct an explicit formula for the simultaneous eigenfunctions of  $\mathbf{H}_0$ ,  $\mathbf{H}_1$  and  $\mathbf{H}_2$  by means of the coordinate Bethe Ansatz and to show that the multiparticle  $S$ -matrices are factorised into 2-particles ones.

#### 4. FOCK SPACE

The canonical operators  $\Psi_m^\dagger(x)$  and  $\Psi_m(x)$  are labelled by the pairs  $(m, x) \in \mathbb{N} \times \mathbb{R}$ .

It is convenient to treat the pair of labels as a single composite entity  $\xi = (m, x)$ , or  $\eta = (n, y)$ . Denoting  $\delta_{\xi\eta} = \delta_{mn}\delta(x-y)$  we can thus rewrite (3.6) as

$$[\Psi_\xi, \Psi_\eta^\dagger] = \delta_{\xi\eta}, \quad \Psi_\xi |0\rangle = 0. \quad (4.1)$$

The bosonic Fock space  $\mathcal{F}$  is decomposed into  $N$ -particle components spanned by the vectors

$$|f\rangle = \sum_{N=0}^{\infty} \frac{1}{N!} \sum_{\mathbf{m} \in \mathbb{N}^N} \int_{\mathbb{R}^N} dx_1 \dots dx_N f_N(\mathbf{m}, \mathbf{x}) \prod_{j=1}^N \Psi_{m_j}^\dagger(x_j) |0\rangle \quad (4.2)$$

defined in terms of the  $N$ -particle wave functions

$$f_N(\mathbf{m}, \mathbf{x}) = f_N \begin{pmatrix} m_1 & m_2 & \dots & m_N \\ x_1 & x_2 & \dots & x_N \end{pmatrix} = f_N(\xi_1, \dots, \xi_N) = f_N(\boldsymbol{\xi}) \quad (4.3)$$

depending on  $N$  discrete indices  $\mathbf{m} = (m_1, \dots, m_N) \in \mathbb{N}^N$  and  $N$  continuous variables  $\mathbf{x} = (x_1, \dots, x_N) \in \mathbb{R}^N$ , and symmetric with respect to permutations of the pairs  $\xi_i = (m_i, x_i)$ . We shall use the notation

$$(|f\rangle)_N(\boldsymbol{\xi}) \equiv f_N(\boldsymbol{\xi}) \quad (4.4)$$

to refer to the  $N$ -particle component of the vector  $|f\rangle$ .

Using the shorthand notation, one can rewrite (4.2) as

$$|f\rangle = \sum_{N=0}^{\infty} \frac{1}{N!} \int d\boldsymbol{\xi}^N f_N(\boldsymbol{\xi}) \prod_{j=1}^N \Psi_{\xi_j}^\dagger |0\rangle, \quad (4.5)$$

where

$$\int d\boldsymbol{\xi}^N = \sum_{\mathbf{m} \in \mathbb{N}^N} \int_{\mathbb{R}^N} dx_1 \dots dx_N. \quad (4.6)$$

The norm of the vector  $|f\rangle$  is

$$\|f\|^2 = \langle f|f\rangle = \sum_{N=0}^{\infty} \frac{1}{N!} \int d\xi^N |f_N(\xi)|^2. \quad (4.7)$$

From (4.1) and (4.5) the action of the canonical operators on the  $N$ -particle wave function can be computed easily:

$$(\Psi_\eta | f\rangle)_N(\xi_1, \dots, \xi_N) = f_{N+1}(\xi_1, \dots, \xi_N, \eta), \quad (4.8)$$

or, simply

$$(\Psi_\eta | f\rangle)_N(\xi) = f_{N+1}(\xi, \eta). \quad (4.9)$$

Respectively,

$$(\Psi_\eta^\dagger | f\rangle)_N(\xi_1, \dots, \xi_N) = \sum_{j=1}^N \delta_{\eta\xi_j} f_{N-1}(\xi_1, \dots, \widehat{\xi_j}, \dots, \xi_N), \quad (4.10)$$

where  $\widehat{\xi_j}$  means omitting  $\xi_j$ .

From (4.9), and (4.10) one easily derives the action of the Hamiltonians  $\mathbf{H}_0$ ,  $\mathbf{H}_1$  and  $\mathbf{H}_2$  on the state  $|f\rangle$  in terms of its  $N$ -particle components.

From (3.7) one computes that

$$(\mathbf{H}_0 | f\rangle)_N \begin{pmatrix} \mathbf{m} \\ \mathbf{x} \end{pmatrix} = |\mathbf{m}| f_N \begin{pmatrix} \mathbf{m} \\ \mathbf{x} \end{pmatrix}, \quad |\mathbf{m}| = m_1 + \dots + m_N, \quad (4.11)$$

measuring thus the total mass of an  $N$ -particle system.

Similarly, from (3.8) one derives that  $\mathbf{H}_1$  is the total momentum operator (generator of infinitesimal translation):

$$(\mathbf{H}_1 | f\rangle)_N \begin{pmatrix} \mathbf{m} \\ \mathbf{x} \end{pmatrix} = -i(\partial_{x_1} + \dots + \partial_{x_N}) f_N \begin{pmatrix} \mathbf{m} \\ \mathbf{x} \end{pmatrix}. \quad (4.12)$$

From (3.9) the action of  $\mathbf{H}_2$  on  $|f\rangle$  takes the form:

$$\begin{aligned} (\mathbf{H}_2 | f\rangle)_N \begin{pmatrix} m_1 & \dots & m_N \\ x_1 & \dots & x_N \end{pmatrix} &= - \left( \frac{1}{m_1} \partial_{x_1}^2 + \dots + \frac{1}{m_N} \partial_{x_N}^2 \right) f_N \begin{pmatrix} m_1 & \dots & m_N \\ x_1 & \dots & x_N \end{pmatrix} \\ + 2 \sum_{1 \leq i_1 < i_2 \leq N} \beta_{m_{i_1} m_{i_2}} f_{N-1} \begin{pmatrix} m_1 & \dots & \widehat{m_{i_1}} & \dots & \widehat{m_{i_2}} & \dots & m_N, & m_{i_1} + m_{i_2} \\ x_1 & \dots & \widehat{x_{i_1}} & \dots & \widehat{x_{i_2}} & \dots & x_N, & x_{i_1} \end{pmatrix} \delta(x_{i_1} - x_{i_2}) \\ + \sum_{k=1}^N \sum_{\substack{n_1, n_2 \in \mathbb{N} \\ n_1 + n_2 = m_k}} \beta_{n_1 n_2} f_{N+1} \begin{pmatrix} m_1 & \dots & \widehat{m_k} & \dots & m_N & n_1 & n_2 \\ x_1 & \dots & \widehat{x_k} & \dots & x_N & x_k & x_k \end{pmatrix} \\ + (\gamma_{m_1} + \dots + \gamma_{m_N}) f_N \begin{pmatrix} m_1 & \dots & m_N \\ x_1 & \dots & x_N \end{pmatrix}. \end{aligned} \quad (4.13)$$

As in (4.10), the hat marks omitted arguments. Due to the symmetry of the wave function, the order of the arguments is irrelevant, so we put the new arguments replacing the omitted ones at the end of the list.

Whereas the operators  $\mathbf{H}_0$  and  $\mathbf{H}_1$  preserve the number  $N$  of particles, the Hamiltonian  $\mathbf{H}_2$  does not do so, due to the exchange terms  $\Psi_{m_1+m_2}^\dagger \Psi_{m_1} \Psi_{m_2}$  and  $\Psi_{m_1}^\dagger \Psi_{m_2}^\dagger \Psi_{m_1+m_2}$ . However, since  $\mathbf{H}_2$  commutes with  $\mathbf{H}_0$ , it preserves the mass  $M = m_1 + \dots + m_N$  instead. The original quantum-field-theoretical model splits thus into a series of quantum-mechanical

ones restricted to the eigenspaces  $\mathcal{F}_M$  of  $\mathbf{H}_0$  which we call mass- $M$  sectors. By (4.13), in each mass- $M$  sector the Hamiltonian  $\mathbf{H}_2$  is represented by a multicomponent differential operator with singular (delta-function) coefficients.

### 5. STRUCTURE OF MASS- $M$ SECTOR

To describe the structure of the mass- $M$  sector of the Fock space in more details we shall need a few definitions from combinatorics [20].

A *composition*  $\mathbf{m}$  of a nonnegative integer  $M \in \mathbb{N}$  is defined as a sequence  $\mathbf{m} = (m_1, \dots, m_N)$  of  $m_i \in \mathbb{N}$  such that  $m_1 + \dots + m_N \equiv |\mathbf{m}| = M$ . The number  $N = \ell(\mathbf{m})$  is called *length* of the composition, and  $M = |\mathbf{m}|$  its *weight*. The number of compositions of  $M$  equals  $2^{M-1}$ .

We introduce a partial order  $\succ$  on the set of compositions:  $\mathbf{m} \succ \widetilde{\mathbf{m}}$  means that  $\ell(\widetilde{\mathbf{m}}) = \ell(\mathbf{m}) - 1$  and  $\widetilde{\mathbf{m}}$  can be obtained from  $\mathbf{m}$  by replacing an adjacent pair  $(m_i, m_{i+1})$  for some  $i = 1, \dots, \ell(\mathbf{m}) - 1$  with  $m_i + m_{i+1}$ , that is

$$\begin{aligned} \mathbf{m} &= (m_1, \dots, m_{i-1}, m_i, m_{i+1}, m_{i+2}, \dots, m_N), \\ \widetilde{\mathbf{m}} &= (m_1, \dots, m_{i-1}, m_i + m_{i+1}, m_{i+2}, \dots, m_N). \end{aligned}$$

The set of compositions  $\mathbf{m}$  of  $M$  becomes then an ordered graph, with vertices  $\mathbf{m}$  and arrows pointing from  $\mathbf{m}$  to  $\widetilde{\mathbf{m}}$  if  $\mathbf{m} \succ \widetilde{\mathbf{m}}$ . The graph is topologically equivalent to an  $(M - 1)$ -dimensional hypercube having  $2^{M-1}$  vertices and  $(M - 1)2^{M-2}$  edges, as exemplified by Fig. 1. The vertex  $(1, \dots, 1)$  is the *source*, having no predecessors. Starting from it and travelling along the arrows one can reach any point of the hypercube in a variety of ways, terminating at the *sink*  $(M)$ .




FIGURE 1. Composition hypercubes

The mass- $M$  sector  $\mathcal{F}_M$  of our Fock space  $\mathcal{F}$  is the eigenspace of the mass operator  $\mathbf{H}_0$  corresponding to the eigenvalue  $M$ . It is spanned by the vectors

$$|f\rangle = \sum_{\mathbf{m}: |\mathbf{m}|=M} \frac{1}{N!} \int_{\mathbb{R}^N} dx_1 \dots dx_N f_N \left( \begin{matrix} \mathbf{m} \\ \mathbf{x} \end{matrix} \right) \prod_{j=1}^N \Psi_{m_j}^\dagger(x_j) |0\rangle \in \mathcal{F}_M, \quad (5.1)$$

with the norm

$$\|f\|^2 = \sum_{\mathbf{m}: |\mathbf{m}|=M} \frac{1}{N!} \int_{\mathbb{R}^N} dx_1 \dots dx_N \left| f_N \left( \begin{matrix} \mathbf{m} \\ \mathbf{x} \end{matrix} \right) \right|^2 \quad (5.2)$$

(here and below we always imply  $N = \ell(\mathbf{m})$ ).

The *Weyl alcove*  $\mathcal{W}_N$  is defined as

$$\mathcal{W}_N = \{ \mathbf{x} \in \mathbb{R}^N : x_1 < x_2 < \dots < x_N \}. \quad (5.3)$$

Due to the symmetry of the wave functions, the terms  $f_N(\boldsymbol{\xi})$  contribute to the sum (5.1) with the multiplicity  $N!$ . Consequently, one can replace the integration over  $\mathbb{R}^N$  in (5.1) and in (5.2) with the integration over  $\mathcal{W}_N$ , having adjusted the combinatorial coefficients:

$$|f\rangle = \sum_{\mathbf{m}: |\mathbf{m}|=M} \int_{\mathcal{W}_N} dx_1 \dots dx_N f_N \left( \begin{matrix} \mathbf{m} \\ \mathbf{x} \end{matrix} \right) \prod_{j=1}^N \Psi_{m_j}^\dagger(x_j) |0\rangle \in \mathcal{F}_M, \quad (5.4)$$

$$\|f\|^2 = \sum_{\mathbf{m}: |\mathbf{m}|=M} \int_{\mathcal{W}_N} dx_1 \dots dx_N \left| f_N \left( \begin{matrix} \mathbf{m} \\ \mathbf{x} \end{matrix} \right) \right|^2. \quad (5.5)$$

As a result,  $\mathcal{F}_M$  splits into the orthogonal sum

$$\mathcal{F}_M = \bigoplus_{\mathbf{m}: |\mathbf{m}|=M} \mathcal{F}_M^{\mathbf{m}}, \quad (5.6)$$

where  $\mathcal{F}_M^{\mathbf{m}} \simeq L^2(\mathcal{W}_N)$ .

The vectors of  $\mathcal{F}_M$  are thus identified with the collection of  $2^{M-1}$  functions  $f_N(\boldsymbol{\xi})$  labelled by the compositions  $\mathbf{m}$ , with arguments  $\mathbf{x} \in \mathcal{W}_N$ . The component  $f_N(\boldsymbol{\xi})$  describes a collection of  $N = \ell(\mathbf{m})$  one-dimensional particles with masses  $m_i$  and coordinates  $x_i$  ordered from left to right.

In the next section we shall rewrite the eigenvalue problem for the differential operator (4.13) with delta-function coefficients on functions  $f_N(\boldsymbol{\xi})$  with  $\mathbf{x} \in \mathbb{R}^N$  as an equivalent system of differential equations and boundary conditions for functions  $f_N(\boldsymbol{\xi})$  with  $\mathbf{x} \in \mathcal{W}_N$ .

## 6. FROM $\delta$ -FUNCTION TO BOUNDARY CONDITIONS

Replacing a delta-function term with boundary conditions is a standard trick, see e.g. [3, 7, 15], for the case of a scalar Bose-gas (quantum nonlinear Schrödinger equation). We only need to adapt the technique to the case of particles of different masses.

Let us analyse first a simple two-particle example. Let a function  $f(x_1, x_2)$  on  $\mathbb{R}^2$  satisfy the Schrödinger equation describing two particles of masses  $m_1$  and  $m_2$  and containing a singular inhomogeneous term (external source)

$$\left[ \left( -\frac{1}{m_1} \partial_{x_1}^2 - \frac{1}{m_2} \partial_{x_2}^2 \right) \right] f(x_1, x_2) + \sigma(x_1) \delta(x_1 - x_2) + \tau(x_1, x_2) = 0, \quad (6.1)$$

where the densities  $\sigma(x)$  and  $\tau(x_1, x_2)$  are assumed to be smooth functions.

Since  $\delta(x_1 - x_2)$  vanishes off the diagonal, one obtains immediately the differential equation “in the bulk”

$$-\left( \frac{1}{m_1} \partial_{x_1}^2 + \frac{1}{m_2} \partial_{x_2}^2 \right) f(x_1, x_2) + \tau(x_1, x_2) = 0, \quad x_1 \neq x_2. \quad (6.2)$$

To derive the boundary conditions on the diagonal  $x_1 = x_2$ , assume that the function  $f$  is piecewise smooth, meaning that it is given by two different expressions  $f^{(+)}(x_1, x_2)$  in the half-plane  $x_1 - x_2 > 0$  and  $f^{(-)}(x_1, x_2)$  in the half-plane  $x_1 - x_2 < 0$ . Furthermore,

both functions  $f^{(\pm)}$  are assumed to be smooth and defined in an open neighbourhood of the cut  $x_1 = x_2$ , the domain of each function extending thus beyond its native half-plane.

Introducing the step function

$$\theta(x) = \begin{cases} 1, & x > 0 \\ 0, & x < 0 \end{cases} \quad (6.3)$$

one can represent  $f$  as

$$f(x_1, x_2) = f^{(+)}(x_1, x_2)\theta(x_1 - x_2) + f^{(-)}(x_1, x_2)\theta(x_2 - x_1) \quad (6.4)$$

(we treat  $f$  as a measurable function defining a distribution, so its values on the zero-measure set  $x_1 = x_2$  are irrelevant and can be left undefined).

Substitute now (6.4) into (6.1) and perform the differentiations, using the identities valid for any smooth function  $\omega(x_1, x_2)$

$$\omega(x_1, x_2)\delta(x_1 - x_2) = \omega(x_1, x_1)\delta(x_1 - x_2),$$

$$\omega(x_1, x_2)\delta'(x_1 - x_2) = (\partial_{x_2}\omega)(x_1, x_1)\delta(x_1 - x_2) + \omega(x_1, x_1)\delta'(x_1 - x_2),$$

so that the coefficients at  $\theta$ - and  $\delta$ -functions in the resulting sum depend only on  $x_1$ . The coefficients at  $\theta(x_1 - x_2)$  and  $\theta(x_2 - x_1)$  then give the ‘‘bulk’’ equation (6.2) for  $f^{(+)}$  and  $f^{(-)}$ , respectively. The coefficient at  $\delta'(x_1 - x_2)$  gives the continuity condition

$$f^{(+)}(x_1, x_1) = f^{(-)}(x_1, x_1). \quad (6.5)$$

The coefficient at  $\delta(x_1 - x_2)$  gives, after a simplification using (6.5), the boundary condition

$$\left[ \frac{1}{m_1}\partial_{x_1}(f^{(+)} - f^{(-)}) + \frac{1}{m_2}\partial_{x_2}(-f^{(+)} + f^{(-)}) \right](x_1, x_1) = \sigma(x_1). \quad (6.6)$$

Let  $g(x \pm 0)$  denote the one-sided limiting values for  $g(x \pm \varepsilon)$  as  $\varepsilon \searrow 0$ . Then, one can rewrite the continuity condition (6.5) as

$$f(x + 0, x - 0) = f(x - 0, x + 0) \equiv f(x, x), \quad (6.7)$$

and the jump-of-transversal-derivative condition (6.6) as

$$\begin{aligned} - [(m_2 \partial_{x_1} - m_1 \partial_{x_2})f](x + 0, x - 0) + [(m_2 \partial_{x_1} - m_1 \partial_{x_2})f](x - 0, x + 0) \\ + m_1 m_2 \sigma(x) = 0. \end{aligned} \quad (6.8)$$

In (6.8), it is assumed that the differential operator is applied first to the function of two variables  $(x_1, x_2)$  and then the limit  $\varepsilon \searrow 0$  is taken in  $(x_1, x_2) = (x \pm \varepsilon, x \mp \varepsilon)$ .

The above argument works also in the multiparticle case since in the neighbourhood of the cut  $x_i = x_j$  the functions depend on the rest of the variables continuously, and we can ignore all remaining variables. Besides, when  $x$ 's are ordered as in (5.3) the only jump conditions to take into account are those for the adjacent particles  $x_i = x_{i+1}$ .

Consider the eigenvalue problem  $\mathbf{H}_2 |f\rangle = \lambda |f\rangle$ . Taking (4.13) and applying (6.2) we then obtain the set of bulk equations labelled by the vertices  $\mathbf{m}$  of the hypercube

$$\lambda f_N(\boldsymbol{\xi}) = \sum_{i=1}^N \left( -\frac{1}{m_i} \partial_{x_i}^2 + \gamma_{m_i} \right) f_N(\boldsymbol{\xi}) + \sum_{\mathbf{m}' \succ \mathbf{m}} \beta_{n_1 n_2} f_{N+1}(\boldsymbol{\xi}'). \quad (6.9)$$

The sum over the compositions  $\mathbf{m}'$  such that  $\mathbf{m}' \succ \mathbf{m}$  is in fact a double sum over the integers  $k, n_1, n_2$  with  $n_1 + n_2 = m_k$  that label the compositions and the associated vectors  $\boldsymbol{\xi}, \boldsymbol{\xi}'$  as

$$\begin{aligned}\boldsymbol{\xi} &= \begin{pmatrix} \mathbf{m} \\ \mathbf{x} \end{pmatrix} = \begin{pmatrix} m_1 & \dots & m_{k-1} & m_k & m_{k+1} & \dots & m_N \\ x_1 & \dots & x_{k-1} & x_k & x_{k+1} & \dots & x_N \end{pmatrix}, \\ \boldsymbol{\xi}' &= \begin{pmatrix} \mathbf{m}' \\ \mathbf{x}' \end{pmatrix} = \begin{pmatrix} m_1 & \dots & m_{k-1} & n_1 & n_2 & m_{k+1} & \dots & m_N \\ x_1 & \dots & x_{k-1} & x_k & x_k & x_{k+1} & \dots & x_N \end{pmatrix}.\end{aligned}\quad (6.10)$$

Note also that in (6.9) one does not need to distinguish the limits  $x_k \pm 0$  owing to the continuity of  $f_{N+1}$ .

Respectively, (6.8) produces the set of jump conditions labelled by the arrows of the hypercube pointing from  $\mathbf{m}$  that is the pairs

$$(m_1, \dots, m_k, m_{k+1}, \dots, m_N) \rightarrow (m_1, \dots, m_k + m_{k+1}, \dots, m_N), \quad k = 1, 2, \dots, N-1.$$

Applying (6.8) to (4.13) we get the jump of transversal derivative on the line  $x_k = x_{k+1}$ :

$$\begin{aligned}& - [(m_{k+1} \partial_{x_k} - m_k \partial_{x_{k+1}}) f_N] \begin{pmatrix} m_1 & \dots & m_{k-1} & m_k & m_{k+1} & m_{k+2} & \dots & m_N \\ x_1 & \dots & x_{k-1} & x_{k+1} + 0 & x_{k+1} - 0 & x_{k+2} & \dots & x_N \end{pmatrix} \\ & + [(m_{k+1} \partial_{x_k} - m_k \partial_{x_{k+1}}) f_N] \begin{pmatrix} m_1 & \dots & m_{k-1} & m_k & m_{k+1} & m_{k+2} & \dots & m_N \\ x_1 & \dots & x_{k-1} & x_k - 0 & x_k + 0 & x_{k+2} & \dots & x_N \end{pmatrix} \\ & + 2m_k m_{k+1} \beta_{m_k, m_{k+1}} f_{N-1} \begin{pmatrix} m_1 & \dots & m_{k-1} & \widehat{m}_k & \widehat{m}_{k+1} & m_{k+2} & \dots & m_N & m_k + m_{k+1} \\ x_1 & \dots & x_{k-1} & \widehat{x}_k & \widehat{x}_{k+1} & x_{k+2} & \dots & x_N & x_k \end{pmatrix} = 0.\end{aligned}\quad (6.11)$$

To express the result as a function of the arguments  $(x_1 < \dots < \widehat{x}_{k+1} < \dots < x_N)$  on a Weyl alcove it remains to use the symmetry of  $f_N$  and  $f_{N-1}$  to swap  $x_k \leftrightarrow x_{k+1}$  in the first term of (6.11) and, respectively, to rearrange the  $x$ 's in the increasing order in the last term. The resulting final form of the jump condition can be recast in a compact form by using the compositions  $\mathbf{m}', \mathbf{m}'' \succ \mathbf{m}$  of  $\mathbf{m}$  and the associated vectors  $\boldsymbol{\xi}', \boldsymbol{\xi}''$  and  $\boldsymbol{\xi}$ :

$$\begin{aligned}\boldsymbol{\xi}' &= \begin{pmatrix} \mathbf{m}' \\ \mathbf{x}' \end{pmatrix} = \begin{pmatrix} m_1 & \dots & m_k & m_{k+1} & \dots & m_N \\ x_1 & \dots & x_k - 0 & x_k + 0 & \dots & x_N \end{pmatrix}, \\ \boldsymbol{\xi}'' &= \begin{pmatrix} \mathbf{m}'' \\ \mathbf{x}'' \end{pmatrix} = \begin{pmatrix} m_1 & \dots & m_{k+1} & m_k & \dots & m_N \\ x_1 & \dots & x_k - 0 & x_k + 0 & \dots & x_N \end{pmatrix}, \\ \boldsymbol{\xi} &= \begin{pmatrix} \mathbf{m} \\ \mathbf{x} \end{pmatrix} = \begin{pmatrix} m_1 & \dots & m_{k-1} & m_k + m_{k+1} & m_{k+2} & \dots & m_N \\ x_1 & \dots & x_{k-1} & x_k & x_{k+2} & \dots & x_N \end{pmatrix}.\end{aligned}\quad (6.12)$$

$$\begin{aligned}[(m_k \partial_{x_k} - m_{k+1} \partial_{x_{k+1}}) f_N](\boldsymbol{\xi}'') + [(m_{k+1} \partial_{x_k} - m_k \partial_{x_{k+1}}) f_N](\boldsymbol{\xi}') \\ + 2m_k m_{k+1} \beta_{m_k, m_{k+1}} f_{N-1}(\boldsymbol{\xi}) = 0.\end{aligned}\quad (6.13)$$

Note that the swapping  $m_k \leftrightarrow m_{k+1}$  produces an identical equation. Also, for  $m_k = m_{k+1}$  the first and the second term in (6.13) are equal.

To conclude, the eigenvalue problem for the Hamiltonian  $\mathbf{H}_2$  in the sector of mass  $M$  is now formulated in terms of a set of functions  $f_N(\boldsymbol{\xi})$  labelled by compositions  $\mathbf{m}$  of  $M$  with length  $N = \ell(\mathbf{m})$  defined on Weyl alcoves  $\mathcal{W}_N$ . The equations for  $f_N(\boldsymbol{\xi})$  are divided into two classes: the bulk differential equations of 2nd order (6.9) labelled by the vertices  $\mathbf{m}$  of the compositions hypercube, and the jump conditions (6.13) for the transversal

derivatives labelled by the edges of the compositions hypercube that correspond to the merging  $(m_k, m_{k+1}) \rightarrow (m_k + m_{k+1})$  of the adjacent particles.

### 7. SOLUTION IN THE SECTOR $M = 2$

The number  $M = 2$  admits two compositions:  $1+1$  and  $2$ , see Fig. 1a. Respectively, the mass-2 sector splits as  $\mathcal{F}_2 = \mathcal{F}_2^{(11)} \oplus \mathcal{F}_2^{(2)}$ , so that any vector  $|f\rangle \in \mathcal{F}_2$  can be represented as

$$|f\rangle = \int_{x_1 < x_2} dx_1 dx_2 f_2 \begin{pmatrix} 1 & 1 \\ x_1 & x_2 \end{pmatrix} \Psi_1^\dagger(x_1) \Psi_1^\dagger(x_2) |0\rangle + \int_{-\infty}^{\infty} dx_1 f_1 \begin{pmatrix} 2 \\ x_1 \end{pmatrix} \Psi_2^\dagger(x_1) |0\rangle, \quad (7.1)$$

$$\|f\|^2 = \int_{x_1 < x_2} dx_1 dx_2 \left| f_2 \begin{pmatrix} 1 & 1 \\ x_1 & x_2 \end{pmatrix} \right|^2 + \int_{-\infty}^{\infty} dx_1 \left| f_1 \begin{pmatrix} 2 \\ x_1 \end{pmatrix} \right|^2. \quad (7.2)$$

The general bulk equation (6.9) produces two bulk equations corresponding to the vertices of the graph in Fig. 1a:

$$\lambda f_2 \begin{pmatrix} 1 & 1 \\ x_1 & x_2 \end{pmatrix} = (-\partial_{x_1}^2 - \partial_{x_2}^2 + 2\gamma_1) f_2 \begin{pmatrix} 1 & 1 \\ x_1 & x_2 \end{pmatrix}, \quad (7.3a)$$

$$\lambda f_1 \begin{pmatrix} 2 \\ x_1 \end{pmatrix} = \left(-\frac{1}{2}\partial_{x_1}^2 + \gamma_2\right) f_1 \begin{pmatrix} 2 \\ x_1 \end{pmatrix} + \beta_{11} f_2 \begin{pmatrix} 1 & 1 \\ x_1 & x_1 \end{pmatrix}. \quad (7.3b)$$

Respectively, equation (6.13) produces the jump condition corresponding to the single arrow  $(11) \rightarrow (2)$  of the graph in Fig. 1a:

$$2[(\partial_{x_1} - \partial_{x_2})f_2] \begin{pmatrix} 1 & 1 \\ x_1 & x_1 \end{pmatrix} + 2\beta_{11} f_1 \begin{pmatrix} 2 \\ x_1 \end{pmatrix} = 0 \quad (7.3c)$$

(the two terms with derivatives coincide due to the symmetry  $m_1 = m_2 = 1$ ).

In the spirit of Bethe Ansatz [3, 7], we look for a solution of the boundary problem (7.3) in the subsector  $\mathbf{m} = (11)$  as a linear combination of plain waves: the incoming one  $e^{i(u_2 x_1 + u_1 x_2)}$  and the scattered one  $e^{i(u_1 x_1 + u_2 x_2)}$ , with the scattering coefficient  $S_{21}$ :

$$f_2 \begin{pmatrix} 1 & 1 \\ x_1 & x_2 \end{pmatrix} = e^{i(u_2 x_1 + u_1 x_2)} + S_{21} e^{i(u_1 x_1 + u_2 x_2)}, \quad x_1 < x_2. \quad (7.4a)$$

The jump condition (7.3c) implies then that the wave function  $f_1$  in the subsector  $\mathbf{m} = (2)$  has to be the exponent  $e^{i(u_1 + u_2)x_1}$ , up to a coefficient  $R$ :

$$f_1 \begin{pmatrix} 2 \\ x_1 \end{pmatrix} = R e^{i(u_1 + u_2)x_1}. \quad (7.4b)$$

Substituting the Ansatz (7.4) into (7.3) we obtain, respectively, the bulk-11 equation:

$$u_1^2 + u_2^2 + 2\gamma_1 = \lambda, \quad (7.5a)$$

the bulk-2 equation:

$$\left(\frac{1}{2}(u_1 + u_2)^2 + \gamma_2\right) R + \beta_{11}(1 + S_{21}) = \lambda R, \quad (7.5b)$$

and the jump  $(11) \rightarrow (2)$  equation:

$$i(u_2 - u_1) + i(u_1 - u_2)S_{21} + \beta_{11}R = 0. \quad (7.5c)$$

The system of three linear equations (7.5) for  $\lambda$ ,  $S_{21}$ ,  $R$  is easily solved. Equation (7.5a) gives immediately the value of  $\lambda$ , and the two remaining equations produce the answer

$$S_{21} = S(u_2 - u_1), \quad S(u) = -\frac{P(iu)}{P(-iu)} \quad (7.6)$$

where  $P$  is the cubic polynomial

$$P(v) = v^3 + (2\gamma_2 - 4\gamma_1)v - 2\beta_{11}^2, \quad (7.7)$$

or substituting  $\beta_{11} = \sqrt{2}\beta$ ,  $\gamma_1 = \gamma$ ,  $\gamma_2 = 8\gamma$  from (3.10),

$$P(v) = v^3 + 12\gamma v - 4\beta^2. \quad (7.8)$$

Respectively,

$$R = \frac{4i\beta_{11}u_{21}}{P(-iu_{21})} = \frac{4\sqrt{2}i\beta u_{21}}{P(-iu_{21})}, \quad u_{21} \equiv u_2 - u_1. \quad (7.9)$$

As befits a Galilei-invariant theory, the  $S$ -matrix is invariant under the simultaneous translations  $u_a \mapsto u_a + c$ ,  $a = 1, 2$ .

Note that the sum of the zeroes of the cubic polynomial  $P$  is 0 due to the absence of the quadratic term. Since  $P$  has real coefficients and negative free term  $-4\beta^2$ , it has exactly one positive root. The two remaining zeroes lie in the left half-plane, and their position is determined by the discriminant  $D = -432(16\gamma^3 + \beta^4)$ . For  $D < 0$  they are complex-conjugated, and for  $D > 0$  they are both real negative, as shown on Fig. 2.

It is tempting to call the case  $D < 0$ , or  $16\gamma^3 + \beta^4 > 0$  the quantum KP-I equation, and  $D > 0$ , or  $16\gamma^3 + \beta^4 < 0$  the quantum KP-II equation. Note that the boundary between the two cases is not  $\gamma = 0$  as in the classical case but  $16\gamma^3 + \beta^4 = 0$  when  $P$  has a double negative zero, the term  $\beta^4$  playing the role of a quantum correction. It remains disputable what to call the *dispersionless* quantum KP: either  $16\gamma^3 + \beta^4 = 0$ , or  $\gamma = 0$  that corresponds to  $P(u) = u^3 - 4\beta^2$ , the zeroes forming an equilateral triangle.

The corresponding scattering coefficient  $S(u)$  given by (7.6) is a rational function having three zeroes and three poles. Their positions ( $\circ$  for zeroes,  $\bullet$  for poles), depending on  $D$ , are shown on Fig. 3





FIGURE 2. Zeroes of  $P(u)$

## 8. BETHE ANSATZ IN SECTOR $\mathcal{F}_M^{(1\dots 1)}$

The known multiparticle integrable models share two common features: the absence of diffraction (preservation of the asymptotic momenta of the particles after collision), and the factorisation of the multiparticle  $S$ -matrix into two-particles factors [3, 7]. For the models with a delta-function interaction, like the quantum nonlinear Schrödinger


 FIGURE 3. Zeroes and poles of  $S(u)$ 

model [2, 15], such behaviour is manifested via the coordinate Bethe Ansatz [3, 7], or the assumption that the eigenfunction can be written as a sum of plane waves with the coefficients differing by a two-particles  $S$ -factor when any pair of momenta permutes.

In this section, we shall describe the Bethe Ansatz for our model in the subsector  $\mathcal{F}_M^{(1\dots 1)}$  of  $\mathcal{F}_M$  containing  $M$  particles of unit mass and corresponding to the composition  $\mathbf{m} = (1, \dots, 1)$  of  $M$ .

Let  $\mathfrak{S}_{[1,M]}$  be the permutation group of  $(1, \dots, M)$ . Let  $\mathbf{u} \equiv (u_1, \dots, u_M)$  be the vector of momenta, and  $\mathbf{v} \equiv i\mathbf{u}$ . Define the action of a permutation  $\mathfrak{s} = (\mathfrak{s}_1, \dots, \mathfrak{s}_M) \in \mathfrak{S}_{[1,M]}$  on functions of  $\mathbf{v}$  by substitutions  $\mathfrak{s} : v_j \mapsto v_{\mathfrak{s}_j}$ . Then, for a plane wave

$$\exp(\mathbf{v} \cdot \mathbf{x}) \equiv \exp(v_1 x_1 + \dots + v_M x_M) \quad (8.1)$$

we have

$$\mathfrak{s}(\exp(\mathbf{v} \cdot \mathbf{x})) = (\exp(\mathfrak{s}(\mathbf{v}) \cdot \mathbf{x})) = \exp(v_{\mathfrak{s}_1} x_1 + \dots + v_{\mathfrak{s}_M} x_M). \quad (8.2)$$

We choose to normalise the Bethe wave function by making the coefficients at the plane waves polynomial in  $v_j$ . Such a normalisation was proposed first for the quantum nonlinear Schrödinger equation in [7], *c.f.* Chapter 4, eq. (4.8), see also [3], Chapter 1, eq. (1.24). Such a choice has the advantage of allowing for algebraic manipulations with polynomials rather than rational functions.

**Conjecture 1** (Bethe Ansatz for  $\mathbf{m} = (1, \dots, 1)$ ). *The  $\mathcal{F}_M^{(1\dots 1)}$  component of the eigenfunction of  $\mathbf{H}_2$  can be chosen for  $\mathbf{x} \in \mathcal{W}_{1\dots 1}$  as*

$$f_M \begin{pmatrix} 1 & \dots & 1 \\ x_1 & \dots & x_M \end{pmatrix} = \sum_{\mathfrak{s} \in \mathfrak{S}_{[1,M]}} \text{sgn}(\mathfrak{s}) \left( \prod_{j < k} P(v_{\mathfrak{s}_k} - v_{\mathfrak{s}_j}) \right) \mathfrak{s}(\exp(\mathbf{v} \cdot \mathbf{x})), \quad (8.3)$$

where  $\text{sgn}(\mathfrak{s})$  is the sign of the permutation  $\mathfrak{s}$ , and the polynomial  $P(u)$  is given by (7.8).

By construction, the Bethe wave function (8.3) is antisymmetric in the momenta  $\mathbf{v}$ . Note that the ratio of the coefficients for two plane waves in (8.3) differing by a transposition of two adjacent momenta  $v_{\mathfrak{s}_j}$  and  $v_{\mathfrak{s}_{j+1}}$  is  $S(v_{\mathfrak{s}_j} - v_{\mathfrak{s}_{j+1}})$  due to (7.6), as expected.

Two more conventional wave functions  $f^{(\text{in})}$  and  $f^{(\text{out})}$  having unitary factors at the plane waves are defined from

$$f_M \begin{pmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_M \end{pmatrix} = f^{(\text{in})} (-1)^{M(M-1)/2} \prod_{j < k} P(v_j - v_k) = f^{(\text{out})} \prod_{j < k} P(v_k - v_j). \quad (8.4)$$

Assuming that  $u_1 < \dots < u_M$  and  $x_1 < \dots < x_M$ , one can interpret  $\mathbf{t}(\exp(i\mathbf{u} \cdot \mathbf{x}))$ , with  $\mathbf{t} \equiv (M, \dots, 1)$ , as the incident wave, and  $\exp(i\mathbf{u} \cdot \mathbf{x})$  as the outgoing scattered wave, corresponding to the ordering of the particles carrying the momenta  $u_i$  as  $t \rightarrow -\infty$  or, respectively,  $t \rightarrow +\infty$ .

The function  $f^{(\text{in})}$  is then normalised by the unit coefficient at the incoming wave  $\mathbf{t}(\exp(i\mathbf{u} \cdot \mathbf{x}))$ , and  $f^{(\text{out})}$  by the unit coefficient at the outgoing wave  $\exp(i\mathbf{u} \cdot \mathbf{x})$ .

The functions  $f^{(\text{in})}$  and  $f^{(\text{out})}$  differ only by the factor (multiparticle  $S$ -matrix)

$$f^{(\text{out})} = \mathbb{S} f^{(\text{in})}, \quad \mathbb{S} = \prod_{1 \leq j < k \leq M} S(u_j - u_k) \quad (8.5)$$

that is factorised into a product of the factors corresponding to all two-particle collisions, in the spirit of Bethe Ansatz.

By antisymmetry in  $\mathbf{v}$ , the whole wave function (8.3) can be restored from a single term containing  $\exp(\mathbf{v} \cdot \mathbf{x})$ . Let  $1 \leq a < b \leq M$ . For a subsegment  $(a, \dots, b) \subset (1, \dots, M)$  define the polynomial

$$\mathbb{P}_{[a,b]}(\mathbf{v}) \equiv \prod_{a \leq j < k \leq b} P(v_k - v_j) \quad (8.6)$$

and the linear operator  $\mathfrak{P}_{[a,b]}$  acting on functions of  $\mathbf{v} = (v_1, \dots, v_M)$  by antisymmetrisation, with the weight factor  $\mathbb{P}_{[a,b]}$ , in respect to the group  $\mathfrak{S}_{[a,b]}$  of permutations of  $(a, \dots, b)$  acting on  $(v_a, \dots, v_b)$

$$\mathfrak{P}_{[a,b]} : g(\mathbf{v}) \mapsto \sum_{\mathfrak{s} \in \mathfrak{S}_{[a,b]}} \text{sgn}(\mathfrak{s}) \mathbb{P}_{[a,b]}(\mathfrak{s}(\mathbf{v})) g(\mathfrak{s}(\mathbf{v})). \quad (8.7)$$

The  $\mathfrak{P}_{[a,b]}$  operator for a subsegment of  $(1, \dots, M)$  will be used only in the Appendices. In the main text we use the abbreviation  $\mathfrak{P} \equiv \mathfrak{P}_{[1,M]}$ .

In terms of the operator  $\mathfrak{P}$ , the formula (8.3) for the Bethe function simplifies to

$$f_M \begin{pmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_M \end{pmatrix} = \mathfrak{P}(\exp(\mathbf{v} \cdot \mathbf{x})). \quad (8.8)$$

## 9. BETHE ANSATZ IN GENERIC SECTOR

The jump conditions (6.13) can be viewed as recurrence relations allowing one to obtain the wave function  $f_N(\boldsymbol{\xi})$  by differentiating the wave functions  $f_{N+1}(\boldsymbol{\xi})$  corresponding to the preceding compositions (in the sense of the relation  $\succ$ ). Thus, starting from the source  $\mathbf{m} = (1, \dots, 1)$  and travelling along the arrows of the composition graph one can in principle obtain the wave functions for all the remaining compositions of  $M$ . The problem is, however, that different paths produce, in principle, different expressions, and one ends with a bunch of *consistency conditions* for the wave function. To show that the Bethe Ansatz works at all one has to prove that those conditions have a joint solution. Besides, there remain the bulk conditions (6.9) that also have to be verified. The differentiation  $\partial_{x_i}$  acting on the exponent in (8.8) are replaced by  $v_i$ , and the resulting consistency equations

take the form of a (very overdetermined) set of algebraic equations for the coefficients of the Bethe wave functions.

To any composition  $\mathbf{m} = (m_1, \dots, m_N)$  of  $M$  of length  $\ell(\mathbf{m}) = N$ , there corresponds a split of the sequence of the momenta  $v_i$

$$\mathbf{v} = (v_1, \dots, v_M) = (\mathbf{w}_1^{\mathbf{m}}; \dots; \mathbf{w}_N^{\mathbf{m}}) \quad (9.1)$$

into the consecutive segments  $\mathbf{w}_j^{\mathbf{m}}$  of respective length  $m_j$ , so that

$$\mathbf{w}_j^{\mathbf{m}} = (v_{m_1+\dots+m_{j-1}+1}, \dots, v_{m_1+\dots+m_j}), \quad j = 1, \dots, N \quad (9.2)$$

or

$$(\mathbf{w}_j^{\mathbf{m}})_i = v_{m_1+\dots+m_{j-1}+i}, \quad i = 1, \dots, m_j. \quad (9.3)$$

At the vertex  $\mathbf{m}$ , the original coordinates  $(x_1, \dots, x_M)$  merge into consecutive groups of length  $m_i$ :

$$(x_1, \dots, x_M) \mapsto \mathbf{X}^{\mathbf{m}} = (\overbrace{x_1, \dots, x_1}^{m_1}, \overbrace{x_2, \dots, x_2}^{m_2}, \dots, \overbrace{x_N, \dots, x_N}^{m_N}), \quad (9.4)$$

and now we set  $\mathbf{x} = (x_1, \dots, x_N)$ .

Let  $\langle \mathbf{w} \rangle$  denote the sum of the components of a vector  $\mathbf{w}$ , e.g.  $\langle \mathbf{v} \rangle = v_1 + \dots + v_M$ . Let also

$$\mathbf{W}^{\mathbf{m}} = (\langle \mathbf{w}_1^{\mathbf{m}} \rangle, \dots, \langle \mathbf{w}_N^{\mathbf{m}} \rangle) \quad (9.5)$$

so that

$$\mathbf{W}^{\mathbf{m}} \cdot \mathbf{x} = \langle \mathbf{w}_1^{\mathbf{m}} \rangle x_1 + \dots + \langle \mathbf{w}_N^{\mathbf{m}} \rangle x_N, \quad (9.6)$$

and

$$\mathbf{v} \cdot \mathbf{X}^{\mathbf{m}} = \mathbf{W}^{\mathbf{m}} \cdot \mathbf{x}. \quad (9.7)$$

**Conjecture 2** (Bethe Ansatz for generic  $\mathbf{m}$ ). *The Bethe eigenfunction in the generic subsector  $\mathcal{F}_M^{\mathbf{m}}$  can be written in the form*

$$f_N \begin{pmatrix} m_1 & \dots & m_N \\ x_1 & \dots & x_N \end{pmatrix} = \mathfrak{P}(Q^{\mathbf{m}}(\mathbf{v}) \exp(\mathbf{W}^{\mathbf{m}} \cdot \mathbf{x})), \quad (9.8)$$

where  $Q^{\mathbf{m}}(\mathbf{v})$  is a polynomial in  $\mathbf{v}$ . In particular,  $Q^{(1\dots 1)}(\mathbf{v}) \equiv 1$ .

If  $\mathfrak{P} : g \mapsto 0$  for some function  $g(\mathbf{v})$  we shall say that  $g(\mathbf{v})$  is  $\mathfrak{P}$ -reducible and write  $g \stackrel{\mathfrak{P}}{\equiv} 0$ . If  $g_1 - g_2 \stackrel{\mathfrak{P}}{\equiv} 0$  we shall say that  $g_1$  and  $g_2$  are  $\mathfrak{P}$ -equivalent and write  $g_1 \stackrel{\mathfrak{P}}{\equiv} g_2$ . As a consequence, all quantities under the sign of  $\mathfrak{P}$  are defined only up to  $\mathfrak{P}$ -equivalence.

Consider the bulk equation (6.9).

Since the vertex  $\mathbf{m} = (1, \dots, 1)$  is the source of the composition graph, *c.f.* Fig. 1, having no predecessors, the corresponding bulk equation (6.9) contains no  $\beta$ -terms and is obviously satisfied by the Ansatz (8.3) producing the eigenvalue  $\lambda$  of  $\mathbf{H}_2$

$$\lambda = -v_1^2 - \dots - v_M^2 + M\gamma_1. \quad (9.9)$$

Let  $K(\mathbf{w})$  be the sum of the squares of the components of a vector  $\mathbf{w}$ . Recalling that  $\gamma_1 = \gamma$ , by (3.10), we have  $\lambda = -K(\mathbf{v}) + M\gamma_1$  or, splitting the sum into groups of size  $m_j$ ,

$$\lambda = \sum_{j=1}^N (m_j \gamma - K(\mathbf{w}_j^{\mathbf{m}})). \quad (9.10)$$

Each  $\partial_{x_j}$  in (6.9) is replaced by  $\langle \mathbf{w}_j^m \rangle$ . Moving the  $\lambda$  term into the right-hand-side we find that the coefficient in front of  $f_N(\boldsymbol{\xi})$  produces the factor

$$\sum_{j=1}^N \tilde{K}(\mathbf{w}_j^m) + (m_j^3 - m_j)\gamma, \quad (9.11)$$

where we use the notation

$$\tilde{K}(\mathbf{w}_j^m) = K(\mathbf{w}_j^m) - \frac{\langle \mathbf{w}_j^m \rangle^2}{m_j} \quad (9.12)$$

for the ‘‘kinetic energy’’ of the cluster  $\mathbf{w}_j^m$  reduced w.r.t. the center-of-mass. Note that  $\tilde{K}(\mathbf{w}_j^m)$  is invariant under translations  $\mathbf{w}_j^m \mapsto \mathbf{w}_j^m + (c, \dots, c)$ , as a manifestation of the Galilei invariance.

Upon summing up over the compositions  $\mathbf{m}' \succ \mathbf{m}$  introduced in (6.10), the bulk equation (6.9) takes finally the form

$$\left[ \left( \sum_{j=1}^N \tilde{K}(\mathbf{w}_j^m) + (m_j^3 - m_j)\gamma \right) Q^m(\mathbf{v}) + \sum_{\mathbf{m}' \succ \mathbf{m}} \beta_{n_1 n_2} Q^{\mathbf{m}'}(\mathbf{v}) \right] \exp(\mathbf{W}^m \cdot \mathbf{x}) \stackrel{\mathfrak{P}}{=} 0. \quad (9.13)$$

After performing differentiations of the exponent, the jump equation (6.13) can be recast in terms of the compositions  $\mathbf{m}', \mathbf{m}''$  of  $\mathbf{m}$  introduced in (6.12)

$$\left[ V_{m_k m_{k+1}}(\mathbf{w}_k^m) \cdot Q^{\mathbf{m}'}(\mathbf{v}) + V_{m_{k+1} m_k}(\mathbf{w}_k^m) \cdot Q^{\mathbf{m}''}(\mathbf{v}) + 2m_k m_{k+1} \beta_{m_k, m_{k+1}} Q^m(\mathbf{v}) \right] \exp(\mathbf{W}^m \cdot \mathbf{x}) \stackrel{\mathfrak{P}}{=} 0. \quad (9.14)$$

Here, given an  $n_1 + n_2$ -dimensional vector  $\mathbf{u} = (u_1, \dots, u_{n_1+n_2})$ , we have defined

$$V_{n_1 n_2}(\mathbf{u}) = n_2(u_1 + \dots + u_{n_1}) - n_1(u_{n_1+1} + \dots + u_{n_1+n_2}). \quad (9.15)$$

**Conjecture 3** (Factorisation property). *The polynomial  $Q^m(\mathbf{v})$  in (9.8) can be chosen, up to a  $\mathfrak{P}$ -equivalent expression, in the factorised form*

$$Q^m(\mathbf{v}) = \prod_{k=1}^N Q^{(m_k)}(\mathbf{w}_k^m). \quad (9.16)$$

A justification of the above conjecture is presented in Appendix A.

## 10. BETHE ANSATZ IN SECTOR $\mathcal{F}_M^{(M)}$

Conjecture 3 suggests that it is sufficient to analyze the consistency equations only for the compositions of unit length  $\mathbf{m} = (M)$ ,  $N = 1$ . In this case, the exponent

$$\exp(\mathbf{W}^m \cdot \mathbf{x}) = \exp((v_1 + \dots + v_M)x_1)$$

becomes completely symmetric and can be factored out from under  $\mathfrak{P}$ . The equations for  $Q^{(M)}(\mathbf{v})$  are thus purely polynomial.

For the bulk equation (9.13) we have now

$$\begin{aligned} \mathbf{m}' &= (n_1, n_2), \quad n_1 + n_2 = M, \quad \beta_{n_1 n_2} = \sqrt{n_1 n_2 M} \beta \\ \mathbf{w}_1^{\mathbf{m}'} &= (v_1, \dots, v_m), \quad \mathbf{w}_2^{\mathbf{m}'} = (v_{m+1}, \dots, v_M), \end{aligned}$$

and the equation takes form

$$\begin{aligned} & (\tilde{K}(\mathbf{v}) + (M^3 - M)\gamma) Q^{(M)}(v_1, \dots, v_M) \\ & + \beta \sum_{\substack{n_1, n_2 \geq 1 \\ n_1 + n_2 = M}} \sqrt{n_1 n_2 M} Q^{(n_1)}(v_1, \dots, v_{n_1}) Q^{(n_2)}(v_{n_1+1}, \dots, v_M) \stackrel{\mathfrak{P}}{=} 0. \end{aligned} \quad (10.1)$$

For the jump equation (9.14) we consider the compositions

$$\mathbf{m}' = (m_1, m_2), \quad \mathbf{m}'' = (m_2, m_1), \quad \mathbf{m} = (m_1 + m_2),$$

and denote  $\mathbf{v} = (v_1, \dots, v_{m_1+m_2})$ , what recasts the equation in the form

$$\begin{aligned} & V_{m_1 m_2}(\mathbf{v}) Q^{(m_1)}(v_1, \dots, v_{m_1}) Q^{(m_2)}(v_{m_1+1}, \dots, v_{m_1+m_2}) \\ & + V_{m_2 m_1}(\mathbf{v}) Q^{(m_2)}(v_1, \dots, v_{m_2}) Q^{(m_1)}(v_{m_2+1}, \dots, v_{m_1+m_2}) \\ & + 2m_1 m_2 \beta_{m_1 m_2} Q^{(m_1+m_2)}(v_1, \dots, v_{m_1+m_2}) \stackrel{\mathfrak{P}}{=} 0. \end{aligned} \quad (10.2)$$

The equations (10.1) and (10.2) together with  $Q^{(1)}(v) = 1$  constitute the complete set of conditions for the polynomials  $Q^{(M)}(v_1, \dots, v_M)$ ,  $M = 1, 2, \dots$ , defined up to  $\mathfrak{P}$ -equivalence. Extensive computer experiments have led us to the following explicit solution to the equations (10.1) and (10.2).

**Conjecture 4** (Solution). *Set  $Q^{(1)}(v) = 1$  and for  $M \geq 2$  define the polynomial  $Q^{(M)}(\mathbf{v})$  as the following homogeneous polynomials of degree  $M - 1$*

$$Q^{(M)}(\mathbf{v}) = \frac{2\sqrt{M}}{M!(M-1)} (2\beta)^{1-M} \sum_{1 \leq i < j \leq M} (-1)^{j-i} \binom{M-1}{j-i-1} (v_i - v_j)^{M-1} \quad (10.3)$$

*invariant under translations  $v_i \mapsto v_i + c$ . Then such  $Q^{(M)}(\mathbf{v})$  satisfy all the equations (10.1) and (10.2).*

Note that  $Q^{(M)}(\mathbf{v})$  do not contain coupling constants  $\beta, \gamma$  that are hidden inside the  $\mathfrak{P}$ -operator.

Conjecture 4 has been confirmed by means of computer algebra for  $M \leq 8$ . In fact, instead of verifying Conjectures 4 literally, we have verified a stronger Conjecture 5, see Appendix B.

## 11. DISCUSSION

As a test of quantum integrability of the system, we have demonstrated consistency of the Bethe Ansatz for  $M \leq 8$ . This is a pretty convincing though not conclusive result. A rigorous proof of Conjecture 4, or superseding Conjecture 5 remains an open problem. The  $\mathfrak{P}$  operator, and the notions of 2- and 3-reducibility introduced in Appendix B seem to be new combinatorial objects that might be of interest for themselves.

An alternative way to establish quantum integrability could be provided through the Algebraic Bethe Ansatz [3] based on quantum Lax operator and  $R$ -matrix. That would also help to identify the underlying quantum algebra. The work in this direction is in progress.

Except for  $M = 2$ , we have not pursued a comprehensive study of the orthogonality and completeness of the Bethe eigenfunctions, neither of the structure of bound states.

In the case of the quantum nonlinear Schrödinger equation (delta-function Bose gas) it is known that the bound states of the quantum model correspond in the classical limit to the solitons of the classical model [13]. It would be interesting to study a similar correspondence for the KP-model.

The model we study is associated with a cubic polynomial  $P$  with zero sum of the roots (7.7) through which the two-particle  $S$ -matrix is expressed (7.6) and, in turn, the factorised multiparticle  $S$ -matrix. The question arises what possible QFT models could be associated with polynomials  $P$  of higher degree, or without the restriction on the roots. In [11] the properties of Bethe equations associated with a generic polynomial  $P$  were studied in an abstract way, without clarifying the nature of the corresponding QFT. In a recent paper [16] a possible example of a model of that class is proposed.

The model we study is nonrelativistic and Galilei invariant. It appears that it corresponds to a nonrelativistic limit of a relativistic integrable model known as affine  $A_{N-1}$  Toda field theory [1, 4] and given by the Lagrangian

$$\mathcal{L} = \frac{1}{2} \sum_{i=1}^N \partial_\mu \varphi_i \cdot \partial^\mu \varphi_i - \frac{2M^2}{\beta^2} \sum_{i=1}^N \exp \left[ \frac{\beta}{\sqrt{2}} (\varphi_i - \varphi_{i+1}) \right]. \quad (11.1)$$

Indeed, the  $S$ -matrix for a pair of main particles of the Toda FT is conjectured in [1] to be

$$S_{11}(\theta) = \frac{\sinh \left( \frac{\theta}{2} + \frac{i\pi}{N} \right) \sinh \left( \frac{\theta}{2} - \frac{i\pi}{N} + i\frac{b}{2} \right) \sinh \left( \frac{\theta}{2} - i\frac{b}{2} \right)}{\sinh \left( \frac{\theta}{2} - \frac{i\pi}{N} \right) \sinh \left( \frac{\theta}{2} + \frac{i\pi}{N} - i\frac{b}{2} \right) \sinh \left( \frac{\theta}{2} + i\frac{b}{2} \right)}. \quad (11.2)$$

Upon carrying out the rescaling

$$\theta = \frac{2\kappa^{-1}\pi u}{N} \quad \text{and} \quad b = \frac{2\kappa^{-1}\tau\pi}{N} \quad (11.3)$$

and then sending  $N \rightarrow +\infty$  one obtains the rational degeneration

$$\lim_{N \rightarrow +\infty} S_{11}(\theta) = \tilde{S}_{11}(u) = \frac{u^3 + u(\kappa^2 + \tau^2 - \kappa\tau) - i\kappa\tau(\kappa - \tau)}{u^3 + u(\kappa^2 + \tau^2 - \kappa\tau) + i\kappa\tau(\kappa - \tau)}. \quad (11.4)$$

Thus choosing  $\tau$  and  $\kappa$  such that  $\kappa^2 + \tau^2 - \kappa\tau = -12\gamma$  and  $\kappa\tau(\kappa - \tau) = 4\beta^2$  one obtains the scalar  $S$ -matrix of qKP, or more precisely, qKP-II, since  $0 < b < 2\pi/N$ :

$$\tilde{S}_{11}(u) = S_{\text{qKP}}(u) = \frac{u^3 - 12\gamma u - 4i\beta^2}{u^3 - 12\gamma u + 4i\beta^2}. \quad (11.5)$$

**Acknowledgements:** We thank Benoît Vicedo for valuable discussions at the initial stages of the project. A.T. thanks the EPSRC for funding under the First Grant project EP/K014412/1, and the STFC for support under the Consolidated Grant project nr. ST/L000490/1.

## APPENDIX A.

In the appendices we shall use again the non-abbreviated notation  $\mathbb{P}_{[a,b]}$  (8.6) and  $\mathfrak{P}_{[a,b]}$  (8.7) for a subsegment  $(a, \dots, b) \subset (1, \dots, M)$ .

Let us state a couple of elementary properties of the operator  $\mathfrak{P}_{[a,b]}$ . It is assumed below that  $\mathbf{v} = (v_1, \dots, v_M)$ .

**Lemma 1.** *If a function  $F(\mathbf{v})$  is  $\mathfrak{P}_{[a,b]}$ -reducible, and a function  $G(\mathbf{v})$  is symmetric under permutations  $\mathfrak{S}_{[a,b]} \subset \mathfrak{S}_{[1,M]}$  then the product  $F(\mathbf{v})G(\mathbf{v})$  is also  $\mathfrak{P}_{[a,b]}$ -reducible.*

*Proof.* Since  $G(\mathbf{v})$  is invariant under  $\mathfrak{S}_{[a,b]}$  it is factored out from the sum over  $\mathfrak{S}_{[a,b]}$  in (8.7).  $\blacksquare$

**Lemma 2.** *If a function  $F(\mathbf{v})$  is  $\mathfrak{P}_{[a,b]}$ -reducible then  $F(\mathbf{v})$  is also  $\mathfrak{P}_{[1,M]}$ -reducible.*

*Proof.* The product  $\mathbb{P}_{[1,M]}$  factorises as  $\mathbb{P}_{[1,M]} = \mathbb{P}_{[a,b]} \cdot \tilde{\mathbb{P}}$  where the complementary factor  $\tilde{\mathbb{P}}$  is  $\mathfrak{S}_{[a,b]}$ -symmetric, hence the product  $F(\mathbf{v})\tilde{\mathbb{P}}$  is  $\mathfrak{P}_{[a,b]}$ -reducible, by Lemma 1.

The sum over the group  $\mathfrak{S}_{[1,M]}$  can be rewritten as the double sum, first over the subgroup  $\mathfrak{S}_{[a,b]} \subset \mathfrak{S}_{[1,M]}$ , then over the coset  $\mathfrak{S}_{[1,M]}/\mathfrak{S}_{[a,b]}$ . The alternating sum over  $\mathfrak{S}_{[a,b]}$  with the weight  $\mathbb{P}_{[a,b]}$  then nullifies  $F(\mathbf{v})\tilde{\mathbb{P}}$ .  $\blacksquare$

Now we can justify Conjecture 3.

**Proposition 1.** *Assume that a sequence of polynomials  $Q^{(m)}(v_1, \dots, v_m)$ ,  $m = 1, 2, \dots$  solve equations (10.1) and (10.2). Then  $Q^{\mathbf{m}}(\mathbf{v})$  given by (9.16) solve equations (9.13) and (9.14).*

*Proof.* Given the composition  $\mathbf{m} = (m_1, \dots, m_{k-1}, m_k, m_{k+1}, \dots, m_N)$ , let  $(a_i, \dots, b_i)$  be the consecutive subsegments of length  $m_i$  of the sequence  $(1, \dots, M)$ , or, explicitly,

$$a_i = m_1 + \dots + m_{i-1} + 1, \quad b_i = m_1 + \dots + m_i, \quad i = 1, \dots, N. \quad (\text{A.1})$$

By using the product structure of  $Q^{\mathbf{m}}$  one can recast the below sum as

$$\begin{aligned} & \left[ \left( \sum_{j=1}^N \tilde{K}(\mathbf{w}_j^{\mathbf{m}}) + (m_j^3 - m_j)\gamma \right) Q^{\mathbf{m}}(\mathbf{v}) + \sum_{\mathbf{m}': \mathbf{m}' \succ \mathbf{m}} \beta_{n_1 n_2} Q^{\mathbf{m}'}(\mathbf{v}) \right] e^{\mathbf{W}^{\mathbf{m}} \cdot \mathbf{x}} \\ & = \sum_{j=1}^N \left\{ \left[ \tilde{K}(\mathbf{w}_j^{\mathbf{m}}) + (m_j^3 - m_j)\gamma \right] Q^{(m_j)}(\mathbf{w}_j^{\mathbf{m}}) \right. \\ & \left. + \sum_{\substack{n_1+n_2 \\ =m_j}} \beta_{n_1 n_2} Q^{(n_1)}(w_{j,1}^{\mathbf{m}}, \dots, w_{j,n_1}^{\mathbf{m}}) Q^{(n_2)}(w_{j,n_1+1}^{\mathbf{m}}, \dots, w_{j,m_j}^{\mathbf{m}}) \right\} \cdot e^{\mathbf{W}^{\mathbf{m}} \cdot \mathbf{x}} \cdot \prod_{\substack{a=1 \\ \neq j}}^N Q^{(m_j)}(\mathbf{w}_a^{\mathbf{m}}). \end{aligned}$$

Here  $\mathbf{x} = (x_1, \dots, x_N)$ ,  $\mathbf{W}^{\mathbf{m}}$  are as defined in (9.5) while the vector  $\mathbf{w}_j^{\mathbf{m}}$  introduced in (9.2) have components

$$\mathbf{w}_j^{\mathbf{m}} = (w_{j,1}^{\mathbf{m}}, \dots, w_{j,m_j}^{\mathbf{m}}).$$

The product outside of the bracket is symmetric in respect to permutations of the coordinates of  $\mathbf{w}_j^{\mathbf{m}}$ , viz. in respect to the action of the permutation group  $\mathfrak{S}_{[a_j, b_j]}$ . By construction, the functions appearing inside of the brackets are  $\mathfrak{P}_{[a_j, b_j]}$ -reducible. Thus  $Q^{\mathbf{m}}(\mathbf{v})$  given by (9.16) solves the bulk equation (9.13) in virtue of Lemma 1.

It thus remains to deal with the gluing conditions issuing from the jump of the transversal derivatives. Here, we introduce the auxiliary compositions  $\mathbf{m}' \succ \mathbf{m}$ ,  $\mathbf{m}'' \succ \mathbf{m}$

$$\begin{aligned} \mathbf{m}' &= (m_1, \dots, m_{k-1}, n_1, n_2, m_{k+1}, \dots, m_N), \\ \mathbf{m}'' &= (m_1, \dots, m_{k-1}, n_2, n_1, m_{k+2}, \dots, m_N) \end{aligned}$$

with  $n_1 + n_2 = m_k$ . Then, it holds

$$\begin{aligned} & \left[ V_{n_1 n_2}(\mathbf{w}_k^m) \cdot Q^{m'}(\mathbf{v}) + V_{n_2 n_1}(\mathbf{w}_k^m) \cdot Q^{m''}(\mathbf{v}) + 2n_1 n_2 \beta_{n_1, n_2} Q^m(\mathbf{v}) \right] \cdot e^{\mathbf{W}^m \cdot \mathbf{x}} \\ &= \left[ 2n_1 n_2 \beta_{n_1, n_2} Q^{(m_k)}(\mathbf{w}_k^m) + V_{n_1 n_2}(\mathbf{w}_k^m) \cdot Q^{(n_1)}(w_{k,1}^m, \dots, w_{k,n_1}^m) Q^{(n_2)}(w_{k,n_1+1}^m, \dots, w_{k,m_k}^m) \right. \\ & \left. + V_{n_2 n_1}(\mathbf{w}_k^m) \cdot Q^{(n_2)}(w_{k,1}^m, \dots, w_{k,n_2}^m) Q^{(n_1)}(w_{k,n_2+1}^m, \dots, w_{k,m_k}^m) \right] \cdot e^{\mathbf{W}^m \cdot \mathbf{x}} \cdot \prod_{\substack{a=1 \\ \neq k}}^N Q^{(m_j)}(\mathbf{w}_a^m). \end{aligned}$$

Again, the product outside of the bracket is symmetric in respect to permutations of the coordinates of  $\mathbf{w}_k^m$ , *viz.*  $\mathfrak{S}_{[a_j, b_j]}$ . By construction, the functions appearing inside of the brackets are  $\mathfrak{P}_{[a_j, b_j]}$ -reducible. Thus  $Q^m(\mathbf{v})$  given by (9.16) solves the jump of transversal derivative condition (9.14) in virtue of Lemma 1. ■

## APPENDIX B.

Checking  $\mathfrak{P}$ -equivalence of polynomials directly is difficult even with computer since it involves summation over  $M!$  permutations, which leads to the exponential growth of the computational complexity with  $M$ . When verifying Conjecture 4, we checked in fact some stronger conditions that we call 2- and 3-reducibility having the advantage of a polynomial complexity.

Let  $P(v)$  be given by (7.8) and  $\mathbf{v} = (v_1, \dots, v_M)$ . Let  $v_{ij} = v_i - v_j$ , and  $P_{ij} = P(v_i - v_j)$ . Assuming  $M \geq 2$ , we shall say that a polynomial  $F(\mathbf{v})$  is 2-reducible and write  $F \stackrel{2}{\equiv} 0$  if  $F(\mathbf{v})$  admits a decomposition

$$F(\mathbf{v}) = \sum_{i=1}^{M-1} P_{i,i+1} G_i(\mathbf{v}) \tag{B.1}$$

with some polynomials  $G_i(\mathbf{v})$  such that  $G_i(\mathbf{v})$  is symmetric under permutation  $v_i \leftrightarrow v_{i+1}$  for each  $i$ . Note that such a decomposition is not necessarily unique.

**Proposition 2.** *If  $F$  is 2-reducible then  $F$  is  $\mathfrak{P}_{[1, M]}$ -reducible.*

*Proof.* Note that  $P_{i,i+1}$  is  $\mathfrak{P}_{[i, i+1]}$ -reducible since  $P_{i,i+1} \mathbb{P}_{[i, i+1]} = P_{i,i+1} P_{i+1, i}$  is  $\mathfrak{S}_{[i, i+1]}$ -symmetric, hence nullified by the antisymmetrisation. Then, by Lemma 1, the  $i$ -th term in (B.1) is  $\mathfrak{P}_{[i, i+1]}$ -reducible, hence  $\mathfrak{P}_{[1, M]}$ -reducible, by Lemma 2. ■

The property of 2-reducibility is not always sufficient to prove the  $\mathfrak{P}$ -reducibility, and we shall also use the notion of 3-reducibility defined below.

**Lemma 3.** *The polynomial  $v_{12} - v_{23} = v_1 - 2v_2 + v_3$  is  $\mathfrak{P}_{[1, 3]}$ -reducible.*

*Proof.* Note that  $P_{12}$  is  $\mathfrak{P}_{[1, 2]}$ -reducible, and  $P_{23}$  is  $\mathfrak{P}_{[2, 3]}$ -reducible, as shown in the proof of Proposition 2. By Lemma 2,  $P_{12}$ ,  $P_{23}$ , and therefore  $P_{12} - P_{23}$  are  $\mathfrak{P}_{[1, 3]}$ -reducible. Now note that the difference

$$P_{12} - P_{23} = v_{12}^3 - v_{23}^3 + 12\gamma(v_{12} - v_{23})$$

factorises into  $v_{12} - v_{23}$  and a quadratic polynomial  $J$  that is  $\mathfrak{S}_{[1, 3]}$ -symmetric:

$$P_{12} - P_{23} = (v_{12} - v_{23})J, \tag{B.2}$$

$$J = v_{12}^2 + v_{12}v_{23} + v_{23}^2 + 12\gamma = v_1^2 + v_2^2 + v_3^2 - v_1v_2 - v_1v_3 - v_2v_3 + 12\gamma. \quad (\text{B.3})$$

Then from the symmetry of  $J$  it follows that

$$0 = \mathfrak{P}_{[1,3]}(P_{12} - P_{23}) = \mathfrak{P}_{[1,3]}((v_{12} - v_{23})J) = J \mathfrak{P}_{[1,3]}(v_{12} - v_{23}) \quad (\text{B.4})$$

and therefore  $\mathfrak{P}_{[1,3]}(v_{12} - v_{23}) = 0$  since  $J \neq 0$ .  $\blacksquare$

By Lemma 2, an immediate corollary is that  $v_{i,i+1} - v_{i+1,i+2}$  is  $\mathfrak{P}_{[1,M]}$ -reducible for any  $M$ , and  $i = 1, \dots, M - 2$ .

Remarkably, the condition that

$$J = \frac{P_{12} - P_{23}}{v_{12} - v_{23}} \quad (\text{B.5})$$

is an  $\mathfrak{S}_{[1,3]}$ -symmetric polynomial fixes the polynomial  $P(v)$  uniquely as a cubic polynomial with zero  $v^2$ -term. The easiest way to prove this is to use the homogeneity and to check the monomials  $v^p$  to see that the solution is  $p \in \{0, 1, 3\}$ .

Assuming  $M \geq 3$ , we shall say that a polynomial  $F(\mathbf{v})$  is 3-reducible and write  $F \stackrel{\text{3}}{\equiv} 0$  if  $F(\mathbf{v})$  admits a decomposition

$$F(\mathbf{v}) = \sum_{i=1}^{M-2} (v_{i,i+1} - v_{i+1,i+2}) J_i(\mathbf{v}) \quad (\text{B.6})$$

with some  $\mathfrak{S}_{[i,i+2]}$ -symmetric polynomials  $J_i(\mathbf{v})$ .

Note that such a decomposition is not necessarily unique.

**Proposition 3.** *If  $F$  is 3-reducible then  $F$  is  $\mathfrak{P}_{[1,M]}$ -reducible.*

*Proof.* For the  $i$ -th term in (B.6) we have

$$\mathfrak{P}_{[i,i+2]}((v_{i,i+1} - v_{i+1,i+2}) J_i(\mathbf{v})) = J_i(\mathbf{v}) \mathfrak{P}_{[i,i+2]}(v_{i,i+1} - v_{i+1,i+2}) = 0,$$

using first the symmetry of  $J_i$ , then Lemma 3. By Lemma (2), each term is  $\mathfrak{P}_{[1,M]}$ -reducible.  $\blacksquare$

The following conjecture supersedes Conjecture 4. It has been verified by means of computer algebra for  $M \leq 8$ .

**Conjecture 5.** *For the polynomials  $Q^{(M)}(\mathbf{v})$  given by (10.3) the left-hand-side of the jump equation (10.2) is in fact 3-reducible, which, by Proposition 3, implies  $\mathfrak{P}_{[1,M]}$ -reducibility.*

*The left-hand-side of the bulk equation (10.1) is respectively a sum of a 2-reducible and a 3-reducible parts, which, by Propositions 2 and 3, implies  $\mathfrak{P}_{[1,M]}$ -reducibility.*

## REFERENCES

- [1] A. Arinshtein, V. Fateyev, and A. B. Zamolodchikov. Quantum  $S$ -matrix of the  $(1+1)$ -dimensional Todd chain. *Phys. Lett. B*, 87(4):389–392, 1979. doi:10.1016/0370-2693(79)90561-6.
- [2] F. Berezin, G. Pokhil, and V. Finkelberg. Schrödinger equation for a system of one-dimensional particles with point interaction (in Russian). *Vestnik MGU*, 1:21–28, 1964.
- [3] N. M. Bogoliubov, A. G. Izergin, and V. E. Korepin. *Quantum Inverse Scattering Method, Correlation Functions and Algebraic Bethe Ansatz*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1993. doi:10.1017/CB09780511628832.
- [4] H. Braden, E. Corrigan, P. Dorey, and R. Sasaki. Affine Toda field theory and exact  $S$ -matrices. *Nucl. Phys. B*, 338(3):689–746, 1990. doi:10.1016/0550-3213(90)90648-W.
- [5] K. Case. Polynomial constants for the quantized NLS equation. *J. Math. Phys.*, 25(7):2306–2314, 1984. doi:10.1063/1.526400.

- [6] B. Davies and V. Korepin. Higher conservation laws for the quantum non-linear Schrödinger equation. Technical report, Centre for Mathematical Analysis of Australian National University CMA-R33-89, 1989. URL: <http://arxiv.org/abs/1109.6604>.
- [7] M. Gaudin. *The Bethe Wavefunction*. Cambridge University Press, Cambridge, 2014. doi:10.1017/CB09781107053885.
- [8] E. Gutkin. Conservation laws for the nonlinear schrödinger equation. In *Annales de l'IHP Analyse non linéaire*, volume 2, pages 67–74, 1985. URL: [http://archive.numdam.org/article/AIHPC\\_1985\\_\\_2\\_1\\_67\\_0.pdf](http://archive.numdam.org/article/AIHPC_1985__2_1_67_0.pdf).
- [9] V. Kac and A. Raina. *Bombay Lectures on Highest Weight Representations of Infinite Dimensional Lie Algebras*, volume 2 of *Advanced Series in Mathematical Physics*. World Scientific, Singapore, 1987.
- [10] B. Konopelchenko. *Introduction to Multidimensional Integrable Equations: The Inverse Spectral Transform in 2 + 1 Dimensions*. Plenum Monographs in Nonlinear Physics. Springer US, 1993. doi:10.1007/978-1-4899-1170-4.
- [11] K. Kozłowski and E. Sklyanin. Combinatorics of Generalized Bethe Equations. *Lett. Math. Phys.*, 103(10):1047–1077, 2013. arXiv:1205.2968, doi:10.1007/s11005-013-0630-9.
- [12] P. Kulish. Quantum nonlinear wave interaction system. *Physica D: Nonlinear Phenomena*, 18(1–3):360–364, 1986. doi:10.1016/0167-2789(86)90197-1.
- [13] P. P. Kulish, S. V. Manakov, and L. D. Faddeev. Comparison of the exact quantum and quasiclassical results for a nonlinear schrödinger equation. *Theor. Math. Phys.*, 28(1):615–620, July 1976. doi:10.1007/BF01028912.
- [14] T. D. Lee. Some special examples in renormalizable field theory. *Phys. Rev.*, 95:1329–1334, Sep 1954. doi:10.1103/PhysRev.95.1329.
- [15] E. H. Lieb and W. Liniger. Exact analysis of an interacting bose gas. I. The general solution and the ground state. *Phys. Rev.*, 130:1605–1616, May 1963. doi:10.1103/PhysRev.130.1605.
- [16] A. V. Litvinov. On spectrum of ILW hierarchy in conformal field theory. *JHEP*, 2013(11):1–14, 2013. doi:10.1007/JHEP11(2013)155.
- [17] S. Manakov and P. Santini. On the solutions of the dKP equation: the nonlinear Riemann-Hilbert problem, longtime behaviour, implicit solutions and wave breaking. *J. Phys. A: Math. Theor.*, 41(5), 2008. doi:10.1088/1751-8113/41/5/055204.
- [18] E. Sklyanin. Quantum version of the method of inverse scattering problem. *J. Soviet Math.*, 19(5):1546–1596, 1982. doi:10.1007/BF01091462.
- [19] E. K. Sklyanin. Quantization of the continuous Heisenberg ferromagnet. *Lett. Math. Phys.*, 15(4):357–368, 1988. doi:10.1007/BF00419595.
- [20] R. P. Stanley and S. Fomin. *Enumerative Combinatorics, vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999. doi:10.1017/CB09780511609589.