

This is a repository copy of *Sums of reciprocals and the three distance theorem*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/121745/>

Version: Submitted Version

Monograph:

Beresnevich, Victor orcid.org/0000-0002-1811-9697 and Leong, Nicol En Ci (2017) Sums of reciprocals and the three distance theorem. Working Paper.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Sums of reciprocals and the three distance theorem

VICTOR BERESNEVICH* NICOL LEONG

Abstract

In this paper we investigate the sums of reciprocals to an arithmetic progression taken modulo one, that is sums of $\{n\alpha - \gamma\}^{-1}$, where α and γ are real parameters and $\{\cdot\}$ is the fractional part of a real number. Bounds for these sums have been studied for a long while in connection with various applications. In this paper we develop an alternative technique for obtaining upper bounds for the sums and obtain new efficient and fully explicit results. The technique uses the so-called three distance theorem.

1 Introduction

Throughout, given an $x \in \mathbb{R}$, we write $[x]$ for its integer part, $\{x\} = x - [x]$ for its fractional part and $\|x\|$ for the distance of x from the nearest integer. Throughout $\alpha, \gamma \in \mathbb{R}$ and α is irrational. The key object of interest of this paper is the following sum

$$T_N(\alpha, \gamma) := \sum_{\substack{0 \leq n \leq N \\ n \neq n'}} \frac{1}{\{n\alpha - \gamma\}} \quad (1)$$

where $N \in \mathbb{N}$ and $0 \leq n' \leq N$ is an integer such that

$$\{n'\alpha - \gamma\} = \min_{0 \leq n \leq N} \{n\alpha - \gamma\}. \quad (2)$$

Note that n' depends on α , γ and N and, since α is irrational, n' is unique. One obvious reason for imposing the condition $n \neq n'$ in (1) is to avoid a vanishing denominator. However, even if $\{n'\alpha - \gamma\}$ does not vanish the term $\{n'\alpha - \gamma\}^{-1}$ cannot be bounded by a function of N and α . This is because, for any given N and α , the parameter γ can be arbitrarily close to one of the points $\{n\alpha\}$ with $0 \leq n \leq N$. Also in this paper, we shall consider the sums with $\{n\alpha - \gamma\}$ replaced by $\|n\alpha - \gamma\|$ and the more general expression $n^a \{n\alpha - \gamma\}^b$ – see the last section.

*Supported by EPSRC grant EP/J018260/1

There are various applications of bounds on $T_N(\alpha, \gamma)$, e.g. for counting lattice points in polygons, in the theory of uniform distribution, metric theory of Diophantine approximation and so on. The above sums have therefore been studied in depth for a long while and also recently, see, for example, [2], [3], [4], [5], [9], [11], [12], [13], [15, pp. 108-110], [16], [17], [19], [24], [26], [32], [33].

The purpose of this paper is to introduce yet another technique for studying the above sums and obtain new results as well as recover known bounds in a more precise form. A detailed overview of previously obtained bounds for the sums of reciprocals can be found in the recent paper [5]. The technique we propose makes use of the so-called Three Distance Theorem, see §2. This theorem relies on the theory of continued fractions. The facts from the theory of continued fractions we use in this paper can be found in any textbook on the topic, for example, [14], [18] or [23].

From now on, unless otherwise stated, $[a_0; a_1, a_2, \dots]$ will denote the simple continued fraction expansion of a fixed real irrational number α , where the a_k are partial quotients. Thus, $a_0 \in \mathbb{Z}$, $a_1, a_2, \dots \in \mathbb{N}$ are the unique integers such that

$$\alpha = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}.$$

Further, $p_k/q_k = [a_0; a_1, a_2, \dots, a_k]$ will denote the k th convergent to α . Recall that the numbers p_k and q_k are coprime integers that satisfy the following recurrence relations:

$$q_{k+1} = a_{k+1}q_k + q_{k-1}, \quad p_{k+1} = a_{k+1}p_k + p_{k-1} \quad (k \geq 0) \quad (3)$$

with $p_0 = a_0$, $p_{-1} = 1$, $q_0 = 1$, $q_{-1} = 0$, and

$$q_{k+1}p_k - q_kp_{k+1} = (-1)^{k+1} \quad (k \geq -1). \quad (4)$$

Our key result concerning $T_N(\alpha, \gamma)$ is the following.

Theorem 1. *Let $N \in \mathbb{N}$, $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, p_k/q_k denote the convergents to α and $K = K(N, \alpha)$ be the largest integer satisfying $q_K \leq N$. Further, let $\gamma \in \mathbb{R}$ and $n' \in [0, N]$ be an integer given by (2). Then*

$$T_N(\alpha, \gamma) < 4N(\log q_K + 1) + 2q_{K+1} \left(\log \left(\frac{N}{q_K} \right) + 1 \right) \quad (5)$$

and furthermore

$$\sum_{\substack{0 \leq n \leq N \\ n \not\equiv n' \pmod{q_K}}} \frac{1}{\{n\alpha - \gamma\}} < 4N(\log q_K + 1). \quad (6)$$

Remark 1. The sequence $(q_k)_{k \geq 1}$ of the denominators of the convergents to α is the unique largest increasing sequence of positive integers such that $\|q_1\alpha\| > \|q_2\alpha\| > \|q_3\alpha\| > \dots$. Since for any positive integer q we have that $\|q(-\alpha)\| = \|q\alpha\|$, we have that $q_k(-\alpha) = q_k(\alpha)$ for all α . Therefore Theorem 1 is applicable to $-\alpha$, resulting in the same bounds. Further for any real number x note that $\|x\| = \min(\{x\}, \{-x\})$ and furthermore that $\frac{1}{2} \leq \max(\{x\}, \{-x\}) \leq 1$. Hence

$$1 < \frac{1}{\{x\}} + \frac{1}{\{-x\}} - \frac{1}{\|x\|} \leq 2. \quad (7)$$

We shall use the above observations to prove the following

Corollary 1. *Let N , α , p_k/q_k and K be the same as in Theorem 1. Further, let $\gamma \in \mathbb{R}$ and $n^* \in [0, N]$ be an integer such that*

$$\|n^*\alpha - \gamma\| = \min_{0 \leq n \leq N} \|n\alpha - \gamma\|.$$

Then

$$\sum_{\substack{0 \leq n \leq N \\ n \neq n^*}} \frac{1}{\|n\alpha - \gamma\|} \leq 8N(\log q_K + 1) + 4q_{K+1} \left(\log \left(\frac{N}{q_K} \right) + 2 \right). \quad (8)$$

Proof. Let n' be the same as in Theorem 1 and similarly let n'' be the unique integer in $[0, N]$ such that $\{n''(-\alpha) + \gamma\} = \min_{0 \leq n \leq N} \{n(-\alpha) + \gamma\}$. Then, by Theorem 1 and inequalities (7), we have that

$$\begin{aligned} \sum_{\substack{0 \leq n \leq N \\ n \neq n', n''}} \frac{1}{\|n\alpha - \gamma\|} &\leq \sum_{\substack{0 \leq n \leq N \\ n \neq n'}} \frac{1}{\{n\alpha - \gamma\}} + \sum_{\substack{0 \leq n \leq N \\ n \neq n''}} \frac{1}{\{-(n\alpha - \gamma)\}} \\ &\leq 8N(\log q_K + 1) + 4q_{K+1} \left(\log \left(\frac{N}{q_K} \right) + 1 \right). \end{aligned} \quad (9)$$

Note that n^* is either n' or n'' . If $n' \neq n''$ then we also have that $\|n'\alpha - n''\alpha\| \geq \|q_K\alpha\|$ since q_K is the largest best approximation $\leq N$ and $1 \leq |n' - n''| \leq N$. Therefore, the $\max\{\|n'\alpha - \gamma\|, \|n''\alpha - \gamma\|\} \geq \frac{1}{2}\|q_K\alpha\| \geq 1/(4q_{K+1})$. Hence

$$\min \left\{ \frac{1}{\|n'\alpha - \gamma\|}, \frac{1}{\|n''\alpha - \gamma\|} \right\} \leq 4q_{K+1}$$

and combining this with (9) completes the proof. \square

2 The Three Distance Theorem

The main question we discuss in this section is the following: *given an $N \in \mathbb{N}$ and $\alpha \in \mathbb{R}$, what can we say about the distribution of the points*

$$\{\alpha\}, \{2\alpha\}, \dots, \{N\alpha\} \quad (10)$$

in the unit interval $[0, 1)$? Equivalently this question can be posed using circle rotations on identifying $[0, 1)$ with the unit circle in the usual way via the map $\alpha \mapsto e^{2\pi i\alpha}$. The following statement conjectured by Hugo Steinhaus is widely known as the three distance theorem or three gaps theorem.

Theorem 2 (The Three Distance Theorem). *For any $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and any integer $N \geq 1$ the points (10) partition $[0, 1]$ into $N + 1$ intervals which lengths take at most 3 different values δ_A , δ_B and δ_C with $\delta_C = \delta_A + \delta_B$.*

There are various generalisations of the above fact and several independent proofs, in particular in [10, 20, 21, 27, 28, 29, 30, 31]. Remarkably, the length of the gaps as well as the number of gaps of every length can be exactly specified using the continued fraction expansion of α . Within this paper we will use an even more refined statement which in addition specifies the order in which the intervals of each length appear. The theorem will require the theory of continued fractions. Some basic notation and facts of this theory have already been recalled in the introduction above. In addition, for $k \geq 0$, we define the following quantities, which measure how the k th convergent approximates α :

$$D_k = q_k\alpha - p_k. \quad (11)$$

By (3) and the definition of D_k , we clearly have that

$$a_{k+1}D_k = D_{k+1} - D_{k-1} \quad (k \geq 1). \quad (12)$$

The following well known statement, that can be found in [18], reveals the approximation properties of the convergents to α :

Lemma 1. *The sequence of even convergents p_{2k}/q_{2k} is strictly increasing, the sequence of odd convergents p_{2k+1}/q_{2k+1} is strictly decreasing. Both converge to α and furthermore we have that*

$$\frac{1}{2q_{k+1}} < \frac{1}{q_{k+1} + q_k} < |D_k| < \frac{1}{q_{k+1}}. \quad (13)$$

Since $q_k \geq 2$ for $k \geq 2$, we have from the above lemma that $D_k = (-1)^k \|q_k\alpha\|$ for all $k \geq 1$. Also we have that D_k alternates the sign, that is

$$D_k D_{k+1} < 0 \quad (k \geq 0). \quad (14)$$

In particular, in view of (12) and (14) we have that

$$a_{k+1}|D_k| + |D_{k+1}| = |D_{k-1}| \quad (k \geq 1). \quad (15)$$

We are now ready to state the full version of the three distance theorem that we will use in this paper.

Theorem 3. *Let $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, $[a_0; a_1, a_2, \dots]$ be the continued fraction expansion of α , p_k/q_k be the convergents to α and $D_k = q_k\alpha - p_k$. Then for any $N \in \mathbb{N}$ there exists a unique integer $k \geq 0$ such that*

$$q_k + q_{k-1} \leq N < q_{k+1} + q_k \quad (16)$$

and unique integers r and s satisfying

$$N = rq_k + q_{k-1} + s, \quad 1 \leq r \leq a_{k+1}, \quad 0 \leq s \leq q_k - 1, \quad (17)$$

such that the points $\{\alpha\}, \{2\alpha\}, \dots, \{N\alpha\}$ split $[0, 1]$ into $N+1$ intervals, namely

$$J_0 := [0, \{n_1\alpha\}], \quad J_1 := [\{n_1\alpha\}, \{n_2\alpha\}], \quad \dots, \quad J_N := [\{n_N\alpha\}, 1], \quad (18)$$

of which

$$N_A := N + 1 - q_k \quad \text{are of length } \delta_A := |D_k|, \quad (19)$$

$$N_B := s + 1 \quad \text{are of length } \delta_B := |D_{k+1}| + (a_{k+1} - r)|D_k|, \quad (20)$$

$$N_C := q_k - s - 1 \quad \text{are of length } \delta_C := \delta_A + \delta_B. \quad (21)$$

Furthermore, the unique permutation (n_1, \dots, n_N) of $(1, \dots, N)$ such that

$$0 < \{n_1\alpha\} < \{n_2\alpha\} < \dots < \{n_{N-1}\alpha\} < \{n_N\alpha\} < 1 \quad (22)$$

can be found by setting $n_0 := 0$ and then defining

$$n_{i+1} := n_i + \Delta_i \quad \text{for } i = 0, \dots, N-1, \quad (23)$$

where

$$\Delta_i := \begin{cases} (-1)^k q_k & \text{if } n_i \in A, \\ (-1)^{k-1} (q_{k-1} + r q_k) & \text{if } n_i \in B, \\ (-1)^{k-1} (q_{k-1} + (r-1) q_k) & \text{if } n_i \in C \end{cases} \quad (24)$$

and

$$\begin{aligned} A &:= \{n \in \mathbb{Z} \cap [0, N] : 0 \leq n + (-1)^k q_k \leq N\}, \\ B &:= \{n \in \mathbb{Z} \cap [0, N] : 0 \leq n + (-1)^{k-1} (q_{k-1} + r q_k) \leq N\}, \\ C &:= \{n \in \mathbb{Z} \cap [0, N] : n \notin (A \cup B)\} \end{aligned} \quad (25)$$

are disjoint subsets of integers; the length of the intervals (18) is given by

$$|J_i| = \begin{cases} \delta_A & \text{if } n_i \in A, \\ \delta_B & \text{if } n_i \in B, \\ \delta_C & \text{if } n_i \in C. \end{cases} \quad (26)$$

Remark 2. Strictly speaking Theorem 3 is not new and can be assembled from published results. The closest versions can be found in [1]. However, given that Theorem 3 underpins our approach, in §2.1 we present a complete yet short proof of the result, which also makes this paper self-contained.

Remark 3. Throughout this paper the length of the interval J_i will be referred to as the i th gap. It is clear from (13) that $|D_{k+1}| < |D_k|$. Hence, $\delta_B < \delta_A$ if and only if $r = a_{k+1}$, and if $r \neq a_{k+1}$, δ_A will be the smallest (of δ_A , δ_B and δ_C) gap.

Remark 4. As α is irrational, δ_A , δ_B and δ_C are distinct. In fact, while the two gaps δ_A and δ_B always appear, δ_C -gaps exist if and only if $s < q_k - 1$. There are infinitely many integers N for which there are only two gaps. The structure and the transformation rules for partitioning intervals are studied in details in [21]. Other in-depth studies involving the two-gaps case can be found in [6, 7, 8, 22, 25].

2.1 Proof of Theorem 3

Observe that $(q_k + q_{k-1})_{k \geq 0}$ is a strictly increasing sequence of integers starting from $q_0 + q_{-1} = 1$, whence the existence of k satisfying (16) readily follows. Next, observe that $q_k \leq N - q_{k-1} < q_{k+1} + q_k - q_{k-1} = (a_{k+1} + 1)q_k$. Therefore, by division with remainder, r and s satisfying (17) exist and are unique.

Now we verify that $A \cap B = \emptyset$. Indeed, if there existed $n \in A \cap B$, then for even k we would have that

$$rq_k + q_{k-1} \stackrel{n \in B}{\leq} n \stackrel{n \in A}{\leq} N - q_k = rq_k + q_{k-1} + s - q_k \stackrel{s < q_k}{<} rq_k + q_{k-1},$$

while for odd k we would have that

$$q_k \stackrel{n \in A}{\leq} n \stackrel{n \in B}{\leq} N - (rq_k + q_{k-1}) = s < q_k.$$

In both instances we would get a contradiction. In addition, note that C is disjoint from A and from B simply by its definition.

Now we show that

$$0 \leq n_i \leq N \quad \text{for all } 0 \leq i \leq N. \quad (27)$$

We use induction on i . Clearly, $n_0 = 0$ satisfies the inequalities. Now assuming that $i < N$ and $0 \leq n_i \leq N$ we shall verify that $0 \leq n_{i+1} \leq N$. Indeed, if $n_i \in A \cup B$ this claim follows immediately from the definitions of A and B and (23). Thus we only need to consider $n_i \in C$. First assume that k is even. Then, since $n_i \notin A$, we have that $n_i > N - q_k$ and therefore

$$\begin{aligned} n_{i+1} &\stackrel{(23)}{=} n_i - (q_{k-1} + (r-1)q_k) \geq N - q_k - (q_{k-1} + (r-1)q_k) \\ &\stackrel{(17)}{=} rq_k + q_{k-1} + s - q_k - (q_{k-1} + (r-1)q_k) = s \geq 0 \end{aligned}$$

while clearly $n_{i+1} < n_i \leq N$ as required. Now assume that k is odd. Then, since $n_i \notin A$, we have that $n_i < q_k$ and therefore

$$n_{i+1} \stackrel{(23)}{=} n_i + (q_{k-1} + (r-1)q_k) < q_k + (q_{k-1} + (r-1)q_k) \stackrel{(17)}{=} N - s \leq N$$

while clearly $0 \leq n_i < n_{i+1}$. This completes the proof of (27). The disjointness of A , B and C together with (27) imply that n_i given by (23) is well defined.

Next, by (19), (20), (21) and (25), we have that

$$N_A = \#A, \quad N_B = \#B, \quad N_C = \#C \quad \text{and} \quad N_A + N_B + N_C = N + 1. \quad (28)$$

Furthermore,

$$\begin{aligned} N_A \delta_A + N_B \delta_B + N_C \delta_C &= (N - s) \delta_A + q_k \delta_B \\ &= (rq_k + q_{k-1}) |D_k| + q_k (|D_{k+1}| + (a_{k+1} - r) |D_k|) \\ &= (a_{k+1} q_k + q_{k-1}) |D_k| + q_k |D_{k+1}| \stackrel{(3)}{=} q_{k+1} |D_k| + q_k |D_{k+1}| \\ &\stackrel{(14)}{=} |q_{k+1} D_k - q_k D_{k+1}| \stackrel{(11)}{=} |-q_{k+1} p_k + q_k p_{k+1}| \stackrel{(4)}{=} 1. \quad (29) \end{aligned}$$

In particular, it means that δ_A , δ_B and δ_C are all strictly less than 1, except when $N = 1$ in which case $N_C = 0$ and $\delta_C = 1$.

Next we will show that for $0 \leq i < N$

$$\{(n_{i+1} - n_i)\alpha\} = \begin{cases} \delta_A & \text{if } n_i \in A, \\ \delta_B & \text{if } n_i \in B, \\ \delta_C & \text{if } n_i \in C. \end{cases} \quad (30)$$

We shall use (23) and (24). In the case $n_i \in A$ we have that

$$\{(n_{i+1} - n_i)\alpha\} = \{(-1)^k q_k \alpha\} = \{(-1)^k (q_k \alpha - p_k)\} = \{|D_k|\} = \{\delta_A\} = \delta_A,$$

since $0 < \delta_A < 1$. In the case $n_i \in B$ we have that

$$\begin{aligned} \{(n_{i+1} - n_i)\alpha\} &= \{(-1)^{k-1} (q_{k-1} + r q_k)\alpha\} \stackrel{(3)}{=} \{(-1)^{k-1} (q_{k+1} - (a_{k+1} - r)q_k)\alpha\} \\ &= \{(-1)^{k-1} (q_{k+1}\alpha - p_{k+1} - (a_{k+1} - r)(q_k\alpha - p_k))\} \\ &= \{|D_{k+1}| + (a_{k+1} - r)|D_k|\} = \{\delta_B\} = \delta_B, \end{aligned}$$

since $0 < \delta_B < 1$. Finally, in the case $n_i \in C$ we have that

$$\begin{aligned} \{(n_{i+1} - n_i)\alpha\} &= \{(-1)^{k-1} (q_{k-1} + (r-1)q_k)\alpha\} \\ &\stackrel{(3)}{=} \{(-1)^{k-1} (q_{k+1} - (a_{k+1} + 1 - r)q_k)\alpha\} \\ &= \{(-1)^{k-1} (q_{k+1}\alpha - p_{k+1} - (a_{k+1} + 1 - r)(q_k\alpha - p_k))\} \\ &= \{|D_{k+1}| + (a_{k+1} + 1 - r)|D_k|\} = \{\delta_C\} = \delta_C, \end{aligned}$$

since in this case $N_C \neq 0$ and so $0 < \delta_C < 1$.

Now, we prove (22). First of all note that $0 < \{n_1\alpha\}$ since α is irrational. The proof continues by induction. Suppose that $1 \leq i < N$ and that $0 < \{n_1\alpha\} < \dots < \{n_i\alpha\}$. This means that n_0, n_1, \dots, n_i are all different. Therefore, by (30) and the disjointness of A , B and C , we get that

$$\sum_{j=0}^i \{(n_{j+1} - n_j)\alpha\} < N_A \delta_A + N_B \delta_B + N_C \delta_C \stackrel{(29)}{=} 1. \quad (31)$$

Therefore,

$$\sum_{j=0}^i \{(n_{j+1} - n_j)\alpha\} = \{\sum_{j=0}^i (n_{j+1} - n_j)\alpha\} = \{(n_{i+1} - n_0)\alpha\} = \{n_{i+1}\alpha\}.$$

Similarly, $\sum_{j=0}^{i-1} \{(n_{j+1} - n_j)\alpha\} = \{n_i\alpha\}$ and thus

$$\{n_{i+1}\alpha\} = \{(n_{i+1} - n_i)\alpha\} + \{n_i\alpha\} > \{n_i\alpha\}. \quad (32)$$

This completes the proof of (22). Furthermore, since $n_0 = 0$, by (22) and (27), we have that (n_1, \dots, n_N) is the required permutation of $(1, \dots, N)$.

Finally, (30) and (32) verify (26) for $0 \leq i \leq N - 1$, while the facts that n_0, \dots, n_N are all different and lie in the disjoint sets A , B and C put together with (29) implies (26) for $i = N$. In turn, (19), (20) and (21) are a consequence of (26) and (28), namely that $N_A = \#A$, $N_B = \#B$, $N_C = \#C$. The proof of Theorem 3 is thus complete. \square

Remark 5. Since $0 \leq n_N \leq N$, one can formally define n_{N+1} using (23) and (24). This number is easy to compute since amongst $\Delta_0, \dots, \Delta_N$ there are clearly N_A values of $(-1)^k q_k$, N_B values of $(-1)^{k-1}(q_{k-1} + r q_k)$ and N_C values of $(-1)^{k-1}(q_{k-1} + (r-1)q_k)$. Hence, n_{N+1} is exactly

$$(-1)^k q_k N_A + (-1)^{k-1}(q_{k-1} + r q_k) N_B + (-1)^{k-1}(q_{k-1} + (r-1)q_k) N_C.$$

Substituting the values for N_A , N_B and N_C into the above expression and also taking into account that $N = r q_k + q_{k-1} + s$ one readily verifies that

$$n_{N+1} = 0.$$

This means that the sequence n_i can be continued for i beyond N and that it will always be well defined and cyclic. Furthermore it will satisfy the property that

$$n_i = n_j \iff i \equiv j \pmod{N+1}.$$

3 Proof of Theorem 1

3.1 Semi-homogeneous case

We begin by establishing Theorem 1 in the case $\{n'\alpha - \gamma\} = 0$. Without loss of generality we may assume that $\gamma = n'\alpha$. In what follows we will use the following basic estimate

$$\sum_{t=1}^T \frac{1}{t} \leq 1 + \int_1^T \frac{dt}{t} = \log T + 1. \quad (33)$$

Let $(n_i)_{i \geq 0}$ be the sequence defined by (23) and (24) extended to all i as described in Remark 5. Then, there exists a unique integer $\ell \in [0, N]$ such that $n_\ell = n'$. By Theorem 3 and Remark 5, $(n_\ell, \dots, n_{\ell+N})$ is the permutation of integers $(0, \dots, N)$ such that

$$0 = \{(n_\ell - n')\alpha\} < \{(n_{\ell+1} - n')\alpha\} < \dots < \{(n_{\ell+N} - n')\alpha\} < 1. \quad (34)$$

Hence

$$\sum_{\substack{0 \leq n \leq N \\ n \neq n'}} \frac{1}{\{(n - n')\alpha\}} = \sum_{j=1}^N \frac{1}{\{(n_{\ell+j} - n')\alpha\}}, \quad (35)$$

where the sum on the right of (35) is of decreasing terms. We shall be considering two cases depending on the relative size of δ_A and δ_B .

Case (i): $\delta_A < \delta_B$. Note that this means that $r \neq a_{k+1}$ and that $K = k$, where k is as in Theorem 3 and K is as in Theorem 1. Given an index j with $\ell \leq j \leq \ell + N$, let h_j denote the largest integer such that $0 \leq h_j \leq \max\{0, \ell + N - j\}$ and

$$\{(n_{j+i} - n')\alpha\} - \{(n_{j+i-1} - n')\alpha\} = \delta_A \quad \text{for all } i \text{ with } 1 \leq i \leq h_j. \quad (36)$$

By (23) and (24), we have that $\{(n_{j+i} - n')\alpha\} - \{(n_{j+i-1} - n')\alpha\} = \delta_A$ if and only if $n_{j+i} - n_{j+i-1} = (-1)^k q_k$. Hence $n_{j+h_j} = n_j + h_j(-1)^k q_k$. Now, since $0 \leq n_{j+h_j}, n_j \leq N$, we must have that

$$h_j = \frac{(-1)^k(n_{j+h_j} - n_j)}{q_k} = \frac{|n_{j+h_j} - n_j|}{q_k} \leq \left\lceil \frac{N}{q_k} \right\rceil =: H. \quad (37)$$

Hence, by (33) and the fact that $\delta_A = |D_k|$, we have that the first H terms on the right of (35) in total give us at most

$$\sum_{j=1}^H \frac{1}{j\delta_A} < \frac{1}{|D_k|} (\log H + 1) \stackrel{(13)}{<} 2q_{k+1} \left(\log \left(\frac{N}{q_k} \right) + 1 \right) \quad (38)$$

which is the right most term in (5), since $k = K$.

Now consider the remaining terms of (35): $\sum_{j=H+1}^N \{(n_{\ell+j} - n')\alpha\}^{-1}$. By (37), amongst any $H + 1$ consecutive gaps in (34), there will be at least one of length δ_B or δ_C . Therefore, writing j by division with remainder as $t(H + 1) + u$ with $0 \leq u \leq H$ we get that

$$\{(n_{\ell+j} - n')\alpha\} = \sum_{i=1}^j \underbrace{\left(\{(n_{\ell+i} - n')\alpha\} - \{(n_{\ell+i-1} - n')\alpha\} \right)}_{\text{gaps}} \geq t(\delta_B + H\delta_A).$$

Observe that

$$t = \left\lfloor \frac{j}{H + 1} \right\rfloor \leq \left\lfloor \frac{N}{[N/q_k] + 1} \right\rfloor \leq q_k$$

and that $t \geq 1$ if and only if $j \geq H + 1$. Then

$$\sum_{j=H+1}^N \frac{1}{\{(n_{\ell+j} - n')\alpha\}} \leq \sum_{t=1}^{q_k} \sum_{u=0}^H \frac{1}{t(\delta_B + H\delta_A)} \leq \frac{H + 1}{\delta_B + H\delta_A} \sum_{t=1}^{q_k} \frac{1}{t}. \quad (39)$$

Note that $r = [(N - q_{k-1})/q_k] \leq H$. Then, using the explicit values for δ_A and δ_B given in Theorem 3 and inequalities (13), we get that

$$\frac{H + 1}{\delta_B + H\delta_A} \leq \frac{H + 1}{(a_{k+1} + H - r)|D_k|} \leq \frac{H + 1}{a_{k+1}|D_k|} \leq \frac{(q_{k+1} + q_k)(H + 1)}{a_{k+1}}. \quad (40)$$

By (3), we have that

$$\frac{q_{k+1} + q_k}{a_{k+1}} = q_k + \frac{1}{a_{k+1}}(q_k + q_{k-1}). \quad (41)$$

Since $r \neq a_{k+1}$ we must have that $a_{k+1} \geq 2$ and since $q_{k-1} < q_k$ we then conclude that (41) is bounded above by $2q_k$. Putting this together with (40) and the facts that $H \leq N/q_k$ and $q_k \leq N$ we get that

$$\frac{H + 1}{\delta_B + H\delta_A} \leq 2q_k(N/q_k + 1) \leq 4N. \quad (42)$$

This together with (39) and (33) gives the left most term on the right of (5) and thus completes the proof of (5).

Case (ii): $\delta_A > \delta_B$. Note that this means that $r = a_{k+1}$ and that $K = k + 1$. Since in this case $\delta_B = |D_{k+1}|$ is the smallest gap, we have that

$$\frac{1}{\{(n_{\ell+1} - n')\alpha\}} \leq \frac{1}{\delta_B} \stackrel{(13)}{\leq} 2q_{k+2} = 2q_{K+1} \leq 2q_{K+1} \left(\log \left(\frac{N}{q_K} \right) + 1 \right), \quad (43)$$

which gives the right most term in (5). Now consider the remaining terms of (35), namely $\sum_{j=2}^N \{(n_{\ell+j} - n')\alpha\}^{-1}$.

Given an index j with $\ell \leq j \leq \ell + N$, let h_j denote the largest integer such that $0 \leq h_j \leq \max\{0, \ell + N - j\}$ and

$$\{(n_{j+i} - n')\alpha\} - \{(n_{j+i-1} - n')\alpha\} = \delta_B \quad \text{for all } i \text{ with } 1 \leq i \leq h_j. \quad (44)$$

By (23) and (24), we have that $\{(n_{j+i} - n')\alpha\} - \{(n_{j+i-1} - n')\alpha\} = \delta_B$ if and only if $n_{j+i} - n_{j+i-1} = (-1)^{k-1} (q_{k-1} + rq_k)$. Hence $n_{j+h_j} = n_j + h_j (-1)^{k-1} (q_{k-1} + rq_k)$. Now, since $0 \leq n_{j+h_j}, n_j \leq N$, we must have that

$$h_j = \frac{(-1)^k (n_{j+h_j} - n_j)}{q_{k-1} + rq_k} = \frac{|n_{j+h_j} - n_j|}{q_{k-1} + rq_k} \leq \left\lfloor \frac{N}{q_{k-1} + rq_k} \right\rfloor \stackrel{(17)}{=} 1. \quad (45)$$

Hence, there may not be two consecutive δ_B -gaps in (34) and every other gap in there is of length at least δ_A . Therefore, writing j by division with remainder as $2t + u$ with $0 \leq u \leq 1$ we get that

$$\{(n_{\ell+j} - n')\alpha\} = \sum_{i=1}^j \underbrace{\left(\{(n_{\ell+i} - n')\alpha\} - \{(n_{\ell+i-1} - n')\alpha\} \right)}_{\text{gaps}} \geq t(\delta_B + \delta_A).$$

Observe that

$$t = \left\lfloor \frac{j}{2} \right\rfloor \leq \left\lfloor \frac{N}{2} \right\rfloor \leq \left\lfloor \frac{q_{k+1} + q_k}{2} \right\rfloor < q_{k+1} = q_K$$

and that $t \geq 1$ if and only if $j \geq 2$. Then

$$\sum_{j=2}^N \frac{1}{\{(n_{\ell+j} - n')\alpha\}} \leq \sum_{t=1}^{q_K} \sum_{u=0}^1 \frac{1}{t(\delta_B + \delta_A)} \leq \frac{2}{\delta_B + \delta_A} \sum_{t=1}^{q_K} \frac{1}{t}. \quad (46)$$

Then, using the explicit values for δ_A and δ_B given in Theorem 3 and inequalities (13), we get that

$$\frac{2}{\delta_B + \delta_A} \leq \frac{2}{|D_k| + |D_{k+1}|} \leq \frac{2}{|D_k|} \stackrel{(13)}{\leq} 2(q_{k+1} + q_k). \quad (47)$$

Since $r = a_{k+1}$ we must have that $N = rq_k + q_{k-1} + s \geq q_{k+1}$. Putting this together with (47) we get that

$$\frac{2}{\delta_B + \delta_A} \leq 4N. \quad (48)$$

This together with (46) and (33) gives the left most term on the right of (5) and thus completes the proof of (5) in Case (ii).

Finally, note that the condition $n \equiv n' \pmod{q_K}$ excludes the terms contributing to the right most term of (5). Hence (6) follows and the proof of Theorem 1 is complete in the semi-homogeneous case.

3.2 The case of arbitrary γ

Without loss of generality assume that $0 < \gamma < 1$ (the case when $\gamma = 0$ is covered by the semi-homogeneous case). Then,

$$\{n\alpha - \gamma\} = \begin{cases} \{n\alpha\} - \gamma & \text{if } \{n\alpha\} \geq \gamma, \\ \{n\alpha\} - \gamma + 1 & \text{if } \{n\alpha\} < \gamma. \end{cases} \quad (49)$$

Let ℓ be the largest integer in the range $0 \leq \ell \leq N$ such that $\{n_\ell\alpha\} < \gamma$. Since $\{0\alpha\} = 0 < \gamma$, ℓ does exist. First assume that $\ell < N$. Set $n' = n_{\ell+1}$. Then, if $\ell + 1 \leq i \leq N$ we get that

$$\{n_i\alpha - \gamma\} \stackrel{(49)}{=} \{n_i\alpha\} - \gamma \geq \{n_i\alpha\} - \{n_{\ell+1}\alpha\} = \{(n_i - n')\alpha\},$$

while if $0 \leq i \leq \ell$ we get that

$$\{n_i\alpha - \gamma\} \stackrel{(49)}{=} \{n_i\alpha\} - \gamma + 1 \geq \{n_i\alpha\} - \{n_{\ell+1}\alpha\} + 1 = \{(n_i - n')\alpha\}.$$

Now assume that $\ell = N$. Define $n' = 0$. Then for all $0 \leq i \leq N$

$$\{n_i\alpha - \gamma\} \stackrel{(49)}{=} \{n_i\alpha\} - \gamma + 1 \geq \{n_i\alpha\} = \{(n_i - n')\alpha\}.$$

In either case we have that $T_N(\alpha, \gamma) \leq T_N(\alpha, \{n'\alpha\})$ with appropriately defined n' and therefore we finish the proof of the general case of Theorem 1 by appealing to the semi-homogeneous case that we have already considered in §3.1.

4 Generalisations

The proof of Theorem 1 given above can be easily generalised to give a bound on the following more general sum:

$$T_N^b(\alpha, \gamma) := \sum_{\substack{0 \leq n \leq N \\ n \neq n'}} \frac{1}{\{n\alpha - \gamma\}^b}.$$

Theorem 4. Let $N \in \mathbb{N}$, $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, p_k/q_k denote the convergents to α and $K = K(N, \alpha)$ be the largest integer satisfying $q_K \leq N$. Further, let $\gamma \in \mathbb{R}$ and $n' \in [0, N]$ be an integer given by (2). Then

$$T_N^b(\alpha, \gamma) \leq \begin{cases} 2^{1+b}\zeta(b)N^b + 2^b\zeta(b)q_{K+1}^b & \text{if } b > 1 \\ \frac{2^{1+b}}{1-b}N + \frac{2^b}{1-b}q_{K+1}^b(N/q_K)^{1-b} & \text{if } b < 1, \end{cases}$$

where $\zeta(b) = \sum_{j=1}^{\infty} j^{-b}$.

Proof. The arguments are similar to the proof of Theorem 1. We therefore only describe the relevant changes in the calculations made in the proof of Theorem 1. To begin with, note that the general case is again reduced to the semi-homogeneous case. First, assume that $b > 1$. Then, in **Case (i)**, inequalities (38) will be replaced by

$$\sum_{j=1}^H \frac{1}{\{(n_{\ell+j} - n')\alpha\}^b} \leq \sum_{j=1}^H \frac{1}{(j\delta_A)^b} < \frac{1}{|D_k|^b} \sum_{j=1}^H j^{-b} < \zeta(b)(2q_{K+1})^b = 2^b\zeta(b)q_{K+1}^b,$$

while (39) will be replaced by

$$\begin{aligned} \sum_{j=H+1}^N \frac{1}{\{(n_{\ell+j} - n')\alpha\}^b} &\leq \sum_{t=1}^{q_K} \sum_{u=0}^H \frac{1}{(t(\delta_B + H\delta_A))^b} \leq \zeta(b) \frac{H+1}{(\delta_B + H\delta_A)^b} \\ &= \zeta(b) \left(\frac{H+1}{(\delta_B + H\delta_A)} \right)^b (H+1)^{1-b} \\ &\stackrel{(42)}{\leq} \zeta(b)(4N)^b (H+1)^{1-b} \\ &\stackrel{(*)}{\leq} \zeta(b)(4N)^b (N/q_K)^{1-b} \\ &= 4^b \zeta(b) N q_K^{b-1} \stackrel{q_K \leq N}{\leq} 4^b \zeta(b) N^b, \end{aligned}$$

where (*) is due to the inequalities $b > 1$ and $H+1 = [N/q_K] + 1 \geq N/q_K$.

In **Case (ii)**, (43) will be replaced by

$$\frac{1}{\{(n_{\ell+1} - n')\alpha\}^b} \leq (2q_{K+1})^b \leq 2^b \zeta(b) q_{K+1}^b,$$

since $\zeta(b) > 1$, while (46) will be replaced by

$$\begin{aligned} \sum_{j=2}^N \frac{1}{\{(n_{\ell+j} - n')\alpha\}^b} &\leq \sum_{t=1}^{q_K} \sum_{u=0}^1 \frac{1}{(t(\delta_B + \delta_A))^b} \leq \zeta(b) \frac{2}{(\delta_B + \delta_A)^b} \\ &= 2^{1-b} \zeta(b) \left(\frac{2}{\delta_B + \delta_A} \right)^b \stackrel{(48)}{\leq} 2^{1-b} \zeta(b) (4N)^b = 2^{1+b} \zeta(b) N^b. \end{aligned}$$

Now assume that $b < 1$. In this case we will use the following inequality instead of (33)

$$\sum_{t=1}^T \frac{1}{t^b} \leq 1 + \int_1^T \frac{dt}{t^b} = \frac{T^{1-b} - b}{1-b} \leq \frac{T^{1-b}}{1-b}.$$

Then, in **Case (i)**, (38) will be replaced by

$$\sum_{j=1}^H \frac{1}{\{(n_{\ell+j} - n')\alpha\}^b} \leq \sum_{j=1}^H \frac{1}{(j\delta_A)^b} = \frac{1}{|D_k|^b} \sum_{j=1}^H j^{-b} < \frac{1}{1-b} (2q_{K+1})^b (N/q_K)^{1-b},$$

while (39) will be replaced by

$$\begin{aligned} \sum_{j=H+1}^N \frac{1}{\{(n_{\ell+j} - n')\alpha\}^b} &\leq \sum_{t=1}^{q_k} \sum_{u=0}^H \frac{1}{(t(\delta_B + H\delta_A))^b} \\ &\leq \left(\frac{H+1}{(\delta_B + H\delta_A)} \right)^b (H+1)^{1-b} \frac{q_K^{1-b}}{1-b} \\ &\leq (4N)^b (2N/q_K)^{1-b} \frac{q_K^{1-b}}{1-b} = \frac{2^{1+b}}{1-b} N. \end{aligned}$$

In **Case (ii)**, (43) will be replaced by

$$\frac{1}{\{(n_{\ell+1} - n')\alpha\}^b} \leq (2q_{K+1})^b \leq \frac{1}{1-b} (2q_{K+1})^b (N/q_K)^{1-b},$$

while (46) will be replaced by

$$\begin{aligned} \sum_{j=2}^N \frac{1}{\{(n_{\ell+j} - n')\alpha\}^b} &\leq \sum_{t=1}^{q_K} \sum_{u=0}^1 \frac{1}{(t(\delta_B + \delta_A))^b} \leq 2^{1-b} \left(\frac{2}{\delta_B + \delta_A} \right)^b \frac{q_K^{1-b}}{1-b} \\ &\leq 2^{1-b} (4N)^b \frac{q_K^{1-b}}{1-b} = \frac{2^{1+b}}{1-b} N^b q_K^{1-b} \leq \frac{2^{1+b}}{1-b} N. \end{aligned}$$

□

Remark 6. Using the same argument as in Corollary 1 one can easily get an analogue of Theorem 4 for sums that involve the distance to the nearest integer. Finally, note that using the partial summation formula (see for example [5]) an interested reader can readily deduce upper bounds for the even more general sums of the form

$$\sum \frac{1}{n^a \{n\alpha - \gamma\}^b} \quad \text{and} \quad \sum \frac{1}{n^a \|n\alpha - \gamma\|^b},$$

where $a \geq 0$ and $b > 0$.

References

- [1] P. ALESSANDRI, V. BERTHÉ, Three distance theorems and combinatorics on words, *Enseignement Mathématique* 44 (1998), 103-132.
- [2] J. BECK, Probabilistic Diophantine approximation I: Kronecker sequences, *Ann. of Math. (2)* 140 (1994), no. 2, 449–502.
- [3] J. BECK, Probabilistic Diophantine approximation: Randomness in Lattice Point Counting, *Springer Monographs in Mathematics*, 2014.
- [4] H. BEHNKE, Zur theorie der diophantischen approximationen, *Abh. Math. Sem. Univ. Hamburg* 3 (1924), no. 1, 261–318.
- [5] V. BERESNEVICH, A. HAYNES, S. VELANI, Sums of reciprocals of fractional parts and multiplicative Diophantine approximation, *Mem. Amer. Math. Soc.* (to appear), arXiv:1511.06862.
- [6] N. CHEVALLIER, Distances dans la suite des multiples d'un point du tore à deux dimensions, *Acta Arith.* 74 (1996), 47-59.
- [7] N. CHEVALLIER, Meilleures approximations d'un élément du tore \mathbb{T}^2 et géométrie de la suite des multiples de cet élément, *Acta Arith.* 78 (1996), 19-35.
- [8] N. CHEVALLIER, Géométrie des suites de Kronecker, *Manuscripta Mathematica* 94, no. 1 (1997), 231-241.
- [9] S. D. CHOWLA, Some problems of diophantine approximation (I), *Math. Z.*, 33 (1931), no. 1, 544-563.
- [10] J. H. HALTON, The distribution of the sequence $\{n\xi\}$ ($n = 0, 1, 2, \dots$), *Proc. Cambridge Phil. Soc.* 71 (1965), 665-670.
- [11] G. HARDY, J. LITTLEWOOD, Some problems of diophantine approximation: The lattice-points of a right-angled triangle, *Proc. London Math. Soc.* (2) 20 (1922), 15-36.
- [12] G. HARDY, J. LITTLEWOOD, Some problems of diophantine approximation: The lattice-points of a right-angled triangle (second memoir), *Abhandlungen aus dem Mathematischen Seminar der Hamburgischen Universität* 9 (1922), 212-249.
- [13] G. HARDY, J. LITTLEWOOD, Some problems of diophantine approximation: a series of cosecants, *Bull. Calcutta Math. Soc.* 20 (1930), 251-266.
- [14] A. Y. KHINTCHINE, Continued fractions. Dover Publications, 1997.
- [15] J. F. KOKSMA, Diophantische Approximationen. (German) Reprint. Springer-Verlag, Berlin-New York, 1974.
- [16] L. KUIPERS, H. NIEDERREITER, Uniform distribution of sequences. Wiley-Interscience, John Wiley & Sons, New York-London-Sydney, 1974.
- [17] A. H. KRUSE, Estimates of $\sum_{k=1}^N k^{-s} \langle kx \rangle^{-t}$. *Acta Arith.* 12 (1966/1967), 229-261.
- [18] S. LANG, Introduction to Diophantine Approximations, Springer Science & Business Media, 1995.

- [19] T. H. LÊ, J. D. VAALER, Sums of products of fractional parts. *Proc. Lond. Math. Soc.* (3) 111 (2015), no. 3, 561-590.
- [20] M. MAYERO, The three gap theorem (Steinhaus Conjecture), *Types for Proofs and Programs, Springer Berlin Heidelberg*, 2000, 162-173.
- [21] T. RAVENSTEIN, The three gap theorem (Steinhaus Conjecture), *J. Australian Math. Soc. (Series A)* 45 (1988), no. 3, 360-370.
- [22] T. RAVENSTEIN, G. WINLEY, K. TOGNETTI, Characteristics and the three gap theorem, *Fibonacci Quarterly* 28 (1990), no. 3, 204-214.
- [23] A. ROCKETT, P. SZÜSZ, Continued Fractions, World Scientific, Singapore, 1992.
- [24] W. M. SCHMIDT, Metrical theorems on fractional parts of sequences, *Trans. Amer. Math. Soc.* 110 (1964), 493-518.
- [25] A. SIEGEL, Théorème des trois longueurs et suites sturmiennes: mots d'agencement des longueurs, *Acta Arith* 97 (2001), no. 3, 195-210.
- [26] Y. SINAI, AND C. ULCIGRAI, Estimates from above of certain double trigonometric sums. *J. Fixed Point Theory Appl.* 6 (2009), no. 1, 93-113.
- [27] N. B. SLATER, Gaps and steps for the sequence $n\theta \bmod 1$, *Proc. Cambridge Phil. Soc.* 73 (1967), 1115-1122.
- [28] V. T. SÓS, On the theory of diophantine approximations. I, *Acta Math. Acad. Sci. Hungar.* 8 (1957), 461-472.
- [29] V. T. SÓS, On the distribution mod 1 of the sequence $n\alpha$, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* 1 (1958), 127-134.
- [30] J. SURÁNYI, Über die Anordnung der Vielfachen einer reellen Zahl mod 1, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* 1 (1958), 107-111.
- [31] S. ŚWIERCKOWSKI, On successive settings of an arc on the circumference of a circle, *Fund. Math.* 46 (1958), 187-189.
- [32] A. WALFISZ, Über einige trigonometrische Summen. (German) *Math. Z.* 33 (1931), no. 1, 564-601.
- [33] A. WALFISZ, Über einige trigonometrische Summen II. (German) *Math. Z.* 35 (1932), no. 1, 774-788.