

This is a repository copy of *Bowing fracture of the inferior angle of the scapula, a difficult diagnosis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/122327/>

Version: Accepted Version

Article:

Miller, Christopher, Grainger, Andrew J, Phillips, Robert S orcid.org/0000-0002-4938-9673 et al. (2 more authors) (2017) Bowing fracture of the inferior angle of the scapula, a difficult diagnosis. *Pediatric radiology*. pp. 1-4. ISSN: 1432-1998

<https://doi.org/10.1007/s00247-017-3945-3>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Dear Author

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style.
- Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s00247-017-3945-3>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

Article Title	Bowing fracture of the inferior angle of the scapula, a difficult diagnosis	
Article Sub-Title		
Article Copyright Year	Springer-Verlag GmbH Germany 2017 (This will be the copyright line in the final PDF)	
Journal Name	Pediatric Radiology	
Author	Family Name	Miller
	Particle	
	Given Name	Christopher
	Suffix	
	Division	Clarendon Wing Radiology Department
	Organization	Leeds Children’s Hospital, Leeds Teaching Hospitals NHS Trust, Leeds Children’s Hospital at The Leeds General Infirmary
	Address	Leeds LS2 9NS, UK
Author	Family Name	Grainger
	Particle	
	Given Name	Andrew
	Suffix	
	Division	Department of Radiology
	Organization	Chapel Allerton Hospital, Leeds Teaching Hospitals NHS Trust
	Address	Leeds, UK
Author	Family Name	Phillips
	Particle	
	Given Name	Robert
	Suffix	
	Division	Department of Paediatric Oncology
	Organization	Leeds Children’s Hospital, Leeds Teaching Hospitals NHS Trust
	Address	Leeds, UK
Author	Family Name	Sabouni
	Particle	
	Given Name	Mohamed
	Suffix	
	Division	Department of Paediatric Orthopaedic Surgery
	Organization	Leeds Children’s Hospital, Leeds Teaching Hospitals NHS Trust
	Address	Leeds, UK
	Family Name	Kraft

Corresponding Author	Particle	
	Given Name	Jeannette K.
	Suffix	
	Division	Clarendon Wing Radiology Department
	Organization	Leeds Children’s Hospital, Leeds Teaching Hospitals NHS Trust, Leeds Children’s Hospital at The Leeds General Infirmary
	Address	Leeds LS2 9NS, UK
	e-mail	Jeannette.Kraft@nhs.net
Schedule	Received	8 April 2017
	Revised	25 June 2017
	Accepted	10 July 2017
Abstract	A 4-year-old boy presented with swelling over the inferior tip of the scapula and an unclear history. Initial radiographic findings were concerning for an aggressive lesion. This case highlights how a multimodality imaging approach was used to relieve uncertainty by diagnosing a paediatric bowing type fracture of the scapular tip.	
Keywords (separated by '-')	Child - Fracture - Magnetic resonance imaging - Radiography - Scapula - Ultrasound	
Foot note information		

Bowing fracture of the inferior angle of the scapula, a difficult diagnosis

Christopher Miller¹ · Andrew Grainger² · Robert Phillips³ · Mohamed Sabouni⁴ · Jeannette K. Kraft¹

Received: 8 April 2017 / Revised: 25 June 2017 / Accepted: 10 July 2017
© Springer-Verlag GmbH Germany 2017

Abstract A 4-year-old boy presented with swelling over the inferior tip of the scapula and an unclear history. Initial radiographic findings were concerning for an aggressive lesion. This case highlights how a multimodality imaging approach was used to relieve uncertainty by diagnosing a paediatric bowing type fracture of the scapular tip.

Keywords Child · Fracture · Magnetic resonance imaging · Radiography · Scapula · Ultrasound

Introduction

Scapula fractures occur infrequently in children and are usually the result of major trauma with multiple injuries. This is because the scapula is well protected by surrounding musculature. Therefore, fractures usually involve the glenoid, coracoid process and acromion. Fractures of the inferior angle of the scapula are very rare in children with only a few case reports in the

literature [1–4]. They usually represent avulsion fractures due to the action of serratus anterior or latissimus dorsi muscles [1].

Case report

A 4-year-old boy presented to the emergency department at an outside institution with swelling over the left scapula noticed by his mother. Earlier that day, he had fallen down stairs and landed on his back without apparent initial sequelae. It was uncertain if the swelling predated the injury. Physical examination revealed a painless lump over the left scapula, with full range of movement at the left shoulder joint. The patient was otherwise well with no significant medical or family history.

Radiographs performed in the emergency department demonstrated an irregular bony mass projecting towards the chest wall from the inferior angle of the scapula (Fig. 1). Routine blood tests including inflammatory markers were normal except for low vitamin D values of 12.7 nmol/L (<30 nmol/L suggests vitamin D deficiency). Blood cultures were negative.

The boy was referred to the paediatric oncology department at our institution as the plain film findings were suspicious for an aggressive bone lesion. A US scan performed 5 days after the initial presentation demonstrated a curved inferior scapular border with an angled cartilaginous tip of the scapula. Associated was an ill-defined mass-like area with increased vascularity and surrounding soft-tissue oedema (Fig. 2). Concerns regarding malignancy triggered further investigations. An MR scan performed 10 days after the initial presentation showed no soft-tissue mass but extensive muscle and soft-tissue oedema surrounding a bony ridge at the inferior angle of the scapula with bone marrow oedema. Post gadolinium marked enhancement was seen in the bone and surrounding tissues (Fig. 3). A CT scan demonstrated a curved scapular tip with surrounding periosteal reaction and early

✉ Jeannette K. Kraft
Jeannette.Kraft@nhs.net

¹ Clarendon Wing Radiology Department, Leeds Children's Hospital, Leeds Teaching Hospitals NHS Trust, Leeds Children's Hospital at The Leeds General Infirmary, Leeds LS2 9NS, UK

² Department of Radiology, Chapel Allerton Hospital, Leeds Teaching Hospitals NHS Trust, Leeds, UK

³ Department of Paediatric Oncology, Leeds Children's Hospital, Leeds Teaching Hospitals NHS Trust, Leeds, UK

⁴ Department of Paediatric Orthopaedic Surgery, Leeds Children's Hospital, Leeds Teaching Hospitals NHS Trust, Leeds, UK



Fig. 1 Lateral radiograph of a 4-year-old boy with scapular bowing fracture. The inferior tip of the scapula appears mass-like with bony irregularity (*arrow*)

with hard callus formation and a well-rounded inferior scapula tip was demonstrated on a radiograph. Clinical examination revealed normal range of shoulder movement.

Discussion

In contrast to the common types of scapula fractures, which usually require high force, inferior angle fractures can be sustained in isolation and with lower levels of trauma. Fractures of the inferior angle of the scapula are very rare. It should be considered that fractures in such an unusual location may be related to non-accidental injury, especially if there is no history of trauma or, as in our case, the causality is not initially apparent. We identified three previously reported cases in children [1–4]. These papers describe the mechanism of injury as avulsion due to the strong action of periscapular muscles such as serratus anterior or latissimus dorsi. In our case, the mechanism of injury is likely direct trauma with a blow to the lower aspect of the scapula that occurred on the day of presentation to the emergency department. Impact on the edge of a step as the boy fell must have caused inwards folding of the scapular tip in a paediatric-type plastic deformation pattern. The diagnostic difficulty arose from the plain radiographic appearances of an aggressive lesion, which is probably related to the radiographic projection and difficulty in depicting the blade of the scapula in a true lateral projection as can be achieved with CT. Initially, the preceding traumatic event was not given enough consideration to suggest an unusual fracture, leading to further investigations and referral to oncology clinic. Therefore, sonography was performed several days after presentation when the injury had started to heal with increased vascularity and granulation tissue suggesting a more aggressive process. As demonstrated in a case report by Szopinski, Adamczyk and Drwiega [2], the cartilaginous part

callus formation (Fig. 4). When compared with the normal right scapula, it was apparent that the inferior tip of the scapula had folded inwards in keeping with a bowing or plastic deformation fracture. The boy was managed conservatively as he was pain free. At follow-up 3 months later, a healing fracture





Fig. 2 Sonograms of a 4-year-old boy with scapular bowing fracture. **a** Long section through the inferior tip of the scapula shows a curved scapular edge (*short arrows*) with angulation just above the cartilaginous tip (*long arrow*). **b** Transverse section demonstrates a

mass-like area (*arrow*) with increase in vascularity on colour Doppler related to healing of the fracture. **c** Long section through the inferior tip of the right scapula is shown for comparison

Fig. 3 MR images of a 4-year-old boy with scapular bowing fracture. **a** Sagittal STIR image of the left scapula demonstrates marked soft tissue and muscle oedema surrounded by the folded inferior tip of the scapula (*arrow*). **b** Axial gadolinium-enhanced T1 fat-saturated image demonstrates prominent enhancement within the soft tissues surrounding the scapular tip



of the scapular tip, which ossifies around puberty, is well-demonstrated with US. Angulation of it in our case probably suggests associated detachment at the bone cartilage interface (Fig. 2). This could be related to direct trauma or an avulsion type component related to the action of the serratus anterior muscle on the inferior aspect of the scapula. Avulsion injuries of the tip of the scapula are rare but have been previously described [1–4]. However, the case presented here is not a simple avulsion injury. Bowing of the scapula was not described in any of the previously published cases we identified and detachment or dysfunction of the serratus anterior muscle would lead to winging of the scapula, which was not observed in our case [1, 5]. Another two cases previously described paediatric-type greenstick fractures in children [6, 7]. Similar to our case, both cases occurred after low-energy trauma. However, these greenstick fractures were associated with significant scapular winging.

In our case, the fracture was stable against the chest wall, which probably explains why the child presented with a painless lump. It is likely that the fracture was initially painful, which was not communicated by the young child. Once a fracture is stabilised, it usually becomes pain free.

MRI is not commonly used to investigate fractures in children. Therefore, an unusual healing fracture in an unusual location is difficult to diagnose by MRI, especially in a child referred from oncology clinic. The marked muscle oedema and enhancement together with early callus formation suggested a more aggressive or inflammatory lesion (Fig. 3). CT, however, provides superior bone detail suitable to demonstrate periosteal reaction and callus formation. CT is usually reserved for complex fractures and preoperative planning, but its multiplanar and surface-rendered 3-D imaging capabilities finally revealed the true nature of the lesion (Fig. 4). A previous case report details a similar situation where a suspicious lesion was seen on plain film and MRI, with a bone biopsy only avoided when a fracture line was identified on the chest CT performed for staging purposes [8].

In contrast to a reported case of avulsion of the cartilaginous tip of the inferior angle of the scapula that was surgically treated, our patient was conservatively managed with no ill effects to his shoulder and scapular function [2]. Chang et al. [1] recently reviewed the literature and identified 10 cases of inferior angle fractures. The review

Fig. 4 CT images of a 4-year-old boy with scapular bowing fracture. **a** Sagittal CT image (bone window) shows angulation of the tip of the scapula with callus formation (*arrow*). **b** 3-D reconstruction demonstrates the folded and rounded tip of scapula (*arrow*) that leads to shortening of the scapular body when compared to the normal right side



139 included 2 children (a 13- and a 17-year-old) who both had
 140 undisplaced fractures successfully managed conservative-
 141 ly. The paper suggests that displaced fractures, however,
 142 should be surgically treated to avoid painful non-union [1].
 143 This case illustrates how multimodality imaging can be
 144 utilised to reach a diagnosis when plain films reveal un-
 145 usual appearances and the history is uncertain. Whilst in
 146 hindsight US and MRI could have been avoided, neither of
 147 these carries a radiation burden. It was important to recog-
 148 nise this rare fracture involving the inferior angle of the
 149 scapula and exclude sinister pathology.

150 **Compliance with ethical standards**

151 **Conflicts of interest** None.
 174

References

1. Chang AC, Phadnis J, Earldy-Harris N et al (2016) Inferior angle of the scapula fractures: a review of literature and evidence-based treatment guidelines. *J Shoulder Elb Surg* 25:1170–1174
2. Szopinski KT, Adamczyk G, Drwiega M (2012) Traumatic detachment of the inferior angle of the scapula in a 5-year-old boy – a sonographic diagnosis. *Skelet Radiol* 41:615–618
3. Heyse-Moore GH, Stoker DJ (1982) Avulsion fractures of the scapula. *Skelet Radiol* 9:27–32
4. Brindle TJ, Coen M (1998) Scapular avulsion fracture of a high school wrestler. *J Orthop Sports Phys Ther* 27:444–447
5. Franco M, Albano L, Blaimont A et al (2004) Spontaneous fracture of the lower angle of scapula. Possible role of cough Joint Bone Spine 71:580–582
6. Tryfonidis M, Reynolds J, Ostlere S et al (2008) Greenstick fracture of the scapular blade; an unusual case of winging of the scapula. *Injury Extra* 39:130–133
7. Bowen TR, Miller F (2006) Greenstick fracture of the scapula: a cause of scapular winging. *J Orthop Trauma* 20:147–149
8. Kim MJ, Rinsky LA, MacKenzie JD (2011) Scapular stress fracture initially masquerading as tumor or infection. *Radiol Case Rep* 6:510

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES.

- Q1. Please check the exact presentation of author names including any initial(s) keeping in mind that this will remain the spelling/format used in any future citation of this paper.
- Q2. Please check if the affiliations are presented/captured correctly.

UNCORRECTED PROOF