



This is a repository copy of *The Typhoid Vaccine Acceleration Consortium (TyVAC): Vaccine effectiveness study designs: Accelerating the introduction of typhoid conjugate vaccines and reducing the global burden of enteric fever. Report from a meeting held on 26-27 October 2016, Oxford, UK.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/122899/>

Version: Published Version

Article:

Meiring, J.E., Gibani, M., Basnyat, B. et al. (23 more authors) (2017) The Typhoid Vaccine Acceleration Consortium (TyVAC): Vaccine effectiveness study designs: Accelerating the introduction of typhoid conjugate vaccines and reducing the global burden of enteric fever. Report from a meeting held on 26-27 October 2016, Oxford, UK. *Vaccine*, 35 (38). pp. 5081-5088. ISSN 0264-410X

<https://doi.org/10.1016/j.vaccine.2017.08.001>

Reuse



This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Conference report

The Typhoid Vaccine Acceleration Consortium (TyVAC): Vaccine effectiveness study designs: Accelerating the introduction of typhoid conjugate vaccines and reducing the global burden of enteric fever. Report from a meeting held on 26–27 October 2016, Oxford, UK

James E. Meiring^{a,*}, Malick Gibani^a, the TyVAC Consortium Meeting Group: Buddha Basnyat^{a,b}, Adwoa D. Bentsi-Enchill^c, John Clemens^d, Thomas C. Darton^{e,f}, Kashmira Date^g, Gordon Dougan^h, Denise Garrettⁱ, Bradford D. Gessner^j, Melita A. Gordon^{k,l}, Robert S. Heyderman^m, Joachim Hombach^c, Karen L. Kotloffⁿ, Myron M. Levineⁿ, Stephen P. Luby^o, Vadrevu Krishna Mohan^p, Anthony A. Marfin^q, Kim Mulholland^r, Kathleen Neuzilⁿ, Virginia E. Pitzer^s, Andrew J. Pollard^e, Firdausi Qadri^d, David Salisbury^t, Anita Zaidi^u

^a Oxford University Clinical Research Unit, Patan Academy of Health Sciences, Kathmandu, Nepal

^b Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, United Kingdom

^c Department of Immunization, Vaccines and Biologicals, World Health Organization, Geneva, Switzerland

^d International Centre for Diarrhoeal Diseases Research, Bangladesh (icddr), Dhaka, Bangladesh

^e Oxford Vaccine Group, Department of Paediatrics, University of Oxford, and the NIHR Oxford Biomedical Research Centre, Oxford, United Kingdom

^f The Hospital for Tropical Diseases, Wellcome Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

^g Global Immunization Division, Center for Global Health, Centers for Disease Control and Prevention (CDC), Atlanta, USA

^h The Wellcome Trust Sanger Institute, Hinxton, Cambridgeshire, United Kingdom

ⁱ Sabin Vaccine Institute, Washington DC, USA

^j Agence de Médecine Préventive, Paris, France

^k Malawi Liverpool Wellcome Trust Clinical Research Programme, University of Malawi College of Medicine, Blantyre, Malawi

^l Institute of Infection and Global Health, University of Liverpool, Liverpool, United Kingdom

^m Division of Infection and Immunity, University College London, London, United Kingdom

ⁿ Center for Vaccine Development, University of Maryland School of Medicine, Baltimore, USA

^o Division of Infectious Diseases and Geographic Medicine, Stanford University, Stanford, CA 94305, United States

^p Bharat Biotech International Limited, Hyderabad, India

^q PATH, Seattle, USA

^r Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK

^s Department of Epidemiology of Microbial Diseases, Yale School of Public Health, Yale University, New Haven, CT, USA

^t Centre on Global Health Security, Chatham House, London, UK

^u Enteric and Diarrheal Diseases, Global Health, Bill and Melinda Gates Foundation, Seattle, USA

^a Oxford Vaccine Group, Department of Paediatrics, University of Oxford, and the NIHR Oxford Biomedical Research Centre, Oxford, United Kingdom

ARTICLE INFO

Article history:

Received 16 June 2017

Received in revised form 31 July 2017

Accepted 1 August 2017

Available online 10 August 2017

Keywords:

Typhoid
Vaccines
Policy
Effectiveness
Clinical trials
Global health

ABSTRACT

Typhoid fever is estimated to cause between 11.9–26.9 million infections globally each year with 129,000–216,510 deaths. Access to improved water sources have reduced disease incidence in parts of the world but the use of efficacious vaccines is seen as an important public health tool for countries with a high disease burden.

A new generation of Vi typhoid conjugate vaccines (TCVs), licensed for use in young children and expected to provide longer lasting protection than previous vaccines, are now available. The WHO Strategic Advisory Group of Experts on Immunization (SAGE) has convened a working group to review the evidence on TCVs and produce an updated WHO position paper for all typhoid vaccines in 2018 that will inform Gavi, the Vaccine Alliance's future vaccine investment strategies for TCVs.

The Typhoid Vaccine Acceleration Consortium (TyVAC) has been formed through a \$36.9 million funding program from the Bill & Melinda Gates Foundation to accelerate the introduction of TCVs into Gavi-eligible countries.

* Corresponding author.

E-mail address: james.meiring@paediatrics.ox.ac.uk (J.E. Meiring).

In October 2016, a meeting was held to initiate planning of TCV effectiveness studies that will provide the data required by policy makers and stakeholders to support decisions on TCV use in countries with a high typhoid burden.

Discussion topics included (1) the latest evidence and data gaps in typhoid epidemiology; (2) WHO and Gavi methods and data requirements; (3) data on TCV efficacy; (4) cost effectiveness analysis for TCVs from mathematical models; (5) TCV delivery and effectiveness study design. Specifically, participants were asked to comment on study design in 3 sites for which population-based typhoid surveillance is underway.

The conclusion of the meeting was that country-level decision making would best be informed by the respective selected sites in Africa and Asia vaccinating children aged from 9-months to 15-years-old, employing either an individual or cluster randomized design with design influenced by population characteristics, transmission dynamics, and statistical considerations.

© 2018 Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Typhoid fever is estimated to cause between 11.9 million–26.9 million cases and 129,000–216,510 deaths annually [1–3]. The burden of disease is largely within low- and middle-income countries primarily throughout Asia and Africa [4,5]. Whilst improvements in drinking water quality have been successful in reducing rates of typhoid fever in certain parts of the world [6–8], control has been hampered elsewhere and typhoid remains a significant public health problem [1,3].

In 2008 the World Health Organization (WHO) recommended the consideration of two licensed typhoid vaccines (Vi-polysaccharide and Ty21a), for programmatic use by countries with high rates of typhoid fever for controlling endemic disease, as well as for use in outbreak settings and for travelers to endemic areas [9]. However, with the expectation of second generation typhoid conjugate vaccines (TCV) becoming available in the near future Gavi, the Vaccine Alliance, deferred decisions on funding support until TCVs were licensed and prequalified by WHO [10,11].

Two Vi tetanus toxoid conjugate vaccines have now been licensed in India, with one manufacturer (Bharat Biotech) applying for WHO prequalification [12,13]. Due to these developments, the WHO Strategic Advisory Group of Experts on Immunization (SAGE) has convened a working group to 'review the scientific evidence and relevant programmatic considerations to formulate updated recommendations on the use of typhoid vaccines' [14]. Specific consideration will be given to estimates of disease burden, vaccine schedules, and economic analyses of vaccination programs. Publication of an updated WHO position paper on typhoid vaccines is scheduled for 2018 [14]. These recommendations will be highly relevant to Gavi, given their prior positive statements on support for TCV.

At this key stage in typhoid vaccine development, the Bill & Melinda Gates Foundation (BMGF) published a request for proposals to accelerate the introduction of TCVs into Gavi-eligible countries and to contribute to the data to support the use of TCVs as means of reducing the global typhoid burden. With \$36.9M from BMGF, the Typhoid Vaccine Acceleration Consortium (TyVAC) was formed in 2016 comprising core partners at the University of Maryland's Center for Vaccine Development, the University of Oxford's Oxford Vaccine Group and PATH to achieve this goal [15].

A meeting of key stakeholders was held in Oxford, UK in October 2016 to discuss some of the critical issues surrounding typhoid fever and the impact on TCV effectiveness study design.

2. Typhoid vaccine acceleration consortium

In her introductory remarks, Dr. Anita Zaidi, Director of Enteric Diseases at BMGF, highlighted the motivations for this program on behalf of the foundation. Firstly, typhoid fever is primarily a disease of the poor which will require coordinated public health interventions to prevent and with the rising concern of antimicrobial

drug resistance the global situation may yet get worse [16]. Secondly, with the recent development of TCVs the case for global typhoid control is compelling and there may be opportunity to demonstrate a dramatic impact through vaccination.

Following this the Director of TyVAC, Professor Kathleen Neuzil, outlined some of the key objectives for the consortium;

- To serve as a coordinating body for typhoid-related research and control activities
- To foster supportive global policies
- To ensure typhoid and TCVs are recognized as global, regional and national health priorities
- To provide data on impact, effectiveness, appropriate vaccine strategies and cost of TCV use
- To support countries in decision-making and preparation for sustained TCV introduction

Dr. Neuzil also reviewed lessons learned from other successful vaccine introduction efforts, and the need for clear goals and stakeholder involvement. Further, as TCVs are licensed, the goal of these studies will be to inform policy and financing decisions – thus the studies must be designed in light of that goal. Current funding only allows these studies to be done in a limited number of settings. Therefore, we must ensure that these settings and designs are sufficiently generalizable to inform non-trial countries in translating results to their local settings.

3. WHO perspectives

Understanding the data required by WHO to recommend TCVs is an important consideration for the consortium. Two central policy issues are currently under review by the WHO, (1) should TCV be recommended over Vi-polysaccharide (ViPS) and Ty21a vaccines for use in persons 2 years of age and older? (2) should TCV be recommended for routine use in children <2 years of age and what should be the lower age limit for use in this group? Recent ad-hoc WHO consultations to discuss the initiation of a SAGE policy pathway for TCVs identified multiple gaps in the data required to inform these policy decisions (Table 1). The design of TCV effectiveness studies should be made with these specific data gaps in mind, to provide the critical data that policy makers at global, regional and country levels will require to adopt and promote these new vaccines.

The SAGE Working Group on Typhoid Vaccines' review of the data to support TCV use will lead to consideration by SAGE for policy recommendations in October 2017.

4. Gavi

Understanding data used by Gavi before investing in TCV introduction is an important consideration for the consortium. In 2008,

Table 1

Proposed data gaps for near-to-medium term research agenda to guide TCV policy and use (Used with permission from WHO; TCV Data Gaps Meeting Report September 2015 http://www.who.int/immunization/research/meetings_workshops/typhoid_conjugate_vaccine_sept15/en/).

Critical 1–3 year research agenda Evidence generation	Critical 1–3 year research agenda Vaccine development	Critical 2–5 year research agenda Typhoid fever control strategies
<ol style="list-style-type: none"> 1. Surveillance projects using standardized methodology (e.g., SETA^a, SEAP^b) – ongoing 2. Disease transmission and carriage – ongoing 3. Cost-effectiveness analysis and modelling – ongoing 4. Diagnostics – different types (Point-of-care, surveillance, environmental) 	<ol style="list-style-type: none"> 1. Assay & reagents development and standardization – immediate 2. Oxford University Controlled Human Infections Models – ongoing 3. Clinical trial endpoints/study design for new vaccines – immediate 4. WHO global consensus on the requirements for licensure 5. WHO global reference laboratories and reagents 	<ol style="list-style-type: none"> 1. Integration with new WaSH^c initiatives 2. Achievement of control and prevention in targeted communities 3. Need new vaccines to achieve licensure 4. Vaccine immunization strategies and schedules 5. Environmental surveillance

^a Severe Typhoid in Africa Program.

^b Surveillance of Enteric Fever in Asia Project.

^c Water, Sanitation and Hygiene.

Gavi instituted a new method to identify vaccines that would be supported in Gavi-eligible countries. In order to evaluate each vaccine in the same objective manner, Gavi proposed evaluation indicators including overall health impact (e.g., childhood mortality, all-age mortality/morbidity); other vaccine impact considerations (e.g., potential to stop epidemics, indirect protection [“herd immunity”]), implementation feasibility (including ease of programmatic integration) and cost-effectiveness. In the future, long-term morbidity, equity, and overall affordability may feature prominently in Gavi investment decisions. Many of these evaluation indicators will benefit from the availability of vaccine effectiveness study data.

Gavi has previously considered typhoid vaccine support. Despite a 2007 SAGE recommendation of live, oral Ty21a vaccine (3–4 dose schedule) and injectable ViPS vaccine (single dose schedule) by typhoid-endemic countries [9], with the expectation of near-term TCV availability, a 2008 Gavi Vaccine Investment Strategy (VIS) Working Group recommended that Gavi should not invest in typhoid vaccination in its 2009–13 VIS. This working group cited the limited duration of protection given by the Ty21a and ViPS vaccines that would necessitate re-vaccination every 3 years and the inability to use these vaccines in children less than 2 years of age as important limitations of those vaccines. In 2008, Gavi expressed a strong preference for TCVs which could theoretically be administered through routine vaccination of infants as part of a national immunization schedule following catch-up campaigns.

At this time, it is not known if Gavi will consider TCV support based on the 2009–2013 VIS or whether it will need to be considered in new discussions related to the 2019–2023 VIS. Although this is not known, the consortium discussed the importance of trial design, campaign strategy, and types of data that are collected to inform VIS working groups.

Whilst providing data to support a Gavi investment is important, it is also important to consider data needed to inform introduction decisions by countries that are not eligible for Gavi support. Engaging with country-level policymakers will help the consortium understand perceived need for typhoid vaccinations and identify the data needed by lower middle-income countries to support the introduction of TCVs. Policymaker surveys conducted as part of the Disease of Most Impoverished (DOMI) program identified that typhoid vaccine introduction would be best facilitated with country-specific disease burden data, efficacy studies in local populations and evidence of potential economic savings from vaccination [17].

5. Burden of disease

Precise estimates of the global burden of disease are difficult to establish owing to the non-specific presentation of disease, the lack of reliable diagnostic facilities in many regions, and the insensitivity of commonly used diagnostic tests [18]. Additionally, the burden of typhoid disease can fluctuate widely depending on geographic context and the occurrence of epidemics. Currently it is estimated that typhoid fever is responsible for between 11.9 million–26.9 million cases and 129,000–216,510 deaths annually [1–3,19,5].

Burden of disease studies performed to date have shown high rates of typhoid fever throughout South and South East Asia. Within these high-burden regions, the epidemiology of typhoid is complicated by marked inter- and intra-country variation. For example, data from the DOMI program highlighted incidence rates varying from 24.2/100,000 in Vietnam, to 493.5/100,000 in India (per 100,000 person-years) [20]. Surveillance throughout Asia has predominantly been undertaken in urban centers with consistently high rates of disease [21–24], but less is known about the rates of disease in rural areas [8].

Historically, there has been less information available regarding disease epidemiology in Africa [19]. Fewer studies have been conducted and the heterogeneity of disease may be higher [25]. Surveillance performed in two sites in Kenya between 2006 and 2009 found that the incidence of blood-culture proven typhoid fever in rural and urban sites varied from 29 up to 247 cases per 100,000 person-years of observation (PYO), although these are crude rates [26]. Recent data from the Typhoid Fever Surveillance in Africa Program (TSAP) [27] highlighted marked differences in incidence rates between sites in Africa with adjusted rates ranging from 0 in Sudan to 383 per 100,000 PYO in Burkina Faso [28]. This study also demonstrated marked intra-country variation with higher rates in rural Ghana compared with an urban setting [29].

The uncertain global disease burden of typhoid fever presents several challenges. Firstly, conclusions drawn from vaccine effectiveness studies need to be generalizable to different epidemiological contexts, so that individual countries can make informed decisions on vaccine introduction. With the heterogeneity of disease between and within countries, policy makers will require applicable data for their context.

Secondly, the design of any effectiveness trial must be robust enough to provide accurate data despite unpredictable rates of disease. This has a direct impact on the design of any trial as data will need to be analyzed concurrently in the intervention and control

arm to minimize the bias produced by any significant change in incidence over time. Finally, disease burden and severity estimates are important to policy makers when considering cost-effectiveness analyses and cases/deaths averted [30].

6. Site-specific data

Specifically, for TyVAC, three sites (See interactive maps for Dhaka, Bangladesh; Kathmandu, Nepal and Blantyre, Malawi) had been evaluated prior to this meeting in which to conduct planned vaccine effectiveness studies. Each site is currently conducting detailed disease burden studies as part of the STRATAA program [31] (Supplemental Table 2), which could be leveraged for these planned studies. As such, there will be recent epidemiological data on the incidence of blood culture confirmed typhoid fever, as well as rates of sub-clinical infection identified by serosurveillance and the likely prevalence of chronic carriers in all three sites.

7. Efficacy of TCVs

Proof-in-principle of TCV efficacy is derived from trials of the US NIH Vi-rEPA vaccine, and data from these trials were reviewed to inform future trial designs. Efficacy of up to 92% at 24 months and 89% at 46 months has been demonstrated in trials of the Vi-rEPA vaccine carried out in Vietnam, when given as a two-dose schedule in 2–5 year olds [32,33]. A post hoc analysis of trial participants administered a single dose of the Vi-rEPA vaccine indicated a protective efficacy of approximately 88% [34]. However, these trials do not address the issues of TCV efficacy in children under the age of 2 years.

Whilst efficacy data are available for Vi-rEPA, it is unclear whether similar results would be observed with TCVs that use other carrier proteins, such as tetanus toxoid. Immuno-bridging from the Vi-rEPA studies would be possible in an ideal situation, and efforts to standardize assays to compare products are ongoing.

Several TCVs are currently in development, including Vi-CRM₁₉₇ and Vi-diphtheria toxoid conjugates, many of which are in Phase 1 and 2 trials [35,13]. There are currently two Vi-tetanus toxoid conjugates licensed on the basis of immunogenicity. One such vaccine, PedaTyph™ (a Vi-polysaccharide tetanus toxoid conjugate manufactured by Bio-Med®) has been trialed in a school-based cluster randomized study in an urban slum setting of Kolkata, India, although published data are currently limited in size to allow for firm conclusions [12].

Clinical and immunogenicity data on the other tetanus-conjugate Vi vaccine Typbar-TCV® was presented by Bharat Biotech. Typbar-TCV® is the first TCV to be submitted for prequalification by WHO. Immunogenicity data were derived from a phase 3 study, comprising an open label arm of 327 children aged 6 months to 2 years, and a randomized control arm of 654 individuals aged 2–45 years who received either Typbar-TCV® or a comparator Vi-polysaccharide vaccine (Typbar Vi). In these studies, Typbar-TCV® induced consistently higher anti-Vi IgG responses than Vi-polysaccharide (anti-Vi IgG 1292–1937 EU/ml Typbar-TCV® vs. 411 EU/ml Typbar Vi) and was noted to be immunogenic in children aged under 2 years. The avidity of anti-Vi IgG following Typbar-TCV vaccination was increased compared with that following Typbar-Vi vaccination. Long term immunogenicity data are available for between three to five years post vaccination, which indicate long term persistence of anti-Vi [13].

8. Cost-effectiveness

A major aim of the TyVAC program will be to generate measures of cost-effectiveness within the study sites that could be scaled up

to aid future country-level decisions on TCV introduction. The consortium includes a group of health economics specialists who will engage at an early stage with the aim of tailoring trial design to capture relevant economic data to undertake a prospective cost-effectiveness analysis.

Currently unpublished modelling data were presented to review the predicted cost-effectiveness of TCVs across different settings with different epidemiological and health-economic characteristics. Routine vaccination at 9 months of age was predicted to result in a substantial decline in disease incidence at all sites, with an additional benefit of catch-up campaigns, both in terms of disease incidence and disability adjusted life years (DALYs) averted. When viewed from a healthcare provider perspective, routine vaccination alone at 9 months of age was predicted to be “cost-effective”, “very cost effective” or “cost saving” in most of the settings modelled, with the exception of areas with low incidence of disease and low cost of illness. Additional cost-effectiveness benefits were predicted to be achieved by one-time catch up campaigns – including routine vaccination at 9 months with catch up to the age of 5 years, 15 years, 25 years or to include all ages.

To undertake a prospective cost-effectiveness analysis, economic endpoints should be included as secondary objectives of effectiveness trials. The cost-effectiveness models presented primarily considered healthcare provider costs, rather than individual and societal costs resulting from typhoid infection. In part, this results from the absence of empirical economic data on these potentially important metrics. It was recognized that within the context of a trial, data on patient-level expenditure and health utilization could be collected with relatively limited additional effort using a well-designed case report form. Such data could subsequently be assigned monetary costs that form the basis of cost-effectiveness analyses and improved modelling estimates.

9. Vaccine delivery strategies

The meeting considered a range of different vaccine delivery strategies, which could be applied either in an effectiveness trial or to provide evidence for future programmatic use. It was recognized that there are distinct benefits and limitations of each delivery strategy, which must be balanced against overall feasibility and the need to fill specific data gaps.

9.1. Mass vaccination

A strategy of routine vaccination starting at 9 months of age and mass vaccination of the population at risk, with no upper age limit, was discussed by the meeting attendees. This approach is likely to be the most impactful strategy, particularly in terms of direct effects of TCVs as well as indirect effects. A mass-vaccination strategy could be a test of the feasibility of eventual disease elimination in a confined area and would increase the likelihood for vaccination of high-risk groups, including food handlers, non-immune adults from non-endemic areas moving into endemic areas as well as those with medical co-morbidities. It is unknown whether countries can afford this approach or external donors such as Gavi will support it.

9.2. EPI vaccination plus catch-up campaigns

Multiple variations of a delivery strategy incorporating TCVs into the routine childhood (“EPI”) vaccination schedule with a concomitant catch-up campaign were discussed. The proposed catch-up campaigns could be used in individuals up to 5 years, 15 years or 45 years of age, as outlined in modelling studies.

Such a strategy is the most likely platform for post-study use in endemic countries. Introduction of TCVs into the existing EPI

schedule is anticipated to result in high vaccine coverage, as was seen with similar catch-up campaigns for other vaccines (e.g. MenAfrivac). Whilst this approach may result in a less pronounced effect on disease burden compared with a mass vaccination campaign discussed above, this might be balanced by increased feasibility.

9.3. School-based vaccination

Vaccination of school-age children (defined as those aged 5–15 years) was also considered as a delivery strategy. An advantage of this approach is that it is a convenient group to access and achieve high coverage depending on the proportion of children in high risk age groups who attend school in the target population. Indirect vaccine effects could also be estimated, particularly in those aged under 2 years, by measuring transmission within households of vaccinated school children.

The cost of delivery of a school-based campaign might be high comparing with integration within the EPI schedule. School-based vaccination strategies could also raise issues of equity, either by missing children not enrolled in schools due to economic issues or gender imbalance, and would require engagement with both public and private school systems.

10. Trial design

The performance of TCVs could, in theory, be studied using several different trial designs, each with a distinct set of benefits and limitations. Competing priorities to be addressed through these trials include the desire to inform financing decisions; define the extent of indirect protection conferred by TCVs; define efficacy in children under 2 years of age; or, simply, feasibility.

Data describing TCV immunogenicity and efficacy has already been published for certain settings [12,13,36]. What is now required for policy makers are effectiveness studies, performed in ordinary conditions providing real-world data on both the protection provided by a vaccine and also the practicalities of implementation.

General issues included the need to design an ethical, unbiased trial which is acceptable to the population at study sites. Blinding of both participants and study staff was identified as a key question to consider, in particular with regards to ascertainment of cases during the follow up period after vaccination, as knowledge of vaccination status might affect care-seeking, or risk-taking behaviour.

It was recognized that different trial designs might be more, or less, appropriate for different study sites. However, introducing different trial designs, or target populations to be enrolled, risks complicating cross-site conclusions as evidenced through the divergent findings generated by cluster randomized trials of ViPS vaccine in Kolkata and Karachi [37,38,49].

Observational studies were rejected as means of evaluating TCV impact due to the high risk of bias with a disease like typhoid because of its epidemic potential and seasonal variation. Whilst no single trial design received unanimous support, the strengths and weaknesses of four designs were proposed to answer the specific questions related to typhoid fever vaccination.

10.1. Individual randomization

In an individually randomized control trial, participants would be randomly allocated to receive either TCV or control vaccine. Large scale, individually randomized controlled trials have been used to determine efficacy of conjugate vaccines for Hib and pneumococcal disease [39,40], as well as for typhoid fever. [32,33] Such a trial design offers several advantages, including feasibility and smaller sample size requirements, compared with other trial

designs discussed below. Importantly, an individually randomized trial could be used to determine the direct efficacy of TCVs in children under 2 years of age, which has been identified as a major knowledge gap. The estimate of direct efficacy would be independent of herd effects, which might vary depending on geographic setting or the age-group targeted. A major drawback of this approach is that an individually randomized trial would not generate data on indirect and overall protection of TCVs.

10.2. Cluster randomized control trials

In a cluster randomized trial (CRT), population-based clusters would be defined and randomized to receive either the intervention vaccine or control vaccine. In this design, a percentage of each individual cluster would be vaccinated, and total protection assessed by comparing the incidence of disease in vaccinated individuals in the vaccinated arm, compared with those who received the control vaccine in control clusters. This design was successfully applied in two large trials of ViPS vaccine in Karachi and Kolkata as part of the DOMI program [37,38,41]. In a separately funded study, plans are underway to evaluate effectiveness of TCV for a public-sector implementation group using a cluster randomized approach in NaviMumbai, India. (Personal communication with CDC.)

One of the main strengths of CRTs is the capacity to produce data on total as well as indirect and overall protection of TCVs under blinded conditions. It, arguably, is more reflective of a 'real-world' situation, than an individually randomized study, where less than 100% of a population will be vaccinated and herd protection can be measured.

Drawbacks of the CRT approach include the need for a larger sample size than an individually randomized study design. With the CRT design, there is a risk of 'contamination' between clusters by movement of persons into, out of, or between clusters. The proposed study sites represent densely populated urban areas with high levels of migration and population mixing between clusters, which could reduce all effectiveness estimates. The risk of "contamination" could be mitigated either by incorporating geographic "buffer zones" between clusters or by performing surveillance and analysis of the inner area of a cluster only (the so-called 'fried-egg' design).

10.3. Step wedge

The step-wedge trial (STW) design represents a type of cluster-randomized trial. In a STW [42] instead of randomization by geographic region as in a cluster randomized control trial, the time of intervention within each cluster is randomized. Clusters are phased in for vaccination in a random order at regular intervals, such that by the end of the study all clusters have crossed over. Vaccine effectiveness is assessed through comparison of the incidence in clusters already vaccinated versus those not yet phased in [43]. One proposed advantage of STWs is the need for smaller sample sizes to demonstrate impact, as each cluster acts as its own control [44]. However, it was noted that a step wedge represents a poorly controlled and usually underpowered CRT. In both step-wedge designs and CRT, there is a need to consider seasonality and potential epidemics from point sources, to avoid an accumulation of cases in one cluster or on one side of the 'wedge'. It is likely these reasons have played a role in STWs rarely being used for vaccine evaluations [43,45].

10.4. Ring vaccination

In a ring-vaccination study, an index positive case would be identified by the vaccination team, who would proceed to identify contacts, and contacts of contacts in a cluster. Individuals within

the single cluster, or the entire cluster, could be randomized to either the active vaccine or a control with the aim of interrupting the transmission cycle. Both groups would be under follow-up for signs of clinical disease [46]. Recent experience of this trial design, taken from the West African Ebola outbreak, was given [47].

The major role of a ring-vaccination approach would be in the measurement of TCV effectiveness in the context of a typhoid epidemic. Such an approach might, theoretically, be relevant for testing typhoid vaccines in settings such as in Fiji, where typhoid frequently occurs in multiple small outbreaks [48]. The predicted high-incidence of cases in each ring around an identified index case reduces the overall sample size requirements compared with other designs. Perhaps the major drawback of the ring-vaccination approach for typhoid relates to uncertainties surrounding disease transmission. If, typhoid transmission is via the long-cycle route, contacts of cases have likely already been exposed to the same environmental source, or don't share the same risk factors as the cases [49,50].

A general consensus at the meeting was reached recommending a cluster randomized trial design, if feasible, to capture both total and overall effects with vaccination starting at 9 months of age and a catch up campaign to 15 years of age. Whilst the modeling data suggested that vaccination up to the age of 45 years is likely to have the largest impact and remain cost-effective, this approach is unlikely to be feasible nor affordable within the context of the trials. It would also be less likely to be supported by financing agencies or individual governments. A nested immunogenicity study with individual randomization in the under 2-year age group was also considered.

11. WASH considerations

TCV effectiveness studies will be undertaken in the context of ongoing water, sanitation and hygiene (WASH) interventions, which might lead to a significant reduction in typhoid disease incidence independent of vaccines. This presents two risks to a typhoid vaccine trial. Firstly, a trial with a small geographic footprint may be undermined by improvements in water supply during the process of the trial. Secondly, the perception of local residents that what is actually required is not vaccines but improvements to infrastructure may affect enrollment and community acceptance.

Mitigation of these risks could be possible through surveillance over a large geographic area and inclusion of a water intervention into an arm of the trial. The focus of this particular program however is on vaccine effectiveness, and introducing water improvement was deemed beyond the scope of the planned trials as well as potentially confounding results.

12. Measuring outcomes

The selection of robust outcome measures will be central to the design of TCV effectiveness studies. The outcome recorded due to the vaccination will affect the generalizability of data generated. Whilst there is no perfect outcome measure, some have clear advantages.

The primary outcome of TCV effectiveness studies should be to measure the impact on the burden of typhoid disease using the incidence of blood-culture positive cases as a fixed endpoint.

Clinically defined endpoints are problematic due to the non-specific case presentation of typhoid and range of alternate diagnoses [48–51]. A careful clinical case definition for enrolment and collection of blood cultures will have to be defined. Exploratory diagnostics (e.g. antibody in lymphocyte supernatant (ALS),

culture-PCR) could be used to broaden the case-definition in some cases, as secondary/exploratory endpoints.

Surveillance at the community level may increase the detection rate of uncomplicated disease and therefore an increased burden, but measuring severe disease averted, intestinal perforation, or hospitalization, for example [52], may be more impactful. With non-specific outcomes, all-cause mortality is unlikely to be attainable to measure with the relatively low mortality rate associated with typhoid requiring a large sample size. Impact could be seen, however, with a reduction in hospital febrile admission or outpatient visits, antimicrobial usage or change in antimicrobial resistance patterns. Where clinical outcomes have been used to measure vaccine preventable disease incidence, much higher estimates of impact have been described [53]. In pneumococcal conjugate vaccine studies, for example, a 10-fold increase in impact was observed when non-specific clinical outcomes were measured rather than an etiology-confirmed endpoint [40]. In the context of typhoid fever where blood culture is estimated to be between 50–75% sensitive [18], there will be a significant proportion of vaccine-preventable but non-etologically-confirmable disease; therefore, measuring non-specific outcomes could be an important factor to build into the program.

13. Conclusion

Typhoid is an important cause of morbidity and mortality in low resource settings. There is an opportunity now to advance the use of newly available conjugate vaccines. Through TyVAC, key stakeholders are now engaged in the process of designing and implementing trials in 3 settings. These trials should complement other important initiatives, such as the planned Navimumbai government introduction in India.

Through the discussions at this meeting, an approach using either a cluster or individually randomized control strategy starting vaccination at 9 months old with a catch-up campaign to 15 years of age has been proposed using a comparator vaccine in the control group. Surveillance for a primary outcome of blood-culture proven typhoid fever, and a broad range of non-specific secondary outcomes will be performed in 3–4 sites over a period of two years.

The use of typhoid vaccines to control typhoid in those regions with the greatest burden could now be a real possibility. This consortium has been created to realize that aim.

14. Competing interests

This report contains the collective views of an international group of experts. The authors alone are responsible for the views expressed and they do not necessarily represent the decisions, policy or views of the World Health Organization.

AJP chairs the UK Department of Health's (DH) Joint Committee on Vaccination and Immunisation (JCVI) and the European Medicines Agency (EMA) Scientific Advisory Group on Vaccines and is a member of WHO's SAGE. The views expressed in this manuscript do not necessarily reflect those of JCVI, DH, EMA or WHO. AJP has previously conducted clinical trials on behalf of the University of Oxford funded by vaccine manufacturers but has no personal financial interests.

Funding

Funding for this meeting was provided by the Bill & Melinda Gates Foundation.

Funding for the activities of the TyVAC consortium has also been provided by the Bill & Melinda Gates Foundation.

Authors contributions

The manuscript was drafted by JM and MG. AM drafted the Gavi section. All authors within the TyVAC Consortium Meeting Group read, critically revised and approved the final manuscript.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.vaccine.2017.08.001>. These data include Google maps of the most important areas described in this article.

References

- [1] Crump JA, Luby SP, Mintz ED. The global burden of typhoid fever. *Bull World Health Organ* 2004;82:346–53.
- [2] Buckle GC, Walker CLF, Black RE. Typhoid fever and paratyphoid fever: Systematic review to estimate global morbidity and mortality for 2010. *J Glob Health* 2012;2:10401.
- [3] Mogasale V, Maskery B, Ochiai RL, et al. Burden of typhoid fever in low-income and middle-income countries: a systematic, literature-based update with risk-factor adjustment. *Lancet Glob Heal* 2014;2:e570–80.
- [4] Mogasale V, Mogasale V, Ramani E, et al. Revisiting typhoid fever surveillance in low and middle income countries: lessons from systematic literature review of population-based longitudinal studies. *BMC Infect Dis* 2016;16:35.
- [5] Antillon M, Warren JL, Crawford FW, et al. The burden of typhoid fever in low and middle-income countries: a meta-regression approach. *PLoS Negl Trop Dis* 2017;11:e0005376.
- [6] Sedgwick WT, Macnutt JS. On the Mills-Reincke Phenomenon and Hazen's theorem concerning the decrease in mortality from diseases other than typhoid fever following the purification of public water-supplies. *J Infect Dis* 1910;7:489–564.
- [7] Cutler D, Miller G. The role of public health improvements in health advances: the twentieth-century United States. *Demography* 2005;42:1–22.
- [8] UN-Water: WHO/UNICEF Joint Monitoring Programme for Water Supply and Sanitation (JMP). <<http://www.unwater.org/publications/jmp/en/>> [accessed March 31, 2017].
- [9] WHO|8 February 2008, vol. 83, 6. WHO; 2012. p. 49–60.
- [10] Steele AD, Hay Burgess DC, Diaz Z, Carey ME, Zaidi AKM. Challenges and opportunities for typhoid fever control: a call for coordinated action. *Clin Infect Dis* 2016;62(Suppl 1):S4–8.
- [11] WHO|Expert consultation to review evidence in support of the use of typhoid conjugate vaccines. WHO; 2014. <http://www.who.int/immunization/research/meetings_workshops/typhoidvaccines_july14/en/> [accessed March 31, 2017].
- [12] Mitra M, Shah N, Ghosh A, et al. Efficacy and safety of vi-tetanus toxoid conjugated typhoid vaccine (PedaTyph™) in Indian children: school based cluster randomized study. *Hum Vaccin Immun* 2016;12:939–45.
- [13] Mohan VK, Varanasi V, Singh A, et al. Safety and immunogenicity of a Vi polysaccharide-tetanus toxoid conjugate vaccine (typbar-TCV) in healthy infants, children, and adults in typhoid endemic areas: a multicenter, 2-cohort, open-label, double-blind, randomized controlled phase 3 study. *Clin Infect Dis* 2015;61:393–402.
- [14] WHO|SAGE Working Group on Typhoid Vaccines (established March 2016). WHO; 2016.
- [15] Typhoid Vaccine Acceleration Consortium; 2017. <<http://www.medschool.umaryland.edu/CVD/Projects/Typhoid-Vaccine-Acceleration-Consortium-TyVAC/>>. doi: <<http://www.medschool.umaryland.edu/CVD/Projects/Typhoid-Vaccine-Acceleration-Consortium-TyVAC/>>.
- [16] Wong VK, Baker S, Pickard DJ, et al. Phylogeographical analysis of the dominant multidrug-resistant H58 clade of *Salmonella* Typhi identifies inter- and intracontinental transmission events. *Nat Genet* 2015;47:632–9.
- [17] DeRoock D, Clemens JD, Nyamete A, Mahoney RT. Policymakers' views regarding the introduction of new-generation vaccines against typhoid fever, shigellosis and cholera in Asia. *Vaccine* 2005;23:2762–74.
- [18] Mogasale V, Ramani E, Mogasale VV, Park J. What proportion of *Salmonella* Typhi cases are detected by blood culture? A systematic literature review. *Ann Clin Microbiol Antimicrob* 2016;15:32.
- [19] Crump JA, Mintz ED. Global trends in typhoid and paratyphoid Fever. *Clin Infect Dis* 2010;50:241–6.
- [20] Ochiai RL, Acosta CJ, Danovaro-Holliday MC, et al. A study of typhoid fever in five Asian countries: disease burden and implications for controls. *Bull World Health Organ* 2008;86:260–8.
- [21] Brooks WA, Hossain A, Goswami D, et al. Bacteremic Typhoid Fever in Children in an Urban Slum, Bangladesh. *Emerg Infect Dis* 2005;11:326–9.
- [22] Siddiqui FJ, Rabbani F, Hasan R, Nizami SQ, Bhutta ZA. Typhoid fever in children: some epidemiological considerations from Karachi, Pakistan. *Int J Infect Dis* 2006;10:215–22.
- [23] Punjabi NH, Agtini MD, Ochiai RL, et al. Enteric fever burden in North Jakarta, Indonesia: a prospective, community-based study. *J Infect Dev Ctries* 2013;7:781–7.
- [24] Karkey A, Thompson CN, Tran Vu Thieu N, et al. Differential epidemiology of *Salmonella* Typhi and Paratyphi A in Kathmandu, Nepal: a matched case control investigation in a highly endemic enteric fever setting. *PLoS Negl Trop Dis* 2013;7:e2391.
- [25] Reddy EA, Shaw AV, Crump JA. Community-acquired bloodstream infections in Africa: a systematic review and meta-analysis. *Lancet Infect Dis* 2010;10:417–32.
- [26] Breiman RF, Cosmas L, Njuguna H, et al. Population-based incidence of typhoid fever in an urban informal settlement and a rural area in Kenya: implications for typhoid vaccine use in Africa. *PLoS One* 2012;7:e29119.
- [27] von Kalckreuth V, Konings F, Aaby P, et al. The Typhoid Fever Surveillance in Africa Program (TSAP): clinical, diagnostic, and epidemiological methodologies. *Clin Infect Dis* 2016;62(Suppl 1):S9–S16.
- [28] Marks F, von Kalckreuth V, Aaby P, et al. Incidence of invasive salmonella disease in sub-Saharan Africa: a multicentre population-based surveillance study. *Lancet Glob Heal* 2017;5:e310–23.
- [29] Cruz Espinoza LM, Nichols C, Adu-Sarkodie Y, et al. Variations of invasive *Salmonella* infections by population size in Asante Akim North Municipal, Ghana. *Clin Infect Dis* 2016;62:S17–22.
- [30] Kallenberg J. Gavi's vaccine investment strategy. <<http://www.gavi.org/>> [accessed Dec 2, 2016].
- [31] Darton TC, Meiring JE, Tonks S, et al. The STRATAA study protocol: a programme to assess the burden of enteric fever in Bangladesh, Malawi and Nepal using prospective population census, passive surveillance, serological studies and healthcare utilisation surveys. *BMJ Open* 2017;7:e016283.
- [32] Lin FY, Ho VA, Khiem HB, et al. The efficacy of a *Salmonella* typhi Vi conjugate vaccine in two-to-five-year-old children. *N Engl J Med* 2001;344:1263–9.
- [33] Mai Ngoc L, Phan Van B, Vo Anh H, et al. Persistent efficacy of Vi conjugate vaccine against typhoid fever in young children. *N Engl J Med* 2003;349:1390–1.
- [34] Szu SC. Development of Vi conjugate – a new generation of typhoid vaccine. *Expert Rev Vaccines* 2013;12:1273–86.
- [35] Bhutta ZA, Capeding MR, Bavdekar A, et al. Immunogenicity and safety of the Vi-CRM197 conjugate vaccine against typhoid fever in adults, children, and infants in south and southeast Asia: results from two randomised, observer-blind, age de-escalation, phase 2 trials. *Lancet Infect Dis* 2014;14:119–29.
- [36] Thiem VD, Lin F-YYCFYC, Canh DG, et al. The Vi conjugate typhoid vaccine is safe, elicits protective levels of IgG anti-Vi, and is compatible with routine infant vaccines. *Clin Vaccine Immunol* 2011;18:730–5.
- [37] Khan MI, Soofi SB, Ochiai RL, et al. Effectiveness of Vi capsular polysaccharide typhoid vaccine among children: a cluster randomized trial in Karachi. *Pakistan. Vaccine* 2012;30:5389–95.
- [38] Sur D, Ochiai RL, Bhattacharya SK, et al. A cluster-randomized effectiveness trial of Vi typhoid vaccine in India. *N Engl J Med* 2009;361:335–44.
- [39] Mulholland K, Hilton S, Adegbola R, et al. Randomised trial of Haemophilus influenzae type-b tetanus protein conjugate vaccine for prevention of pneumonia and meningitis in Gambian infants. *Lancet* 1997;349:1191–7.
- [40] Cutts FT, Zaman SMA, Enwere G, et al. Efficacy of nine-valent pneumococcal conjugate vaccine against pneumonia and invasive pneumococcal disease in The Gambia: randomised, double-blind, placebo-controlled trial. *Lancet (London, England)* 2002;365:1139–46.
- [41] Ochiai RL, Acosta CJ, Agtini M, et al. The use of typhoid vaccines in Asia: the DOMI experience. *Clin Infect Dis* 2007;45(Suppl 1):S34–8.
- [42] Hemming K, Haines TP, Chilton PJ, Girling AJ, Lilford RJ. The stepped wedge cluster randomised trial: rationale, design, analysis, and reporting. *BMJ* 2015;350:h391.
- [43] Beard E, Lewis JJ, Copas A, et al. Stepped wedge randomised controlled trials: systematic review of studies published between 2010 and 2014. *Trials* 2015;16:353.
- [44] The Gambia Hepatitis Intervention Study. The gambia hepatitis study group. *Can Res* 1987;47:5782–7.
- [45] Mdege ND, Man M-S, Taylor (nee Brown) CA, Torgerson DJ. Systematic review of stepped wedge cluster randomized trials shows that design is particularly used to evaluate interventions during routine implementation. *J Clin Epidemiol* 2011;64:936–48.
- [46] Kucharski AJ, Eggo RM, Watson CH, Camacho A, Funk S, Edmunds WJ. Effectiveness of ring vaccination as control strategy for ebola virus disease. *Emerg Infect Dis* 2016;22:105–8.
- [47] Henao-Restrepo AM, Longini IM, Egger M, et al. Efficacy and effectiveness of an rVSV-vectored vaccine expressing Ebola surface glycoprotein: interim results from the Guinea ring vaccination cluster-randomised trial. *Lancet* 2015;386:857–66.
- [48] Thompson CN, Kama M, Acharya S, et al. Typhoid fever in Fiji: a reversible plague? *Trop Med Int Health* 2014;19:1284–92.
- [49] Pitzer VE, Bowles CC, Baker S, et al. Predicting the impact of vaccination on the transmission dynamics of typhoid in South Asia: a mathematical modeling study. *PLoS Negl Trop Dis* 2014;8:e2642.

- [50] Baker S, Holt KE, Clements ACA, et al. Combined high-resolution genotyping and geospatial analysis reveals modes of endemic urban typhoid fever transmission. *Open Biol* 2011;1.
- [51] Prasad N, Murdoch DR, Reyburn H, Crump JA. Etiology of severe febrile illness in low- and middle-income countries: a systematic review. *PLoS One* 2015;10: e0127962.
- [52] Mogasale V, Desai SN, Mogasale VV, Park JK, Leon Ochiai R, Wierzba TF. Case fatality rate and length of hospital stay among patients with typhoid intestinal perforation in developing countries: a systematic literature review. *PLoS One* 2014;9. <http://dx.doi.org/10.1371/journal.pone.0093784>.
- [53] Feikin DR, Scott JAG, Gessner BD. Use of vaccines as probes to define disease burden. *Lancet* 2014;383:1762–70.