



This is a repository copy of *Vitamin D in obesity*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/123619/>

Version: Accepted Version

Article:

Walsh, J.S. orcid.org/0000-0002-7122-2650, Bowles, S. and Evans, A.L. (2017) Vitamin D in obesity. *Current Opinion in Endocrinology, Diabetes and Obesity*, 24 (6). pp. 389-394. ISSN 1752-296X

<https://doi.org/10.1097/MED.0000000000000371>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Vitamin D in Obesity

Jennifer S Walsh, Simon Bowles, Amy L Evans

Mellanby Centre for Bone Research, University of Sheffield, UK

Corresponding author:

Dr Jennifer S Walsh

Senior Clinical Lecturer, University of Sheffield, UK

Metabolic Bone Centre

Northern General Hospital

Herries Road

Sheffield

UK

S5 7AU

+44 144 2714705

j.walsh@sheffield.ac.uk

Abstract

Purpose of review: Vitamin D is essential for bone health, and may also have important functions in immunity and other systems. Vitamin D deficiency is common, and testing and supplementation is increasing. Serum vitamin D is lower in obese people; it is important to understand the mechanism of this effect and whether it indicates clinically significant deficiency.

Recent findings: Vitamin D is fat soluble, and distributed into fat, muscle, liver and serum. All of these compartments are increased in volume in obesity, so the lower vitamin D likely reflects a volumetric dilution effect and whole body stores of vitamin D may be adequate. Despite lower serum vitamin D, obese adults do not have higher bone turnover or lower bone mineral density (BMD). Patients undergoing bariatric surgery do have bone loss, and ensuring vitamin D sufficiency in these patients may help to attenuate bone loss.

Summary: Lower vitamin D in obese people is a consistent finding across age, ethnicity and geography. This may not always reflect a clinical problem. Obese people need higher loading doses of vitamin D to achieve the same serum 25OHD as normal weight.

Keywords vitamin D, obesity, bone,

Introduction

The main source of vitamin D in humans is production from the action of UVB light on cholesterol in skin. There is a small contribution from dietary sources such as oily fish and fortified foods. There are some geographical variations, for example in Scandinavia, dietary fish is a major source, and in the USA there is more food fortification. Vitamin D deficiency is very prevalent in the northern hemisphere; most of the UK population are vitamin D deficient in winter ¹.

There is ongoing debate about the fundamental principles of defining vitamin D deficiency. The most commonly used measure of vitamin D status is total 25-hydroxyvitamin D (25OHD) which captures 25OHD₃ and 25OHD₂. 25OHD has a long serum half-life and is a technically easy measurement, but it has been proposed that free (non-protein-bound) 25OHD is a better measurement because it more closely reflects biologically available 25OHD status. There are variations in the sufficiency threshold recommended by different advisory bodies, with recommendations generally between 50 and 75 nmol/l ^{2,3}.

Vitamin D is a pre-hormone. It is converted to the active hormone 1,25 dihydroxyvitamin D (1,25(OH)₂D) by 25-hydroxylation in the liver and 1-hydroxylation in the kidney. The final step of 1-hydroxylation is tightly regulated to prevent hypercalcaemia.

The main circulating form of vitamin D is 25OHD. It is protein-bound in circulation (about 85% to vitamin D binding protein (VDBP), and about 15% to albumin). 25OHD is fat soluble, and distributed into fat, muscle and liver, with smaller amounts into other tissues. 25OHD is cleared by 24-hydroxylation to 24,25(OH)₂D or other inactive metabolites.

Active vitamin D, 1,25(OH)₂D is required for gut absorption of dietary calcium. Severe vitamin D deficiency causes osteomalacia in adults and in children. Less severe insufficiency causes increased bone turnover and is associated with increased fracture risk. Vitamin D deficiency has been associated with a wide range of other disorders, particularly auto-immune disorders and cancers, but a causative role in these conditions is not yet well established.

Obesity is inversely correlated with serum 25OHD. This is a consistent finding in adults and children of different ethnicities in a range of geographic locations. This observation raises several questions: Is the low serum 25OHD a consequence or cause of obesity? What is the cause of low serum 25OHD in obesity? Does the low serum 25OHD in obesity have clinical consequences for bone or other systems? Do obese people need higher doses of vitamin D supplements? Does vitamin D supplementation in obesity ameliorate any of the metabolic consequences of obesity? What happens to vitamin D status when weight changes rapidly after weight loss surgery?

Relationship between body weight and vitamin D

Obese people have lower serum 25OHD than normal weight people, and serum 25OHD is inversely correlated with body weight, BMI and fat mass. This has been shown in adults and children in northern and southern Europe, Australia and New Zealand, Saudi Arabia, Latin America and in White, Black and Hispanic groups in the USA ⁴⁻¹². Serum 25OHD is about 20% lower in obese people than normal weight ^{4,5,13}, and the prevalence of 25OHD deficiency is greater in obese people, reported at between 40-80% ^{4,9,12}.

Other measures of vitamin D status (free 25OHD and 1,25(OH)₂D) are also lower in obesity ^{4,14}.

Parathyroid hormone is often used as an indicator of vitamin D status. PTH tends to be higher in obesity¹⁵, but the relationship between serum calcium and PTH is left-shifted in obesity¹⁶, so it is difficult to interpret the clinical significance of higher PTH.

It is likely that low serum 25OHD is a consequence of obesity, rather than the cause of obesity. A large genetic study found that high BMI and genes that predispose to obesity decrease serum 25OHD, while low 25OHD and genes associated with low 25OHD have very little effect on obesity¹⁷. In meta-analysis, vitamin D supplementation has no effect on body weight or fat mass¹⁸.

Causes of low 25OHD in obesity

There are a number of possible mechanisms that could cause low 25OHD in obesity. There could be lower input due to lower dietary intake, lower sunlight exposure or impaired skin synthesis of vitamin D. Alterations in protein binding or faster metabolic clearance in obesity could lead to lower serum 25OHD. Or the lower serum 25OHD could be due to distribution of 25OHD into a larger whole body tissue volume, particularly if 25OHD was actively sequestered in other tissues.

Dietary vitamin D intake did not differ between obese and normal weight adults in a UK population⁴. Obese people may have less sunlight exposure than normal weight people¹⁹ in some geographic regions, although in two UK studies, sunlight exposure did not vary with BMI^{4,20}. When exposed to UVB, normal-weight and obese people have a similar cutaneous synthesis of vitamin D²¹. Clearly, diet and sunlight behaviours will vary between different geographic and cultural groups, and may be a contributory factor to lower vitamin D in some groups.

Serum VDBP and albumin concentrations affect the total 25OHD measurement, and there are genetic variations in VDBP which cause variation in the binding affinity of DBP. However, VDBP and albumin do not differ between obese and normal weight people, and the distribution of VDBP genotypes is similar in obese and normal weight groups^{4,22}.

If metabolic clearance rate of 25OHD were faster in obesity, the half-life of 25OHD would be shorter and circulating 25OHD would be lower. Measurements using a stable isotope 25OHD tracer show that metabolic clearance is similar in obese and normal weight adults⁴, so this is unlikely to be the explanation for lower serum 25OHD.

The most important mechanism is probably volumetric dilution of 25OHD into greater tissue volume in obese people. 25OHD is distributed into serum, fat, muscle, liver and a small amount into other tissues²³, and all of these compartments are increased in obesity. The difference in serum 25OHD between obese and normal weight groups is greater in summer than in winter because the summer rise in serum 25OHD is less in obese people^{4,24,25}. Obese people get similar sunlight exposure to normal weight people, and produce the same amount of vitamin D in response to sunlight exposure, but the synthesised vitamin is distributed into a larger volume, so the amount distributed into serum is less. There is a similar difference between obese and normal weight people in response to oral vitamin D dosing, with a smaller serum 25OHD rise in obesity²⁶. The dose-response relationship of vitamin D supplementation in obese adults is BMI-dependent and curvilinear, consistent with volumetric dilution²⁷ (**Figure 1**). The clinical implication of this volumetric dilution effect is that obese people will need larger loading doses of vitamin D than normal weight to achieve the same serum level of vitamin D, but on cessation of supplementation may have a slower decrease in serum vitamin D due to redistribution from other tissue stores. Conversely, people with low body weight would have a large rise in serum vitamin D after dosing (due to the smaller volume of distribution),

but would become deficient again quickly if supplementation was stopped (due to less tissue storage).

It has been proposed that vitamin D is actively retained (sequestered) in fat but the modelling of the dose-response doesn't seem consistent with that hypothesis. There is experimental evidence that vitamin D concentration in abdominal subcutaneous fat increases with vitamin D supplementation^{28,29}, and one in vitro study suggests that adipocytes from insulin-resistant obese people might have impaired release of vitamin D³⁰. Adipocytes express 1-hydroxylase, but also 24-hydroxylase, so can activate and inactivate 25OHD. The activity of the enzymes may be altered in obesity, which could affect the amount of 25OHD available for release into serum³¹. However the distribution and concentration of vitamin D and 25OHD in subcutaneous and omental fat does not differ between obese and normal weight people, and the correlation between 25OHD in subcutaneous fat and serum is similar in obese and normal weight³².

Does lower 25OHD have clinical consequences in obesity?

Usually, low total 25OHD, free 25OHD and 1,25(OH)₂D would lead to lower dietary calcium absorption, and increased bone turnover with lower bone mineral density (BMD). However, obese adults have lower bone turnover than normal weight, and higher BMD with thicker, denser cortices and greater trabecular number³³. It is important to note that in contrast, obesity in children has adverse effects on bone strength³⁴.

This lack of adverse effects on bone may indicate that obese people are not truly vitamin D deficient; although serum 25OHD is lower due to volumetric dilution, their whole body vitamin D stores are greater due to the reservoir in their fat tissue, which maintains an equilibrium with serum 25OHD and a sufficient supply.

An alternative explanation is that obese people are vitamin D deficient but that other effects of obesity compensate for the effects of vitamin D deficiency. For example, greater skeletal loading or the action of hormones such as leptin, adiponectin or oestrogen are known to have positive effects on bone mass.

If obese people are truly vitamin D deficient, there may be implications for systems other than bone. Vitamin D deficiency has been associated with a large number of disorders, such as auto-immunity, cancer, neurodegenerative disease and metabolic syndrome. However, there is not yet clear evidence for a causative role of vitamin D deficiency in many of these conditions³⁵. Obesity does increase the risk for several of these disorders but there are other possible mechanisms than low vitamin D for these associations, and the interaction of vitamin D and obesity in causation has not yet been clearly characterised³⁶.

In the US NHANES population study, low serum 25OHD was associated with higher all-cause mortality in postmenopausal women with normal waist circumference; the hazard ratio for the lowest versus the highest serum vitamin D quartile (< 36.5 nmol/l vs > 65.4 nmol/l) was 1.85 (95% CI 1.00 to 3.44). In women with abdominal obesity there was no association between serum 25OHD quartile and all-cause mortality (HR 0.96, 95% CI 0.52 to 1.76)³⁷. This result suggests that obesity does modulate the relationship between serum vitamin D and health outcomes.

Does vitamin D supplementation improve the adverse metabolic profile in obesity?

A recent meta-analysis found only four or five good quality intervention studies for each of the metabolic parameters they considered (fat mass, blood pressure, lipids and glucose tolerance). There were some positive effects of vitamin D supplementation on fat mass, triglycerides, HDL cholesterol and oral glucose tolerance, but adverse effects on LDL cholesterol and blood pressure. However, due to the small numbers of studies and significant heterogeneity of methodology, the current evidence is insufficient to draw conclusions ³⁸.

Inflammatory markers do not seem to be higher in vitamin D deficient obese people than in vitamin D replete obese people ³⁹ and vitamin D supplementation has little effect on circulating inflammatory markers in overweight or obese people ⁴⁰⁻⁴².

Vitamin D with weight loss surgery

Obesity surgery leads to substantial weight loss in most patients, often about 30% of their body weight. The two main types of surgery are gastric volume reduction (e.g. sleeve gastrectomy) or malabsorptive (most commonly Roux-en-Y gastric bypass (RYGB), and less commonly biliopancreatic diversion with gastric switch). As expected, obese people have low serum 25OHD pre-operatively. Most studies find that serum 25OHD increases in the first month after surgery ⁴³, which may reflect the smaller volume of dilution with weight loss. One study tried to determine whether the vitamin D stored in fat was mobilised into serum during weight loss, but found no association between vitamin D concentration in fat at time of surgery and subsequent change in serum 25OHD ⁴⁴. With biliopancreatic diversion, serum 25OHD tends to decline immediately after surgery, which is probably due to the greater bile and fat malabsorption with this procedure ⁴⁵.

In the longer term, vitamin D decreases. Five years after RYGB, with calcium and vitamin D supplementation in most patients, they are just vitamin D sufficient at about 50 nmol/l, but have parathyroid hormone above the reference range ⁴⁶. Bone density and microarchitecture deteriorates for at least two years after RYGB, beyond stabilisation of body weight, despite vitamin D supplementation which maintained serum 25OHD well into the replete range ⁴⁷. The bone loss is likely to be multifactorial, with decreased loading, changes in adipokines, gut hormones, insulin and sex hormones as well as other nutritional factors. Maintaining vitamin D sufficiency after bariatric surgery may require higher doses than standard supplementation, but in combination with calcium and protein supplements and physical activity may attenuate bone loss ⁴⁸.

Optimum vitamin D dosing for BMI

As explained above, the observed serum 25OHD of about 20% below the normal weight population in obese people may not indicate true deficiency or a clinical problem. However, despite higher total body stores, obese people can become vitamin D deficient for the same reasons as any other group. If obese people are treated to the same serum 25OHD repletion threshold as normal weight, higher loading doses will be required due the greater volume of distribution. Obese people may need repeated loading courses to achieve repletion. (Conversely, people with below-normal BMI may need lower doses, but more frequently). Drincic ²⁷ describes the formula to calculate the required vitamin D dose based on BMI and the required increment in serum 25OHD. Because there are no alterations in protein binding or metabolic clearance rate in obesity, maintenance dose requirements should be similar to normal weight (although as above, higher doses may be need after bariatric surgery).

Summary

Obesity is associated with lower serum vitamin D, assessed as total 25OHD, free 25OHD or 1,25(OH)₂D. This is likely due to volumetric dilution into the greater volumes of serum, fat, muscle and liver. The observed lower serum vitamin D in obesity doesn't seem to have adverse consequences for bone health. There is less available information about consequences for other systems, ~~and but no clear evidence of adverse effects.~~ Vitamin D supplementation has not yet been clearly shown to benefit the adverse metabolic profile in obesity. [More information on other system effects is needed before it can be determined what the serum 25OHD target for repletion should be in obese people.](#) Vitamin D does decrease after bariatric surgery, and careful attention to vitamin D repletion might help to prevent bone loss in these patients. ~~Obese people will need greater loading doses of vitamin D to achieve the same serum 25OHD threshold as normal weight. Obese people will need higher loading doses than normal weight to achieve the same serum level of vitamin D, but because the metabolic clearance rate is not affected by obesity, the maintenance dose should be the similar.~~

Acknowledgements: Thanks to Richard Eastell, who is principal investigator and senior author on the Sheffield vitamin D and obesity study (AJCN 2017, ref no. 4). That study was funded by the Department of Health Policy Research Programme. The views expressed in this article are those of the authors and not necessarily those of the Department of Health. SB is funded by the ARC/MRC Centre for Integrated study of Musculoskeletal Ageing (CIMA).

Conflicts of interest: JW: Speaker's honoraria from Eli Lilly, grant funding from Alexion and Immunodiagnostic Systems, donations of drug from Eli Lilly, Prostrakan (Kyowa Kirin) and Consilient for clinical studies, donation of assay kits from Biomedica, consulting fees from Shire and Mereo Biopharma. SB: donation of drug from Consilient for clinical study. ALE: none.

References

1. Hypponen E, Power C. Hypovitaminosis D in British adults at age 45 y: nationwide cohort study of dietary and lifestyle predictors. *AmJClinNutr* 2007;85:860-8.
2. Institute of Medicine. Dietary reference intakes for calcium and vitamin D. The National Academies Press; Washington DC. 2011.
3. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *The Journal of clinical endocrinology and metabolism* 2011;96:1911-30.
4. Walsh JS, Evans AL, Bowles S, et al. Free 25-hydroxyvitamin D is low in obesity, but there are no adverse associations with bone health. *The American journal of clinical nutrition* 2016;103:1465-71.
5. Macdonald HM, Mavroeidi A, Barr RJ, Black AJ, Fraser WD, Reid DM. Vitamin D status in postmenopausal women living at higher latitudes in the UK in relation to bone health, overweight, sunlight exposure and dietary vitamin D. *Bone* 2008;42:996-1003.
6. Ardawi MS, Sibiany AM, Bakhsh TM, Qari MH, Maimani AA. High prevalence of vitamin D deficiency among healthy Saudi Arabian men: relationship to bone mineral density, parathyroid hormone, bone turnover markers, and lifestyle factors. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 2012;23:675-86.
7. Shea MK, Houston DK, Tooze JA, et al. Correlates and prevalence of insufficient 25-hydroxyvitamin D status in black and white older adults: the health, aging and body composition study. *JAmGeriatrSoc* 2011;59:1165-74.
8. Rajakumar K, de las HJ, Chen TC, Lee S, Holick MF, Arslanian SA. Vitamin D status, adiposity, and lipids in black American and Caucasian children. *JClinEndocrinolMetab* 2011;96:1560-7.
9. Lagunova Z, Porojnicu AC, Lindberg F, Hexeberg S, Moan J. The dependency of vitamin D status on body mass index, gender, age and season. *Anticancer Res* 2009;29:3713-20.
10. McKinney K, Breikopf CR, Berenson AB. Association of race, body fat and season with vitamin D status among young women: a cross-sectional study. *ClinEndocrinol(Oxf)* 2008;69:535-41.
11. Palacios C, Gil K, Perez CM, Joshipura K. Determinants of vitamin D status among overweight and obese Puerto Rican adults. *Annals of nutrition & metabolism* 2012;60:35-43.
12. Samuel L, Borrell LN. The effect of body mass index on optimal vitamin D status in U.S. adults: the National Health and Nutrition Examination Survey 2001-2006. *Annals of epidemiology* 2013;23:409-14.
13. Ardawi MS, Qari MH, Rouzi AA, Maimani AA, Raddadi RM. Vitamin D status in relation to obesity, bone mineral density, bone turnover markers and vitamin D receptor genotypes in healthy Saudi pre- and postmenopausal women. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 2011;22:463-75.
14. Lagunova Z, Porojnicu AC, Vieth R, Lindberg FA, Hexeberg S, Moan J. Serum 25-hydroxyvitamin D is a predictor of serum 1,25-dihydroxyvitamin D in overweight and obese patients. *JNutr* 2011;141:112-7.
15. Snijder MB, van Dam RM, Visser M, et al. Adiposity in relation to vitamin D status and parathyroid hormone levels: a population-based study in older men and women. *JClinEndocrinolMetab* 2005;90:4119-23.
16. Hultin H, Edfeldt K, Sundbom M, Hellman P. Left-shifted relation between calcium and parathyroid hormone in obesity. *JClinEndocrinolMetab* 2010;95:3973-81.
17. Vimalleswaran KS, Berry DJ, Lu C, et al. Causal relationship between obesity and vitamin D status: bi-directional Mendelian randomization analysis of multiple cohorts. *PLoS medicine* 2013;10:e1001383.

18. Chandler PD, Wang L, Zhang X, et al. Effect of vitamin D supplementation alone or with calcium on adiposity measures: a systematic review and meta-analysis of randomized controlled trials. *Nutrition reviews* 2015;73:577-93.
19. Kull M, Kallikorm R, Lember M. Body mass index determines sunbathing habits: implications on vitamin D levels. *InternMedJ* 2009;39:256-8.
20. Macdonald HM, Mavroei A, Aucott LA, et al. Skin color change in Caucasian postmenopausal women predicts summer-winter change in 25-hydroxyvitamin D: findings from the ANSAViD cohort study. *JClinEndocrinolMetab* 2011;96:1677-86.
21. Wortsman J, Matsuoka LY, Chen TC, Lu Z, Holick MF. Decreased bioavailability of vitamin D in obesity. *AmJClinNutr* 2000;72:690-3.
22. Winters SJ, Chennubhatla R, Wang C, Miller JJ. Influence of obesity on vitamin D-binding protein and 25-hydroxy vitamin D levels in African American and white women. *Metabolism* 2009;58:438-42.
23. Heaney RP, Horst RL, Cullen DM, Armas LA. Vitamin D3 distribution and status in the body. *Journal of the American College of Nutrition* 2009;28:252-6.
24. Bolland MJ, Grey AB, Ames RW, et al. The effects of seasonal variation of 25-hydroxyvitamin D and fat mass on a diagnosis of vitamin D sufficiency. *AmJClinNutr* 2007;86:959-64.
25. Blum M, Dolnikowski G, Seyoum E, et al. Vitamin D(3) in fat tissue. *Endocrine* 2008;33:90-4.
26. Gallagher JC, Yalamanchili V, Smith LM. The effect of vitamin D supplementation on serum 25(OH)D in thin and obese women. *The Journal of steroid biochemistry and molecular biology* 2013;136:195-200.
27. Drincic A, Fuller E, Heaney RP, Armas LA. 25-hydroxyvitamin d response to graded vitamin d3 supplementation among obese adults. *The Journal of clinical endocrinology and metabolism* 2013;98:4845-51.
28. Didriksen A, Burild A, Jakobsen J, Fuskevag OM, Jorde R. Vitamin D3 increases in abdominal subcutaneous fat tissue after supplementation with vitamin D3. *European journal of endocrinology / European Federation of Endocrine Societies* 2015;172:235-41.
29. Heaney RP, Recker RR, Grote J, Horst RL, Armas LA. Vitamin D(3) is more potent than vitamin D(2) in humans. *JClinEndocrinolMetab* 2011;96:E447-E52.
30. Di Nisio A, De Toni L, Sabovic I, et al. Impaired release of vitamin D in dysfunctional adipose tissue: new cues on vitamin D supplementation in obesity. *The Journal of clinical endocrinology and metabolism* 2017; doi: 10.1210/jc.2016-3591. [Epub ahead of print]
31. Wamberg L, Christiansen T, Paulsen SK, et al. Expression of vitamin D-metabolizing enzymes in human adipose tissue-the effect of obesity and diet-induced weight loss. *International journal of obesity* 2013;37:651-7.
32. Carrelli A, Bucovsky M, Horst R, et al. Vitamin D Storage in Adipose Tissue of Obese and Normal Weight Women. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research* 2017;32:237-42.
33. Evans AL, Paggiosi MA, Eastell R, Walsh JS. Bone density, microstructure and strength in obese and normal weight men and women in younger and older adulthood. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research* 2015;30:920-8.
34. Dimitri P, Bishop N, Walsh JS, Eastell R. Obesity is a risk factor for fracture in children but is protective against fracture in adults: A paradox. *Bone* 2012;50:457-66.
35. Autier PB, M.; Pizot, C.; Mullie, P. Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol* 2014;2:76-89.
36. Shanmugalingam T, Crawley D, Bosco C, et al. Obesity and cancer: the role of vitamin D. *BMC cancer* 2014;14:712.
37. Eaton CB, Young A, Allison MA, et al. Prospective association of vitamin D concentrations with mortality in postmenopausal women: results from the Women's Health Initiative (WHI). *The American journal of clinical nutrition* 2011;94:1471-8.

38. Manousopoulou A, Al-Daghri NM, Garbis SD, Chrousos GP. Vitamin D and cardiovascular risk among adults with obesity: a systematic review and meta-analysis. *European Journal of Clinical Investigation* 2015;45:1113-26.
39. Vilarrasa N, Vendrell J, Maravall J, et al. Is plasma 25(OH) D related to adipokines, inflammatory cytokines and insulin resistance in both a healthy and morbidly obese population? *Endocrine* 2010;38:235-42.
40. Beilfuss J, Berg V, Sneve M, Jorde R, Kamycheva E. Effects of a 1-year supplementation with cholecalciferol on interleukin-6, tumor necrosis factor-alpha and insulin resistance in overweight and obese subjects. *Cytokine* 2012;60:870-4.
41. Wamberg L, Kampmann U, Stodkilde-Jorgensen H, Rejnmark L, Pedersen SB, Richelsen B. Effects of vitamin D supplementation on body fat accumulation, inflammation, and metabolic risk factors in obese adults with low vitamin D levels - results from a randomized trial. *European journal of internal medicine* 2013;24:644-9.
42. Jorde R, Sneve M, Torjesen PA, Figenschau Y, Goransson LG, Omdal R. No effect of supplementation with cholecalciferol on cytokines and markers of inflammation in overweight and obese subjects. *Cytokine* 2010;50:175-80.
43. Himbert C, Ose J, Delphan M, Ulrich CM. A systematic review of the interrelation between diet- and surgery-induced weight loss and vitamin D status. *Nutr Res* 2017;38:13-26.
44. Pramyothin P, Biancuzzo RM, Lu Z, Hess DT, Apovian CM, Holick MF. Vitamin D in Adipose Tissue and Serum 25-Hydroxyvitamin D After Roux-en-Y Gastric Bypass. *Obesity(SilverSpring)* 2011;19:2228-34.
45. Aasheim ET, Bjorkman S, Sovik TT, et al. Vitamin status after bariatric surgery: a randomized study of gastric bypass and duodenal switch. *The American journal of clinical nutrition* 2009;90:15-22.
46. Switzer NJ, Marcil G, Prasad S, et al. Long-term hypovitaminosis D and secondary hyperparathyroidism outcomes of the Roux-en-Y gastric bypass: a systematic review. *Obesity reviews : an official journal of the International Association for the Study of Obesity* 2017;18:560-6.
47. Shanbhogue VV, Stoving RK, Frederiksen KH, et al. Bone structural changes after gastric bypass surgery evaluated by HR-pQCT: a two-year longitudinal study. *European journal of endocrinology / European Federation of Endocrine Societies* 2017;176:685-93.
48. Muschitz C, Kocijan R, Haschka J, et al. The Impact of Vitamin D, Calcium, Protein Supplementation, and Physical Exercise on Bone Metabolism After Bariatric Surgery: The BABS Study. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research* 2016;31:672-82.

Key points

Serum vitamin D is low in obese people across age, ethnicity and geography.

The low vitamin D is due to greater volumetric dilution, and may not necessarily indicate vitamin D deficiency.

Despite lower serum vitamin D, obese people have lower bone turnover and higher bone density than normal weight people.

Bariatric surgery does cause bone loss, and vitamin D supplementation may be important in this group of patients.

Obese people need higher loading doses of vitamin D than normal weight people to achieve a similar increment in serum vitamin D.

Figure legend

Figure 1 Serum response to oral vitamin D supplementation is BMI- and dose-dependent.

The time course of the 25(OH)D increment in the obese 1,000-IU group (○), the obese 5,000-IU group (△), and the obese 10,000-IU group (□) in the current study, contrasted with a historical comparison from normal-weight males who were taking 5,000 IU (▲) ~~daily~~ and 10,000 IU (■) daily. The error bars are 1 SEM.

Reproduced with permission from Drincic et al, J Clin Endocrinol Metab 2013;98(12):4845-4851