



**UNIVERSITY OF LEEDS**

This is a repository copy of *Large emissions from floodplain trees close the Amazon methane budget*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/123683/>

Version: Accepted Version

---

**Article:**

Pangala, SR, Enrich-Prast, A, Basso, LS et al. (12 more authors) (2017) Large emissions from floodplain trees close the Amazon methane budget. *Nature*, 552. pp. 230-234. ISSN 0028-0836

<https://doi.org/10.1038/nature24639>

---

© 2017 Macmillan Publishers Limited, part of Springer Nature. This is an author produced version of a paper published in *Nature*. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## 1        **Large emissions from floodplain trees close the Amazon methane budget**

2        **Sunitha R. Pangala<sup>1\*</sup>, Alex Enrich-Prast<sup>2,3</sup>, Luana S. Basso<sup>4</sup>, Roberta Bittencourt Peixoto<sup>3</sup>,**  
3        **David Bastviken<sup>2</sup>, Edward Hornibrook<sup>5,6</sup>, Luciana V. Gatti<sup>4,7</sup>, Humberto Marotta Ribeiro<sup>8,9</sup>,**  
4        **Luana Silva Braucks Calazans<sup>3</sup>, Cassia Mônica Sakuragui<sup>3</sup>, Wanderley Rodrigues Bastos<sup>10</sup>,**  
5        **Olaf Malm<sup>11</sup>, Emanuel Gloor<sup>12</sup>, John Miller<sup>13</sup>, Vincent Gauci<sup>1\*</sup>**

6        <sup>1</sup> School of Environment, Earth and Ecosystem Sciences, The Open University, Walton Hall,  
7        Milton Keynes, MK7 6AA, UK.

8        <sup>2</sup> Department of Thematic Studies – Environmental Change, Linköping University, Linköping  
9        SE-581 83, Sweden.

10       <sup>3</sup> Department of Botany, Institute of Biology, University Federal of Rio de Janeiro, Rio de  
11       Janeiro, Brazil.

12       <sup>4</sup> Instituto de Pesquisas Energéticas e Nucleares (IPEN)–Comissão Nacional de Energia  
13       Nuclear (CNEN)–Atmospheric Chemistry Laboratory, 2242 Avenida Professor Lineu Prestes,  
14       Cidade Universitária, São Paulo CEP 05508-000, Brazil.

15       <sup>5</sup> School of Earth Sciences, The University of Bristol, Wills Memorial Building, Queen’s Road,  
16       Bristol, BS8 1RJ, UK

17       <sup>6</sup> Earth, Environmental and Geographic Sciences, Irving K. Barber School of Arts and  
18       Sciences, The University of British Columbia, 1177 Research Road, Kelowna, BC, V1V 1V7,  
19       Canada.

20       <sup>7</sup> National Institute for Space Research (INPE), Center for Earth System Science (CCST),  
21       Greenhouse Gas Laboratory (LaGEE), Av. Dos Astronautas, 1758, São José dos Campos, CEP  
22       12227-010, Brazil.

23       <sup>8</sup> Ecosystems and Global Change Laboratory (LEMG-UFF) / International Laboratory of Global  
24       Change (LINCGlobal). Biomass and Water Management Research Center (NAB-UFF).  
25       Graduated Program in Geosciences (Environmental Geochemistry). Universidade Federal  
26       Fluminense (UFF), Av. Edmundo March, s/nº – Zip Code: 24210-310, Niterói/RJ- Brazil.

27       <sup>9</sup> Sedimentary and Environmental Processes Laboratory (LAPSA-UFF). Department of  
28       Geography. Graduated Program in Geography. Universidade Federal Fluminense (UFF), Av.  
29       Gal. Milton Tavares de Souza, s/nº - Zip Code: 24210-346, Niterói/RJ- Brazil.

30       <sup>10</sup> Environmental Biogeochemistry Laboratory, Federal University of Rondônia, Rondônia,  
31       Brazil.

32 <sup>11</sup> Radioisotopes Laboratory, Biophysics Institute, Federal University of Rio de Janeiro (UFRJ),  
33 Rio de Janeiro, Brazil.

34 <sup>12</sup> School of Geography, University of Leeds, Woodhouse Lane, Leeds LS9 2JT, UK.

35 <sup>13</sup> Global Monitoring Division, Earth System Research Laboratory, National Oceanic and  
36 Atmospheric Administration, 325 Broadway, Boulder, CO 80305, USA.

37 \*Authors for correspondence: Sunitha.Pangala@open.ac.uk, Vincent.Gauci@open.ac.uk.

38 **Keywords:** tropical wetlands, methane, tree stem methane emissions, Amazon wetlands.

39 **Wetlands are the largest global source of atmospheric methane (CH<sub>4</sub>)<sup>1</sup>, a potent greenhouse gas.**  
40 **However, methane emission inventories from the Amazon floodplain<sup>2,3</sup>, the largest natural**  
41 **geographic source of CH<sub>4</sub> in the tropics, consistently underestimate the atmospheric burden of CH<sub>4</sub>**  
42 **determined via remote sensing and inversion modelling<sup>4,5</sup>, pointing to a major gap in our**  
43 **understanding of the contribution of these ecosystems to CH<sub>4</sub> emissions. Here we report CH<sub>4</sub>**  
44 **fluxes from the stems of 2357 individual Amazonian floodplain trees from 13 locations across the**  
45 **central Amazon basin. We find that egress of soil gas through wetland trees is the dominant**  
46 **source of regional CH<sub>4</sub> emissions. Amazon tree stem fluxes were up to 150-200 times larger than**  
47 **emissions reported for temperate wet forests<sup>6</sup> and tropical peat swamp forests<sup>7</sup>, representing the**  
48 **largest non-ebullitive wetland fluxes observed. Tree emissions had an average δ<sup>13</sup>C-CH<sub>4</sub> value of -**  
49 **66.2±6.4‰ consistent with a soil biogenic origin. We estimate that floodplain trees emit 15.1 ± 1.8**  
50 **to 21.2 ± 2.5 Tg CH<sub>4</sub> yr<sup>-1</sup>, in addition to 20.5±5.3 Tg CH<sub>4</sub> yr<sup>-1</sup> emitted regionally from other sources.**  
51 **Furthermore, we provide a top-down regional estimate of CH<sub>4</sub> emissions of 42.7±5.6 Tg CH<sub>4</sub> yr<sup>-1</sup> for**  
52 **the Amazon basin based on regular vertical lower troposphere CH<sub>4</sub> profiles covering the period**  
53 **2010-13. We find close agreement between our 'top-down' and combined 'bottom-up' estimates,**  
54 **indicating that large CH<sub>4</sub> emissions from trees adapted to permanent or seasonal inundation can**  
55 **account for the missing emission source required to close the Amazon CH<sub>4</sub> budget.**

56 Wetlands are the single largest global source of atmospheric methane (CH<sub>4</sub>), emitting an estimated  
57 160 to 210 Tg of CH<sub>4</sub> each year to the troposphere<sup>1</sup>. Wetlands are concentrated globally in two  
58 broad latitudinal bands; one rich in peatlands spanning the boreal and subarctic zones and a second  
59 in the tropics and sub-tropics containing vast swamps and seasonally inundated floodplains<sup>1</sup>. Low  
60 latitude wetlands are notably prolific sources of CH<sub>4</sub> because of their substantial net primary  
61 productivity (NPP) and high seasonal temperatures<sup>2</sup>. However, relative to northern wetlands, flux  
62 measurements from Amazon floodplain ecosystems are comparatively sparse and have focussed  
63 mainly on soil and water surfaces, and gas exchange mediated by aquatic macrophytes<sup>8,9</sup>.

64 Integration of these emission sources across the lowland Amazon basin based upon remotely sensed  
65 wetland distributions, yields an estimated flux of 26 to 29 Tg CH<sub>4</sub> yr<sup>-1 2,3</sup>. In contrast, estimates  
66 derived from atmospheric transport inversion modelling using *in-situ* CH<sub>4</sub> concentrations measured  
67 at surface sites remote from Amazonia and satellite greenhouse gas measurements (the so-called  
68 'top-down' approaches) are considerably greater at 44 to 52 Tg yr<sup>-1 4,10</sup> and consistent with estimates  
69 of CH<sub>4</sub> flux determined from modelling heterotrophic anaerobic respiration of regional NPP<sup>10</sup>.

70 Results of these global inversions should be treated with some caution. This is because the surface

71 air sampling sites are minimally sensitive to the Amazon and the number of total column CH<sub>4</sub>  
72 estimates from space likely suffer from both temporal sampling bias (data are concentrated in the  
73 early dry season between seasons of smoke and clouds) and measurement biases<sup>11</sup>. In contrast *in-*  
74 *situ* measured vertical profile data capture directly the surface flux signals and discern the boundary  
75 layer signal from the free troposphere signal<sup>12</sup>. New measurements are therefore required to resolve  
76 the discrepancy between bottom-up inventories and top-down estimates which cannot be  
77 reconciled via contributions from other currently reported CH<sub>4</sub> sources from the Amazon region e.g.,  
78 biomass burning, termites and ruminants<sup>5,13</sup> nor UV-induced aerobic emissions from plants<sup>14</sup> and  
79 tank bromeliads<sup>15</sup>. Further, the regional stable carbon isotope composition (i.e., <sup>13</sup>C/<sup>12</sup>C ratio  
80 expressed as a δ<sup>13</sup>C value) of atmospheric CH<sub>4</sub> indicates unequivocally that the ‘missing’ Amazonian  
81 CH<sub>4</sub> source is derived from microbial metabolism of C<sub>3</sub> photosynthate<sup>16</sup>. Consequently, the most  
82 likely scenario is that surface-based flux measurements have either missed intense but perhaps  
83 spatially disaggregated CH<sub>4</sub> emission sources or they have overlooked an important pathway for  
84 egress of soil-produced CH<sub>4</sub>.

85 Trees subjected to permanent or periodic inundation develop adaptive features such as enlarged  
86 lenticels and hollow aerenchyma tissue to enhance oxygenation of their root systems<sup>17,18</sup>. The  
87 internal conduits that enable air to move downwards also facilitate upward escape of soil CH<sub>4</sub> to the  
88 atmosphere<sup>7,17,18</sup>. Tree-mediated gas emission has been shown to dominate ecosystem CH<sub>4</sub>  
89 emissions in tropical peat swamp forest where aerobic CH<sub>4</sub>-oxidizing bacteria form a highly effective  
90 barrier to diffusive flux through peat soil<sup>7</sup>. Total CH<sub>4</sub> emission rates are relatively modest in Borneo  
91 peat swamps<sup>1,7</sup>; however, the capacity for trees to emit CH<sub>4</sub> at higher rates is determined largely by  
92 rates of soil CH<sub>4</sub> production and supply<sup>18</sup>. Tree-mediated transport of CH<sub>4</sub> has not been investigated  
93 to date in the seasonally flooded, dense forests of the Amazon floodplains although ongoing efforts  
94 continue to extend the database of flux measurements quantifying CH<sub>4</sub> emission from soil, emergent  
95 macrophytes<sup>8,9</sup>, and open water<sup>8,19,20</sup>.

96 We measured CH<sub>4</sub> fluxes at 13 floodplain locations in the central Amazon River basin (Fig.1a),  
97 quantifying emissions from all known transport pathways, including forested floodplain soil, aquatic  
98 surfaces, and floating herbaceous macrophytes as well as stem and leaf surfaces of mature and  
99 young trees. At each floodplain site, a 50 × 80 m plot was established that encompassed four  
100 transects in which water table depth varied from ~1 m below the soil surface to ~10 m above the soil  
101 surface. Nine of the 12 sites sampled in 2014 included an area of exposed floodplain soil in which  
102 large hummocks occupied <13.5% of the total surface area. The relative contribution of emissions  
103 from individual pathways was determined relative to total ecosystem CH<sub>4</sub> flux (Table 1). Methane  
104 emissions from tree stems and aquatic surfaces were the dominant egress pathways (Fig. 1; Table 1).

105 All trees studied released substantial quantities of CH<sub>4</sub>. Emission rates for mature and young trees  
106 ranged from 0.33 to 337 mg m<sup>-2</sup> stem h<sup>-1</sup> and 0.39 to 581 mg m<sup>-2</sup> stem h<sup>-1</sup>, respectively. Methane flux  
107 from tree stems exceeded CH<sub>4</sub> emissions from all other pathways in the study plots (Fig. 1b-f; Table  
108 1). Moreover, CH<sub>4</sub> emission rates from Amazon floodplain trees were ~150 times larger than stem  
109 flux rates reported for southeast Asian peat swamp forests<sup>7</sup> where less CH<sub>4</sub> is released owing to low  
110 soil pH, high CH<sub>4</sub> oxidation rates and recalcitrant carbon impeding rates of methanogenesis. Fewer  
111 than 4% of wood cores extracted from tree stems at 20 and 130 cm above the soil or water surface

112 displayed capacity for CH<sub>4</sub> production (Table 2) and stem cores from sampled trees displayed no  
113 visual sign of wood rot. These observations suggest that CH<sub>4</sub> emitted from the tree stems originated  
114 in the floodplain soil.

115 The  $\delta^{13}\text{C}$  values of tree-mediated CH<sub>4</sub> flux ranged from -76.3 to -59.1‰, averaging  $-66.2 \pm 6.4\%$  (n =  
116 18; Table 3) consistent with the stable carbon isotope composition of CH<sub>4</sub> in soil water (range -70.8  
117 to -54.5‰; Table 3) in the study plots. The  $\delta^{13}\text{C}$  values are typical for wetland CH<sub>4</sub> albeit more  
118 negative than values generally attributed to tropical wetlands<sup>21</sup>.

119 Young tree leaves emitted small but significant quantities of CH<sub>4</sub> (Fig.1b-f; Table 1). Methane  
120 emission from mature leaves, if present, was below the instrument detection limit of c. 2 ppbv.  
121 Similar to temperate<sup>6</sup> and other tropical<sup>7</sup> trees, stem CH<sub>4</sub> flux rates decreased either linearly or  
122 exponentially with increasing stem height sampling position.

123 We pursued two approaches to scaling fluxes to the entire Amazon basin. Firstly, the measured CH<sub>4</sub>  
124 emission rates and areas of emission surfaces (Supplementary Table 3) were used to estimate the  
125 contribution of each transport pathway to total ecosystem CH<sub>4</sub> flux estimated for each 50 × 80 m  
126 study plot and then averaged for the river type. Emissions from tree stems and leaves collectively  
127 were the dominant source of CH<sub>4</sub> evasion from Amazon floodplain soil (44 to 65 %; Table 1). The  
128 contribution from aquatic surfaces was the second most significant source, accounting for 27 to 41%  
129 of total CH<sub>4</sub> flux. Soil surfaces, which were corrected for tree basal areas, emitted 2.5 to 15.7% of  
130 ecosystem CH<sub>4</sub> flux (Table 1). Conservative scaling of stem emission (considering only 0-140 cm of  
131 tree stem emissions) to the central Amazon basin<sup>22</sup> yields an annual source strength of  $15.1 \pm 1.8$  Tg  
132 CH<sub>4</sub> yr<sup>-1</sup> for tree-mediated flux (Table 4). Inclusion of tree emissions to 2.3-5 m stem height,  
133 estimated using the relationship between stem CH<sub>4</sub> flux and stem height intervals, yields an annual  
134 source strength of  $21.2 \pm 2.5$  Tg CH<sub>4</sub> yr<sup>-1</sup>, which is equivalent to current bottom-up inventories of  
135 total CH<sub>4</sub> emissions for Amazonian wetlands ( $26.2 \pm 9.8$  Tg yr<sup>-12,3</sup>; Table 4) that exclude tree  
136 emissions. Further, while recent evidence suggests the potential for non-wetland trees to emit CH<sub>4</sub><sup>23-</sup>  
137 <sup>25</sup>, no robust measurements of upland tree emission have been reported in the region and those few  
138 flux measurements reported elsewhere have been several orders of magnitude smaller than our  
139 wetland tree observations, so in keeping with our conservative approach to regional upscaling we  
140 have excluded upland tree fluxes pending further evidence.

141 Secondly, during the period 2010 to 2013 we also established top-down regional estimates of CH<sub>4</sub>  
142 emissions based upon novel regularly measured *in-situ* atmospheric CH<sub>4</sub> profiles from the surface to  
143 4.5 km height above sea level using an air-column budgeting approach. Profiles were measured at  
144 four locations in the Amazon basin (Alta Floresta (ALF), Rio Branco (RBA), Santarém (SAN) and  
145 Tabatinga (TAB)). Flux estimates determined using this approach integrate CH<sub>4</sub> emissions from  
146 regions upwind of the sampling sites, covering an increasing area the farther west a site is located in  
147 the basin. Based on the envelope of back-trajectory ensembles we estimate the regions of influence  
148 to be 2.53 million km<sup>2</sup> for TAB, 3.67 million km<sup>2</sup> for RBA, 0.59 million km<sup>2</sup> for SAN and 1.31 million  
149 km<sup>2</sup> for ALF. The total Amazon basin area is 6.7 million km<sup>2</sup>. The upwind regions of all four sites  
150 during all four years were a significant source of CH<sub>4</sub> to the atmosphere with emission rates varying  
151 from  $11.4 \pm 4.5$  to  $15.9 \pm 2.2$  mg CH<sub>4</sub> m<sup>-2</sup> day<sup>-1</sup> at ALF,  $11.4 \pm 1.6$  to  $15.4 \pm 3.2$  mg CH<sub>4</sub> m<sup>-2</sup> day<sup>-1</sup> at RBA,

152 11.1 ± 4.7 to 18.9 ± 3.2 mg CH<sub>4</sub> m<sup>-2</sup> day<sup>-1</sup> at TAB and 48.4 ± 7.6 to 60.9 ± 6.3 mg CH<sub>4</sub> m<sup>-2</sup> day<sup>-1</sup> at SAN.  
 153 We observed substantially larger mean annual fluxes at SAN relative to the other three sites, which  
 154 is consistent with spatial differences observed in CH<sub>4</sub> emission rates within our 13 floodplain study  
 155 plots. The SAN area of influence includes the Tapajós River where we measured the largest CH<sub>4</sub>  
 156 fluxes from trees and other sources among the 13 floodplain study plots (T10, T11, T12; Fig. 1a).

157 Extrapolation of inversion results to the whole of the Amazon basin using an area-weighted average

158  $(F = \bar{F} \cdot A_{basin} \text{ with } \bar{F} = \sum_{i=1}^4 (\frac{A_i}{\sum_{n=1}^4 A_n}) \cdot F_i, A_{basin} = 6.7 \cdot 10^6 \text{ km}^2)$

159  $(F = \bar{F} \times A_{basin} \text{ with } \bar{F} = \sum_{i=1}^4 (\frac{A_i}{\sum_{n=1}^4 A_n}) \times F_i, A_{basin} = 6.7 \times 10^6 \text{ km}^2)$  yields a mean total CH<sub>4</sub> flux of

160 42.7 ± 5.6 Tg CH<sub>4</sub> yr<sup>-1</sup> for the four-year period, which is the equivalent of ~8% of global CH<sub>4</sub>  
 161 emissions. The uncertainty of 5.6 Tg CH<sub>4</sub> yr<sup>-1</sup> is the standard deviation (1σ) of the four annual  
 162 emission estimates. In an earlier study<sup>26</sup>, we used the 2010-2011 vertical profile data and a simple  
 163 Bayesian synthesis inversion approach constrained by both prior flux estimates and atmospheric  
 164 profile data to obtain a net flux estimate of 37 ± 5.9 Tg yr<sup>-1</sup>. For all inversions and periods considered,  
 165 the estimated fluxes exceeded the prior flux estimates with wetland prior fluxes based either on the  
 166 JULES land surface model or the model of Bloom *et al.*<sup>2</sup>. While these earlier estimates are somewhat  
 167 smaller than the estimates reported here, this is expected because the presence of the prior flux  
 168 estimates biases the estimates low. The combinations of floodplain tree emissions (15.1 ± 1.8 - 21.2  
 169 ± 2.5 Tg CH<sub>4</sub> yr<sup>-1</sup>) and CH<sub>4</sub> emission from other transport pathways (20.5 ± 5.3 Tg yr<sup>-1</sup>) yields a total  
 170 that agrees well with our estimate of regional CH<sub>4</sub> emissions determined from inversion modelling of  
 171 atmosphere CH<sub>4</sub> profiles. Thus, inclusion of tree-mediated CH<sub>4</sub> fluxes reconciles current disparities  
 172 between ‘bottom-up’ and ‘top down’ approaches effectively closing the Amazonian CH<sub>4</sub> budget.

173 Our results demonstrate that exceptionally large emissions from Amazon floodplain trees alone are  
 174 equivalent in size to the entire Arctic CH<sub>4</sub> source and account for ~15% of the global wetland CH<sub>4</sub>  
 175 source. Together with already understood emission pathways, our findings demonstrate that the  
 176 Amazon, in contributing up to a third of the global wetland CH<sub>4</sub> source, is a far larger source of CH<sub>4</sub>  
 177 than inventories previously acknowledged and is therefore likely to exert greater influence over  
 178 global atmospheric CH<sub>4</sub> concentration variability than was previously thought. Given this increased  
 179 influence over atmospheric CH<sub>4</sub> there is a need to quantify the controls on soil CH<sub>4</sub> production and  
 180 tree emission variability within the biodiverse, hydrologically dynamic and geochemically  
 181 heterogeneous Amazon basin while re-appraising representation of CH<sub>4</sub> transport mechanisms in  
 182 process-based wetland models if global models are to possess the capacity to accurately predict  
 183 changes in CH<sub>4</sub> flux resulting from climate change or other human perturbations such as the planned  
 184 construction of hydroelectric dams across the basin<sup>27</sup>. Finally, given that tropical forested wetlands  
 185 spanning the Congo and southeast Asia experience either seasonal or permanent inundation,  
 186 wetland-adapted trees may be responsible for a similar proportion of CH<sub>4</sub> flux in those regions,  
 187 pointing to potential gross underestimates in bottom-up CH<sub>4</sub> inventories across globally important  
 188 regions using current approaches that exclude trees.

189 **References**

190 1 Kirschke, S. *et al.* Three decades of global methane sources and sinks. *Nature Geoscience* **6**, 813-823, doi:10.1038/ngeo1955 (2013).

191 2 Bloom, A. A., Palmer, P. I., Fraser, A. & Reay, D. S. Seasonal variability of tropical wetland CH<sub>4</sub> emissions: the role of the methanogen-available

192 carbon pool. *Biogeosciences* **9**, 2821-2830, doi:10.5194/bg-9-2821-2012 (2012).

193 3 Melack, J. M. *et al.* Regionalization of methane emissions in the Amazon Basin with microwave remote sensing. *Global Change Biology* **10**, 530-

194 544, doi:10.1111/j.1529-8817.2003.00763.x (2004).

195 4 Bergamaschi, P. *et al.* Inverse modeling of global and regional CH<sub>4</sub> emissions using SCIAMACHY satellite retrievals. *Journal of Geophysical*

196 *Research-Atmospheres* **114**, doi:10.1029/2009jd012287 (2009).

197 5 Frankenberg, C. *et al.* Pressure broadening in the 2 ν<sub>3</sub> band of methane and its implication on atmospheric retrievals. *Atmospheric Chemistry*

198 *and Physics* **8**, 5061-5075 (2008).

199 6 Pangala, S. R., Hornibrook, E. R. C., Gowing, D. J. & Gauci, V. The contribution of trees to ecosystem methane emissions in a temperate forested

200 wetland. *Global Change Biology* **21**, 2642-2654, doi:10.1111/gcb.12891 (2015).

201 7 Pangala, S. R., Moore, S., Hornibrook, E. R. C. & Gauci, V. Trees are major conduits for methane egress from tropical forested wetlands. *New*

202 *Phytologist* **197**, 524-531, doi:10.1111/nph.12031 (2013).

203 8 Devol, A. H., Richey, J. E., Forsberg, B. R. & Martinelli, L. A. Seasonal dynamics in methane emissions from the Amazon River floodplain to the

204 troposphere. *Journal of Geophysical Research-Atmospheres* **95**, 16417-16426, doi:10.1029/JD095iD10p16417 (1990).

205 9 Bartlett, K. B. *et al.* Methane flux from the central Amazonian floodplain. *Journal of Geophysical Research: Atmospheres* **93**, 1571-1582,

206 doi:10.1029/JD093iD02p01571 (1988).

207 10 Ringeval, B. *et al.* Methane emissions from floodplains in the Amazon Basin: challenges in developing a process-based model for global

208 applications. *Biogeosciences* **11**, 1519-1558, doi:10.5194/bg-11-1519-2014 (2014).

209 11 Webb, A. J. *et al.* CH<sub>4</sub> concentrations over the Amazon from GOSAT consistent with in situ vertical profile data. *Journal of Geophysical Research-*

210 *Atmospheres* **121**, 11006-11020, doi:10.1002/2016jd025263 (2016).

211 12 Chou, W. W. *et al.* Net fluxes of CO<sub>2</sub> in Amazonia derived from aircraft observations. *Journal of Geophysical Research-Atmospheres* **107**,

212 doi:10.1029/2001jd001295 (2002).

213 13 Frankenberg, C., Meirink, J. F., van Weele, M., Platt, U. & Wagner, T. Assessing methane emissions from global space-borne observations. *Science*

214 **308**, 1010-1014, doi:10.1126/science.1106644 (2005).

215 14 Bloom, A. A. *et al.* Global methane emission estimates from ultraviolet irradiation of terrestrial plant foliage. *New Phytologist* **187**, 417-425,

216 doi:10.1111/j.1469-8137.2010.03259.x (2010).

217 15 Martinson, G. O. *et al.* Methane emissions from tank bromeliads in neotropical forests. *Nature Geoscience* **3**, 766-769, doi:10.1038/ngeo980

218 (2010).

219 16 Beck, V. *et al.* Methane airborne measurements and comparison to global models during BARCA. *JOURNAL OF GEOPHYSICAL RESEARCH-*

220 *ATMOSPHERES* **117**, doi:10.1029/2011JD017345 (2012).

221 17 Rusch, H. & Rennenberg, H. Black alder (*Alnus glutinosa* (L.) Gaertn.) trees mediate methane and nitrous oxide emission from the soil to the

222 atmosphere. *Plant and Soil* **201**, 1-7, doi:10.1023/a:1004331521059 (1998).

223 18 Pangala, S. R., Gowing, D. J., Hornibrook, E. R. C. & Gauci, V. Controls on methane emissions from *Alnus glutinosa* saplings. *New Phytologist* **201**,

224 887-896, doi:10.1111/nph.12561 (2014).

225 19 Sawakuchi, H. O. *et al.* Methane emissions from Amazonian Rivers and their contribution to the global methane budget. *Global Change Biology*

226 **20**, 2829-2840, doi:10.1111/gcb.12646 (2014).

227 20 Belger, L., Forsberg, B. R. & Melack, J. M. Carbon dioxide and methane emissions from interfluvial wetlands in the upper Negro River basin, Brazil.

228 *Biogeochemistry* **105**, 171-183, doi:10.1007/s10533-010-9536-0 (2011).

229 21 Schaefer, H. *et al.* A 21st-century shift from fossil-fuel to biogenic methane emissions indicated by 13CH<sub>4</sub>. *Science* **352**, 80-83 (2016).

230 22 Hess, L. L. *et al.* Wetlands of the lowland Amazon basin: extent, vegetative cover, and dual-season inundated area as mapped with JERS-1

231 Synthetic Aperture Radar. *Wetlands* **35**, 745-756, doi:10.1007/s13157-015-0666-y (2015).

232 23 Covey, K. R., Wood, S. A., Warren, R. J., Lee, X. & Bradford, M. A. Elevated methane concentrations in trees of an upland forest. *Geophysical*

233 *Research Letters* **39**, doi:10.1029/2012gl052361 (2012).

234 24 Pitz, S. & Megonigal, J. P. Temperate forest methane sink diminished by tree emissions. *New Phytologist* **214**, 1432-1439, doi:10.1111/nph.14559

235 (2017).

- 236 25 Wang, Z. P. *et al.* Methane emissions from the trunks of living trees on upland soils. *New Phytologist* **211**, 429-439, doi:10.1111/nph.13909  
237 (2016).
- 238 26 Wilson, C. *et al.* Contribution of regional sources to atmospheric methane over the Amazon Basin in 2010 and 2011. *Global Biogeochemical Cycles*  
239 **30**, 400-420, doi:10.1002/2015gb005300 (2016).
- 240 27 Latrubesse, E. M. *et al.* Damming the rivers of the Amazon basin. *Nature* **546**, 363-369, doi:10.1038/nature22333 (2017).

## 241 **Supplementary information**

242 This file contains supplementary tables (1-5) and supplementary figures (1-2).

## 243 **Acknowledgements**

244 The overall bottom-up measurement study was funded by the following grants: UK Natural  
245 Environment Research Council to VG (PI) and EH (grant no: NE/J010928/1); several CNPq (Brazil  
246 National Council of Research and Development) and FAPERJ (Science Foundation from the State of  
247 Rio de Janeiro) grants to AEP; Swedish Research Council VR (grant no: 2012-00048) to DB; and a  
248 Swedish-Brazil exchange grant from STINT – Capes to DB and AEP (grant no: 2012-2085).  
249 Atmospheric CH<sub>4</sub> vertical profile measurements were funded by the following agencies: NERC  
250 consortium AMAZONICA led by EG (grant no: NE/F005806/1) with the consortium including LVG and  
251 JBM, FAPESP (State of Sao Paulo Science Foundation) via the 'Carbon Tracker' and (FAPESP-NERC) via  
252 the 'ACO' project (08/58120-3; 11/51841-0), the EU via the 7th grant framework GEOCARBON  
253 project (283080) and CNPq (403241/2012-3) awarded to LVG. NASA, NOAA, IPEN and INPE made  
254 large contributions to the construction and maintenance of the GHG laboratory in Brazil. We also  
255 acknowledge support from NERC FAPESP ACO project which funded a workshop where bottom up  
256 and top down data were brought together. VG and EG would like to thank the NERC consortium  
257 MOYA (grant no: NE/N015606/1) and DB, the Swedish Research Council VR for salary contributions.  
258 VG and SRP acknowledge support from the AXA Research Fund. We thank Tiago for assistance with  
259 tree species identification and André Breves, Eliane Cristina, Fausto Machado Silva, Juliana Valle,  
260 João Paulo Felizardo, Laís Rodrigues, Luciene Valladares, Paula Souto, Ricardo Pollery and Viviane  
261 Figueiredo for field work assistance. Ani Dwarakanath is thanked for data analysis and Paul  
262 Monaghan and Adam McAleer for performing  $\delta^{13}\text{C}$ -CH<sub>4</sub> analyses.

## 263 **Author contributions**

264 SRP, VG, AP and DB conceived and designed the bottom-up measurement study. The Brazil  
265 expeditions (bottom-up measurements) in 2013 and 2014 were planned and organised by AP, OM,  
266 WRB, RBP and SRP, which was carried out by SRP, RBP, HMR, LSBC and WRB. EH was responsible for  
267  $\delta^{13}\text{C}$ -CH<sub>4</sub> analysis and interpretation of those data. LSBC and CMS identified the tree species in the  
268 2014 Brazil expedition. The top down measurement study was designed and carried out by LSB, LVG,  
269 JM and EG. VG coordinated integration of the various elements of the study. SRP, VG, LB, EG, DB, EH  
270 and AP all contributed to writing of the manuscript.

## 271 **Author information**

272 Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). Authors declare no  
273 competing financial interests. Correspondence and requests for materials should be addressed to  
274 SRP (Sunitha.Pangala@open.ac.uk) and VG ([Vincent.Gauci@open.ac.uk](mailto:Vincent.Gauci@open.ac.uk)).

## 275 **Main table legends**

276 **Table 1:** Methane fluxes and estimated ecosystem contributions from five major rivers in the central  
277 Amazon basin.  
278 **Table 2:** Methane production potentials measured from the wood cores extracted.  
279 **Table 3:**  $\delta^{13}\text{C}$  values of tree  $\text{CH}_4$  flux and porewater  $\text{CH}_4$ .  
280 **Table 4:** Estimated annual  $\text{CH}_4$  emissions from the Amazon basin using bottom up and top down methods.

281 **Main figure legends**

282 **Figure 1: Sampling site locations and  $\text{CH}_4$  flux distributions.** a) Map showing the location of the 13  
283 sampling sites within the central Amazon River basin, Brazil. (×) and (●) represent the sites sampled  
284 in 2013 and 2014, respectively. Sampling sites are labelled: S1, S2 (River Solimões); N3, N4, N5, N6  
285 (River Negro); A7, A8, A9 (River Amazon); T10, T11, T12 (River Tapajós) and M13 (River Madeira).  
286 Box and whisker plots showing the distribution of  $\text{CH}_4$  fluxes measured from all  $\text{CH}_4$  emitting  
287 pathways from river b) Negro, c) Madeira, d) Amazon, e) Solimões and f) Tapajós. Box plots  
288 represents  $\text{CH}_4$  fluxes measured from mature tree stem surfaces (M.stems), young tree stem  
289 surfaces (Y.stems), young tree leaf surfaces  $\times 10^{-2}$  (Y.leaves), emergent macrophytes (MAC), aquatic  
290 surfaces where the water table was 0-10 m above the soil surface and soil surfaces where the water  
291 table was 0-1 m below the soil surfaces. Stem  $\text{CH}_4$  fluxes for mature trees were measured at four 30  
292 cm intervals between 20 and 140 cm and young trees at 10 cm intervals between 15 and 135 cm.  
293 The box plot represents the averaged flux value between the 20 to 140 cm stem portion for mature  
294 trees and 15 to 135 cm for young trees.  $\text{CH}_4$  fluxes ( $\text{mg m}^{-2} \text{hr}^{-1}$ ) are expressed per unit area of the  
295  $\text{CH}_4$  emitting surface measured.

296 **Methods**

297 **Ecosystem scale measurements**

298 Thirteen temporary plots (50 × 80 m) were set up in the floodplains (várzeas and Igapó) of the five  
299 major rivers of the central Amazon basin, Brazil. During 2013, sampling was conducted at the Cuniã  
300 ecological field station (Rondônia) a floodplain fed by the River Madeira (Fig. 1). During 2014, all  
301 sampling locations ( $n = 12$ ) were within the 1.77 million km<sup>2</sup> reference quadrant of the central  
302 Amazon basin previously characterised in detail with Synthetic Aperture Radar (SAR) imagery<sup>3,28</sup>. The  
303 12 sampling locations consisted of four sampling locations in River Negro (black water), two in River  
304 Solimões (white water), three in River Amazon (white water), and three in River Tapajós (clear  
305 water). Methane sampling was conducted in the flooded forests (Supplementary Table 1) and  
306 sample locations S1, S2, A7, A8 and M13 were comprised of várzeas with white waters, neutral pH,  
307 and high sediment load from the Andean and pre-Andean regions. Sample plots N3, N4, N5, N6, T10,  
308 T11 and T12 consisted of igapós with black water (N3, N4, N5 and N6) or clear water (T10, T11 and  
309 T12), having a pH ranging from 4 to 5.5 and 4.4 to 7, respectively. Our measurements across the 13  
310 sites ensured that any differences between the distinct water types (clear, white and black)  
311 characteristic of the Amazon River and attributed mostly to its channel morphology and geology  
312 were captured.

313 Within each study plot, stem CH<sub>4</sub> flux from mature trees (diameter at breast height; DBH = 6-74 cm;  
314 tree height = 5-22 m;  $n = 1759$  trees; Supplementary Table 2) was measured at 30 cm intervals  
315 between 20 and 140 cm height and for young trees (tree height ≤ 5 m; DBH ≤ 6 cm;  $n = 598$  trees) at  
316 10 cm intervals between 15 and 135 cm above the soil/water surface. CH<sub>4</sub> emissions from young and  
317 mature trees were measured across the plot, split into four transects within which the water table  
318 depths ranged from wet (0-10 m above the soil surface) to dry (0 – 1 m below the soil surface)  
319 conditions. Methane emissions from stems of mature and young trees were measured using static  
320 chambers as described by Pangala *et al.*<sup>7,18</sup> and Siegenthaler *et al.*<sup>29</sup>. Methane emissions ( $n = 207$ )  
321 were measured from aquatic surfaces within each plot, inside the flooded forests using floating  
322 chambers (Supplementary Figure 1) deployed for 24 hours as described by Bastviken *et al.*<sup>30</sup>. Floating  
323 chambers were deployed in four transects within each plot, where the water table depths ranged  
324 from 0 to 10 m above the soil surface. These transects also extended into the raised hummocks  
325 where the water-table was below the soil surface and in these areas soil CH<sub>4</sub> fluxes ( $n = 380$ ) were  
326 measured using cylindrical static chambers (30 × 30 cm; diameter × height; Supplementary Figure 1).  
327 'Aquatic surfaces' refers to the water body within the flooded forest and does not include 'open  
328 waters' outside the flooded forest with no vegetation.

329 Floating chambers (1 × 1 × 1.5 m; height × width × length) were used to measure CH<sub>4</sub> emissions from  
330 emergent floating macrophytes ( $n = 80$ ). The chambers were constructed of gas-impermeable  
331 fluorinated ethylene propylene film (Adtech Ltd., Gloucestershire, UK) wrapped around a pipe  
332 frame. Floats were attached to the bottom of the frame. Emergent macrophytes were absent in  
333 study locations in the River Negro catchment probably due to low nutrient concentrations in the  
334 acidic black waters. Due to receding water table levels, floating macrophytes were absent in River  
335 Madeira. Therefore, CH<sub>4</sub> fluxes from emergent floating macrophytes were measured only in Rivers

336 Solimões, Amazon and Tapajós. Rooted macrophytes were absent in all sampling locations during  
337 our study period.

338 Leaf emissions were measured from leaf surfaces of young trees ( $n = 260$  trees) and mature trees  
339 (when accessible;  $n = 180$  trees) using static chambers as described by Pangala *et al.*<sup>18</sup>. The  
340 chambers, which enclosed four different branches per tree, were deployed for 10 minutes during  
341 each flux measurement. In the 2014 campaign, we measured CH<sub>4</sub> emissions from tree stem and leaf  
342 surfaces in the flooded forest and emergent macrophytes in real-time by cavity-ring down laser  
343 spectroscopy as described in Pangala *et al.*<sup>18</sup>. However, on days with heavy rainfall, gas sampling  
344 and analysis were conducted as described in Pangala *et al.*<sup>7</sup> i.e. collection via syringes and later  
345 analysis for CH<sub>4</sub> content. Methane emissions from tree stems and leaf surfaces from trees with  
346 water table below the soil surface in the 2014 campaign and all measurements in the 2013  
347 campaigns were performed as described in Siegenthaler *et al.*<sup>29</sup> and Pangala *et al.*<sup>7</sup>, respectively.  
348 Gas samples from chambers enclosing soil and aquatic surfaces were extracted using a syringe and  
349 then transferred to glass vials for CH<sub>4</sub> analysis by modified cavity ring down laser spectroscopy<sup>6,7</sup>. CH<sub>4</sub>  
350 fluxes are expressed per unit surface area enclosed within the corresponding static chambers and  
351 fluxes therefore reported as mg m<sup>-2</sup> h<sup>-1</sup> correspond to mg m<sup>-2</sup> soil h<sup>-1</sup> for soil fluxes, mg m<sup>-2</sup> stem h<sup>-1</sup>  
352 for mature and young stem fluxes, mg m<sup>-2</sup> leaf h<sup>-1</sup> for leaf fluxes, mg m<sup>-2</sup> aquatic h<sup>-1</sup> for aquatic fluxes  
353 and mg m<sup>-2</sup> MAC h<sup>-1</sup> for macrophytes fluxes. Two sets of wood cores were extracted diagonally at 20  
354 and 130 cm stem height above the forest floor/water surface for 67% and 73%, respectively, of  
355 mature trees investigated for stem CH<sub>4</sub> fluxes. The wood cores were incubated to investigate CH<sub>4</sub>  
356 production potential as described by Covey *et al.*<sup>23</sup>.

357 Gas samples were collected from flux chambers and porewater (head space equilibration method)  
358 for  $\delta^{13}\text{C}$ -CH<sub>4</sub> analysis using gas-tight syringes and then transferred to evacuated ( $10^{-3}$  bar) 125 ml  
359 Wheaton® vials fitted with Bellco® stoppers and crimp seals. Vials were over-pressured by  $\sim 0.5$  bar  
360 to ensure ingress of air did not occur as a result of pressure or temperature changes during transport  
361 to the laboratory. The  $\delta^{13}\text{C}$  values of CH<sub>4</sub> were measured using a ThermoFinnigan® Delta XP stable  
362 isotope ratio mass spectrometer. Methane in the glass vials was purified and combusted to CO<sub>2</sub>  
363 using a ThermoFinnigan PreCon®, which was modified to house a 6.4 mm stainless steel combustion  
364 reactor containing palladium on quartz wool heated to 780°C<sup>31</sup> and a Sofnocat® reagent trap  
365 operated at room temperature to remove carbon monoxide. The instrument was calibrated using  
366 BOC alpha-gravimetric and Isometric Ltd standards (ISO-B, ISO-H, ISO-L and ISO-T)<sup>32</sup>. Analysis  
367 precision based upon replicate measurements of standards containing 2 ppmv CH<sub>4</sub> was  $\pm 0.1\%$ . The  
368  $\delta^{13}\text{C}$  values and mixing ratios of CH<sub>4</sub> in the chamber headspace measured either three or four times  
369 during each 30 minute deployment were used to determine the  $\delta^{13}\text{C}$  value of CH<sub>4</sub> flux via Keeling  
370 regression analysis.

371 The locations of trees were mapped in each of the 13 study plots along with the area occupied by  
372 emergent macrophytes and water-table depths (measured within 1 m of all trees) along the  
373 boundary of the plot and within four internal transects. Tree height, DBH, stem diameter at 10 cm  
374 intervals between 0 and 200 cm stem height, and basal diameter were measured for all trees in each  
375 plot. The floodplain on River Madeira site sampled in 2013 was comprised of non-flooded forest  
376 because of receding water-table levels. Várzeas in the region had shrunk to small ponds with trees

377 around the edges, which were subjected to water-table levels at or below the soil surface. In all the  
378 study plots, the edge of the floodplain where floating macrophytes ceased to exist was regarded as  
379 the plot boundary and open water beyond that point, which contained no vegetation, was excluded  
380 from the ecosystem contribution estimations but was later included in the regional upscaling using  
381 the literature values<sup>8</sup>. Nine of the 12 sites investigated during 2014 contained both flooded and non-  
382 flooded portions (<13.5%) of floodplain, three sites were fully flooded. Area occupied by aquatic  
383 surfaces, soil surfaces and mature and young trees were mapped for each study site and the  
384 corresponding surface areas were calculated.

385 Using ArcGIS, a polygon map for each of the sampling sites was developed, which contained water  
386 table depth information and locations of trees across the transects. A spatial distribution model  
387 developed from the information collected during the campaign was used to estimate macrophyte  
388 surface area, aquatic surface area and soil surface areas after deducting tree basal area  
389 (Supplementary Table 3). Methane fluxes from soil and water surfaces, and macrophytes were  
390 estimated using CH<sub>4</sub> emission rates measured during the campaign and emission surfaces estimated  
391 using the spatial distribution model. The leaf surface area of the young trees were estimated using  
392 the methods described by Santiago *et al.*<sup>33</sup> which was multiplied by measured leaf CH<sub>4</sub> flux rates to  
393 determine total ecosystem leaf CH<sub>4</sub> emissions. Using the stem diameter measured between 20 and  
394 140 cm stem height, stem surface area was estimated and multiplied by the corresponding stem CH<sub>4</sub>  
395 flux rate to obtain stem emissions for each tree. Stem CH<sub>4</sub> emissions for individual trees measured  
396 along the length of trees were then estimated based upon relationships between stem CH<sub>4</sub> flux rates  
397 and stem sampling position at 30 cm tree stem height intervals. Approximately 42% of trees  
398 measured displayed a linear relationship ( $R^2 > 0.95$ ;  $P < 0.0001$ ) between stem sampling height and  
399 stem CH<sub>4</sub> flux rate. Trees exhibiting such a relationship had stem CH<sub>4</sub> flux rates equal to zero at stem  
400 height between 2.3 and 3.5 m. The remaining trees studied exhibited an exponential relationship  
401 between stem CH<sub>4</sub> flux rate and stem height. Although regression models based on exponential  
402 relationships suggested the possibility of the entire tree emitting CH<sub>4</sub>, we set stem CH<sub>4</sub> emissions to  
403 zero when the percentage difference between the ratios of stem CH<sub>4</sub> flux at two consecutive 30 cm  
404 stem height intervals was  $\geq 0.1\%$ . In such cases, stem CH<sub>4</sub> flux rate was equal to zero at stem heights  
405 ranging between 3.8 and 5 m. Using the stem diameter measured at 10 cm intervals between 20 and  
406 200 cm stem height, a relationship was established (exponential and/or power function relationship)  
407 to estimate stem circumference and surface area for each tree up to 5 m. Total CH<sub>4</sub> emission up to  
408 2.3 - 5 m length of the individual trees based upon the relationship each tree followed, was  
409 estimated by multiplying measured and/or estimated CH<sub>4</sub> flux rates and corresponding stem surface  
410 areas (Supplementary Table 3). Average stem CH<sub>4</sub> flux per tree was estimated by dividing total stem  
411 emissions measured by the number of trees studied, within each study plot. The average flux rate  
412 per tree subsequently was multiplied by the total number of trees within each plot to obtain total  
413 ecosystem CH<sub>4</sub> contribution from trees for each study site.

414 To estimate total annual CH<sub>4</sub> contributions from the entire lowland Amazon basin, we averaged CH<sub>4</sub>  
415 emissions across 13 sites for each individual pathways studied, assumed the estimated fluxes are  
416 representative of basin-wide fluxes and then applied the fluxes to the entire Amazon basin area,  
417 which was estimated using surface area data obtained from Melack *et al.*<sup>34</sup> and Hess *et al.*<sup>22</sup>  
418 (Supplementary Table 5). Monthly area coverage for open water, flooded forest and macrophytes in

419 1.77 million km<sup>2</sup> of the central Amazon basin were obtained from Melack *et al.*<sup>34</sup> and the percentage  
 420 decrease in water-table depths relative to October data (lowest water-table month reported for  
 421 most land cover classes by Melack *et al.*<sup>34</sup>) and percentage increase in water-table depths relative to  
 422 May data (highest water-table month reported for most land cover classes in Melack *et al.*<sup>34</sup>) was  
 423 estimated. The percentage increases/decreases were applied to the high and low water surface area  
 424 for flooded forest, open water and macrophyte area within the Amazon basin wetland area (8.4 ×  
 425 10<sup>5</sup> km<sup>2</sup>) reported in Hess *et al.*<sup>22</sup> and surface areas for the remaining months were estimated. Soil  
 426 surface area at the peak of the wet season was considered to be zero and for the remaining 11  
 427 months, soil surface area was estimated by subtracting the subsequent month flooded-forest  
 428 surface area and tree basal area from the flooded forest area during the peak of the wet season. Our  
 429 work suggests that up to 13.5% of the flooded forest was comprised of exposed soil and raised  
 430 hummocks in May, hence it is estimated that the soil surface area reached zero in June and  
 431 thereafter the water table receded. This observation was applied to soil surface area calculations.  
 432 Aquatic surface area was estimated by subtracting tree basal area from flooded-forest area.  
 433 Estimated monthly surface areas are listed in Supplementary Table 5. Tree-mediated CH<sub>4</sub> flux, similar  
 434 to other CH<sub>4</sub> emission pathways, was averaged across all 13 sites and was estimated to be 1350 ±  
 435 553 g ha<sup>-1</sup> d<sup>-1</sup> and 98 ± 47 g ha<sup>-1</sup> d<sup>-1</sup> for mature and young tree stem emissions between 0-140 cm  
 436 stem heights above the forest floor/water surface. However, when 0 to 5 m stem height was  
 437 considered the fluxes increased to 1927 ± 793 g ha<sup>-1</sup> d<sup>-1</sup> and 104 ± 49 g ha<sup>-1</sup> d<sup>-1</sup> for mature and young  
 438 trees, respectively. Open water CH<sub>4</sub> fluxes outside/beyond the edges of the flooded-forest were not  
 439 measured in our study. Fluxes from macrophytes were measured in some plots but the macrophytes  
 440 tended to be floating at the edges rather than inside the flooded-forest. Rooted macrophytes were  
 441 absent in all the plots. Thus CH<sub>4</sub> flux data for open water and macrophytes from Devol *et al.*<sup>8</sup> were  
 442 used to estimate these components for the entire Amazon basin. Uncertainties expressed as  
 443 standard deviation (SD) of means in CH<sub>4</sub> fluxes from all pathways were estimated using a  
 444 bootstrapping method (10,000 iterations).

#### 445 Aircraft measurements

446 To estimate CH<sub>4</sub> fluxes ( $F$ ) based on atmospheric CH<sub>4</sub> vertical profile measurements we apply a  
 447 simple air column budgeting technique following Miller *et al.*<sup>35</sup>:

$$448 \quad F = \int_{z=0 \text{ (agl)}}^{4.4 \text{ km}} \frac{\Delta CH_4(z)}{t(z)} dz$$

$$449 \quad F = \int_{z=0 \text{ (agl)}}^{4.4 \text{ km}} \frac{CH_4(z')}{t(z')} dz'$$

450 where  $\Delta CH_4 = CH_{4,site} - CH_{4,bg}$  is the difference between CH<sub>4</sub> mass per volume measured *in situ* at a site  
 451 inside the basin and background (*bg*) air entering the basin from the Atlantic,  $z$  is height above  
 452 ground (*agl*) and  $t(z)$  air-mass trajectory travel time from the coast to height  $z$  at the site. The CH<sub>4</sub>  
 453 concentration of background air is estimated from atmospheric SF<sub>6</sub> measured at the site and  
 454 compared with NOAA background stations Barbados (*RGB*, 7.92°S, 14.42°W) and Ascension (*ASC*,  
 455 7.92°S, 14.42°W) respectively, using a linear mixing model:

$$456 \quad CH_{4,bg} = f_{ASC} \cdot CH_{4,ASC} + (1 - f_{ASC}) \cdot CH_{4,RPB} \quad \text{with } f_{ASC} = \frac{SF_{6,site} - SF_{6,RPB}}{SF_{6,ASC} - SF_{6,RPB}}$$

$$CH_{4,bg} = f_{ASC} \times CH_{4,ASC} + (1 - f_{ASC}) \times CH_{4,RPB} \quad \text{with } f_A = \frac{S_{6,S} F_{S_{6,R}} I}{S_{6,A} F_{S_{6,R}} I}$$

SF<sub>6</sub> is suited for this purpose because it has virtually no sources in the Amazon Basin and atmospheric SF<sub>6</sub> concentration is substantially higher in the northern compared to the southern hemisphere. Air mass travel times are estimated using back trajectories calculate d using the HYSPLIT model<sup>36</sup> ([http://ready.arl.noaa.gov/HYSPLIT\\_traj.php](http://ready.arl.noaa.gov/HYSPLIT_traj.php)).

We applied this method to vertical air profiles sampled roughly bi-weekly from 2010 to 2013 at four sites in the Brazilian Amazon located along the main airstream: at Alta Floresta (ALF; 8.80°S, 56.75°W), Rio Branco (RBA; 9.38°S, 67.62°W), Santarém (SAN; 2.86°S; 54.95°W) and Tabatinga (TAB; 5.96°S, 70.06°W). Concomitantly, carbon monoxide (CO) also was measured which allowed us to determine the CH<sub>4</sub> component derived from fires during the dry season of each site. Air samples were collected using a two-component portable semi-automatic collection system, consisting of a first unit with two compressors and rechargeable batteries and a second unit with 17 (at SAN) and 12 (at ALF, RBA and TAB) 700 mL boro-silicate glass flasks connected by tubing and valves, which are opened and closed by a microprocessor. The samples were generally taken between noon and 1 PM local time, when the boundary layer tends to be well mixed. After sampling, the unit containing the air flasks was transported to the high-precision greenhouse gas laboratory at IPEN (Instituto de Pesquisas Energeticas e Nucleares) in Sao Paulo, where CH<sub>4</sub> and CO concentrations in air were quantified. The accuracy and precision (1.5 ppb) of our greenhouse gas analysis system in Brazil is similar to the system of the bottom up of NOAA (National Oceanic and Atmospheric Administration, USA)<sup>35</sup>.

## References

- 28 Hess, L. L., Melack, J. M., Novo, E., Barbosa, C. C. F. & Gastil, M. Dual-season mapping of wetland inundation and vegetation for the central Amazon basin. *Remote Sensing of Environment* **87**, 404-428, doi:10.1016/j.rse.2003.04.001 (2003).
- 29 Siegenthaler, A., Welch, B., Pangala, S. R., Peacock, M. & Gauci, V. Technical Note: Semi-rigid chambers for methane gas flux measurements on tree stems. *Biogeosciences* **13**, 1197-1207, doi:10.5194/bg-13-1197-2016 (2016).
- 30 Bastviken, D., Cole, J. J., Pace, M. L. & de Bogert, M. C. Fates of methane from different lake habitats: Connecting whole-lake budgets and CH<sub>4</sub> emissions. *JOURNAL OF GEOPHYSICAL RESEARCH-BIOGEOSCIENCES* **113**, doi:10.1029/2007JG000608 (2008).
- 31 Fisher, R., Lowry, D., Wilkin, O., Sriskantharajah, S. & Nisbet, E. G. High-precision, automated stable isotope analysis of atmospheric methane and carbon dioxide using continuous-flow isotope-ratio mass spectrometry. *Rapid Communications in Mass Spectrometry* **20**, 200–208 (2006).
- 32 Keppler, F. *et al.* Measurements of <sup>13</sup>C/<sup>12</sup>C methane from anaerobic digesters: comparison of optical spectroscopy with continuous-flow isotope ratio mass spectrometry. *Environmental Science & Technology* **44**, 5067–5073 (2010).
- 33 Santiago, L. S., Goldstein, G., Meinzer, F. C., Fownes, J. H. & Mueller-Dombois, D. Transpiration and forest structure in relation to soil waterlogging in a Hawaiian montane cloud forest. *Tree Physiology* **20**, 673-681 (2000).
- 34 Melack, J. M. *et al.* LBA-ECO LC-07 Monthly mean flooded wetlands habitat, Central Amazon Basin: 1979-1996. *Data set. Available on-line* [<http://daac.ornl.gov>] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. , doi:10.3334/ORNLDAAC/1049 (2011).
- 35 Miller, J. B. *et al.* Airborne measurements indicate large methane emissions from the eastern Amazon basin. *Geophysical Research Letters* **34**, doi:10.1029/2006gl029213 (2007).
- 36 Draxler, R. R. & Hess, G. D. An overview of the HYSPLIT\_4 modelling system for trajectories, dispersion, and deposition. *Australian Meteorological Magazine* **47**, 295-308 (1998).

## Data availability statement

499 Our aircraft CO<sub>2</sub> and CH<sub>4</sub> measurement data is available at <http://www.ccst.inpe.br/projetos/lagee/>.  
500 CH<sub>4</sub> flux data from the bottom up study are available from SRP on request.

501 **Supplementary table legends**

502 **Table 1:** Additional information for all sampling sites (50 × 80 m) in this study.

503 **Table 2:** Tree species identified within our 13 plots across the central Amazon basin.

504 **Table 3:** Surface area (m<sup>2</sup>) used to estimate ecosystem contributions from all CH<sub>4</sub> emitting pathways in each  
505 sampling plot.

506 **Table 4:** Coefficient of variation (%) for surface areas used in the ecosystem contribution  
507 estimations.

508 **Table 5:** Estimated surface areas for the entire lowland Amazon basin (km<sup>2</sup>)<sup>a</sup>.

509 **Supplementary figure legends**

510 **Figure 1:** Photographs depicting one of the study sites, a typically inundated flooded forest (a), soil  
511 flux (b), mature tree stem flux (c) and aquatic flux (d) measurements.

512 **Figure 2:** Frequency distribution of stem CH<sub>4</sub> fluxes from 20-50 cm of stem height from mature trees  
513 measured from river a) Negro, b) Madeira, c) Amazon, d) Solimões and e) Tapajós.

**Table 1:** Methane fluxes and estimated ecosystem contributions from five major rivers in the central Amazon basin.

Methane emitting pathways	River Negro		River Madeira		River Amazon		River Solimões		River Tapajós	
	Fluxes ± SD <sup>a</sup>	Ecosystem contribution	Fluxes ± SD	Ecosystem contributions						
	mg m <sup>-2</sup> h <sup>-1</sup>	g ha <sup>-1</sup> d <sup>-1</sup> (%)	mg m <sup>-2</sup> h <sup>-1</sup>	g ha <sup>-1</sup> d <sup>-1</sup> (%)	mg m <sup>-2</sup> h <sup>-1</sup>	g ha <sup>-1</sup> d <sup>-1</sup> (%)	mg m <sup>-2</sup> h <sup>-1</sup>	g ha <sup>-1</sup> d <sup>-1</sup> (%)	mg m <sup>-2</sup> h <sup>-1</sup>	g ha <sup>-1</sup> d <sup>-1</sup> (%)
<b>Mature tree stem emissions<sup>b</sup></b>	474 ± 151 (58.3)		836±323 (52.3)		823±214 (43.6)		1874±477 (53)		2866±759 (41.5)	
20-50 cm	30.2 ± 20.7		33.2±26		46.4 ± 33.7		83.2±42.8		141±71.4	
50-80 cm	22.2 ± 15.3		27.5±23.1		34.5 ± 25.6		62.4±32.4		106±54.5	
80-110 cm	15.4 ± 10.7		24.8±22.7		24.5 ± 18.3		44.2±23.1		73.5±38.4	
110-140 cm	10.7 ± 7.6		20.1± 19.4		16.7 ± 13.1		31.9±17.2		51.8±29.1	
<b>Young tree stem emissions<sup>b</sup></b>	47.4±11 (5.8)		83±33.2 (5.2)		50.3±13.3 (2.7)		157±40.5 (4.4)		181±56.1 (2.6)	
15-45 cm	59±28.2		50.2±32.9		103±44.9		150±67.4		271±109	
45-75 cm	41.9±20.2		42.5±32.3		73.5±32.8		108±49.9		180±74.1	
75-105 cm	29.1±14.1		35.4±31.7		50.6±23.4		77.6±36.2		125±54.1	
105-135 cm	18.9±9.7		28.5±25.7		32.8±16.4		49.1±24.2		77.83±38.3	
<b>Young tree leaf emissions<sup>c</sup></b>	0.016±0.04	3.86±4.6 (0.5)	0.019±0.04	5.07±4.8 (0.317)	0.038±0.07	5.93±7.3 (0.3)	0.051±0.09	13.5±13.1 (0.4)	0.09±0.11	17.3±15.7 (0.2)
<b>Macrophytes</b>	-	-	-	-	7.29±10.8	190±745 (10)	6.62±8.9	134±261 (3.8)	39±41.9	966±2105 (13.9)
<b>Aquatic emissions</b>	1.51±3.2	219±544 (27)	7.34±2.59	423±148 (26.5)	6.1±14.7	768±1792 (40.7)	4.37±5.77	1269±1111 (35.9)	25.7±29.8	2426±2898 (35.1)
<b>Soil emissions</b>	1.06±0.8	67.7±56 (8.3)	1.33±1.57	251±289 (15.7)	2.73±2.62	49±179 (2.6)	4.27±4.3	88.6±108 (2.5)	10.6±7.7	456±564 (6.6)

<sup>a</sup> The fluxes are per unit area of the corresponding CH<sub>4</sub> emitting surface area and SD are estimated using bootstrapping methods; <sup>b</sup> Ecosystem contributions from young and mature tree stems were estimated using the measured stem CH<sub>4</sub> fluxes between 15-20 and 135-140 cm stem height above the soil/water surface at 30 cm stem height intervals and multiplied by the corresponding stem surface area. Contributions between 0-20 cm stem height were assumed to be the same as the 20-50 cm stem CH<sub>4</sub> flux and was included in the ecosystem contributions; <sup>c</sup> young tree leaf CH<sub>4</sub> fluxes are the average of four different branches per tree (*n* = 260). No CH<sub>4</sub> emissions were detected from mature tree leaves (*n* = 180).

**Table 2:** Methane production potentials measured from the wood cores extracted.

No of trees sampled	Percentage trees showing evidence of CH <sub>4</sub> production potential (%)	CH <sub>4</sub> production potential rates ± SD (µg CH <sub>4</sub> h <sup>-1</sup> m <sup>-3</sup> vol of wood) <sup>a</sup>
At 20 cm above the soil/water surface		
<i>n</i> = 1232	1.3	158 ± 274
At 130 cm above the soil/water surface		
<i>n</i> = 1343	3.7	440 ± 579

<sup>a</sup> CH<sub>4</sub> production potential was measured by incubating the stem cores for 12 hrs in 35 ml Wheaton vials flushed with N<sub>2</sub><sup>23</sup>.

**Table 3:**  $\delta^{13}\text{C}$  values of tree  $\text{CH}_4$  flux and porewater  $\text{CH}_4$ .

	Flux			Porewater	
	$\delta^{13}\text{C}(\text{CH}_4)^a$	SD	n <sup>b</sup>	$\delta^{13}\text{C}(\text{CH}_4)^c$	N
	(‰)	(‰)		(‰)	
River Negro					
N3	-76.3	0.9	4	-	-
N6	-64.6	3.2	5	-	-
River Amazon					
A7	-65.4	2.2	4	-58.5/-54.5	2
A9	-61.8	3.3	3	-70.8/-63.3	3
River Tapajós					
T11	-59.1	0.4	3	-55.6	1

<sup>a</sup> Mean  $\delta^{13}\text{C}$  values are reported for  $\text{CH}_4$  flux; <sup>b</sup> n represents one chamber deployment from which three or four pairs of  $\text{CH}_4$  concentration and  $\delta^{13}\text{C}(\text{CH}_4)$  values were used to determine a  $\delta^{13}\text{C}$  value for  $\text{CH}_4$  flux via Keeling regression analysis; <sup>c</sup> The range of  $\delta^{13}\text{C}$  values are reported for porewater  $\text{CH}_4$ .

**Table 4:** Estimated annual CH<sub>4</sub> emissions from the Amazon basin using bottom up and top down methods.

Approach: bottom up (BU) top-down (TD)	CH <sub>4</sub> emitting pathways	CH <sub>4</sub> fluxes ± SD (g ha <sup>-1</sup> d <sup>-1</sup> )	Annual emissions ± SD (Tg CH <sub>4</sub> yr <sup>-1</sup> ) <sup>a</sup>	Study
	Mature tree stems	1350 ± 553 - 1927 ± 793 <sup>b</sup>	14 ± 1.8 - 20 ± 2.5 <sup>b</sup>	This study
	Young tree stems	98 ± 46.8 - 104 ± 49.2 <sup>b</sup>	1.02 ± 0.15 - 1.08 ± 0.16 <sup>b</sup>	This study
	Young tree leaf emissions	9.5 ± 15.9	0.099 ± 0.05	This study
<b>BU</b>			<b>15.1 ± 1.8 - 21.2 ± 2.5<sup>b</sup></b>	This study
	Aquatic surfaces	1033 ± 1622	9.7 ± 5.2	This study
	Soil surfaces	170 ± 299	1.1 ± 0.7	This study
	Macrophytes	3245 ± 721 – 1229 ± 334 <sup>c</sup>	8 ± 0.6 <sup>d</sup>	3,8
	Open water	270 ± 80.1	1.2 ± 0.05 <sup>d</sup>	8
	River channel		0.4 - 0.6 <sup>e</sup>	19
<b>BU</b>		Total surface emissions (including trees)	<b>35.6 ± 5.6 – 41.7 ± 5.9<sup>b</sup></b>	This study
<b>BU</b>		Total surface emissions (no trees)	20.5 ± 5.3	This study
<b>BU</b>		Total surface emissions (no trees)	29.4	3
<b>BU</b>		Total surface emissions (no trees)	26.2 ± 9.8	2
<b>TD</b>	Biomass burning (non-wetland source)		4.1 ± 0.7	This study
<b>TD</b>	All		<b>42.7 ± 5.6</b>	This study
<b>TD</b>	All		44 ± 4.8	10
<b>TD</b>	All		40.2 - 52	4
<b>TD</b>	All		37 ± 5.9	26

<sup>a</sup> Surface area used to estimate regional CH<sub>4</sub> contributions reported in Supplementary Table 5; <sup>b</sup> The upper range represents the inclusion of stem CH<sub>4</sub> emissions estimated for up to 5 m of the stem height for mature trees and 1.85 m for young trees using the relationship between stem CH<sub>4</sub> flux and stem height positions; <sup>c</sup> Aquatic macrophyte CH<sub>4</sub> emissions from high and low water season estimated and reported by Devol *et al.*<sup>8</sup> and Melack *et al.*<sup>3</sup>; <sup>d</sup> CH<sub>4</sub> fluxes to estimate emissions from macrophytes and open water were obtained from Devol *et al.*<sup>8</sup> and Melack *et al.*<sup>3</sup>; <sup>e</sup> total annual CH<sub>4</sub> emission estimates from river channels in the Amazon basin obtained from Sawakuchi *et al.*<sup>19</sup>.