

This is a repository copy of *Prescribers' views and experiences of assessing the appropriateness of prescribed medications in a specialist addiction service*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/123741/>

Version: Accepted Version

---

**Article:**

Oluyase, Adejoke Obirenjeyi, Raistrick, Duncan, Hughes, Elizabeth et al. (1 more author) (2017) Prescribers' views and experiences of assessing the appropriateness of prescribed medications in a specialist addiction service. *International Journal of Clinical Pharmacy*. pp. 1248-1255.

<https://doi.org/10.1007/s11096-017-0541-4>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# International Journal of Clinical Pharmacy

## Prescribers' views and experiences of assessing the appropriateness of prescribed medications in a specialist addiction service

--Manuscript Draft--

|  |   |                |
|--|---|----------------|
| <b>Manuscript Number:</b>                            | IJCP-D-16-00597R3   |                |
| <b>Full Title:</b>                                   | Prescribers' views and experiences of assessing the appropriateness of prescribed medications in a specialist addiction service   |                |
| <b>Article Type:</b>                                 | Research Article  |                |
| <b>Keywords:</b>                                     | appropriateness; specialist addiction service; medications; medical prescribers; nurse prescribers; United Kingdom  |                |
| <b>Corresponding Author:</b>                         | Adejoke Obirenjeyi Oluyase, Ph.D<br>University of York<br>York, UNITED KINGDOM  |                |
| <b>Corresponding Author Secondary Information:</b>   |   |                |
| <b>Corresponding Author's Institution:</b>           | University of York  |                |
| <b>Corresponding Author's Secondary Institution:</b> |   |                |
| <b>First Author:</b>                                 | Adejoke Obirenjeyi Oluyase, Ph.D  |                |
| <b>First Author Secondary Information:</b>           |   |                |
| <b>Order of Authors:</b>                             | Adejoke Obirenjeyi Oluyase, Ph.D  |                |
|  | Duncan Raistrick, MBChB, M.Phil, FRCPsych   |                |
|  | Elizabeth Hughes, Ph.D  |                |
|  | Charlie Lloyd, M. Phil  |                |
| <b>Order of Authors Secondary Information:</b>       |   |                |
| <b>Funding Information:</b>                          | NIHR Collaboration for Leadership in Applied Health Research and Care   | Not applicable |
|  |   |                |
| <b>Abstract:</b>                                     | <p>Background: Mental and physical health problems are common in people with substance misuse problems and medications are often required in their management. Given the extent of prescribing for service users who attend specialist addiction services, it is important to consider how prescribers in this setting assess the appropriateness of service users' prescribed medications.</p> <p>Objective: To explore prescribers' views and experiences of assessing the appropriateness of medications prescribed for service users coming in for treatment as well as the differences between prescribers.</p> <p>Setting: A specialist addiction service in the North of England.</p> <p>Method: A phenomenological approach was adopted. Semi-structured interviews were conducted with four nurse prescribers and eight doctors. Data were analysed using thematic framework analysis.</p> <p>Main outcome measure: Prescribers' views and experiences of assessing the appropriateness of prescribed medications.</p> <p>Results: Assessment of the appropriateness of prescribed medications involved reviewing medications, assessing risk, history-taking, involvement of service users, and comparing guideline adherence and 'successful' prescribing. Doctors and nurse prescribers assessed the appropriateness of medications they considered to be within their competency. Doctors provided support to nurse prescribers and general practitioners (GPs) when dealing with issues around prescribing.</p> <p>Conclusion: Assessment of the appropriateness of prescribed medications is complex. The recent reduction in medical expertise in specialist addiction services may negatively impact on the clinical management of service users. It appears that there is a need for further training of nurse prescribers and GPs so they can provide optimal care to service users.</p> |                |

|                               |   |
|-------------------------------|---|
|                               | <p>Impact of findings on practice statement</p> <p>1. Nurse prescribers and doctors in this specialist addiction service differed in the types of medications they reviewed but appeared to be working within their competency.</p> <p>2. Given that doctors in this specialist addiction service appeared to provide support to nurse prescribers around medication-related issues, decreasing medical expertise in addictions may pose a threat to quality decision-making by nurse prescribers.</p> <p>3. The decreasing availability of medical expertise in addiction services also presents a challenge to the management of complex service users by GPs.</p> <p>4. There is a need to provide training and support to nurse prescribers and GPs so that they can provide optimal care to service users.</p> |
| <b>Response to Reviewers:</b> | <p>Response to the comments of reviewers</p> <p>Reviewer 1:<br/>Comment: I am happy with the changes made</p> <p>Reviewer 2:<br/>Comment: The additional quotations and sentence have added a very small degree of depth to the analysis, however I acknowledge the authors' comments that the journal word limit constrained the ability to increase the depth of analysis any further.</p> <p>One minor formatting change is recommended on line 156, for consistency the words within brackets should not be italicized.</p> <p>Response: The italics has now been removed from line 156 (please see line 157 for the quotation). Thank you for highlighting this.</p>   |

**Prescribers' views and experiences of assessing the appropriateness of prescribed  
medications in a specialist addiction service**

Adejoke Obirenjeyi Oluyase, Ph.D. <sup>a</sup>:aobirenjeyi@yahoo.co.uk, 01904 321912. Elizabeth Hughes, Ph.D. <sup>b</sup>:e.c.hughes@hud.ac.uk, Duncan Raistrick, MBChB, M.Phil, FRCPsych<sup>c</sup>:d.raistrick@nhs.net, Charlie Lloyd, M.Phil. <sup>a</sup>:charlie.lloyd@york.ac.uk.

<sup>a</sup>Department of Health Sciences, Room 208b, Area 4 ARRC Building, University of York, Heslington, YO10 5DD, York.

<sup>b</sup>School of Health and Human Sciences, Harold Wilson Building, University of Huddersfield, Queensgate, Huddersfield, HD1 3DH.

<sup>c</sup>Leeds and York Partnership NHS Foundation Trust, 19 Springfield Mount, LS2 9NG, Leeds.

## 1 Introduction

2 People with substance misuse problems often have co-existing physical and/or mental health  
3 conditions [1,2], and are prescribed a large number of medications which may sometimes not  
4 be justified [3]. Service users who seek treatment in specialist addiction clinics are more  
5 likely to have higher levels of dependence and complex needs that include social problems,  
6 functional impairment, comorbidities and use of multiple medications when compared with  
7 those who do not seek help [4,5]. These complex needs may influence prescribing decisions  
8 made for this population [6]. For instance, prescribing may be targeted at maintaining  
9 equilibrium in the lives of service users, which may lead to prescribing outside of guideline  
10 recommendations. Furthermore, service users may want certain medications such as opioids  
11 and benzodiazepines prescribed for non-medical reasons [7,8].

12

13 Opioids used in pain treatment and benzodiazepines for mental health problems have been  
14 implicated in the occurrence of adverse events in people with substance misuse problems.  
15 Benzodiazepines, antidepressants, antipsychotics and substances such as alcohol have often  
16 been found to be used in combination with opioids such as dihydrocodeine and oxycodone in  
17 opioid-related overdose and fatalities [9-11]. Antidepressant prescriptions, especially tricyclic  
18 antidepressants (hereafter TCAs), have also been linked to heroin overdose [12,13].

19

20 The large number of people entering specialist addiction services with complex needs and  
21 multiple prescriptions provides an important opportunity for exploring addiction service  
22 prescribers' views and experiences of assessing the appropriateness of medications prescribed  
23 for service users coming in for treatment as well as the differences between the various types

of prescribers. Prescribers included in this study were medical and non-medical. The non-medical prescribers (NMP) were independent nurse prescribers who could assess and also devise a treatment plan that may include prescribing for service users [14]. NMPs prescribe within their areas of competence [15]. For instance, nurse prescribers working in addiction medicine are able to prescribe substitute opioids, relapse prevention medications, medications for detoxification and vitamin supplements.

Assessment of the clinical appropriateness of non-medical prescribing, including nurse prescribing, have concluded that NMPs generally make clinically appropriate prescribing decisions [16,17]. However, history taking, assessment and diagnosis skills have been highlighted as areas for further attention.

Service users visiting the service could self-refer or be referred from a range of sources such as general practitioners, psychiatrists, hospital, social services, drug services and the criminal justice system. Consequently, this study explored specialist addiction service prescribers' views and experiences of assessing the appropriateness of medications prescribed by others.

## Aim

This study explored specialist addiction service prescribers' views and experiences of assessing the appropriateness of medications prescribed for service users coming in for treatment as well as the differences between prescribers. Appropriateness was considered to involve maximising effectiveness, minimising risks and costs, and respecting the patient's choice [18].

## 49 Ethics approval

50 The study was approved by the University of York's Research Governance Committee and  
51 the National Research Ethics Service (NRES) Committee Yorkshire & The Humber.  
52 Reference 12/YH/0325.

53

## 54 Methods

### 55 Study design and setting

56 A phenomenological approach was taken to explore individual views and experiences of  
57 assessing the appropriateness of medications prescribed for service users. Semi-structured  
58 qualitative interviews were carried out with prescribers comprising nurse prescribers and  
59 medical doctors working at the specialist addiction service. This service is located in a city in  
60 the North of England and is a statutory NHS specialist service that provides tier 3 level  
61 interventions to adults who misuse alcohol and/or drugs. Tier 3 interventions generally  
62 involve the provision of care-planned interventions following a comprehensive community-  
63 based assessment [19]. One-on-one interviews were used because it would be very difficult to  
64 get time-pressed clinicians together for a focus group discussion. In addition, group  
65 interviews may be prohibitive for some prescribers.

66

### 67 Participants

68 Twelve prescribers took part in this study, comprising four nurse prescribers and eight  
69 medical doctors. In line with qualitative research inquiry, the aim of the sampling strategy  
70 adopted was to recruit respondents who could provide valuable insight into the topic and also  
71 to provide a broad overview of the perspectives of different prescribers. Consequently, all the

fourteen prescribers working at the specialist addiction service during the period of this study were provided with the study details by the chief investigator (A. O). This was followed by a meeting with each prescriber to discuss the study in detail after which written informed consent was sought. Twelve of the fourteen eligible prescribers were interviewed. Participants included five females, three of whom were nurse prescribers and two medical doctors and seven males, of whom one was a nurse prescriber and six medical doctors. The medical doctors had different levels of seniority and included one senior house officer (hereafter SHO), one locum doctor, three specialist registrars (hereafter SpR) and three consultant addiction psychiatrists. Generally, the prescribers represented a broad range of qualifications and experience in the addiction field.<sup>1</sup> Nurse prescribers' ages ranged from 34 to 55 years while doctors were between 31 and 65 years.

#### Data collection

Data were collected by the first author, A. O. All the interviews were conducted at a time convenient for participants at the specialist addiction service and lasted on average 48 mins (range: 36 to 74 mins). The topic guide was informed by knowledge of the literature on prescribing and advice from the project advisory group (which included one consultant addiction psychiatrist). The topic guide was piloted with a consultant addiction psychiatrist and covered the following areas: definition of inappropriate prescribing, classes of medications assessed and how assessment is carried out. The interviews were audio recorded (with permission) and transcribed verbatim.

---

<sup>1</sup> Nurse prescribers had practiced in addiction specialty for between one and five years and as nurses for between five and twenty-two years. Two doctors had no prior experience in addiction specialty while the remaining had between six months and thirty-five years of experience. Consultant psychiatrists in particular, had between three and 35 years of experience in addiction specialty. Doctors had between six and 41 years of clinical experience.



## 94 Data analysis

95 Data were analysed using thematic framework analysis [20]. Familiarisation involved  
96 repeated reading of transcripts alongside listening to the audio-recordings and was followed  
97 by a period of descriptive and interpretive coding facilitated by Atlas ti (v 6.0). This inductive  
98 approach enabled a deeper understanding of the data [21]. As new themes emerged, they  
99 were added to the coding framework. Broader themes were subsequently generated and  
100 frequently reviewed while comparing data from participants that supported the themes and  
101 also looking for explanations of any differences of viewpoints within the data. Numbers  
102 rather than names were allocated to participants in order to ensure anonymity and  
103 confidentiality. Trustworthiness of the data was ensured through an audit trail kept by A.O  
104 which detailed how data were collected, how themes were formed and how decisions were  
105 made during the research process. Furthermore, the interpretation of the data was discussed  
106 in-depth with two of the authors (C.L and E.H), who reflected on the plausibility of the  
107 themes and the depth of the analysis. A. O has a pharmacy background while C. L and E.H  
108 have criminology and nursing backgrounds respectively. D.R is a consultant addiction  
109 psychiatrist.

110

## 111 **Results**

112 The following themes emerged in response to how prescribers assessed the appropriateness of  
113 prescribed medications: review of medications, assessing risk, guideline adherence versus  
114 successful prescribing, history-taking and involvement of service users. There were some  
115 areas of differences in nurse prescribers and medical doctors' approaches and also among the  
116 different types of medical doctors. These differences are highlighted in the text.

117    Review of medications

118    The classes of medications reviewed varied among prescribers with three of them (all  
119    doctors) with the longest years of prescribing experience stating that they reviewed all of  
120    service users' medications for their appropriateness. One of these three prescribers had 41  
121    years of experience in prescribing and made the following statement:

122    *So I'd look at the list of drugs prescribed and see how they matched up to what I thought the*  
123    *person was showing in terms of addiction illness, physical illness and mental illness [P3,*  
124    *consultant].*

125    The remaining prescribers consisting of other doctors and nurse prescribers described a more  
126    limited remit. These doctors considered their scope of practice to encompass medications for  
127    mental health illnesses, addictions and sometimes opioids for pain relief while nurse  
128    prescribers described a focus on medications used for treating addiction problems. This quote  
129    captures a nurse prescriber's view:

130    *So I don't really see, with psychiatric medication, that that would be within my remit really.*  
131    *If somebody came and they were prescribed 100mgs of methadone and they couldn't even*  
132    *open their eyes then, I would be assessing the appropriateness of the dosage and making*  
133    *necessary adjustments to things like that [P10, NP].*

134    Nurse prescribers further described involving doctors at the specialist addiction service or  
135    service users' general practitioners (hereafter GPs) if they had particular concerns about  
136    medications. There was an underlying feeling of cautiousness characterised by their  
137    perceptions of their competency. This was captured by the quote below:

138 *As I say, if I was particularly concerned about someone's mood or I have particular concerns*  
139 *about the medication I would defer to a medic. You know, it's not an area I feel strongly*  
140 *confident on [P6, NP].*

141 Doctors at the specialist addiction service were a valuable source of support to nurse  
142 prescribers in prescribing-related issues. There was also particular reliance on the expertise of  
143 consultant addiction psychiatrists by both nurse prescribers and doctors who were not  
144 consultants. A doctor described contacting a GP concerning an inappropriate medication and  
145 the support of her consultant in providing expert advice when needed:

146 *For the example I started with [patient with schizophrenia on supra-BNF dose of olanzapine],*  
147 *I wrote to the GP saying, you know, Mr So-and-So is stable and is relatively symptom free on*  
148 *this but I'm worried about this monitoring [olanzapine monitoring] but generally if I think*  
149 *something's really inappropriate and I'm in a position to contact the original prescriber I'll*  
150 *try to do that, but I'd always discuss a case with my consultant and make a decision about*  
151 *whether or not I need to do something imminently [P12, SHO].*

152 It appears that prescribers at this specialist addiction service provided a 'safety net' function  
153 to other prescribers such as GPs:

154 *If I find something that's maybe been overlooked or prescribed wrongly, then I will let the*  
155 *GP know about it [P5, Locum].*

156

157 *I'd probably look at it [medication appropriateness] at the initial assessment and if there's*  
158 *anything that comes up or that was sort of glaringly obvious I'd refer to the GP and ask the*  
159 *GP to review, if they're prescribing [P11, NP].*

160

161 Specialist addiction service prescribers further described GPs' varying responses to the need  
162 for review of service users' medications:

163 *Yeah. that has happened on a couple of times where I've written to the GP to ask them to*  
164 *review... there have been a couple of scenarios where I've written and the GP hasn't*  
165 *responded or the GP has written back saying, I don't feel I'm the best person to do this,*  
166 *would you refer to a specialist service or would you basically will you deal with it [P12,*  
167 *SHO].*

168

169 They also described sometimes taking over prescribing of psychiatric medications from GPs:

170

171 *But in general I'd like to take over all of the psychoactive drugs that somebody gets, at least*  
172 *until the point that we're sure that the drugs are appropriate and we've got some sort of*  
173 *stable situation [P3, Consultant].*

174

175 Assessing risk

176 The evaluation of risk is a theme that was highlighted by all prescribers as a means through  
177 which they assess the appropriateness of service users' medications. All the twelve  
178 prescribers said they considered the risk posed by a medication. Some of the quotations  
179 captured this:

180 *Well if it's going to do, first of all, less harm than the actual substance, not more harm, so the*  
181 *actual prescription can be worse than doing nothing [P5, locum doctor].*

182 One prescriber described a service user who she felt had an inappropriate and high risk  
183 prescription of olanzapine (an antipsychotic). The service user was an elderly man who was  
184 being prescribed olanzapine (25mg) at a dose higher than that stated in the British National  
185 Formulary (BNF) without monitoring by a psychiatrist:

186 *I have a patient who has a very old diagnosis of paranoid schizophrenia dating from his late*  
187 *teens, and for this he's prescribed a very high dose of medication called olanzapine and he's*  
188 *prescribed over the limit in the BNF and he's not under the supervision of a specialist. So I*  
189 *would label that as an inappropriate prescription because (a) he's elderly, which means that*  
190 *he's more prone to cardiac disease, and the drug can cause diabetes which can lead to heart*  
191 *disease. It can cause arrhythmias, he's not being monitored regularly with regards to that,*  
192 *and he's not being monitored with regards to his clinical symptoms, which, are actually, from*  
193 *a psychosis point of view, negligible [P12, SHO].*

194 The SHO described contacting the service user's GP concerning the antipsychotic  
195 medication. His GP refused to alter it due to the service user's stability on the dose for a  
196 prolonged period. The GP and SHO differed in their views concerning the antipsychotic.  
197 There was no change made to the antipsychotic.

198

199 Guideline adherence versus successful prescription

200 The need to assess if prescribing is in line with guidelines was highlighted. Some prescribers  
201 further acknowledged that the need to individualise prescribing and ensure optimal  
202 functioning may lead to prescribing outside guideline recommendations. The need to consider  
203 the context of prescribing was emphasised by a nurse prescriber:

204 *And I think any comment about any prescribing should only be made when you know about*  
205 *the circumstances in which the decision was made. For example, we prescribe very high*  
206 *doses of some drugs, now some people say that you shouldn't prescribe at those levels, but*  
207 *they are appropriate if you know about the circumstances [P1, NP].*

208 A consultant addiction psychiatrist also expressed similar views and contrasted guideline  
209 adherence with successful prescribing:

210 *Prescribing is something of an art as well as a science, so prescribers will sometimes*  
211 *prescribe things that they know are not really indicated but with the aim of achieving a*  
212 *particular goal [P3, Consultant].*

213

214 History-taking

215 All prescribers identified history-taking as a part of their assessment of the appropriateness of  
216 service users' medications. The prescribers described enquiring about service users' medical  
217 and medication history:

218 *Looking at the history of their substance use, history of any physical health problems, mental*  
219 *health history, and current mental state as well so I'd get the full history and I think then you*  
220 *can kind of gauge whether something might be inappropriately prescribed [P11, NP].*

221 Despite prescribers routinely obtaining a medical/medication history from service users, most  
222 reiterated that it was not within their remit to explore the appropriateness of all prescribed  
223 medications:

224 *...I would, in as much as part of the assessment, I would ask the service user ...are they on*  
225 *any medications. If they are, what it is, what dose, what's it prescribed for and are they*

226 *taking it. That would be the total sum of my assessment. I wouldn't move to beyond exploring*  
227 *that condition or whether that was appropriate, I don't think that's my place [P6, NP].*

228 All prescribers further described some challenges with self-report when obtaining service  
229 users' histories. These include problems with the reliability of information provided by  
230 service users as some of them may withhold information. This may lead to prescribing of  
231 unnecessary medications. Prescribers also described service users who do not know details of  
232 their medications such as the name and reason for medication use. Some may be cognitively  
233 impaired by substances and therefore unable to provide necessary information. Prescribers  
234 may have to contact GPs concerning needed information. There was however an  
235 acknowledgment that contacting GPs for information was not always routine practice as  
236 prescribers tended to rely on information obtained from service users.

237

238 Involvement of service users

239 This theme was described by all prescribers. It involved discussing with service users in order  
240 to understand their views concerning the appropriateness of their prescribed medications:

241 *Well, firstly I discuss with the patient to see what the patient's view is, and explain what I*  
242 *think, which are the reasons for this inappropriateness [P13, SpR].*

243 Prescribers also highlighted the fact that lack of engagement by service users may affect  
244 prescribing decisions. For instance, service users' medications may need to be stopped due to  
245 repeated non-attendance of clinic appointments.

246

247

## 248 Discussion

249 The evidence from this study shows that the assessment of the appropriateness of prescribed  
250 medications is a complex judgment. Besides a few more experienced doctors, all other  
251 prescribers (doctors and nurse prescribers) tended to review only the subset of medications  
252 which they saw as within their competency. It has been recommended that doctors and nurse  
253 prescribers adhere to their areas of competency for safe practice [22,23]. Nurse prescribers  
254 and doctors appeared to be working within their competency.

255

256 Published evidence suggests non-medical prescribers generally make clinically appropriate  
257 prescribing decisions with the need for further improvement in assessment, diagnosis and  
258 history-taking skills [16,17]. Nurse prescribers described referring service users who they had  
259 concerns about their medications to doctors at the specialist addiction service or service  
260 users' GPs. Specialist addiction service doctors particularly represented a valuable source of  
261 support to nurse prescribers when dealing with issues around prescribing. The more junior  
262 doctors (non-consultants) also relied on their senior colleagues, especially consultant  
263 addiction psychiatrists, for expert advice on medications. There was further evidence that  
264 prescribers were a sort of 'safety net' against medication-related risks as they intervened and  
265 contacted GPs if they found serious problems with service users' medications.

266

267 Service users pose particular challenges in terms of complexity and risk issues. They often  
268 have complex needs including severe comorbid mental and physical health problems [24-29].  
269 In order to meet these needs, Public Health England (2014) has recommended that addiction  
270 specialist doctors such as consultant psychiatrists work alongside non-medical prescribers



and other doctors in a multidisciplinary team [23]. The drug and alcohol treatment system has however undergone some changes in commissioning in recent years. This has involved a move from mainly NHS service provision to a more mixed economy of service providers [23]. These changes have led to a decrease in the number of doctors including consultant addiction psychiatrists in treatment systems [23], with nurses taking on more prescribing roles. Consequently, there is a reduction in the capacity of these new treatment systems for specialist expertise and complex case management.

It appears that there is a possibility of reduction in the quality of prescribing and decision-making as a result of these changes as nurse prescribers and GPs may not have ready access to support and specialist knowledge when required. The potential for specialists to provide clinical supervision that will support nurse prescribers in making clinically appropriate decisions when needed is also hampered. It appears future prescribing practice in alcohol and drug treatment systems will mostly involve nurse prescribers. This raises concerns about the future review practices of psychiatric medications in addiction services if nurse prescribers are not further strengthened to work with service users, including complex clients. In addiction service users, psychiatric comorbidity is highly prevalent [25-28] and medications used in their management have often been implicated in overdose and fatalities [11-13]. Pharmacists' support could be enlisted to guide prescribing decisions for service users with complex comorbidity. This approach may assist in improving medicines management among service users.

There is the need to equip nurse prescribers to work with service users, especially complex cases. Given that assessment, diagnosis and history-taking skills are pre-requisites for

295 undertaking the nurse prescribing qualification, these skills may well be further developed  
296 through training to enable nurse prescribers manage complex service users, especially those  
297 with comorbid mental disorders. Practice should include regular supervision of nurse  
298 prescribers by an experienced doctor or nurse prescriber to ensure that they are making  
299 optimal clinical decisions.

300

301 The relationship between healthcare professionals and service users have changed over the  
302 years from a predominantly paternalistic model to one in which service users have  
303 increasingly become active partners whose views are important [30,31]. Involving service  
304 users assists the prescriber in eliciting their views and is useful in decision-making  
305 concerning treatment [32]. There is evidence that building a positive relationship can lead to  
306 positive client and treatment outcomes [33]. Despite these potential benefits, prescribers  
307 identified problems that may occur when trying to involve service users in decision-making.  
308 The quality of information provided by service users may be poor as a result of cognitive  
309 impairment or even deliberate withholding of information. When service users are actively  
310 misusing substances, prescribers lose access to the most fundamental tool in medicine, the  
311 patient's self-report [34]. While some prescribers described contacting service users' GPs for  
312 further information concerning medications, this was not done by all prescribers.

313

314 Depending on information obtained from only service users in assessing appropriateness  
315 implies that medications which are potentially inappropriate may not be identified if service  
316 users fail to mention them. There is the possibility that different prescribers may go ahead to  
317 prescribe undisclosed medications such as multiple central nervous system depressants. In  
318 addiction medicine, there should be careful consideration of self-report and collateral

information should be sought where possible [34]. Shared medical records [35] and good communication among different service providers are essential in obtaining accurate medical/medication histories and reducing the potential for multiple prescribing, drug interactions, overdose incidents and conflicting treatment plans [34].

The limited applicability of guidelines to service users was also recognised by prescribers. Guidelines often have a disease-specific focus and limited applicability to the varying needs of individual patients [36]. Although prescribing outside guideline recommendations carries its own risks including the potential for greater severity of unwanted side effects [37], there needs to be a weighing of such risks against more pragmatic outcomes that may be of great importance to service users.

#### Strengths and limitations

To the knowledge of the authors, this is the first study to explore the views and experiences of specialist addiction service prescribers when assessing the appropriateness of prescribed medications among service users coming to this setting. Owing to the fact that the interviews were conducted with prescribers after they had taken part in an earlier study in which the appropriateness of opioids and psychiatric medications were assessed using a modified form of the Medication Appropriateness Index [38], it is possible that participation in this initial study may have influenced some of their responses to the different areas explored in the interviews. Consequently, prescribers' responses might be different if they were interviewed before taking part in this initial study.

The findings may lack generalisability to prescribers in other addiction services, especially given the changes that have occurred in drug and alcohol treatment services in the UK. There has been an increase in the number of third sector organisations (non-statutory service providers and the private sector) providing drug and alcohol services. Availability of medical expertise has also diminished in these services. Further research should involve multiple sites (including services run by the NHS and third sector organisations), to establish if the findings of this study are applicable. Given the reducing levels of medical expertise among staff in specialist addiction services, an important area to explore will be the role and scope of nurse prescribers: including their views on the changing drug treatment landscape, management of service users (especially those with complex needs), the support available to nurse prescribers and their training needs. Similarly, there may well be need to interview GPs on these areas since it was evident that specialist addiction service prescribers provided some level of support to them.

Furthermore, data collection was by a single researcher. There is the possibility that the researcher's own perspectives may have affected interpretations that were made. However, the conduct, analysis and interpretation of data were overseen by two of the authors in addition to A.O.

## Conclusion

Assessment of the appropriateness of prescribed medications appeared to be a complex judgment. Optimal assessment of prescribing appropriateness should involve a balance between guideline recommendations, risks and benefits of prescribing, and the context. Nurse

prescribers and medical doctors differed in their approach to reviewing medications but appeared to be working within their competency, with doctors providing support to nurse prescribers when needed. Prescribers were a sort of ‘safety net’ against medication-related risks to GPs. Recent changes in the UK drug and alcohol field have led to diminishing availability of medical expertise and an increasing reliance on non-medical prescribing. These changes have the potential to affect the quality of decision-making around medications. It appears there is a need to further empower non-medical prescribers and GPs to effectively manage service users with comorbidity.

Acknowledgments: The authors are grateful to all those that participated in this study, and also for the helpful comments on an earlier draft of this paper by two reviewers.

Funding: This study is part of independent research funded by the National Institute of Health Research (NIHR) through the NIHR Collaboration for Leadership in Applied Health Research and Care for Leeds, York and Bradford.

Conflicts of interest: The authors declare that there are no conflicts of interest.

## References

1. McLellan T. Revisiting the past for a look toward future research: a final editorial. *J Subst Abuse Treat.* 2009; 36(4):352-4.
2. Mortlock KS, Deane FP, Crowe TP. Screening for mental disorder comorbidity in Australian alcohol and other drug residential treatment settings. *J Subst Abuse Treat.* 2011; 40(4):397-404.
3. Oluyase AO, Raistrick D, Abbasi Y, Dale V, Lloyd, C. A study of the psychotropic prescriptions of people attending an addiction service in England. *Adv Dual Diagn.* 2013; 6(2):54-65.
4. Cunningham JA, Breslin FC. Only one in three people with alcohol abuse or dependence ever seek treatment. *Addict Behav.* 2004; 29(1):221–3.

5. Tucker JA, Simpson CA. The recovery spectrum: from self-change to seeking treatment. *Alcohol Res Health*. 2011; 33(4):371-9.
6. Hughes E. Guidelines for working with mental health-substance use. In: Cooper DB, editor. *Developing services in mental health-substance use*. United Kingdom: Radcliffe Publishing Ltd; 2011.
7. Inciardi JA, Surratt HL, Kurtz SP, Cicero TJ. Mechanisms of prescription drug diversion among drug-involved club- and street-based populations. *Pain Med*. 2007; 8(2):171-83.
8. Savage SR, Kirsh KL, Passik SD. Challenges in using opioids to treat pain in persons with substance use disorders. *Addict Sci Clin Pract*. 2008; 4(2):4-25.
9. Boscarino JA, Kirchner HL, Pitcavage JM, Nadipelli VR, Ronquest NA, Fitzpatrick MH, et al. Factors associated with opioid overdose: a 10-year retrospective study of patients in a large integrated health care system. *Subst Abuse Rehabil*. 2016; 77:131-41.
10. Darke S, Duflou J, Torok M. Toxicology and characteristics of fatal oxycodone toxicity cases in New South Wales, Australia 1999-2008. *J Forensic Sci*. 2011; 56(3):690-3.
11. Zamparutti G, Schifano F, Corkery JM, Oyefeso A, Ghodse AH. Deaths of opiate/opioid misusers involving dihydrocodeine, UK, 1997-2007. *Br J Clin Pharmacol*. 2011; 72(2):330-7.
12. Darke S, Hall W. Heroin overdose: research and evidence-based intervention. *J Urban Health*. 2003; 80(2):189-200.
13. Darke S, Ross J. The use of antidepressants among injecting drug users in Sydney, Australia. *Addiction*. 2000; 95(3):407-17.
14. Department of Health. Review of prescribing, supply and administration of medicines: final report (Crown II). London: Department of Health; 1999.
15. Cope LC, Abuzour AS, Tully MP. Non-medical prescribing: where are we now? *Ther Adv Drug Saf*. 2016; 7(4):165-72.
16. Latter S, Smith A, Blenkinsopp A, Nicholls P, Little P, Chapman S. Are nurse and pharmacist independent prescribers making clinically appropriate prescribing decisions? An analysis of consultations. *J Health Serv Res Policy*. 2012; 17(3):149-56.

17. Naughton C, Drennan J, Hyde A, Allen D, O'Boyle K, Felle P, et al. An evaluation of the appropriateness and safety of nurse and midwife prescribing in Ireland. *J Adv Nurs*. 2013; 69(7):1478–88.
18. Lexchin J. Improving the appropriateness of physician prescribing. *Int J Health Serv*. 1998; 28(2):253-67.
19. National Treatment Agency for Substance Misuse. Models of care for alcohol misusers (MoCAM). London: National Treatment Agency for Substance Misuse; 2006.
20. Miles MB, Huberman AM. *Qualitative data analysis: a sourcebook of new methods*. Beverly Hills, CA: Sage Publications; 1994.
21. Silverman D. *Doing qualitative research: a practical handbook*. London: Sage Publications; 2013.
22. Public Health England. Non-medical prescribing in the management of substance misuse. London: Public Health England; 2014. <http://www.nta.nhs.uk/uploads/nmp-in-the-management-of-substance-misuse.pdf>. Accessed 23 Jul 2015.
23. Public Health England. The role of addiction specialist doctors in recovery orientated treatment systems. A resource for commissioners, providers and clinicians. London: Public Health England; 2014. <http://www.nta.nhs.uk/uploads/the-role-of-addiction-specialist-doctors.pdf>. Accessed 23 Jul 2015.
24. Virgo N, Bennett G, Higgins D, Bennett I, Thomas P. The prevalence and characteristics of co-occurring serious mental illness (SMI) and substance abuse or dependence in the patients of adult mental health and addictions services in eastern Dorset. *J Ment Health*. 2001; 10(2):175-88.
25. Strathdee G, Manning V, Best D, Keaney F, Bhui K, Witton J, et al. Dual diagnosis in a primary care group (PCG), (100,000) population locality: a step-by-step epidemiological needs assessment and design of a training and service response model. *Drugs: Educ Prev Polic*. 2005; 12(1):119-23.
26. Weaver T, Madden P, Charles V, Stimson G, Renton A, Tyrer P, et al. Comorbidity of substance misuse and mental illness in community mental health and substance misuse services. *Br J Psychiatry*. 2003; 183(4):304-13.
27. Delgadillo J, Godfrey C, Gilbody S, Scott P. Depression, anxiety and comorbid substance use: association patterns in outpatient addictions treatment. *Mental Health and Substance Use*. 2013; 6(1):59-75.

28. Walsh K, Copello A. Severe and enduring mental health problems within an established substance misuse treatment partnership. *Psychiatr Bull*. 2014; 38(5):216-9.
29. Dickey B, Normand SL, Weiss RD, et al. Medical morbidity, mental illness, and substance use disorders. *Psychiatr Serv*. 2002; 53(7):861-7.
30. Elwyn G, Edwards A, Kinnersley P. Shared decision-making in primary care: the neglected second half of the consultation. *Br J Gen Pract*. 1999; 49(443):477-82.
31. Bezreh T, Laws MB, Taubin T, Rifkin DE, Wilson IB. Challenges to physician-patient communication about medication use: a window into the skeptical patient's world. *Patient Prefer Adherence*. 2012; 6:11-8.
32. Barnett PB. Rapport and the hospitalist. *Am J Med*. 2001; 111(9B):31S-35S.
33. Leach MJ. Rapport: a key to treatment success. *Complement Ther Clin Pract*. 2005; 11(4):262-5.
34. Action on Addiction. The management of pain in people with a past or current history of addiction. London: Action on Addiction; 2013. <http://www.actiononaddiction.org.uk/Documents/The-Management-of-Pain-in-People-with-a-Past-or-Cu.aspx>. Accessed 23 Jul 2015.
35. Ghitza UE, Sparenborg S, Tai B. Improving drug abuse treatment delivery through adoption of harmonised electronic health record systems. *Subst Abuse Rehabil*. 2011; 2: 125-31.
36. Farquhar CM, Kofa EW, Slutsky JR. Clinicians' attitudes to clinical practice guidelines: a systematic review. *Med J Aust*. 2002; 177(9):502-6.
37. Ali SI, Ajmal SR. When is off-label prescribing appropriate? *Curr Psychiatr*. 2012; 11(7):23-7.
38. Hanlon JT, Schmader KE, Samsa GP, Weinberger M, Uttech KM, Lewis IK, et al. A method for assessing drug therapy appropriateness. *J Clin Epidemiol*. 1992; 45(10):1045-51.



## **Response to the comments of reviewers**

### **Reviewer 1:**

**Comment:** I am happy with the changes made

### **Reviewer 2:**

**Comment:** The additional quotations and sentence have added a very small degree of depth to the analysis, however I acknowledge the authors' comments that the journal word limit constrained the ability to increase the depth of analysis any further.

One minor formatting change is recommended on line 156, for consistency the words within brackets should not be italicized.

**Response:** The italics has now been removed from line 156 (please see line 157 for the quotation). Thank you for highlighting this.



[Click here to access/download](#)

**Electronic Supplementary Material**

Topic guide for interviews with prescribers.docx

